Dissipative quantum oscillator with two competing heat baths

Heiner Kohler†and Fernando Sols‡

Abstract. We study the dissipative dynamics of a harmonic oscillator which couples linearly through its position and its momentum to two independent heat baths at the same temperature. We argue that this model describes a large spin in a ferromagnet. We find that some effects of the two heat baths partially cancel each other. This leads to unexpected features such as underdamped oscillations and long relaxation times in the strong coupling regime. Such a partial frustration of dissipation can be ascribed to the canonically conjugate character of position and momentum. We compare this model to the scenario where a single heat bath couples linearly to both the position and the momentum of the central oscillator. In that case less surprising behavior occurs for strong coupling. The dynamical evolution of the quantum purity for a single and a double wave packet is also investigated.

PACS numbers: 03.65.Yz, 05.40.-a

† Institut für theoretische Physik, Philosophenweg 19 Universität Heidelberg, Germany

‡ Departamento de Física de Materiales , Universidad Complutense de Madrid, E-28040 Madrid, Spain

E-mail: kohler@tphys.uni-heidelberg.de,f.sols@fis.ucm.es

1. Introduction

The dissipative harmonic oscillator has for long attracted considerable interest as a prototype of open quantum system. The early work of Magalinskii [1] and Ullersma [2] focusing mainly on an Ohmic environment and the weak coupling limit has been later extended in many different aspects [3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14] as, for instance, to strong coupling [11], to non–Markovian noise [9, 12] or to nonlinear coupling [3]. Much activity was boosted by the work of Caldeira and Leggett [15] on the dissipative mechanics of a macroscopic quantum variable. Quantum decoherence has also been addressed in Refs. [16, 17, 18]. A comprehensive review can be found in the textbook by Weiss [19].

In the above context relatively little attention has been paid to effects arising from the coupling to different system variables. The system variable which couples to the heat bath is most often assumed to be the position q. In the sequel we refer to this model as the q-oscillator. The choice of q as the coupling variable was favoured in Ref. [15]. There it was argued that a complex dissipative environment can be modelled by a bath of harmonic oscillators, with the couplings parameters chosen to yield a Langevin equation for the q variable in the semiclassical limit. It was shown later [20] (see also Ref. [21]) that, in a superconducting Josepshon juntion, the oscillator bath model can be derived microscopically from the coupling of the phase variable to the quasiparticle bath, with the phase playing the role of position. At that time the possibility of having a second bath coupled to the momentum p variable was ruled out. However, in a superconducting weak link, the electromagnetic field couples to the relative number variable, which in the above scheme plays the role of momentum [21, 22]. If the coupling to the phase is neglected, the semiclassical behavior of the particle number is governed by an Abraham– Lorentz equation [21, 23]. The coupling to the momentum has been also been considered by Leggett [24] and by Cuccoli *et al.* [25]. A systematic study of the combined role of the electromagnetic and quasiparticle fields is presented in Ref. [21]. Thus Josephson junctions provide a scenario where the distinction between the different baths and their different coupling mechanisms becomes imperious.

Our study has also been motivated by the work of Castro Neto, Novais and coworkers [26]. To investigate the problem of a spin $\frac{1}{2}$ impurity in an antiferromagnetic environment, they studied a generalized spin-boson problem using renormalization group techniques. The localized spin couples with two components to the environment. They found a crossover from incoherent to coherent transitions in the spin correlation functions in certain parameter ranges. There the term *quantum frustration* was coined, which refers to the reduction of the effective interaction with the environment due to the non-commutativity of the coupled operators. In this article, we argue that a harmonic oscillator coupled through position and momentum to two independent baths is a suitable representation of a large spin impurity in a ferromagnetic system. We find a weaker cancellation of the effects of the two baths. Since that situation corresponds to a large (quasiclassical) spin, we have referred to this weaker form of frustration as

quasiclassical frustration [27].

Here we wish to investigate systematically the scenario where a particle couples through its position and momentum to two baths, which are independent but have the same temperature. As a paradigmatic system we choose a harmonic oscillator. This choice is motivated not only by mathematical convenience. A more fundamental reason is that the symmetrical roles played by position and momentum makes the harmonic oscillator the natural ground to study the difference and the interplay (when both are present) between position and momentum coupling.

An explanatory comment is in order. As is well known, a coupling to q can be converted into a coupling to p through a suitable canonical transformation. However, the two representations bear an important difference. Only in one of them is it possible to represent the system–bath interaction as the coupling of the quantum variable to a system of otherwise independent oscillators. A characteristic example is provided by a particle of charge e in an electromagnetic field, where a "velocity– coupling model" [28] seems to apply. Minimal coupling $(\mathbf{p} \rightarrow \mathbf{p} - e\mathbf{A}/c)$ generates not only an interaction term $\mathbf{p} \cdot \mathbf{A}$ but also a diamagnetic term $\propto \mathbf{A}^2$ which can be rightly interpreted as an interaction between the effective oscillators. A unitary transformation $U = \exp(ie\mathbf{q} \cdot \mathbf{A}/c)$, with $[q_i, p_j]_- = i\delta_{ij}$, acting on $\mathbf{p} - (e/c)\mathbf{A}$, removes the coupling $\mathbf{p} \cdot \mathbf{A}$. This happens at the expense of generating a coupling $\propto \mathbf{q} \cdot \mathbf{E}$ between the position and the electric field \mathbf{E} . Importantly, in this new representation no quadratic field term is left, i.e. the charge couples to a set of independent photons. Thus, in the precise language which we propose here, a charged particle couples to the electromagnetic field through its position \mathbf{q} .

In Sec. 2 we study the model of an oscillator coupled to two different baths. The general formalism is sketched in Sec. 2.1. In Sec. 2.2 we investigate the case where the two independent Ohmic baths couple linearly to position and momentum. A related feature is the possibility of inducing a crossover from overdamped to underdamped oscillations by *increasing* the damping coefficient. Surprisingly, we obtain underdamped equilibrium oscillations for arbitrarily strong coupling. Sec. 2.3 is devoted to the properties of spectral functions under rather general combinations of double–bath systems. Time dependent phenomena and implications for quantum decoherence are discussed in Sec. 3 and Sec. 4 for the case of a double Ohmic bath. In Sec. 5 we consider the situation where an oscillator couples to a single bath with the most general form of linear coupling to both position and momentum. We find that the most prominent feature of the double–bath model, namely, the existence of underdamped oscillations for arbitrarily strong coupling, disappears in the case of single–bath dissipation.

2. Two independent baths

The general form of a Hamiltonian describing an oscillator coupled to two independent baths is [21]

$$H = \frac{\omega_p}{2} (p + \delta p)^2 + \frac{\omega_q}{2} (q + \delta q)^2 + \sum_k \omega_k a_{qk}^{\dagger} a_{qk} + \sum_k \omega_k a_{pk}^{\dagger} a_{pk} , \qquad (1)$$

where $[q, p]_{-} = i$ and all operators are dimensionless and $\hbar = 1$. The form of the Hamiltonian highlights the symmetry between q and p. The notion of mass is avoided by $m = 1/\omega_p$. For the fluctuating pieces we assume that they are linear in the bath variables and independent of p and q,

$$\delta q = i \sum_{k} \mu_k \left(a_{pk}^{\dagger} - a_{pk} \right)$$

$$\delta p = i \sum_{k} \lambda_k \left(a_{qk}^{\dagger} - a_{qk} \right) , \qquad (2)$$

After two unitary transformations $U_p = \exp(ip\delta q)$ and $U_q = \exp(iq\delta p)$ one arrives at the Hamiltonian

$$H = \frac{\omega_q}{2}q^2 + \sum_k \omega_k \left| a_{qk} + \frac{\lambda_k}{\omega_k} q \right|^2 + \frac{\omega_p}{2}p^2 + \sum_k \omega_k \left| a_{pk} + \frac{\mu_k}{\omega_k} p \right|^2 , \qquad (3)$$

where the short-hand notation $a^{\dagger}a = |a|^2$ is used. The model is a Caldeira–Leggett– type–of Hamiltonian [15]. It describes a harmonic oscillator with momentum p and position q, each variable being coupled to a different oscillator bath. The frequency of the central oscillator is $\omega_0 = (\omega_p \omega_q)^{1/2}$. The baths are described by the spectral densities

$$J_q(\omega) = 2\sum |\lambda_k|^2 \delta(\omega - \omega_k) , \qquad J_p(\omega) = 2\sum |\mu_k|^2 \delta(\omega - \omega_k) .$$
 (4)

Although one could have started directly with Eq. (3), we prefer to present the Hamiltonian Eq. (1) as the starting point in order to provide a natural justification for the renormalization terms $\sum_{k} |\mu_{k}|^{2} \omega_{k}^{-1} p^{2}$ and $\sum_{k} |\lambda_{k}|^{2} \omega_{k}^{-1} q^{2}$ in Eq. (3) which otherwise would have to be introduced *ad hoc*.

For low-lying excitations, and thus for low temperatures, the Hamiltonian (3) becomes equivalent to that of a large spin s in a ferromagnetic environment. This can be seen as follows: The Hamiltonian for a spin in a large magnetic field along the z direction is

$$H' = -\mu_B (S_z B_z + S_x \delta B_x + S_y \delta B_y) + \mathbf{B}^2, \tag{5}$$

where the fluctuating terms model the low-lying bosonic (magnon) excitations of the ferromagnet at the site of the large spin. The fluctuations in the z direction are neglected since they are quadratic in S_x and S_y with S_x , $S_y \ll S_z$. In other words S_z is approximately a constant of motion [29]. For $s \to \infty$ the first term in (5) is a

harmonic oscillator $S_z = \hbar |a|^2 - \hbar s$. The action of S_i on the eigenstates of this harmonic oscillator is

$$\begin{aligned} S_{z}|n\rangle &= (n-s)|n\rangle \\ S_{+}|n\rangle &= \hbar\sqrt{2s-n}\sqrt{n+1}|n+1\rangle \\ &\approx \hbar\sqrt{2s}\sqrt{n+1}|n+1\rangle &\equiv \hbar\sqrt{2s}a^{\dagger}|n\rangle \\ S_{-}|n\rangle &= \hbar\sqrt{2s-n+1}\sqrt{n}|n-1\rangle \\ &\approx \hbar\sqrt{2s}\sqrt{n}|n-1\rangle &\equiv \hbar\sqrt{2s}a|n\rangle , \end{aligned}$$

and $[S_-, S_+] = 2\hbar^2 s$. This amounts to keeping only the leading order term in the Holstein–Primakoff transform of S_+ , S_- [30]. The rescaled Hamiltonian H'/2s becomes formally identical to Eq. (3) in the limit $s \to \infty$, with $\omega_0 = \mu_B B_z/2s$ and the fluctuating pieces scaling as $\delta B_x = \sqrt{s} \sum \lambda_k (a_{qk} + a_{qk}^{\dagger})$ and $\delta B_y = \sqrt{s} \sum \mu_k (a_{pk} + a_{pk}^{\dagger})$. Such large spins have been observed in magnetic particles [31].

In the following we assume a power-law behavior at $\omega = 0$ for both spectral densities $J_q(\omega) = 2\gamma_q \omega^{\alpha_q} / (\omega_{\rm ph}^{\alpha_q-1}\pi)$ and $J_p(\omega) = 2\gamma_p \omega^{\alpha_p} / (\omega_{\rm ph}^{\alpha_p-1}\pi)$. We introduce the "phononic" frequency $\omega_{\rm ph}$ and the dimensionless coupling constants γ_n , with n = q, p. Moreover we introduce a cutoff frequency Ω_n for both baths. Unless stated otherwise, we assume $\omega_{\rm ph} = \Omega_q = \Omega_p \equiv \Omega$.

2.1. General Results

Elimination of the bath variables yields the Heisenberg equations of motion for q and p

$$\dot{q}(t) = \omega_p p(t) + \int^t ds \, K_p(t-s)\dot{p}(s) + F_p(t) -\dot{p}(t) = \omega_q q(t) + \int^t ds \, K_q(t-s)\dot{q}(s) + F_q(t).$$
(6)

The response kernel is defined as

$$K_n(t) \equiv \int_0^\infty \frac{J_n(\omega)}{\omega} \cos(\omega t) d\omega \ , \ n = q, p \tag{7}$$

and the force operator $F_q(t) = \sum \lambda_k a_{qk} \exp(-i\omega_k t) + \text{H.c.}$, with $F_p(t)$ defined accordingly. In Fourier space, Eq. (6) reads

$$\left[\widetilde{J}_{q}(\omega) - \omega_{q}\right]q + i\omega p = F_{q}(\omega)$$
(8)

$$-i\omega q + \left[\widetilde{J}_p(\omega) - \omega_p\right] p = F_p(\omega) , \qquad (9)$$

where $\widetilde{J}_n(\omega)$ is the symmetrized Riemann transform [32]:

$$\widetilde{f}(\omega) = \omega^2 \mathcal{P} \int_0^\infty \frac{f(\omega')}{\omega' (\omega'^2 - \omega^2)} d\omega' - i \operatorname{sgn}(\omega) \frac{\pi}{2} f(|\omega|) .$$
(10)

The oscillation modes are given by the zeros of the function

$$\chi^{-1}(\omega) = \omega_0^2 - \omega^2 - \omega_q \widetilde{J}_p(\omega) - \omega_p \widetilde{J}_q(\omega) + \widetilde{J}_q(\omega) \widetilde{J}_p(\omega)$$
(11)

where $\chi(\omega)$ is the generalized susceptibility.

For the two baths at the same temperature the symmetrized correlation functions for the position $C_{qq}^{(+)}(t) \equiv \frac{1}{2} \langle [q(t), q(0)]_+ \rangle$ and the momentum $C_{pp}^{(+)}(t) \equiv \frac{1}{2} \langle [p(t), p(0)]_+ \rangle$ are obtained from $\chi(\omega)$ as follows

$$C_{qq}^{(+)}(t) = \frac{1}{\pi} \int_0^\infty |\chi(\omega)|^2 \cos(\omega t) \coth(\beta \omega/2)$$
$$\operatorname{Im}\left(\tilde{J}_q(\omega) \left|\omega_p + \tilde{J}_p(\omega)\right|^2 + \omega^2 \tilde{J}_p(\omega)\right) d\omega . \tag{12}$$

The corresponding expression for $C_{pp}^{(+)}(t)$ is obtained by interchanging $p \leftrightarrow q$ in Eq. (12). The antisymmetrized correlation functions for position and momentum $C_{nn}^{(-)}(t) \equiv \langle [n(t), n(0)]_{-} \rangle$ (with n = q, p) are obtained from Eq. (12) in the standard way [19], i.e. by substituting $\sin(\omega t)$ for $\cos(\omega t) \coth(\beta \omega/2)$.

We wish to emphasize that, within the dissipation model studied here, the equations of motion (6) are symmetric in q and p for arbitrary coupling strength. Thus the situation is different from that of a q-oscillator when described within the rotating-wave approximation, which is known to yield a model symmetric in q and p but which holds only in the weak coupling limit [33].

2.2. Ohmic coupling

First we focus on the important special case that both spectral densities are Ohmic, $\alpha_q = \alpha_p = 1$. Then the real parts of $\tilde{J}_q(\omega)$ and $\tilde{J}_p(\omega)$ vanish to lowest order in Ω^{-1} . The susceptibility has poles at the roots of the quadratic polynomial [5]

$$\omega_0^2 - i(\omega_q \gamma_p + \omega_p \gamma_q)\omega - (1 + \gamma_q \gamma_p)\omega^2 , \qquad (13)$$

which are

$$\omega_{\pm} = \frac{\omega_0}{\left(1 + \gamma_q \gamma_p\right)^{1/2}} \left(-i\kappa \pm \sqrt{1 - \kappa^2}\right) \equiv -i\tau^{-1} \pm \zeta , \qquad (14)$$

where

$$\kappa \equiv \frac{\omega_p \gamma_q + \omega_q \gamma_p}{2\omega_0 \left(1 + \gamma_q \gamma_p\right)^{1/2}}.$$
(15)

The solutions of Eq. (13) are either purely imaginary or a pair of complex conjugates depending on whether κ is greater or smaller than 1. Thus,

$$\kappa < 1$$
 Criterion A (16)

is the commonly accepted criterion to distinguish between underdamped and overdamped oscillations. The underdamped region lies in a stripe of width $\Delta = 4\eta (1 + \eta^4)^{-1/2}$, with $\eta \equiv (\omega_q/\omega_p)^{1/2}$, limited by the graphs of the functions

$$f(\gamma_q) = \frac{\gamma_q}{\eta^2} \pm \frac{2}{\eta} . \tag{17}$$

In Fig. 1 the stripes of underdamped oscillations, marked in the (γ_q, γ_p) plane are plotted for three different values of η . For large $\eta = m\omega_0$, corresponding to a large mass of the central oscillator, the stripe of underdamped oscillations becomes smaller. In the limit $\eta \to \infty$, the introduction of an infinesimal coupling to the momentum can induce a transition from overdamped to underdamped oscillations. However the range of values of γ_p allowing for underdamped oscillations also becomes increasingly small as $\propto \eta^{-1}$.

The behavior of the system as a function of γ_p for fixed γ_q is also interesting. We set $\eta = 1$. The inverse damping time τ^{-1} is a monotonously increasing function of γ_p for $\gamma_q < 2$ and a monotonously decreasing function for $\gamma_q > 2$. Therefore an additional bath coupling to p has opposite effects on the damping time τ depending on whether the original q-oscillator starts in the underdamped ($\gamma_q < 2$) or in the overdamped ($\gamma_q > 2$) regime. In the first case, the additional bath always reduces the damping time, leading to infinitely strong damping in the limit $\gamma_p \to \infty$. However, in the regime $\gamma_q > 2$ we can drive the system from the overdamped into the underdamped regime by *increasing* the coupling strength γ_p . What is more, we can take $\gamma_p = \gamma_q = \gamma$, which corresponds to a completely symmetric Hamiltonian in p and q. At this point the system is *always* in the underdamped regime and for large γ the oscillator frequency is close to its maximum. We find for the inverse damping time

$$\tau^{-1} = \frac{\omega_0 \gamma}{1 + \gamma^2} \ . \tag{18}$$

Therefore we are led to the paradoxical situation that the higher the friction coefficient γ the larger τ becomes. In particular, for $\gamma \to \infty$ one gets $\omega_0 \tau \to \infty$. This time dilatation is in itself a remarkable effect, since it contrasts with the behaviour of the *q*-oscillator, for which $\tau \to 0$ as $\gamma_q \to \infty$ [see Eq. (14)].

In the general, non-symmetric case, we note that for a given finite γ_q the damping time τ cannot be made to acquire arbitrary values by tuning γ_p . Rather, it is bounded between $2\gamma_q/\omega_0$ and $2/\gamma_q\omega_0$, which is higher depending on $\gamma_q \ge 1$.



Figure 1. Left: Stripes of underdamped oscillations in the (γ_q, γ_p) plane for three different values of the parameter $\eta = (\omega_q/\omega_p)^{1/2} = m\omega_0$, according to criterion A [Eq. 16]. Right: Regions of underdamped oscillations with criterions B [Eq. (19)] and C [Eq. (21)], for $\eta = 1$.

However, from these striking observations one should not conclude that in the limit of infinite γ the central oscillator recovers the dynamics of a free oscillator. It rather leads to a dilatation of all time scales. For instance, the renormalized oscillator frequency ζ vanishes even faster than the inverse damping time, since $\zeta = \omega_0/(1+\gamma^2)$. An analysis of the correlator in Eq. (12) shows that for $t \ll \tau$ the particle behaves as a free ballistic (though very slow) particle, $C_{qq}^{(+)}(t) \propto t/\tau$.

Instead of using criterion A, we may focus on $D_q(\omega) \equiv \text{Im}C_{qq}^{(-)}(\omega)/\omega$. As a spectral function, $\text{Im}C_{qq}^{(-)}(\omega)$ is independent of temperature. Of special interest is the slope of $D_q(\omega)$ near $\omega = 0$. Since $\lim_{\omega \to \infty} D_q(\omega) = 0$, the condition for the existence for $D_q(\omega)$ displaying a maximum can be written as

$$D'_a(0) > 0$$
 Criterion B, (19)

which may be viewed as an indicator of underdamped oscillations or, equivalently, coherent transitions. It is often employed for the spin–boson model [24, 26]. For Ohmic damping, one finds

$$D_q(\omega) = \frac{\gamma_q \omega_p^2 + \gamma_p (1 + \gamma_q \gamma_p) \omega^2}{[(1 + \gamma_q \gamma_p) \omega^2 - \omega_0^2]^2 + (\gamma_q \omega_p + \gamma_p \omega_q)^2 \omega^2} , \qquad (20)$$

and the critical curve for γ_p is thus given by the relation $\gamma_p^{\text{crit}} = \gamma_q (\gamma_q^2/\eta^2 - 2) /\eta^2$, so that criterion B is satisfied for $\gamma_p > \gamma_p^{\text{crit}}$.

Both being based on exact expressions, we observe that criterion B differs substantially from criterion A, see Fig. 1. Surprisingly, even for the q-oscillator there is a region $\sqrt{2} < \gamma_q < 2$ where criterion A and criterion B are different. In Fig. 2 $D_q(\omega)$ is plotted for different coupling strengths γ_q , γ_p . For large couplings γ_p , we observe that, although we are (according to criterion B) in the underdamped region, the maxima of



Figure 2. The spectral function $D_q(\omega)$ for different coupling strengths γ_p , γ_q . On the left, the marginal case of the *q*-oscillator, at $(\gamma_q, \gamma_p) = (\sqrt{2}, 0)$, (- - -), develops a maximum for $(\gamma_q, \gamma_p) = (\sqrt{2}, \sqrt{2})$, (----). On the right, for a value of $\gamma_q = 3$ (overdamped regime), when the values of γ_p are 0 (- - -), 21 (----) (marginal case), and 40 (....).

		symmetric	only γ_q	only γ_p
$\kappa < 1$	(A)	always	$\gamma_q < 2\eta$	$\gamma_p < 2\eta^{-1}$
$D_q(\omega)$ has peak	(B)	$\gamma < \sqrt{3}$	$\gamma_q < \sqrt{2}\eta$	always
$ \mathrm{Im}\omega_{\pm} < \mathrm{Re}\omega_{\pm} $	(C)	$\gamma < 1$	$\gamma_q < \sqrt{2}\eta$	$\gamma_p < \sqrt{2}\eta^{-1}$

Table 1. Condition for coherent dynamics according to three different criteria (left column, labeled A, B, C), for three particular cases (upper row). The symmetric limit includes the assumption $\eta = 1$. General case is given in main text.

the curves are not very pronounced. This is related to the fact that the renormalized oscillator frequency ζ vanishes for large γ_p faster than the inverse damping time τ^{-1} .

The foregoing discussion suggests the introduction of a third, more restrictive criterion for underdamped oscillations, namely [27],

$$\zeta > \tau^{-1}$$
 Criterion C. (21)

According to this condition, the region of underdamped oscillations is convex, i.e. by increasing the coupling strength no transition from overdamped to underdamped oscillations can be realized. Table I summarizes the different criteria for coherent, underdamped dynamics as they are applied to three prototypical cases.

For high temperatures $(k_B T \gg \omega_0)$ the integral for the coordinate autocorrelation function Eq. (12) can easily be solved yielding the classical solutions of the Heisenberg equations of motion. For $\kappa < 1$ they read

$$C_{qq}^{(+)}(t)/\langle q^2 \rangle = \left\{ \cos\left(\frac{\omega_0\sqrt{1-\kappa^2}}{\sqrt{1+\gamma_q\gamma_p}}t\right) + \frac{\omega_p\gamma_q - \omega_q\gamma_p}{2\omega_0\sqrt{(1-\kappa^2)}(1+\gamma_q\gamma_p)}\sin\left(\frac{\omega_0\sqrt{1-\kappa^2}}{\sqrt{1+\gamma_q\gamma_p}}t\right) \right\} \\ \exp\left(\frac{-\omega_0\kappa|t|}{\sqrt{1+\gamma_q\gamma_p}}\right).$$
(22)

For $\kappa > 1$, Eq. (22) describes the corresponding non-oscillating solution. Analytical expressions for $C_{qq}^{(+)}(t)$ are also available for T = 0. In this case the integral in Eq. (12) can be expressed in terms of exponential integrals [19]. We do not show the result here and limit ourselves to note that for long times only the lowest order term, i. e. the term linear in ω , contributes in the numerator of the r.h.s. of Eq. (12). Thus the long time behaviour is the same as for the q-oscillator, namely, we have

$$C_{qq}^{(+)}(t) = -\frac{\omega_p^2 \gamma_p}{2\pi} \frac{1}{\omega_0 t^2} + \mathcal{O}(t^{-3}) , \quad \text{for} \quad t \to \infty.$$
(23)

For the classical solution Eq. (22) criterion A [Eq. (16)] is the only natural criterion to differentiate between underdamped and overdamped oscillations, since it distinguishes solutions with infinitely many zeros from solutions with no zeros at all. For zero temperature the number of zeros of $C_{qq}^{(+)}$ is always finite and criterion A is less informative.

The zero temperature mean squares $\langle q^2\rangle,\,\langle p^2\rangle$ can be calculated exactly. They are given by

$$\langle q^2 \rangle = \frac{f(\kappa)}{4\kappa(1+\gamma_q\gamma_p)} \left[\frac{\gamma_q}{\eta^2} + \gamma_p \left(1-2\kappa^2\right) \right] + \frac{\gamma_p}{\pi} \left(1+\gamma_p\gamma_q\right) \ln \frac{\Omega_p \sqrt{1+\gamma_q\gamma_p}}{\omega_0} + \mathcal{O}(\Omega_p^{-1}, \Omega_q^{-1}),$$

$$f(\kappa) = \frac{1}{\pi\sqrt{\kappa^2-1}} \ln \frac{\kappa+\sqrt{\kappa^2-1}}{\kappa-\sqrt{\kappa^2-1}}.$$

$$(24)$$

The corresponding expressions for $\langle p^2 \rangle$ are obtained by interchanging $p \leftrightarrow q$ in Eqs. (24) and (15). For small γ_q, γ_p we have $f(\kappa) = 1 - 2\kappa/\pi + \mathcal{O}(\kappa^2)$ and the position mean square becomes

$$\langle q^2 \rangle = \frac{1}{2\eta} - \frac{\gamma_q}{2\eta^2} + \gamma_p \left(\ln \frac{\Omega_p}{\omega_0} - \frac{1}{2} \right) + \mathcal{O}(\gamma_q, \gamma_p) , \qquad (25)$$

and, correspondingly,

$$\langle p^2 \rangle = \frac{\eta}{2} - \frac{\eta^2}{2} \gamma_p + \gamma_q \left(\ln \frac{\Omega_q}{\omega_0} - \frac{1}{2} \right) + \mathcal{O}(\gamma_q, \gamma_p) , \qquad (26)$$

For $\gamma_q \neq 0$, $\gamma_p = 0$ we recover the results for the *q*-oscillator, with the characteristic logarithmic dependency of $\langle p^2 \rangle$ on the cutoff [19]. In the general case, the Heisenberg uncertainty product diverges as $\langle p^2 \rangle \langle q^2 \rangle \propto \ln \Omega_p \ln \Omega_q$. Thus the reduced density matrix at equilibrium becomes approximately the identity, its off-diagonal elements being essentially zero in both the position and the momentum representation.

2.3. Other spectral Densities

For general spectral densities α_q and α_p are arbitrary positive real numbers. While we always have

$$\operatorname{Im}\widetilde{J}_{n}(\omega) = \frac{\pi}{2}\operatorname{sgn}(\omega)J_{n}(|\omega|) , \qquad (27)$$

we find

$$\operatorname{Re}\widetilde{J}_{n}(\omega) = \begin{cases} 0, & \text{if } \alpha_{n} < 2, \\ \frac{\gamma_{n}\omega^{2}}{\pi\omega_{\mathrm{ph}}} \ln \frac{\Omega_{n}^{2}}{\omega^{2}}, & \text{if } \alpha_{n} = 2, \\ \frac{2\gamma_{n}\Omega_{n}^{\alpha_{n}-2}}{\pi(\alpha_{n}-2)\omega_{\mathrm{ph}}^{\alpha_{n}-1}} \omega^{2}, & \text{if } \alpha_{n} > 2, \\ \end{cases}$$
(28)

First, we briefly recall the behavior of $\langle q^2 \rangle$ and $\langle p^2 \rangle$ as functions of α_q and γ_q in the case of the *q*-oscillator. For $\langle q^2 \rangle$ it may be summarized by the formula [19]

$$\frac{d\langle q^2 \rangle}{d\gamma_q} \begin{cases} < 0 & \text{for } \alpha_q < 2\\ \propto -\frac{1}{\Omega} & \text{for } \alpha_q \ge 2 \end{cases}$$
(29)

A measurement of the position, i.e. a diagonalization of the density matrix in the position basis takes place only for $\alpha_q < 2$. The behaviour of $\langle p^2 \rangle$ is opposed to that

of $\langle q^2 \rangle$ in such a way that the product $\langle q^2 \rangle \langle p^2 \rangle$ is always $\geq 1/4$, as required by the Heisenberg relation. For $\langle p^2 \rangle$ we may write

$$\frac{d\langle p^2 \rangle}{d\gamma_q} \begin{cases} \propto \ln\left(\frac{\Omega}{\omega_0}\right) & \text{for } \alpha_q \le 2\\ > 0 & \text{for } \alpha_q > 2 \end{cases}$$
(30)

In the forthcoming discussion we focus on $\langle q^2 \rangle$. We have a contribution from $J_q(\omega)$, which reduces $\langle q^2 \rangle$, and another contribution from $J_p(\omega)$ enhancing $\langle q^2 \rangle$ [see Eq. (25)]. From the results for the *q*-oscillator we expect that the $J_p(\omega)$ contribution should always dominate for $\alpha_p < 2$ and the $J_q(\omega)$ contribution should be negligible for $\alpha_q \geq 2$. The opposite regime $\alpha_q < 2$ and $\alpha_p > 2$ is more interesting (the case $\alpha_q = 1$ and $\alpha_p = 3$ has been studied in Ref. [21]). There we have a sum of two terms of order 1 in Ω^{-1} with opposite sign. In the following we focus on that case.

We assume the susceptibility $\chi(\omega)$ to be an algebraic function. This entails rational exponents $\alpha_n = \beta_n/m$ with integer β_n and m. The integrand in Eq. (12) has $r \equiv \max(\beta_q + \beta_p, 2m)$ poles on the m Riemann sheets which split into two classes. For $\alpha_q, \alpha_p > 1$ there are 2m poles close to the oscillator frequency $\pm \omega_0$ on each sheet. In addition there are $r_{\Omega} \equiv \max(\beta_q + \beta_p - 2m, 0)$ poles of the order of $i\Omega$. If either α_q or α_p becomes equal or smaller one, the 2m poles detach from $\pm \omega_0$ moving into the complex plane. The equilibrium mean square can be written as a sum of the contributions from the two types of poles.

$$\langle q^2 \rangle = \mathcal{C}_{\omega_0} + \mathcal{C}_{\Omega} \ . \tag{31}$$

For the second term we find $(\gamma_q, \gamma_p \text{ small})$

$$C_{\Omega} = \frac{\gamma_p}{\pi} \frac{1}{\alpha_p - 1} \Theta(\alpha_p - 1) + \mathcal{O}(\gamma_p, \gamma_q) .$$
(32)

For a sketch of the derivation of Eq. (32) see Appendix A. Therefore, the high energy poles yield, to first order in γ_p , a finite contribution to the position mean square. This contribution is positive for superohmic coupling to the momentum but vanishes for Ohmic and subohmic coupling. It has a singularity for $\alpha_p = 1$ which marks the transition to the logarithmic dependence on the cutoff frequency, see Eq. (24). We note that the right hand side of Eq. (32) does not depend on the oscillator frequency ω_0 , i. e. it does not depend on the properties of the system itself. The contribution from the poles close to ω_0 , \mathcal{C}_{ω_0} , can in principle be calculated in a similar way as \mathcal{C}_{Ω} for arbitrary exponents α_q , α_p ; however, the calculations become increasingly messy. A general treatment is also hampered by the wide variety of casuistic behavior, with different regimes defined by the conditions $\alpha_n \geq 1$, $\alpha_n \geq 2$ and $\alpha_q + \alpha_p \geq 2$, $\alpha_q + \alpha_p \geq 4$.

We focus on the specific case $\alpha_q \leq 1$ and $\alpha_p > 2$, which is the most relevant, since most baths occuring in nature are either Ohmic (Markovian approximation, electron gas [44]) or superohmic (photon or acoustic phonon baths). In that case, the low energy poles are essentially determined by the lower spectral exponent (here, α_q). This reflects the general property of dissipative quantum systems that an environment becomes



Figure 3. Left: Zero temperature mean square $\langle q^2 \rangle$ as a function of γ_q and γ_p for $\alpha_q = 1$ (Ohmic) and $\alpha_p = 3$ (superohmic). Right: Zero temperature mean square $\langle q^2 \rangle$ as a function of γ_q for $\alpha_q = 1$, $\gamma_p = 1$ and different spectral exponents $\alpha_p = 3$ (- - -), $\alpha_p = 2.2$ (- -) and $\alpha_p = 1.8$ (· · · ·). The full line corresponds to the q-oscillator $\gamma_p = 0$. The cutoff frequency is $\Omega/\omega_0 = 200$ in both figures. The case $\alpha_p = 2$ is avoided because of its singular character [see Eq. (28)].

increasingly efficient with decreasing spectral exponent α_n . To leading order in γ_q , we find

$$C_{\omega_0} = \frac{1}{2} - \frac{\gamma_q}{\pi \alpha_q^2} \left[\frac{1}{2} + 3\alpha_q \left(\alpha_q - 1 \right) \right] + \mathcal{O}(\gamma_q^2) .$$
(33)

If we compare C_{ω_0} with Eq. (32), we observe that the two contributions have opposite sign. In particular, the contributions of C_{ω_0} and C_{Ω} cancel each other to first order in γ_q and γ_p provided that

$$\gamma_p = \frac{\alpha_p - 1}{\alpha_q^2} \left[\frac{1}{2} + 3\alpha_q \left(\alpha_q - 1 \right) \right] \gamma_q .$$
(34)

In Fig. 2.3 $\langle q^2 \rangle$ is plotted against the coupling strengths γ_p and γ_q for $\alpha_q = 1$ and $\alpha_p = 3$. $\langle q^2 \rangle$ is monotonously increasing as a function of γ_p . For $\gamma_p = 0$ it is a monotonously decreasing function of γ_q . For the parameters chosen in Figure 2.3, Equation (34) holds on the diagonal $\gamma_q = \gamma_p$. There $\langle q^2 \rangle$ remains close to its unperturbed value 1/2 if $\gamma_q, \gamma_p \lesssim 1$.

Finally, on the right hand side of Figure 2.3 the position mean square $\langle q^2 \rangle$ is plotted as a function of γ_q for different spectral exponents α_p . For small γ_q the enhancement of $\langle q^2 \rangle$ due to the coupling to the momentum is larger for smaller values of the spectral exponent α_p . This is the expected behaviour. This behavior, however, is inverted when γ_q increases. Then the relative enhancement of $\langle q^2 \rangle$ as compared with the *q*-oscillator is bigger for higher spectral exponent α_p . A crossing occurs always for some value of the coupling constant γ_q , although some times this can be very large.

3. Time evolution

To study the nonequilibrium properties and in particular the loss of coherence for an initially pure state, the calculation of $C_{qq}^{(+)}(t)$ is not sufficient. Instead, one should perform a full nonequilibrium calculation with inclusion of specific initial conditions by means of a Laplace transformation. The position-momentum symmetry of the oscillator Hamiltonian suggests the use of the reduced Wigner-function W(q, p, t) instead of the reduced density matrix. It was shown in Ref. [34] for the q oscillator that the time evolution of W(q, p, t) can only for certain initial conditions be described by an exact master equation. Here we consider decoupled initial conditions $W_0 = W_S W_B^{(q)} W_B^{(p)}$, which fall into this class. In this case we are able to derive not only an exact Master equation for the reduced Wigner function but a solution in terms of a two-fold convolution integral.

Generally W(q, p, t) can be expressed as

$$W(q, p, t) = \langle \delta(q - q(t))\delta(p - p(t)) \rangle_0 , \qquad (35)$$

where the bracket denotes the average over initial conditions [11]

$$\langle \ldots \rangle_0 \equiv \int dq' dp' \prod_{k,k'} da_{qk} da_{pk'} da_{qk}^* da_{pk'}^* (\ldots) W_S(q', p', \{\mathbf{a}_q, \mathbf{a}_p, \mathbf{a}_q^*, \mathbf{a}_p^*\}) , \qquad (36)$$

and q(t), p(t) are the classical solutions of the equations of motion Eq. (6). The Wigner function of the thermalized baths is given by

$$W_B^{(n)} = \prod_k \frac{i}{\pi \coth(\beta \omega_k/2)} \exp\left(-\frac{2a_{nk}^* a_{nk}}{\coth(\beta \omega_k/2)}\right) , \qquad (37)$$

the system itself being in an arbitrary pure state characterized by W_S . In this case one can express Eq. (35) as a twofold convolution integral

$$W(q, p, t) = \int \frac{dk}{2\pi} \frac{dk'}{2\pi} e^{ikp+ik'q} \mathcal{W}_S(k\phi_0(t) + k'\phi_p(t), k'\phi_0(t) - k\phi_q(t))$$

$$\exp\left(-k^2 X_{pq}(t) - k'^2 X_{p0}(t) + 2kk' Y_{pq}(t)\right)$$

$$\exp\left(-k'^2 X_{qp}(t) - k^2 X_{q0}(t) - 2kk' Y_{qp}(t)\right) , \qquad (38)$$

where \mathcal{W}_S is the Fourier transform of W_S in both arguments. We have introduced the auxiliary functions

$$\phi_n(t) \equiv \frac{1}{2\pi i} \int d\omega \chi(\omega) \left(\omega_n + \widetilde{J}_n(\omega) \right) e^{i\omega t} , \qquad n = q, p$$

$$\phi_0(t) \equiv \frac{1}{2\pi i} \int d\omega \omega \chi(\omega) e^{i\omega t} , \qquad (39)$$

and the temperature-dependent quantities

$$X_{pq}(t) = \frac{1}{2} \int_{0}^{\infty} d\omega J_{p}(\omega) \coth(\beta\omega/2) \left| \int_{0}^{t} dt \phi_{q}(t-t') e^{i\omega t} \right|^{2}$$

$$X_{p0}(t) = \frac{1}{2} \int_{0}^{\infty} d\omega J_{p}(\omega) \coth(\beta\omega/2) \left| \int_{0}^{t} dt \phi_{0}(t-t') e^{i\omega t} \right|^{2}$$

$$Y_{pq}(t) = \frac{1}{2} \int_{0}^{\infty} d\omega J_{p}(\omega) \coth(\beta\omega/2)$$

$$\operatorname{Re} \left[\int_{0}^{t} dt \phi_{0}(t-t') e^{i\omega t} \int_{0}^{t} dt \phi_{q}(t-t') e^{-i\omega t} \right].$$
(40)

The functions X_{q0} , X_{qp} and Y_{qp} are defined accordingly by interchanging p and q. For a better understanding of the forthcoming discussion in Sec. 4 we notice that the functions defined in Eq. (39) are the time-dependent coefficients of the solutions q(t), p(t) of the initial value problem Eq. (6). For instance

$$q(t) = \phi_0(t)q(0) + \phi_p(t)p(0) + \int_0^t \phi_0(t-t')F_p(t')dt' + \int_0^t \phi_p(t-t')F_q(t')dt' .$$
(41)

This necessarily requires $\phi_0(0) = 1$ and $\phi_n(0) = 0$. Finally, we note that W fulfills a Fokker–Planck type of equation [19],

$$\dot{W}(q, p, t) = \nabla \left[\mathbf{g}(q, p, t) + \mathbf{B}(t) \nabla \right] W(q, p, t) , \qquad (42)$$

where $\nabla \equiv (\partial/\partial q, \partial/\partial p)$, with $\mathbf{g} = (g_q, g_p)$ the phase-space drift term,

$$g_n(q, p, t) = -\sum_{m=q,p} G_{nm}(t)m$$
, (43)

$$\mathbf{G}(t) \equiv \frac{\begin{pmatrix} \dot{\phi}_0 \phi_0 + \dot{\phi}_p \phi_q & \dot{\phi}_p \phi_0 - \dot{\phi}_0 \phi_p \\ \dot{\phi}_0 \phi_q - \dot{\phi}_q \phi_0 & \dot{\phi}_0 \phi_0 + \dot{\phi}_q \phi_p \end{pmatrix}}{\phi_0^2 + \phi_q \phi_p} , \qquad (44)$$

and with (the 2×2 matrix) $\mathbf{B}(t)$ the state-independent phase-space diffusion term,

$$B_{pp}(t) = \dot{X}_{pq} + \dot{X}_{q0} - 2F_{pp}(X_{pq} + X_{q0}) + 2F_{pq}(Y_{pq} - Y_{qp})$$
(45)

$$B_{qq}(t) = \dot{X}_{qp} + \dot{X}_{p0} - 2F_{qq}(X_{qp} + X_{p0}) - 2F_{qp}(Y_{qp} - Y_{pq})$$

$$B_{pq}(t) = \dot{Y}_{qp} - \dot{Y}_{pq} - (F_{pp} + F_{qq})(Y_{qp} - Y_{pq})$$

$$-F_{pq}(X_{p0} + X_{qp}) - F_{qp}(X_{pq} + X_{q0}).$$
(46)

 $B_{qp}(t)$ is obtained from $B_{pq}(t)$ by exchanging q and p. To derive the coefficients (44) and (45) we have employed the ansatz Eq. (42) for the Fokker–Planck operator. Acting with yet undefined functions f_n, B_{nm} (n, m = p, q) on the r.h.s. of Eq. (42), and comparing with its time derivative, yields conditions for the functions f_n, B_{nm} . An exact Fokker– Planck equation in the form of Eq. (42) for the q-oscillator has been first derived in Ref. [11] and later by a different method in Refs. [13, 16].

3.1. Purity

A convenient quantity to measure the degree of global decoherence is the purity $\mathcal{P}(t)$, defined as the average of the density matrix itself [17],

$$\mathcal{P}(t) \equiv \langle \rho(t) \rangle = \operatorname{tr} \rho^2(t) , \qquad (47)$$

which is basis independent. Of special interest is the equilibrium value $\mathcal{P}_{\beta} \equiv \lim_{t\to\infty} \mathcal{P}(t)$ which measures the efficiency of the environment in destroying quantum coherence. Here we implicitly assume ergodic behavior which, as we shall see, applies in the presence of Ohmic baths. For a harmonic oscillator in thermal equilibrium, the reduced Wigner function is [19]

$$W_{\beta}(q,p) = \frac{1}{2\pi \left[\langle q^2 \rangle_{\beta} \langle p^2 \rangle_{\beta}\right]^{1/2}} \exp\left(-\frac{q^2}{2\langle q^2 \rangle_{\beta}} - \frac{p^2}{2\langle p^2 \rangle_{\beta}}\right),\tag{48}$$

which leads to

$$\mathcal{P}_{\beta} = \frac{\hbar}{2\left[\langle q^2 \rangle_{\beta} \langle p^2 \rangle_{\beta}\right]^{1/2}} \,. \tag{49}$$

4. Coherence decay for Ohmic damping

Although the equilibrium decoherence (as measured by the product $\langle q^2 \rangle_\beta \langle p^2 \rangle_\beta$) is enhanced by the additional noise term, one may wonder whether for low temperatures the decoherence time becomes larger than for the damped *q*-oscillator. To answer this question exhaustively one would have to calculate the time evolution of the purity for an arbitrarily pure initial condition. We consider two cases. First, we choose a coherent (Gaussian) state as initial state. This case should present the greatest robustness against decoherence [17, 35]. Second, we choose a superposition of two Gaussian wave packets. Then a new aspect of decoherence comes into play, namely, the fast vanishing of the relative coherence between the two Gaussian wave packets. We distinguish the two manifestations of decoherence by introducing a new quantity, which we call *relative purity*.

4.1. Decoherence for a Gaussian wave packet as initial state

We consider here the case where the system starts in a coherent state at t = 0. The Wigner function for a Gaussian wave packet is

$$W_S(q,p) = \frac{1}{\pi} \exp\left[-\eta \left(q - q_0\right)^2 - \frac{1}{\eta} \left(p - p_0\right)^2\right] , \qquad (50)$$

where $p_0 = \sqrt{2\eta} \text{Im}(\alpha)$ and $q_0 = \sqrt{2/\eta} \text{Re}(\alpha)$ are defined in terms of the complex eigenvalue α (amplitude) of the coherent state (recall $\eta = \sqrt{\omega_q/\omega_p}$). Since $W_S(q, p, 0)$ is Gaussian, the convolution integral of Eq. (38) can be performed. The integrals become

particularly simple when both baths are equal $(\gamma_p = \gamma_q = \gamma \text{ and } \eta = 1)$, corresponding to the completely symmetric case. Then we have $\phi_p = \phi_q \equiv \phi_1$ and, by the same token, $X_{pq} = X_{qp} \equiv X_1$ in Eq. (39) and also $X_{p0} = X_{q0} \equiv X_0$ in Eq. (40). The crossed terms in Eq. (38) vanish and the Wigner function becomes a product of two Gaussians for all times

$$W(q, p, t) = \frac{1}{4\pi} \frac{1}{\phi_0^2 + \phi_1^2 + 4X_0 + 4X_1} \exp\left[-\frac{(p - p_0\phi_0 + q_0\phi_1)^2 + (q - q_0\phi_0 + p_0\phi_1)^2}{\phi_0^2 + \phi_1^2 + 4X_1 + 4X_0}\right].$$
(51)

For the purity one obtains the simple expression

$$\mathcal{P}(t) = \frac{1}{\phi_0^2 + \phi_1^2 + 4X_0 + 4X_1} \,. \tag{52}$$

The remaining task is to calculate the quantities ϕ_0 , ϕ_1 , X_0 , X_1 from their definition in Eqs. (39) and (40). Using the naïve form $(\Omega \to \infty)$ of the spectral function of an Ohmic bath as given in Sec. 2, we would find $\phi_0(0) = 1/(1 + \gamma_p \gamma_q) \neq 1$ leading to inconsistencies [see the discussion after Eq. (40)]. The reason for this lies in an initial slippage caused by the somewhat unphysical character of the decoupled initial condition [36, 37, 38, 39]. Technically it stems from the fact that for $J(\omega) \propto \omega$ the integrals in Eqs. (39) do not converge at t = 0 [11]. This problem is overcome by regularizing the Ohmic spectral functions, i.e. by reintroducing a finite cutoff Ω . The explicit calculation with a Drude regularized spectral function is sketched in Appendix C. Here we state only the main result: At T = 0 the purity decays on two time scales, given by Ω^{-1} and τ in (18),

$$\mathcal{P}(t) \simeq \begin{cases} e^{-\Omega t}, & \text{for } 0 \le t \lesssim \Omega^{-1} \\ \left[\mathcal{P}_{\beta} + \frac{1}{\omega_0^2 t^2} \frac{2\gamma}{1+\gamma^2} \cos(\Lambda t) e^{-t/\tau} \right], & \text{for } \Omega^{-1} \ll t \to \infty \end{cases}$$
(53)

with $\mathcal{P}_{\beta}^{-1} = 2\langle q^2 \rangle_{\beta}$ and the oscillator frequency $\Lambda = \omega_0/(1 + \gamma^2)$. Coherence is reduced immediately after the start of the coupling. Although afterwards it decreases more slowly, on a time scale τ , for larger couplings the curves of $\mathcal{P}(t)$ for different values of γ never cross, i.e., $\mathcal{P}(t)$ is a monotonously decreasing function of γ for all t.

An initial slip similar to that discussed in this section also occurs for the q-oscillator [11]. However in that case its effect on purity is much less severe. Specifically, out of the set of functions $\phi_q(t)$, $\phi_p(t)$ and $\phi_0(t)$ defined in Eq. (39), only $\phi_q(t)$ is affected by the initial slip. After a time Ω^{-1} it becomes $\phi_q(t) \sim \gamma_p \neq 0$ [11]. However, we note that ϕ_q appears in the temperature-dependent terms X_{qp} , X_{pq} , Y_{qp} , Y_{pq} of Eq. (40) only in combination with the spectral density of the momentum coupling $J_p(\omega)$, which vanishes for the q-oscillator by definition. We thus reach the important conclusion that the purity evolution of the q-oscillator is insensitive to the initial slip stemming from the use of decoupled initial conditions: At $t \sim \Omega^{-1}$ the purity is still approximately unity, decreasing afterwards at a rate $\propto \gamma_q$.

Isar and coworkers [40] studied in detail the decay of purity for the q-oscillator. In particular they found constraints that must be satisfied by the bath if the purity is to

remain constant and close to unity during the whole time evolution of the oscillator. Such a high purity is not possible in the present case due to the initial slip [see Eq. (53].

Another possible preparation of the initial state is in a constrained equilibrium. The expectation values $\langle q(0) \rangle = q_0$ and $\langle p(0) \rangle = p_0$ are held fixed with the bath equilibrated around those values. Then the Wigner function is [11]

$$W(q, p, t) = \frac{1}{2\pi \left[\langle q^2 \rangle_\beta \langle p^2 \rangle_\beta \right]^{1/2}} \exp\left(-\frac{(q - \langle q(t) \rangle)^2}{2\langle q^2 \rangle_\beta} - \frac{(p - \langle p(t) \rangle)^2}{2\langle p^2 \rangle_\beta} \right)$$
(54)

while the purity is given by its equilibrium value Eq. (49). A more detailed discussion of the purity decay of the q-oscillator, together with the general formulae, is given in Appendix C.

4.2. Decoherence of two Gaussian wave packets

Formula (38) also allows us to investigate more complicated initial conditions such as as, for example, the superposition of two Gaussian wave packets. This case has been studied for a single bath by Caldeira and Leggett [41]. It is an interesting case study because it displays two different aspects of decoherence. On the one hand, there is the decoherence which either wave packet would experience alone. This part is essentially described by $\mathcal{P}(t)$, which we will call *Gaussian purity* [see Eqs. (52) and (53)]. On the other hand there is the decoherence due to the spatial separation of the two packets. This second contribution is expected to become increasingly important when the distance *a* between the two packets becomes large. As initial wave function we choose

$$\psi(x) = \frac{1}{c\sqrt[4]{2\pi\sigma^2}} \left(e^{-\frac{(x+a/2)^2}{4\sigma^2}} + e^{-\frac{(x-a/2)^2}{4\sigma^2}} \right) , \qquad (55)$$

which translates into an initial Wigner function

$$W_S(q,p) = \frac{1}{\pi c^2} e^{-\frac{q^2}{2\sigma^2} - 2\sigma^2 p^2} \sum_{k=\pm 1} \left(e^{-\frac{a^2}{8} - \frac{kaq}{2\sigma^2}} + e^{-ikap} \right) .$$
(56)

Here $c \equiv \sqrt{2} \left[1 + \exp(-a^2/8\sigma^2)\right]^{1/2}$ is a normalization constant. We assume the most symmetric case and set $\sigma = 1/\sqrt{2}$. Plugging (56) into (38) we find that the purity can be expressed in terms of the Gaussian purity as

$$P(t) = \frac{\mathcal{P}(t)}{2} \left\{ 1 + \frac{\cosh^2 \left[\frac{a^2}{4} (\phi(t)\mathcal{P}(t) - \frac{1}{2}) \right]}{\cosh^2 \left(\frac{a^2}{8} \right)} \right\}.$$
 (57)

The function $\phi(t)$ is given by

$$\phi(t) = \phi_0^2(t) + \phi_1^2(t) , \qquad (58)$$

where ϕ_0 and ϕ_1 are defined in (39) and computed in (B.2) for the symmetric case. $\phi(t)$ evolves from $\phi(0) = 1$ to $\lim_{t\to\infty} \phi(t) = 0$. We define the *relative purity* as the ratio

$$P_{\rm rel}(t) = P(t)/\mathcal{P}(t) \tag{59}$$

As expected, $P_{\rm rel}(t) \to 1$ as $a \to 0$, and $P_{\rm rel}(t) \to 1/2$ as $a \to \infty$.

Interestingly, the structure of (57) is such that, as time passes and $\phi(t)\mathcal{P}(t)$ evolves from 1 to 0, the ratio $P_{\rm rel}(t)$ starts at unity, as corresponds to a pure state, then decreases and finally, at long times, goes back to unity. When *a* is large $P_{\rm rel}(t)$ decays rapidly to 1/2, on a timescale $\sim 1/4a^2\gamma$. There it stays for a time which increases with distance as $\sim \gamma^{-1} \ln a$. Afterwards it returns to one. The ratio 1/2 can be rightly interpreted as resulting from the incoherent mixture of the two wave packets. Thus it comes as a relative surprise that $P_{\rm rel}(t)$ becomes unity again at long times, as if coherence among the two wave packets were eventually recovered. The physical explanation lies in the ergodic character of the long time evolution, with both wave packets evolving towards the equilibrium configuration [see Eq. (24) and the subsequent discussion].

5. Single bath with linear coupling to both position and momentum

It is instructive to compare Eq. (3) with the Hamiltonian of a harmonic oscillator interacting with a single heat bath in the most general form of linear coupling,

$$H = \frac{\omega_q}{2}q^2 + \frac{\omega_p}{2}p^2 + \sum \omega_k \left| a_k + \frac{\lambda_k}{\omega_k}q + \frac{\mu_k}{\omega_k}p \right|^2 , \qquad (60)$$

with complex parameters λ_k, μ_k . This model is different from that discussed in the previous sections in that here time reversal invariance is broken. By this we mean the following. In the models described by Eq. (3), the bath modes might describe for example a magnetic field coupled to the momentum of a charged particle, which clearly would break time reversal invariance. However in that case, such a symmetry breaking is somewhat fictitious since, due to the linear nature of the coupling and to the modelling of the bath as a set of harmonic oscillators, one can always find a unitary transformation which restores time reversal invariance, i.e. which renders all parameters in the Hamiltonian real quantities. It is easy to see that, for general (complex) λ_k and μ_k , such a unitary transformation cannot be found for Eq. (60).

By an analysis similar to that of Section 2 one finds general expressions for the symmetrized equilibrium correlation functions:

$$C_{qq}^{(+)}(t) = \frac{1}{\pi} \int_{0}^{\infty} d\omega |\chi(\omega)|^{2} \cos(\omega t) \coth(\hbar \beta \omega/2) \left\{ \operatorname{Im} \widetilde{J}_{q}(\omega) \left[\omega_{p} + \operatorname{Re} \widetilde{J}_{p}(\omega) \right]^{2} + \operatorname{Im} \widetilde{J}_{p}(\omega) \left[\left(\omega + \operatorname{Re} \widehat{J}_{-}(\omega) \right)^{2} + \left(\operatorname{Re} \widetilde{J}_{+}(\omega) \right)^{2} \right] - \left[\omega_{p} + \operatorname{Re} \widetilde{J}_{p}(\omega) \right] \operatorname{Im} \left[2\omega \widehat{J}_{-}(\omega) - \widehat{J}_{-}^{2}(\omega) - \widetilde{J}_{+}^{2}(\omega) \right] \right\}.$$
(61)

The same expression applies for $C_{pp}^{(+)}(t)$ with the indexes p, q interchanged everywhere. As before, we define a generalized "susceptibility" for the system which now reads

$$\chi^{-1}(\omega) = \left[\omega_q - \widetilde{J}_q(\omega)\right] \left[\omega_p - \widetilde{J}_p(\omega)\right] - \omega^2 - \widetilde{J}_+^2(\omega) - \widehat{J}_-^2(\omega) + 2\omega \widehat{J}_-(\omega) .$$
(62)

Eqs. (61) and (62) involves four different spectral functions. $\tilde{J}_q(\omega)$ and $\tilde{J}_p(\omega)$ are defined as in Eqs. (4) and (10). Now we introduce the spectral functions

$$J_{+}(\omega) = \sum (\lambda_{k}\mu_{k}^{*} + \lambda_{k}^{*}\mu_{k})\delta(\omega - \omega_{k})$$

$$= 2\sum |\lambda_{k}||\mu_{k}|\cos\theta_{k}\delta(\omega - \omega_{k})$$

$$J_{-}(\omega) = i\sum (\lambda_{k}\mu_{k}^{*} - \lambda_{k}^{*}\mu_{k})\delta(\omega - \omega_{k})$$

$$= 2\sum |\lambda_{k}||\mu_{k}|\sin\theta_{k}\delta(\omega - \omega_{k}) , \qquad (63)$$

which reflect the mixing of the bath modes. The "hat" symbol denotes the transformation

$$\widehat{f}(\omega) \equiv \omega \mathcal{P} \int_0^\infty \frac{f(\omega')}{{\omega'}^2 - \omega^2} d\omega' + i\frac{\pi}{2} f(|\omega|) \quad , \tag{64}$$

In Eq. (64) the real part is an antisymmetric function and the imaginary part is symmetric. This is exactly reverse to the "tilde" (Riemann) transform, defined in Eq. (10).

Before we calculate Eq. (61) for a specific case, we analyze some generic features. First we notice that, by setting $\hat{J}_{-}(\omega)$ and $\tilde{J}_{+}(\omega)$ to zero, we recover the autocorrelation function for two independent baths [see Eq. (12)]. On the other hand, the two spectral densities $J_{+}(\omega)$ and $J_{-}(\omega)$ are not independent from $J_{p}(\omega)$ and $J_{q}(\omega)$; rather, they satisfy

$$J_{+}^{2}(\omega) + J_{-}^{2}(\omega) = J_{p}(\omega)J_{q}(\omega) .$$
(65)

Actually this relation has already been used to simplify the integrand of Eq. (61). We note that it holds for the spectral densities themselves but in general *not* for their Riemann transforms $\tilde{J}_{+}(\omega)$ and $\tilde{J}_{-}(\omega)$. Apart from the above constraint one can, at least in principle, freely choose three of the four spectral functions. If, in addition, we make the physically reasonable assumption that the "mixing angle" in Eq. (63) is frequency independent ($\theta_k = \theta$), condition (65) fixes $J_{+}(\omega)$ and $J_{-}(\omega)$ completely.

$$J_{+}(\omega) = \left[J_{q}(\omega)J_{p}(\omega)\right]^{1/2}\cos\theta$$

$$J_{-}(\omega) = \left[J_{q}(\omega)J_{p}(\omega)\right]^{1/2}\sin\theta .$$
(66)

Therefore, in the important case where both $J_q(\omega)$ and $J_p(\omega)$ have the same spectral exponent ($\alpha_q = \alpha_p = \alpha$), J_+ and J_- also obey the same power law. Using Eq. (65) we observe that the term (Im \widetilde{J}_q Im \widetilde{J}_p) implicit in Eq. (62) drops out. We recall that this interference term has been responsible for the non trivial "phase–space" diagram Fig. 1. What is more, only Re \widetilde{J}_+ appears in Eqs. (61) and (62). That means, according to Eq. (28), that for $\alpha_q + \alpha_p < 4$ the symmetric spectral function J_+ does not contribute at all. The relation between the "hat" and the "tilde" transformation is

$$\widehat{f}(\omega) = \widetilde{\omega f(\omega)} , \qquad (67)$$

from which all properties of \hat{f} can easily be deduced. For example from Eq. (67) and Eqs. (28) and (27) it follows that $\operatorname{Re} \widehat{J}_{-}(\omega) = 0$ only for $\alpha_q + \alpha_p < 2$.

It is illustrative to look at the "generalized" susceptibility Eq. (62) for the case $\alpha_q + \alpha_p < 2$, where all real parts of the spectral functions vanish. We obtain

$$\chi^{-1}(\omega) = \omega_0^2 - \omega^2 - \omega_q \widetilde{J}_p(\omega) - \omega_p \widetilde{J}_q(\omega) + 2\omega \widehat{J}_-(\omega)$$
$$= \omega_0^2 - \omega^2 - \operatorname{sgn}(\omega) \frac{i\pi\omega_p}{2} \left[J_q(|\omega|) + \eta^2 J_p(|\omega|) - \frac{2|\omega|}{\omega_p} J_-(|\omega|) \right] .$$
(68)

This is exactly the suceptibility of a q-oscillator coupled to a single bath with an additive spectral function

$$J(\omega) = J_q(\omega) + \eta^2 J_p(\omega) + \frac{2\omega J_-(\omega)}{\omega_p}.$$
(69)

We conclude that, when only one bath is involved, the particular structure of the coupling of the bath to the position or momentum variable can always be modelled by a q-oscillator with an appropriately chosen spectral function. In this context the combination $\omega J_{-}(\omega)$ can be considered as stemming from an effective additional noise source that couples to position. Note however, that this is not the same as saying that a single-bath coupling to q and p can always be unitarily transformed to a coupling only to q (q-oscillator), since this is not true in general. For one thing, the noise properties, which are not uniquely given by $\chi(\omega)$, would be different.

We also wish to emphasize that a double-bath dissipative oscillator cannot be described in terms of an oscillator coupled to an effective single bath. In particular, it can never be modelled by a susceptibility like (69). This is the reason why the physics explored in Section 2 is so different from that which could be found in any possible single-bath scenario.

We do not repeat here the analysis of Section 2 for arbitrary spectral functions but focus instead on the case of Ohmic coupling ($\alpha_q = \alpha_p = 1$) in order to compare with the results obtained for two independent baths. As $\operatorname{Re}\widetilde{J}_n(\omega) = \operatorname{Re}\widetilde{J}_+(\omega) = 0$ vanish [see Section 2.3], we obtain

$$C_{qq}^{(+)}(t) = \frac{1}{\pi} \int_0^\infty d\omega |\chi(\omega)|^2 \cos(\omega t) \coth(\hbar\beta\omega/2)$$

$$\operatorname{Im} \left\{ \omega_q^2 \widetilde{J}_q(\omega) + \widetilde{J}_p(\omega) \left[\omega - \operatorname{Re} \widehat{J}_-(\omega) \right]^2 - 2\omega_p \widehat{J}_-(\omega) \left[\omega - \operatorname{Re} \widehat{J}_-(\omega) \right] \right\}.(70)$$

The form of the susceptibility varies depending on whether the mixing angle is a multiple of π or not. We discuss the two cases separately.

5.1. Mixing angle $\theta = 0, \pi$

For $\theta = 0$ or $\theta = \pi$ one can bring, by an appropriate redefinition of a_k, a_k^{\dagger} , the interaction part of the Hamiltonian Eq. (60) into the form

$$H_I \propto \sum q_k (|\mu_k| p + |\lambda_k| q) , \qquad (71)$$

so that the bath couples to the main oscillator either through the position of their oscillators or through their momentum, but not through both. From Eq. (66) it can be seen that the antisymmetric spectral function vanishes identically. Apart from the



Figure 4. Regions of underdamped oscillations for a single bath coupling to position and momentum (dark region) and for two independent baths coupling to position and momentum (dark and bright region). $\eta = 1$.

susceptibility, the integrand of Eq. (70) becomes equal to that of Eq. (12). Thus all results of Sec. 2.2, in particular Eqs. (22) and (24), carry formally over. The only but crucial difference lies in the poles of the generalized susceptibility $\chi(\omega)$. The set of zeros can be written as

$$\omega_{\pm} = \omega_0 \left(-i\kappa/2 \pm \sqrt{1-\kappa^2} \right) , \qquad (72)$$

with κ defined as

$$\kappa = \frac{\eta}{2}\gamma_p + \frac{1}{2\eta}\gamma_q \ . \tag{73}$$

According to criterion A, Eq. (16), the regions of overdamped and underdamped oscillations in the (γ_q, γ_p) plane are separated by a straight line with negative slope given as before by the condition $\kappa = 1$ [see Fig. 4]. Therefore, a crossover from overdamped to underdamped oscillations by increasing one of the coupling constants is now impossible. We also note that, in the symmetric case $(\gamma_q = \gamma_p = \gamma)$, the relaxation time is the same as for the q-oscillator [$\tau^{-1} = \omega_0 \gamma/2$]. Increasing γ_q or γ_p leads inevitably to an enhancement of dissipation. This is in blatant *contrast* to the results obtained previously for the double-bath model and one of the most remarkable results of this work. It best illustrates the importance of the specific structure of the bath and justifies a posteriori the detailed study of a model with two independent baths.

5.2. Mixing angle $\theta \neq 0, \pi$

We briefly consider the case of non-vanishing mixing angle. The effects should be largest for $\theta = \pi/2$, corresponding to an interaction Hamiltonian

$$H_I \propto \sum (|\mu_k| p_k p + |\lambda_k| q_k q) . \tag{74}$$

This form of coupling is called amplitude coupling in [43]. Now $J_{-}(\omega)$ is not longer zero but we have

$$\omega \widehat{J}_{-}(\omega) = \sqrt{\gamma_q \gamma_p} \omega^2 \left[\frac{2}{\pi} \ln \left(1 + \frac{\Omega^2}{\omega^2} \right) + i \operatorname{sgn}(\omega) \right]$$
(75)

which is exactly the spectral function of a superohmic bath with exponent $\alpha = 2$. The logarithm in Re \hat{J}_{-} inhibits further analytic treatment of Eq. (70) even in the high temperature limit; thus we limit ourselves to a qualitative analysis. For simplicity, we omit the terms containing logarithms in the generalized succeptibility Eq. (62). The contributions to $C_{qq}^{(+)}$ stemming from those terms should decay on a time scale $\sim \Omega^{-1}$. The remaining terms define a succeptibility which is identical to the succeptibility of a q-oscillator coupled linearly to one bath with spectral density

$$J(\omega) = (\gamma_q + \eta^2 \gamma_p)\omega + \frac{2\sqrt{\gamma_q \gamma_p}}{\omega_p} \omega^2 .$$
(76)

It is well known [19] that, in the case of a polynomial spectral function, the term with the lowest exponent dominates [see also the discussion in Sec. 2.3]. Therefore we may conclude that the crossover diagram Figure 4 remains unchanged by a mixing angle $\theta \neq 0$.

6. Conclusions

We have discussed the behavior of a quantum Brownian particle in a harmonic potential subject to two independent noise sources, one of which couples to its position and the other one to its momentum. In the symmetric case where both baths are Ohmic and their coupling strength is the same, we find underdamped oscillations of the central oscillator for all coupling strengths. This indicates that the two baths partially cancel each other. The effect is due to the mutually conjugate character of position and momentum. It was first noted in Ref. [26] for the (cylindrically) symmetric spin-boson model with spin $s = \frac{1}{2}$, i.e. in the deep quantum regime. "Quantum frustration" of the spin can pictorially be described as the result of having two observers attempting to measure simultaneously, with equal efficiency, two non-commuting components of the spin. Because of the uncertainty principle, both of them fail to measure anything. Here we have investigated the analogous effect for a quantum oscillator coupled to two independent baths. We have found a moderate form of cancellation which we have labelled "quasiclassical frustration" because our dissipative oscillator may describe a large spin impurity coupled to the magnon bath in a ferromagnetic medium. The most notable features of quasiclassical frustration become manifest in the strong coupling limit, where underdamped dynamics survives and all time scales diverge. It remains to be investigated whether the occurrence of frustration in the classical regime is a general property or an artifact of the harmonic oscillator.

We have compared the double-bath model with the case where a single bath couples linearly to the position and momentum of the oscillator. In the latter case the "phase–space" diagram is simpler in that transitions from overdamped to underdamped oscillations can never occur. Comparison of the two models indicates that bath correlations can qualitatively change the behaviour of a dissipative system.

A point of caution is needed in the interpretation of our results. We have definitely ruled out the at first sight enticing but at closer inspection unphysical conjecture that the effects of the two observers cancel each other completely and the particle is not affected by the environment. Indeed we have seen in the specific example of a decoupled initial state that, in destroying quantum coherence, two baths are always more efficient than one. This is true for both the overdamped and the underdamped regime. This demonstrates that, at least for the harmonic oscillator, underdamped dynamics is by no means a reliable signature of high global coherence, which here we identify with purity [see Sec. 4]. Our work provides further evidence that decoherence and dissipation are not necessarily correlated. By dissipation we mean here the net transfer of energy from the central oscillator to the thermal baths. It is characterized by the classical equation of motion or, equivalently, by the properties of the spectral function. We have seen that, depending on the situation, an increase of dissipation can be accompanied by either a reduction or an increase of decoherence. Vice versa, a source of decoherence may or may not lead to dissipation. This is the case in the so called pure dephasing (not considered here), where the interaction part of the Hamiltonian commutes with the system Hamiltonian and no dissipation occurs at all, see i. e. [43].

The fact that we have focused on a quantum oscillator coupled linearly to oscillator baths has allowed us to investigate analytically the equilibrium and dynamical properties. The question of frustration is also raised in other dissipative quantum systems, such as a small spin coupled to a boson [26] or a fermion [44] bath, which are not amenable to an exact analytical treatment. The extension of the present study to less tractable physical scenarios provides a theoretical challenge.

Acknowledgments

We are indebted to F. Guinea for useful discussions. This work has been supported by Ministerio de Ciencia y Tecnología (Spain) under Grants No. BFM2001-0172 and FIS2004-05120, and by the Ramón Areces Foundation. One of us (H.K.) acknowledges financial support from the RTN Network of the European Union under Grant No. HPRN–CT–2000-00144.

Appendix A. Calculation of Eq. (32)

We calculate the high frequency contribution C_{Ω} to the position mean square in Eq. (31). This contribution vanishes for $\alpha_q + \alpha_p < 2$ but yields a finite contribution otherwise. We assume again $\alpha_q = \beta_q/m$ and $\alpha_p = \beta_p/m$, $(\beta_q, \beta_p, m \in \mathbb{Z})$ to be rational numbers. Then this contribution can be written in a determinant form

$$C_{\Omega} = \frac{1}{\gamma_p \gamma_q^2} \frac{m}{\pi \Delta_{r_{\Omega}}(\lambda^2)} \begin{vmatrix} f(\lambda_1^2) & \lambda_1^{2r-4} & \dots & 1\\ \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots \end{vmatrix}$$
(A.1)

Here we use the Vandermonde determinant $\Delta_N(\lambda^2) = \prod_{i < j}^N (\lambda_i^2 - \lambda_j^2)$. We recall that r is the total number of poles, while $r_{\Omega} < r$ is the number of high energy poles. The

Dissipative quantum oscillator with two competing heat baths

function f(x) is given by

$$f(x) = \Phi(x, \beta_p + 3m - 1, r) + \gamma_q \gamma_p \frac{4\Theta(\alpha_p - 2)}{\pi^2(\alpha_p - 2)^2} \Phi(x, 5m + \beta_q - 1, r) + \gamma_q \gamma_p \Phi(x, 2\beta_p + \beta_q + m - 1, r) , \qquad (A.2)$$

where the function $\Phi(x, s, r)$ is a rather complicated function involving a special function called Lerch transcendent [42]. It can, for s odd, be expressed in terms of a logarithm plus a polynomial

$$\Phi(x,s,r) = \frac{x^{s-1}}{2} \left[\ln\left(1 + \frac{1}{x^2}\right) - \sum_{n=1}^{(s+1)/2-r} \frac{(-1)^n}{nx^{2n}} \right] .$$
(A.3)

In order to use this formula we assume m and β_n odd. Consequently we formally exclude α_n even. However, that case can be included by considering the limit $m \to \infty$ with $\beta_n = 2nm-1$. The argument of f(x) is the square of one of the r roots of the characteristic polynomial

$$\left[\frac{2\Theta(\alpha_q-2)\lambda^m}{\pi(\alpha_q-2)} - i\lambda^{\beta_q-m}\right] \left[\frac{2\Theta(\alpha_p-2)\lambda^m}{\pi(\alpha_p-2)} - i\lambda^{\beta_p-m}\right] - \frac{1}{\gamma_q\gamma_p} = 0.$$
(A.4)

The asymptotic behavior of C_{Ω} for $\gamma_q, \gamma_p \ll 1$ can be derived by expanding Φ in Eq. (A.3). For $\alpha_p > 1$ we find to lowest order

$$\mathcal{C}_{\Omega} = \frac{\gamma_p}{\pi} \frac{1}{\alpha_p - 1} + \gamma_q \gamma_p \frac{4\Theta(\alpha_p - 2)}{\pi^2(\alpha_p - 2)^2} \frac{1}{\alpha_q + 1} + \frac{\gamma_q \gamma_p}{2\alpha_p + \alpha_q - 3} + \mathcal{O}(\gamma_q^2, \gamma_p^2) .$$
(A.5)

Keeping only the first term for small γ_q and small γ_p yields Eq. (32).

Appendix B. Derivation of Eq. (53)

We regularize the Ohmic spectral functions by a Drude cutoff

$$J_n(\omega) = \frac{2\gamma_n}{\pi} \frac{\Omega_n^2 \omega}{\Omega_n^2 + \omega^2} , \quad \widetilde{J}_n(\omega) = \frac{\gamma_n \omega \Omega_n}{\omega - i\Omega_n} , \quad n = q, \ p .$$
(B.1)

Assume $\Omega_q = \Omega_p = \Omega$, $\gamma_q = \gamma_p = \gamma$. For ϕ_0 and ϕ_1 we obtain

$$\phi_0(t) = \frac{1}{1+\gamma^2} \left[\cos\left(\Lambda t\right) - \gamma \sin\left(\Lambda t\right) \right] e^{-t/\tau} + \frac{\gamma^2}{1+\gamma^2} \left[\cos(\gamma^2 \Omega t) + \frac{1}{\gamma} \sin(\gamma^2 \Omega t) \right] e^{-\Omega t} \phi_1(t) = \frac{1}{1+\gamma^2} \left[\gamma \cos\left(\Lambda t\right) + \sin\left(\Lambda t\right) \right] e^{-t/\tau} + \frac{\gamma^2}{1+\gamma^2} \left[-\cos(\gamma^2 \Omega t) + \gamma \sin(\gamma^2 \Omega t) \right] e^{-\Omega t} , \qquad (B.2)$$

where no ambiguity is left $[\Lambda = \omega_0 (1+\gamma^2)^{-1}$ and, we recall, $\tau^{-1} = \gamma \Lambda]$. In a time of order Ω^{-1} after the connection (with $\Omega \to \infty$), both functions have dropped on average to a

value $(1 + \gamma^2)^{-1}$. This is the value one would have obtained by using directly $J(\omega) \propto \omega$. After this short initial period, ϕ_0 and ϕ_1 decay much more slowly, on a time scale τ . In the expression (52) for the purity decay after the initial slip, we may safely neglect the first two terms in the denominator. The functions ϕ_0 , ϕ_1 however still govern through X_0 , X_1 the time evolution of the purity $\mathcal{P}(t)$ [see Eq. (40)]. We notice that both X_0 and X_1 are zero at t = 0. However in the initial time interval of order $1/\Omega$ they both increase rapidly. After the initial slip, they settle to a value $\approx \langle q^2 \rangle_{\beta} \gamma^2 / (1 + \gamma^2)$ and increase much more slowly afterwards. This can be seen if we write for instance $X_0(t)$ as follows

$$X_0(t) = \frac{1}{8} \int d\omega J(\omega) \coth\left(\beta\omega/2\right) |\chi(\omega)|^2 \omega^2 \left|1 - g_0(\omega, t)e^{-i\omega t}\right|^2 , \qquad (B.3)$$

where $g_0(\omega, t)$ is the principal value integral

$$g_0(\omega,t) = \frac{1}{2\pi i} \frac{1}{\chi(\omega)\omega} \mathcal{P} \int \frac{\chi(\omega')\omega'}{\omega'-\omega} e^{i\omega't} d\omega' .$$
(B.4)

Eq. (B.4) can be evaluated straightforwardly using the residue theorem. However for our purposes the most important point is that $g_0(\omega, 0) = 1$ and that $g_0(\omega, t)$ behaves in time essentially in the same way as $\phi_0(t)$ in Eq. (B.2). Thus we may write

$$g_{0}(\omega,t) = \frac{1}{\omega(1+\gamma^{2})} \left[\omega \cos\left(\Lambda t\right) - i\left(\omega_{0} + \widetilde{J}(\omega)\right) \sin\left(\Lambda t\right) \right] e^{-t/\tau} + \frac{\gamma^{2}}{1+\gamma^{2}} \left[A_{\Omega}(\omega) \cos(\gamma^{2}\Omega t) + B_{\Omega}(\omega) \sin(\gamma^{2}\Omega t) \right] e^{-\Omega t} .$$
(B.5)

Here A_{Ω} , B_{Ω} are some algebraic functions of ω whose exact form is not important for the present discussion, since after the initial slip of time Ω^{-1} the second line of Eq. (B.5) can be neglected. Proceeding in the same way with $X_1(t)$ we arrive at

$$2X_0(t) + 2X_1(t) = \langle q^2 \rangle_\beta \left(1 + \frac{e^{-2t/\tau}}{(1+\gamma^2)^2} \right) + \frac{2C_{qq}^{(+)}(t)}{1+\gamma^2} \cos\left(\Lambda t\right) e^{-t/\tau} , \qquad (B.6)$$

where $C_{qq}^{(+)}(t)$ is given in Eq. (12). The temperature dependence is encoded in $\langle q^2 \rangle_{\beta}$ and $C_{qq}^{(+)}(t)$. Of course the r.h.s. of Eq. (B.6) is not zero any more at t = 0 due to the neglected initial transient. At zero temperature the autocorrelation function decays algebraically $\propto -\omega_0^2 t^{-2}$. Then we reproduce Eq. (53).

Appendix C. Purity decay of the *q*-oscillator and general formulae

Here we state the general formulae for the purity decay of a coherent (Gaussian) initial state. To calculate the purity decay for two Ohmic baths, both with decoupled initial conditions but with different coupling strengths, we proceed as in Sec. 4.1. Performing a twofold Fourier transform of Eq. (50) and plugging the result into Eq. (38) yields for the reduced Wigner function

$$W(q, p, t) = \frac{1}{\pi \left(A_q A_p - 4A_{qp}^2\right)^{1/2}} \exp\left(\frac{4A_{qp}qp - A_q q^2 - A_p p^2}{A_q A_p - 4A_{qp}^2}\right) .$$
(C.1)

Here,

$$A_{q} \equiv \phi_{0}^{2}\eta + \frac{\phi_{q}^{2}}{\eta} + 4X_{pq} + 4X_{q0}$$

$$A_{p} \equiv \frac{\phi_{0}^{2}}{\eta} + \phi_{p}^{2}\eta + 4X_{qp} + 4X_{p0}$$

$$A_{qp} \equiv \frac{\eta\phi_{0}\phi_{p}}{2} - \frac{\phi_{0}\phi_{q}}{2\eta} - 2Y_{pq} + 2Y_{qp} .$$
(C.2)

For the purity we have with the above definitions

$$\mathcal{P}(t) = \left(A_q A_p - 4A_{qp}^2\right)^{-1/2}$$
 (C.3)

As expected, this expression reduces to Eq. (52) for $J_q = J_p$ and $\eta = 1$. For $J_p = 0$ we have the case of the *q*-oscillator. From the definitions (40) we get $X_{pq} = X_{p0} = Y_{pq} = 0$ and obtain a closed expression for the purity decay of the *q*-oscillator,

$$\mathcal{P}(t) = \left[\left(\phi_0^2 \eta + \frac{\phi_q^2}{\eta} + 4X_{q0} \right) \left(\frac{\phi_0^2}{\eta} + \phi_p^2 \eta + 4X_{qp} \right) - 4 \left(\frac{\eta \phi_0 \phi_p}{2} - \frac{\phi_0 \phi_q}{2\eta} + 2Y_{qp} \right)^2 \right]^{-1/2}$$
(C.4)

The explicit form of the functions $\phi_0(t)$, $\phi_n(t)$ with Drude regularized Ohmic spectral function can be found in Ref. [11]. Here we give only their form after taking the limit $\Omega \to \infty$, i. e. for $t \gg \Omega^{-1}$.

$$\phi_0(t) = \left[\cos(\zeta t) - \frac{1}{\zeta\tau}\sin(\zeta t)\right] e^{-t/\tau}$$

$$\phi_p(t) = \frac{\omega_p}{\zeta}\sin(\zeta t)e^{-t/\tau}$$

$$\phi_q(t) = -\gamma_q - \gamma_q \left[\cos(\zeta t) - \frac{1}{\zeta\tau}\sin(\zeta t)\right]e^{-t/\tau}$$
(C.5)

As in the main text ζ is the renormalized oscillator frequency. With these functions we can calculate $X_{q0}(t)$, $X_{qp}(t)$ and $Y_{qp}(t)$. We obtain,

$$X_{q0}(t) = \frac{1}{2} \langle p^2 \rangle_\beta \left[1 + \phi_0^2(t) \right] - \phi_0(t) C_{qq}^{(+)}(t) + \frac{\dot{\phi}_0^2(t)}{2\omega_p^2} \langle q^2 \rangle_\beta + \frac{\dot{\phi}_0^2}{\omega_p^2} \dot{C}_{qq}^{(+)}(t)$$

$$X_{qp}(t) = \frac{1}{2} \langle q^2 \rangle_\beta \left[1 + \phi_0^2(t) \right] - \phi_0(t) C_{qq}(t) + \frac{1}{2} \phi_p^2(t) \langle p^2 \rangle_\beta - \frac{\phi_p(t)}{\omega_p} \dot{C}_{qq}^{(+)}(t)$$

$$Y_{qp}(t) = \frac{\dot{\phi}_0(t)}{2\omega_p} \left[\phi_0(t) \langle q^2 \rangle_\beta - C_{qq}^{(+)}(t) \right] + \frac{\phi_p}{2} \left[\langle p^2 \rangle_\beta \phi_0(t) - C_{pp}^{(+)}(t) \right] \quad (C.6)$$

We notice that

$$Y_{qp}(0) = X_{q0}(0) = X_{qp}(0) = \phi_p(0) = 0$$
. (C.7)

Therefore we find from Eq. (C.4), $\mathcal{P}(0) = 1$, i.e., the purity is *unaffected* by the initial slip.

References

- [1] V. B. Magalinskii, Soviet. Phys. JETP 9, (1959) 1381.
- [2] P. Ullersma, Physica (Utrecht) 32 (1966) 27; *ibid.* 56; *ibid.* 74; *ibid.* 90.
- [3] B. R. Mollow, Phys. Rev. A 2 (1970) 1477.
- [4] H. Dekker, Phys. Rev. A 16 (1977) 2126.
- [5] P. Talkner, Z. Phys. B 41 (1981) 365.
- [6] A. Schmidt, J. Low Temp. Phys. 49 (1982) 609.
- [7] H. Grabert, P. Talkner, Phys. Rev. Lett. 50 (1983) 1335.
- [8] H. Grabert, U. Weiss, P. Talkner, Z. Phys. B 55 (1984) 87.
- [9] K. Lindeberg, B. J. West, Phys. Rev. A 30 (1984) 568.
- [10] P. S. Riseborough, P. Hänggi, U. Weiss, Phys. Rev. A 31 (1985) 471.
- [11] F. Haake, R. Reibold, Phys. Rev. A 32 (1985) 2462.
- [12] H. Grabert, P. Schramm, G. L. Ingold, Phys. Rep. 168 (1988) 115.
- [13] W. G. Unruh, W. Zurek, Phys. Rev. D 40 (1989) 1071.
- [14] E. G. Harris, Phys. Rev. A 42 (1990) 3685.
- [15] A. O. Caldeira, A. J. Leggett, Ann. Phys. (New York) 149 (1983) 374.
- [16] B. L. Hu, J. P. Paz, Y. Zhang, Phys. Rev. D 45 (1992) 2843.
- [17] W. H. Zurek, S. Habib, J. P. Paz, Phys. Rev. Lett. 70 (1993) 1187.
- [18] L. D. Romero, J. P. Paz, Phys. Rev. A 55 (1997) 4070.
- [19] U. Weiss, Quantum Dissipative Systems, 2nd Edition, World Scientific, Singapore, 1999.
- [20] U. Eckern, V. Ambegaokar, G. Schön, Phys. Rev B 30 (1984) 6419.
- [21] H. Kohler, F. Guinea, F. Sols, Ann. Phys. 310 (2004) 127.
- [22] F. Sols, I. Zapata, New Developments on Fundamental Problems in Quantum Dynamics, Kluwer Academic, Dordrecht, 1997.
- [23] J. D. Jackson, *Classical Electrodynamics*, 2nd Edition, Wiley, New York, 1977.
- [24] A. J. Leggett, Phys. Rev. B 30 (1984) 1208.
- [25] A. Cuccoli, A. Fubini, V. Tognetti, R. Vaia, Phys. Rev. E 64 (2001) 066124.
- [26] A. H. Castro Neto, E. Novais, L. Borda, G. Zarand, I. Affleck, Phys. Rev Lett. 91 (2003) 096401;
 E. Novais *et. al.*, Phys. Rev. B 72, (2005) 014417.
- [27] H. Kohler, F. Sols, Phys. Rev. B 72 (2005) 180404.
- [28] G. W. Ford, J. T. Lewis, R. F. Connell, Phys. Rev. A 37 (1988) 4419.
- [29] C. Kittel, Introduction to Solid State Physics, Wiley, New York, 1976.
- [30] C. Kittel, Quantum Theory of Solids, Wiley, New York, 1963.
- [31] E. M. Chudnovsky and J. Tejada, Macroscopic Quantum Tunneling of the Magnetic Moment, Cambridge U. Press, Cambridge, 1998.
- [32] M. Abramowitz, I. A. Stegun, Handbook of Mathematical Functions, 9th Edition, Dover, New York, 1972.
- [33] C. W. Gardiner, P. Zoller, *Quantum Noise*, 2nd Edition, Berlin, Springer, 2000.
- [34] R. Karrlein and H. Grabert, Phys. Rev. E 55, (1997) 153.
- [35] S. Kohler and F. Sols, Phys. Rev. A 63, (2001) 053605.
- [36] A. J. Leggett *et al.*, Rev. Mod. Phys. 59 (1987) 1.
- [37] W. Bez, Z. Physik B 39 (1980) 319.
- [38] P. Hänggi, Stochastic Dynamics Lecture Notes in Physics Vol. 484, Springer, Berlin, 1997.
- [39] J. Sánchez-Cañizares, F. Sols, Physica A 212 (1994) 181.
- [40] A. Isar, A. Sandulescu and W. Scheid, Phys. Rev. E 60, (1999) 6371
- [41] A. O. Caldeira, A. J. Leggett, Phys. Rev. A 31 (1984) 1059.
- [42] I. S. Gradshteyn, I. M. Ryzhik, Table of Integrals, Series and Products, Academic Press, New York, 2000.
- [43] T. Gorin, T. Prosen, T. H. Seligman, and W. T. Strunz Phys. Rev. A 70, 042105 (2004)
- [44] F. Sols, F. Guinea, Phys. Rev. B 36 (1987) 7775.