Integrable $1/r^2$ Spin Chain with Reflecting End

Takashi Yamamoto^{*}

Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606, Japan

Osamu Tsuchiya[†]

Department of Pure and Applied Sciences, University of Tokyo, Komaba, Meguro-ku, Tokyo 153, Japan

Abstract

A new integrable spin chain of the Haldane-Shastry type is introduced. It is interpreted as the inverse-square interacting spin chain with a *reflecting end*. The lattice points of this model consist of the square roots of the zeros of the Laguerre polynomial. Using the "exchange operator formalism", the integrals of motion for the model are explicitly constructed.

Typeset using REVT_{EX}

^{*}*E-mail address* : yam@yukawa.kyoto-u.ac.jp

[†]*E-mail address* : otutiya@hep1.c.u-tokyo.ac.jp

Studies of the Calogero-Sutherland model [1], the Haldane-Shastry spin chain [2] and their variants [3] have provided many new links with other areas of physics and mathematics. In particular, these models provide exactly solvable models in which the ideas of the fractional exclusion statistics can be tested [4,5].

In ref. [6], with a view to proving the quantum integrability of the Calogero-Sutherland model and, its rational version, the Calogero-Moser model confined in a harmonic potential (we call the Calogero model), Polychronakos had proposed the so-called *exchange operator formalism*. His clever formalism could be applicable not only to the continuum models but to the lattice models and has become a standard technique to study the integrability and the spectrum of the inverse-square interacting systems [7–12]. Within the exchange operator formalism, all of the inverse-square interacting lattice models can be related to the appropriate continuum inverse-square interacting models with the *internal degrees of freedom (spin)*. More precisely, the lattice models are obtained by freezing out the kinematic degrees of freedom in the corresponding continuum models, and lattice sites lie at the classical static-equilibrium positions of the continuum models [13–15]. For example [8,16], the Haldane-Shastry model is related to the spin Calogero-Sutherland model [17,7,18] whose classical equilibrium positions form a regular lattice on the circle.

Polychronakos [16] has applied his formalism to constructing the new lattice model related to the spin Calogero model [7,10,11,19,20]. We call this model the Polychronakos-Frahm (PF) model [21,22]. The lattice sites of the PF model are positioned at the zeros of the Hermite polynomial, *i.e.*, the spins are no more equidistant. Against this unusual property, the spectra of the PF model are equally spaced and therefore simpler than those of the Haldane-Shastry model. Thus the fractional exclusion statistics for the elementary excitations of the PF model is more tractable than one of the Haldane-Shastry model [22].

On the other hands, in ref. [23,24], an another generalization of the spin chain model, the Haldane-Shastry model with open boundary conditions (BC_N -type Haldane-Shastry model), has been introduced. This model is related to the BC_N -type spin Calogero-Sutherland model [25,26]. It is now well known that such BC_N -type models can be applicable to analyzing the physics with boundaries [27–30]. In particular, one of the authors and his collaborators have shown that the above models possess the properties of the chiral Tomonage-Luttinger liquids [30].

The aim of this letter is twofold. The first is to prove the integrability of the B_N type spin Calogero model [31] within the exchange operator formalism. The second is to construct the new integrable lattice model related to the B_N -type spin Calogero model. This lattice model is thought of as the "intersection" of the PF model and the BC_N -type Haldane-Shastry model.

Before turning to the explicit calculation, we shall briefly mention this new integrable spin chain. The Hamiltonian is given by,

$$\mathcal{H}_{PF} = \sum_{1 \le j \ne k \le N} \left[\frac{1}{(x_j - x_k)^2} P_{jk} + \frac{1}{(x_j + x_k)^2} \overline{P}_{jk} \right] + \gamma \sum_{j=1}^N \frac{1}{x_j^2} P_j, \tag{1}$$

where N is the number of sites and $\gamma \in \mathbf{R}$ is a parameter. In the above Hamiltonian we have introduced the B_N -type spin exchange operators for the ν -component spin variables [24,31]; the operator P_{jk} exchanges the spins at the sites j and k, the operator P_j is defined by the condition $P_j^2 = 1$ and thus is regarded as a reflection operator of the spin at the site j, and finally the operator \overline{P}_{jk} is defined by $\overline{P}_{jk} = P_j P_k P_{jk}$. Also it will be shown that, from the integrability condition of the model, lattice points x_j 's lie at the square roots of the zeros of the Laguerre polynomial $L_N^{(|\gamma|-1)}(y)$ (see, for the notation, ref. [32]). It is well known that the Laguerre polynomial $L_N^{(\alpha)}(y)$ with $\alpha > -1$ (resp. = -1) has N distinct roots, $0 < y_1 < y_2 < \cdots < y_N$ (resp. $0 = y_1 < y_2 < \cdots < y_N$) [32]. Therefore the lattice of the model is well defined and does not contain negative sites. For example, in the case $N = 4, \gamma = 2$, the model has the lattice (0.86, 1.60, 2.39, 3.31).

There are several points which should be noticed in (1). Clearly, the Hamiltonian (1) is not translationally invariant because the lattice is not uniform. Even if we suppose that the lattice is uniform, the terms $\overline{P}_{jk}/(x_j + x_k)^2$ and P_j/x_j^2 in (1) break the translational invariance. The term $\overline{P}_{jk}/(x_j + x_k)^2$ represents the interaction between the *j*-th. spin and the "mirror-image" of the *k*-th. spin. With an appropriate choice of the representation of the operator P_j , the last term of (1) can be regarded as magnetic fields whose magnitudes are proportional to the inverse-square of the positions of the sites. From these observations, the origin x = 0 can be regarded as a reflecting end of the system. Then we call the model with Hamiltonian (1) the PF model with reflecting end or the B_N -PF model (if $\gamma = 0$, we call the D_N -PF model).

Consider now the integrability of the B_N -type spin Calogero model. We first recall the B_N -type spin Calogero model. The Hamiltonians of the B_N -type spin Calogero-Moser model and the B_N -type spin Calogero model are respectively given by [31],

$$\bar{H}_{CM} = \sum_{j=1}^{N} \left[-\partial_j^2 + \frac{1}{x_j^2} \beta_1 (\beta_1 - M_j) \right] \\ + \sum_{1 \le j \ne k \le N} \left[\frac{1}{(x_j - x_k)^2} \beta(\beta - M_{jk}) + \frac{1}{(x_j + x_k)^2} \beta(\beta - \overline{M}_{jk}) \right],$$
(2)

$$\bar{H}_C = \bar{H}_{CM} + \omega^2 \sum_{j=1}^N x_j^2,$$
(3)

where $\beta, \beta_1 \in \mathbf{R}$ and $\omega \in \mathbf{R}_{\geq 0}$ are coupling constants, and $\partial_j = \frac{\partial}{\partial x_j}$. In (2), we have already introduced the operators M_j, M_{jk} and $\overline{M}_{jk} (= M_j M_k M_{jk})$ which are called the B_N -type (coordinate) exchange operators, and are defined by the action on the coordinates x_j ;

$$M_{jk}x_j = x_k M_{jk}, \ M_j x_j = -x_j M_j.$$
 (4)

It is easy to see that these operators satisfy the following relations;

$$M_j{}^2 = M_{jk}{}^2 = \overline{M}_{jk}{}^2 = 1, (5)$$

$$M_{jk} = M_{kj}, \ \overline{M}_{jk} = \overline{M}_{kj}, \tag{6}$$

$$M_j M_k = M_k M_j, \tag{7}$$

$$M_{jk}M_j = M_k M_{jk}, \ \overline{M}_{jk}M_j = M_k \overline{M}_{jk} = M_{jk}M_k,$$
(8)

$$M_{jk}M_{kl} = M_{kl}M_{jl} = M_{jl}M_{jk}, (9)$$

$$M_{jk}\overline{M}_{kl} = \overline{M}_{jl}M_{jk} = \overline{M}_{kl}\overline{M}_{jl}.$$
(10)

Remark that the B_N -type spin exchange operators P_j , P_{jk} and \overline{P}_{jk} also satisfy the above relations [31].

The Hamiltonians (2) and (3) does not contain the terms related directly to the spin. The spin degrees of freedom are introduced as follows. Let $\Omega^s = C^{\infty}(\mathbb{C}^N) \otimes V$ where V denotes the space of spins, for example, $(\mathbb{C}^{\nu})^{\otimes N}$. Then operators M_{jk}, M_j, P_{jk} and P_j naturally act on this space, and clearly M_{jk} and M_j commute with P_{jk} and P_j . Next we introduce a projection π which respectively replaces every occurrence of M_{jk} and M_j by P_{jk} and P_j after M_{jk} and M_j have been moved to the right of the expression. Consider the B_N -type "bosonic" subspace

$$\widetilde{\mathbf{\Omega}}^{s} = \{ f \in \mathbf{\Omega}^{s} \mid (M_{jk} - P_{jk}) f = 0, \ (M_{j} - P_{j}) f = 0 \}.$$
(11)

For any operator $\overline{\mathcal{O}}$, the projection π leads to a unique operator \mathcal{O} which satisfies $\overline{\mathcal{O}}\widetilde{\Omega}^s = \mathcal{O}\widetilde{\Omega}^s$ and does not contain the coordinate exchange operators. The Hamiltonians with the spin degrees of freedom are thus given by the operators $\pi(\overline{H}_{CM})$ and $\pi(\overline{H}_C)$. Also, the spinless, *i.e.*, the one-component case can be considered by putting $P_{jk} = 1, P_j = 1$. In this case, the conditions in (11) are nothing but the conditions for the B_N -invariance of the wavefunctions.

First of all, we introduce the operators \mathcal{D}_j for later use;

$$\mathcal{D}_j = \sum_{k \neq j} \left[\frac{1}{x_j - x_k} M_{jk} + \frac{1}{x_j + x_k} \overline{M}_{jk} \right] + \frac{\beta_1}{\beta} \frac{1}{x_j} M_j.$$
(12)

It is easy to show that

$$M_j \mathcal{D}_j = -\mathcal{D}_j M_j, \ M_{jk} \mathcal{D}_j = \mathcal{D}_k M_{jk}, \tag{13}$$

$$[\mathcal{D}_j, \mathcal{D}_k] = 0, \tag{14}$$

$$\left[\mathcal{D}_{j}, x_{k}\right] = \delta_{jk} \left(-\sum_{l \neq j} (M_{jl} + \overline{M}_{jl}) - 2\frac{\beta_{1}}{\beta}M_{j}\right) + (1 - \delta_{jk})(M_{jk} - \overline{M}_{jk}).$$
(15)

Next, we define the B_N -type Dunkl operators [33,26,34] D_j by

$$D_j = \partial_j - \beta \mathcal{D}_j. \tag{16}$$

Using $[\partial_j, x_j] = \delta_{jk}$, $M_j \partial_j = -\partial_j M_j$, $M_{jk} \partial_j = \partial_k M_{jk}$, etc., we can show that the B_N -type Dunkl operators D_j together with the coordinates x_j satisfy the following relations,

$$M_j D_j = -D_j M_j, \ M_{jk} D_j = D_k M_{jk}, \tag{17}$$

$$[D_j, D_k] = 0, \ [x_j, x_k] = 0, \tag{18}$$

$$[D_j, x_k] = \delta_{jk} \left(1 + \beta \sum_{l \neq j} (M_{jl} + \overline{M}_{jl}) + 2\beta_1 M_j \right) - (1 - \delta_{jk})\beta (M_{jk} - \overline{M}_{jk}).$$
(19)

Finally we introduce another type of the B_N -type Dunkl operators,

$$D_j^{\pm} = D_j \mp \omega x_j \tag{20}$$

which satisfy the similar relations among D_j 's and x_j 's;

$$M_j D_j^{\pm} = -D_j^{\pm} M_j, \ M_{jk} D_j^{\pm} = D_k^{\pm} M_{jk},$$
 (21)

$$[D_i^{\pm}, D_k^{\pm}] = 0, \tag{22}$$

$$[D_j^+, D_k^-] = 2\omega[D_j, x_k].$$
(23)

In fact, if we redefine D_j^{\pm} by $D_j^{\pm}/\sqrt{2\omega}$, then $\{D_j, x_j\}$ and $\{D_j^+, D_j^-\}$ have the same algebraic structure.

Remark that we can lead to the similar results starting with the gauge transformed versions of the D_j and D_j^{\pm} ;

$$\widehat{D}_{j} = \Delta(x)^{-1} D_{j} \Delta(x) = D_{j} + \beta \sum_{k \neq j} \left[\frac{1}{x_{j} - x_{k}} + \frac{1}{x_{j} + x_{k}} \right] + \beta_{1} \frac{1}{x_{j}},$$
(24)

$$\widehat{D}_j^{\pm} = \widetilde{\Delta}(x)^{-1} D_j^{\pm} \widetilde{\Delta}(x) = \widehat{D}_j - (\omega \pm \omega) x_j, \qquad (25)$$

where $\Delta(x) = \prod_{j < k} (x_j^2 - x_k^2)^{\beta} \prod_l x_l^{\beta_1}$ and $\widetilde{\Delta}(x) = \Delta(x) \exp(-\frac{\omega}{2} \sum_j x_j^2)$.

As the ordinary case [7,16], the integrals of motion for the B_N -type (spin) Calogero-Moser model and the B_N -type (spin) Calogero model can be constructed by using the Dunkl operators D_j and D_j^{\pm} , respectively. Moreover, under an appropriate transformation of the coordinates, the integrals of motion for the B_N -type (spin) Calogero-Sutherland model are related to the operators $x_j D_j$. Then we shall unify the construction of these integrals of motion following ref. [35]¹. For this purpose, we introduce the operators,

¹Precisely speaking, this treatment is not convenient to the case of the B_N -type (spin) Calogero-Moser model, because the involutiveness of integrals is clear from its definition.

$$\boldsymbol{\Xi}_j = (pD_j + qx_j)(p'D_j + q'x_j), \qquad (26)$$

where $p, p', q, q' \in \mathbf{C}$. They satisfy the relations

$$M_j \Xi_j = \Xi_j M_j, \ M_{jk} \Xi_j = \Xi_k M_{jk}, \tag{27}$$

$$[\Xi_j, \Xi_k] = (pq' - p'q)\beta(\Xi_j - \Xi_k)(M_{jk} + \overline{M}_{jk}).$$
(28)

From the above formulae we can show the key formula,

$$[\Xi_{j}^{n}, \Xi_{k}^{m}] = (pq' - p'q)\beta \sum_{a=1}^{m} \Xi_{k}^{m-a} (\Xi_{j}^{n} - \Xi_{k}^{n}) \Xi_{j}^{a-1} (M_{jk} + \overline{M}_{jk}).$$
(29)

Let us consider the quantities

$$\Upsilon_n = \sum_{j=1}^N \Xi_j^n. \tag{30}$$

Then the involutiveness of Υ_n 's is clear if pq' - p'q = 0. On the other hand, in general, using the formula (29) and then explicitly antisymmetrizing in the index, we can prove the involutiveness of Υ_n 's as follows;

$$\begin{split} &[\Upsilon_n,\Upsilon_m] \\ &= (pq'-p'q)\beta \sum_{j,k=1}^N \sum_{a=1}^m [\Xi_k^{m-a}(M_{jk} + \overline{M}_{jk})\Xi_k^{n+a-1} - \Xi_k^{n+m-a}(M_{jk} + \overline{M}_{jk})\Xi_k^{a-1}] \\ &= (pq'-p'q)\beta \sum_{j,k=1}^N \left(\sum_{a=1}^m -\sum_{a=n+1}^{n+m}\right) \Xi_k^{m-a}(M_{jk} + \overline{M}_{jk})\Xi_k^{n+a-1} \\ &= \frac{(pq'-p'q)\beta}{2} \sum_{j,k=1}^N \left(\sum_{a=1}^m -\sum_{a=n+1}^{n+m} -\sum_{a=1}^n +\sum_{a=m+1}^{n+m}\right) \Xi_k^{m-a}(M_{jk} + \overline{M}_{jk})\Xi_k^{n+a-1} = 0. \end{split}$$

Moreover, from the B_N -symmetry of Υ_n , *i.e.*, $[M_{jk}, \Upsilon_n] = [M_j, \Upsilon_n] = 0$, the projections $\pi(\Upsilon_n)$ are also involutive.

Specializing the parameters p, p', q and q', we define the two sets of the involutive operators $\{I_n^{CM}\}$ and $\{I_n^C\}$ corresponding to the B_N -type spin Calogero-Moser model and the B_N -type spin Calogero model, respectively;

$$I_n^{CM} = \Upsilon_n \big|_{\substack{p=p'=1\\q=q'=0}} = \sum_{j=1}^N (D_j)^{2n},$$
(31)

$$I_n^C = \Upsilon_n \Big|_{\substack{p=p'=1\\-q=q'=\omega}} = \sum_{j=1}^N (D_j^+ D_j^-)^n.$$
(32)

Note that, in contrast to the ordinary (spin) Calogero-Moser model, the integrals I_n^{CM} depend only on D_j^2 . This fact reflects the absence of the translational invariance in the

Hamiltonian (2). Note also that $I_n^{CS} = \Upsilon_n |_{q=0,p'=1}$ are related to the BC_N -type spin Calogero-Sutherland model.

The Hamiltonian \bar{H}_C (resp. \bar{H}_{CM}) is expressed by the operator I_1^{CM} (resp. I_1^C);

$$\bar{H}_{CM} = -I_1^{CM},\tag{33}$$

$$\bar{H}_C = -I_1^C + \mathcal{E}_N^{(0)}, \tag{34}$$

where $\mathcal{E}_N^{(0)} = \omega [N + 2\beta \sum_{j < k} (M_{jk} + \overline{M}_{jk}) + 2\beta_1 \sum_j M_j]$. It remains to show that I_n^{CM} 's (resp. I_n^C 's) commute with \overline{H}_{CM} (resp. \overline{H}_C). These can be checked by using the formulae;

$$[\bar{H}_{CM}, D_j] = 0,$$
 (35)

$$[\bar{H}_C, D_j^{\pm}] = \pm 2\omega D_j^{\pm}. \tag{36}$$

Hence the B_N -type spin Calogero-Moser model and the B_N -type spin Calogero model are integrable. As mentioned, using the projection π , we can obtain the corresponding integrals of motion which depend on the spin variables.

Let us now turn to the lattice model related to the B_N -type spin Calogero model. We apply the standard technique due to Polychronakos [16] (see also [36,37]). That is, we consider the strong coupling limit $\beta \to \infty$ in the Hamiltonian (3). Since the repulsion between particles and also between particles and mirror-image particles become dominant in the strong coupling limit, particles are enforced to localize with the positions x_j which are taken to minimize the potential,

$$V(x) = \beta^2 \tilde{\omega}^2 \sum_{j=1}^N x_j^2 + \beta^2 \sum_{1 \le j \ne k \le N} \left[\frac{1}{(x_j - x_k)^2} + \frac{1}{(x_j + x_k)^2} \right] + \beta^2 \gamma^2 \sum_{j=1}^N \frac{1}{x_j^2}.$$
 (37)

Here we rescaled the the coupling constant ω of the harmonic potential in order for the system to have a nontrivial limit. Also we rescaled $\beta_1 = \beta \gamma$. Note that $\tilde{\omega}$ can be absorbed into the definition of x_j 's. Then we put $\tilde{\omega} = 1$. From $\partial_j V(x) = 0$, we can obtain that such x_j 's satisfy the condition

$$2\sum_{k\neq j} \left[\frac{1}{(x_j - x_k)^3} + \frac{1}{(x_j + x_k)^3} \right] + \gamma^2 \frac{1}{x_j^3} = x_j.$$
(38)

The above formula is equivalent to the condition that $y_j = x_j^2$ are zeros of the Laguerre polynomial $L_N^{(|\gamma|-1)}(y)$ [15].

In the strong coupling limit $\beta \to \infty$, the elastic modes decouple from the internal degrees of freedom (the latter constitute the desired spin chain model);

$$\bar{H}_C \longrightarrow H_{ela} - \beta \bar{\mathcal{H}}_{PF}.$$
 (39)

Here H_{ela} represents the Hamiltonian for the elastic degrees of freedom and $\bar{\mathcal{H}}_{PF}$ is the Hamiltonian which is obtained by replacing P_{jk} and P_j respectively with M_{jk} and M_j in (1), *i.e.*, $\mathcal{H}_{PF} = \pi(\bar{\mathcal{H}}_{PF})$.

Let us define the operators

$$\mathcal{D}_j^{\pm} = \mathcal{D}_j \pm x_j,\tag{40}$$

$$\Xi_j = \mathcal{D}_j^+ \mathcal{D}_j^- = \mathcal{D}_j^2 - x_j^2 - \sum_{k \neq j} (M_{jk} + \overline{M}_{jk}) - \gamma M_j.$$
(41)

The operators \mathcal{D}_{j}^{\pm} can be thought of as the large- β limit of the operators D_{j}^{\pm} . Thus we expect that the operators $\mathcal{I}_{n}^{PF} = \sum_{j=1}^{N} \Xi_{j}^{n}$ are the integrals of motion for the B_{N} -PF model. We can show the involutiveness of the operators \mathcal{I}_{n}^{PF} along the same argument as those for the B_{N} -type spin Calogero model. The remaining task is to show the commutativity of \mathcal{I}_{n}^{PF} with $\overline{\mathcal{H}}_{PF}$. Clearly, it suffices to show $[\overline{\mathcal{H}}_{PF}, \Xi_{j}] = 0$. This can be proved as follows. We recall the relation (35),

$$[\bar{H}_{CM}, D_j] = 0 \iff [-\sum_l \partial_l^2 - \beta \bar{\mathcal{H}}_{PF} + \beta^2 \mathcal{P}, \partial_j - \beta \mathcal{D}_j] = 0,$$
(42)

where

$$\mathcal{P} = \sum_{1 \le j \ne k \le N} \left[\frac{1}{(x_j - x_k)^2} + \frac{1}{(x_j + x_k)^2} \right] + \gamma^2 \sum_{j=1}^N \frac{1}{x_j^2}.$$
(43)

Let us consider the expansion of the relation (42) in the power of β . Since this relation holds for all β , each term must separately vanish. Thus the term of the order β^2 gives,

$$[\bar{\mathcal{H}}_{PF}, \mathcal{D}_j] = [\partial_j, \mathcal{P}] = -4\sum_{k\neq j} \left[\frac{1}{(x_j - x_k)^3} + \frac{1}{(x_j + x_k)^3} \right] - 2\gamma^2 \frac{1}{x_j^3}.$$
 (44)

Also the direct calculation show that

$$[\bar{\mathcal{H}}_{PF}, x_j] = -2\mathcal{D}_j. \tag{45}$$

Using the above two formulae (44), (45) and the properties $[\bar{\mathcal{H}}_{PF}, M_{jk}] = [\bar{\mathcal{H}}_{PF}, M_j] = 0$, we obtain,

$$[\bar{\mathcal{H}}_{PF}, \Xi_j] = ([\bar{\mathcal{H}}_{PF}, \mathcal{D}_j] + 2x_j)\mathcal{D}_j + \mathcal{D}_j([\bar{\mathcal{H}}_{PF}, \mathcal{D}_j] + 2x_j).$$
(46)

If x_j 's are chosen to take values in the set of square roots of the zeros of the Laguerre polynomial $L_N^{(|\gamma|-1)}(y)$, then we have $[\bar{\mathcal{H}}_{PF}, \mathcal{D}_j] + 2x_j = 0$ (\Leftrightarrow (38)), hence $[\bar{\mathcal{H}}_{PF}, \Xi_j] = 0$.

Therefore we proved the integrability of the B_N -PF model and obtained the integrals of motion $\pi(\mathcal{I}_n^{PF})$ for this model. For example, $\pi(\mathcal{I}_1^{PF})$ is given by,

$$\pi(\mathcal{I}_1^{PF}) = -E_N - \left[\sum_{1 \le j \ne k \le N} (P_{jk} + \overline{P}_{jk}) + 2\gamma \sum_{j=1}^N P_j\right],\tag{47}$$

where

$$E_N = \sum_{j=1}^N x_j^2 + \sum_{1 \le j \ne k \le N} \left[\frac{1}{(x_j - x_k)^2} + \frac{1}{(x_j + x_k)^2} \right] + \gamma^2 \sum_{j=1}^N \frac{1}{x_j^2}.$$
 (48)

Finally, we would like to make some comments on algebraic interpretations of the presented results. Our construction naturally leads to the algebra of integrals of motion. For example, the Virasoro-like structure is given by

$$[J_n, J_m] = 0, (49)$$

$$[L_n, J_m] = -mJ_{n+m}, (50)$$

$$[L_n, L_m] = (n - m)L_{n+m},$$
(51)

where

$$J_n = I_n^{CM}, \text{ (or } I_n^C/(2\omega)^n), \tag{52}$$

$$L_n = \frac{1}{2} \sum_{j=1}^N x_j D_j^{2n+1}, \quad \left(\text{or } \frac{1}{2} \sum_{j=1}^N D_j^- (D_j^+)^{2n+1} / (2\omega)^{n+1} \right).$$
(53)

For the proof, we used the formula,

$$\begin{bmatrix} D_{j}^{n}, x_{k} \end{bmatrix}$$

$$= \delta_{jk} \left[n D_{j}^{n-1} + \beta \sum_{l \neq j} \left(P^{-}(D_{j}, D_{l}) M_{jl} + P^{+}(D_{j}, D_{l}) \overline{M}_{jl} \right) + (1 - (-1)^{n}) \beta_{1} D_{j}^{n-1} M_{j} \right]$$

$$- (1 - \delta_{jk}) \beta \left[(P^{-}(D_{j}, D_{k}) M_{jk} - P^{+}(D_{j}, D_{k}) \overline{M}_{jk} \right],$$
(54)

where the polynomials $P^{\pm}(X, Y)$ are defined by $P^{\pm}(X, Y) = (X^n \pm Y^n)/(X \pm Y)$. Notice that in (53) the total degree of the operator is always even as the polynomial of x_j and D_j (or D_j^- and D_j^+), this fact reflects the B_N -symmetry. We can also construct the algebra of integrals of motion related to the W_{∞} algebra.

Another important futures are relations to the spectrum generating algebras and the Yangian symmetries. One of the authors has shown that the spectrum of the B_N -type spin Calogero model is equally spaced [31]. It is easy to see that the same is true for the B_N -PF model. This is caused by the existence of the spectrum generating algebras (36) and

$$[\bar{\mathcal{H}}_{PF}, \mathcal{D}_j^{\pm}] = \mp 2\mathcal{D}_j^{\pm}.$$
(55)

Moreover the numerical studies show that the B_N -PF model possesses the "supermultiplet" structure. The algebra underlying this structure is Yangian [38,39,22,35]. The Yangian symmetries of the B_N -type spin Calogero model and the B_N -PF model are easily see from the transfer matrices of these systems which can be constructed by the Dunkl operators D_j^{\pm} and \mathcal{D}_j^{\pm} . The details will be appeared in [40].

Acknowledgments

We would like to thank H. Awata, N. Kawakami, Y. Matsuo, S. Odake and S.-K. Yang for discussions. T.Y. was supported by the COE (Center of Excellence) researchers program of the Ministry of Education, Science and Culture, Japan.

REFERENCES

- [1] F. Calogero, J. Math. Phys. 10 (1969) 2197;
 B. Sutherland, J. Math. Phys. 12 (1970) 246, 251, Phys. Rev. A 4 (1971) 2019, A 5 (1992) 1372.
- [2] F. D. M. Haldane, Phys. Rev. Lett. **60** (1988) 635;
 B. S. Shastry, Phys. Rev. Lett. **60** (1988) 639.
- [3] For reviews, see following papers and references therein:
 N. Kawakami, Prog. Theor. Phys. 91 (1994) 189;
 F. D. M. Haldane, in *Correlation Effects in Low Dimensional Electron Systems*, eds.
 A. Okiji and N. Kawakami (Springer, 1994).
- [4] F. D. M. Haldane, Phys. Rev. Lett. 66 (1991) 1529.
- [5] D. Bernard and Y.-S. Wu, in New Developments of Integrable Systems and Long-Ranged Interaction Models, eds. M.-L. Ge and Y.-S. Wu (World Scientific, 1995).
- [6] A. P. Polychronakos, Phys. Rev. Lett. **69** (1992) 703.
- [7] J. A. Minahan and A. P. Polychronakos, Phys. Lett. B **302** (1993) 265.
- [8] M. Fowler and J. A. Minahan, Phys. Rev. Lett. **70** (1993) 2325.
- [9] L. Brink, T. H. Hansson and M. A. Vasiliev, Phys. Lett. B 286 (1992) 109.
- [10] L. Brink and M. A. Vasiliev, Mod. Phys. Lett. A 8 (1993) 3583.
- [11] O. V. Dodlov, S. E. Konstein and M. A. Vasiliev, JETP Lett. 58 (1993) 855.
- [12] H. Ujino and M. Wadati, J. Phys. Soc. Jpn. **64** (1995) 4121.
- [13] F. Calogero, Lett. Nouvo Cimento 19 (1977) 505, 20 (1977) 251, 489, in Nonlinear Phenomena, Lecture Notes in Physics 189 (Springer, 1983).
- [14] A. M. Perelomov, Ann. Inst. Henri Poincaré XXVIII (1978) 407.
- [15] S. Ahmed, M. Brushi, F. Calogero, M. A. Olshanetsky and A. M. Perelomov, IL Nouvo Cimento 49 B (1979) 173.
- [16] A. P. Polychronakos, Phys. Rev. Lett. **70** (1993) 2329, Nucl. Phys. **B419** (1994) 553.
- [17] Z. N. C. Ha and F. D. M. Haldane, Phys. Rev. B 46 (1992) 9359.
- [18] K. Hikami and M. Wadati, J. Phys. Soc. Jpn. 62 (1993) 4203.

- [19] K. Hikami and M. Wadati, J. Phys. Soc. Jpn. 62 (1993) 469, Phys. Lett. A 173 (1993) 263.
- [20] K. Vacek, A. Okiji and N. Kawakami, J. Phys. A 27 (1994) L201, Phys. Rev. B 49 (1994) 4635.
- [21] H. Frahm, J. Phys. A **26** (1993) L437.
- [22] K. Hikami, Nucl. Phys. **B441** (1995) 530.
- [23] B. D. Simons and B. L. Altshuler, Phys. Rev. B 50 (1994) 1102.
- [24] D. Bernard, V. Pasquier and D. Serban, Europhys. Lett. **30** (1995) 301.
- [25] M. A. Olshanetsky and A. M. Perelomov, Phys. Rep. 71 (1981) 313, 94 (1983) 313.
- [26] I. Cherednik, Adv. Math. **106** (1994) 65.
- [27] A. Kapustin and S. Skorik, Phys. Lett. A **196** (1994) 47.
- [28] C. W. J. Beenakker and B. Rejaei, Phys. Rev. B 49 (1994) 7499.
- [29] M. Caselle, Phys. Rev. Lett. **74** (1995) 2776.
- [30] T. Yamamoto, N. Kawakami and S.-K. Yang, J. Phys. A **29** (1996) 317.
- [31] T. Yamamoto, Phys. Lett. A **208** (1995) 293.
- [32] G. Szegö, Orthogonal Polynomials, Am. Math. Soc. Colloquium Publishing 23 (Am. Math. Soc., 1939).
- [33] C. F. Dunkl, Trans. Am. Math. Soc. **311** (1989) 167.
- [34] V. B. Kuznetsov, *Hidden symmetry of the quantum Calogero-Moser system*, preprint (1995) solv-int/9509001.
- [35] D. Bernard, K. Hikami and M. Wadati, in New Developments of Integrable Systems and Long-Ranged Interaction Models, eds. M.-L. Ge and Y.-S. Wu (World Scientific, 1995).
- [36] B. Sutherland and B. S. Shastry, Phys. Rev. Lett. **71** (1993) 5.
- [37] Y. Kato and Y. Kuramoto, Phys. Rev. Lett. **74** (1995) 1222.
- [38] F. D. M. Haldane, Z. N. C. Ha, J. C. Talstra, D. Bernard and V. Pasquier, Phys. Rev. Lett. 69 (1992) 2021.
- [39] D. Bernard, M. Gaudin, F. D. M. Haldane and V. Pasquier, J. Phys. A 26 (1993)

5219.

 $\left[40\right]$ O. Tsuchiya and T. Yamamoto, in preperation.