Smoluchowski's equation for cluster exogenous growth

Stéphane Cueille and Clément Sire

Laboratoire de Physique Quantique (UMR C5626 du CNRS), Université Paul Sabatier

31062 Toulouse Cedex, France.

(March 23, 2018)

We introduce an extended Smoluchowski equation describing coagulation processes for which clusters of mass s grow between collisions with $\dot{s} = As^{\beta}$. A physical example, dropwise condensation is provided, and its collision kernel K is derived. In the general case, the gelation criterion is determined. Exact solutions are found and scaling solutions are investigated. Finally we show how these results apply to nucleation of discs on a plane.

Aggregation processes are of practical interest in many fields of science and technology, including chemistry, material sciences, heat transfer engineering, atmosphere sciences, biology, astronomy, among others [1,2]. In a generic aggregation process, clusters of "mass" s, with mass distribution N(s,t), can encounter collisions with other clusters. Whenever two clusters collide, they can stick to form a new cluster with mass conservation. Although simple numerical models [3–5] have proved to be a very successful tool in the study of these phenomena, much of our theoretical understanding is still to a large extent based on Smoluchowski's equation [6],

$$\partial_t N(s,t) = \frac{1}{2} \int_0^s N(s_1,t) N(s-s_1,t) K(s_1,s-s_1) \, ds_1 - N(s,t) \int_0^{+\infty} N(s_1,t) K(s,s_1) \, ds_1,$$
(1)

where the collision kernel $K(s_1, s_2)$, is the collision rate between clusters of mass s_1 and s_2 , and contains the physics of the aggregation process. Smoluchowski's equation is obtained in mean-field, neglecting density fluctuations, and is valid above a model-dependent upper critical dimension d_c .

Smoluchowski's equation can be used to study the scaling properties of a given aggregation model. Dynamic scaling corresponds to the fact, observed both in experiments and in numerical models [7], that the size distribution is asymptotically scale-invariant at large time, $N(s,t) \sim S(t)^{-\theta} f(s/S(t))$, where the typical size S(t)usually diverges as t^z . Smoluchowski's equation can be used to determine z and θ , and the profile of the scaling function.

Van Dongen and Ernst [8,9] have extensively studied the scaling solutions of Smoluchowski's equation. They have shown that collision kernels could be characterized by two exponents λ and μ ,

$$K(bx, by) = b^{\lambda} K(x, y), \qquad (2)$$

$$K(x,y) \sim x^{\mu} y^{\lambda-\mu} \ (y \gg x), \tag{3}$$

which determine the small x behavior of f(x). If $\mu < 0$, f is bell-shaped, while for $\mu \ge 0$, $f(x) \sim x^{-\tau}$. For $\mu > 0$, $\tau = 1 + \lambda$, while for $\mu = 0$, τ is nontrivial and $\tau < 1 + \lambda$. In the latter case, such a crucial quantity as the exponent describing the decay of the total number of clusters $(n(t) \propto t^{-z'})$ is given by $z' = z(2 - \tau)$ if $\tau > 1$, and thus depends on τ . The determination of τ was very difficult until a very effective variational method was recently introduced [10].

Despite this reasonably satisfactory state of the art, standard Smoluchowski's equation does not describe all aggregation processes, and some extensions have been provided to include for instance fragmentation of clusters [11], or source and sink terms [12–14]. In this Letter, we would like to consider another feature, the fact that for certain aggregation phenomena, clusters do not grow solely due to collisions with other clusters, but also collect some mass from the "outside" between collisions events. A practical example is *dropwise condensation*, a daily life phenomenon (water condensation on a mirror) which has also important consequences in material sciences or heat transfer engineering, which motivated most of the early studies. Since the seminal work of Beysens and Knobler [15], the resulting fascinating droplets patterns, or "breath figures", have attracted much interest, and simple computer models have been introduced to study the kinetics of droplet nucleation [16–19], the asymptotic surface (or line) coverage [20], or the time evolution of the "dry" fraction (the surface fraction which has never been touched by any droplet) [21] (also see the review by Meakin [19]).

In dropwise condensation on a *d*-dimensional substrate, individual droplets form *D*-dimensional hyperspherical caps, and grow by absorption from the vapor phase, which leads to a growth law for the droplets radius, $\dot{r} \propto r^{\omega}$, ($\omega = -2$ for water on a plane [19]), or, in terms of the droplet mass $s = r^D$, $\dot{s} = As^{\beta}$, with $\beta = (D + \omega - 1)/D$. When two droplets overlap, they coalesce due to surface tension, to form a new hyperspherical droplet, with mass conservation. Thus, dropwise condensation can be understood as a simple aggregation phenomenon, where droplets grow between collisions (*exogenous growth*), as well as through collisions. Such a picture is the basis of most simplified computer models, such as the droplets growth and coalescence model of Family and Meakin [17].

Now, what is the corresponding mean-field kinetic equation ? If there were no collisions, the equation would simply be a continuity equation expressing the conservation of the number of droplets,

$$\partial_t N(s,t) + \partial_s(\dot{s}N)(s,t) = 0, \qquad (4)$$

with $\dot{s} = s^{\beta}$ (in the following, we set A = 1). Then, the collision rate of two growing droplets of size s_1 and s_2 is, in mean-field, just proportional to the time derivative of the cross section $\sigma(s_1, s_2) \propto (s_1^{1/D} + s_2^{1/D})^d$, thus $K(s_1, s_2) \propto (\dot{s_1} s_1^{1/D-1} + \dot{s_2} s_2^{1/D-1})(s_1^{1/D} + s_2^{1/D})^{d-1}$, and we obtain the following equation,

$$\partial_t N(s,t) + \partial_s (s^\beta N)(s,t) = \frac{1}{2} \int_0^s N(s_1,t) N(s-s_1,t) K(s_1,s-s_1,t) ds_1 \\ - N(s,t) \int_0^{+\infty} N(s_1,t) K(s,s_1,t) ds_1,$$
(5)

with,

$$K(x,y) = (x^{\frac{\omega}{D}} + y^{\frac{\omega}{D}})(x^{\frac{1}{D}} + y^{\frac{1}{D}})^{d-1}.$$
 (6)

Multiplicative constants were set to 1 by a rescaling of t. The expression for K(x, y) is in agreement with an early work of Vincent [22], who found a Smoluchowski equation for $\omega = -2$, d = 2 and D = 3. However, the left hand side of his equation is erroneous since it does not conserve the number of particles. The kernel has the homogeneity $\lambda = (d+\omega-1)/D$, as predicted in [17] from scaling arguments. Eq. (5) for a generic collision kernel actually describes a general irreversible aggregation process with the exogenous growth law $\dot{s} = s^{\beta}$, and we would like to study the properties of such an equation.

Gelation criterion - A very important issue in aggregation problems is the possible occurrence of gelation. Gelation corresponds to the formation of an infinite cluster at a *finite time* (in the thermodynamic limit). Common applications are found in food industry for instance. Another application in material sciences is the formation of fractal aerogels with intriguing physical properties. In terms of Smoluchowski's equation, gelation is the phase transition associated to the breaking of the mass conservation through collisions. Adapting standard arguments [9], let us consider the net mass flux from clusters with $s \leq L$ towards clusters with s > L, $J_L(t)$. From Eq. (5), it is easily seen that,

$$J_{L}(t) = L^{1+\beta}N(L,t) + \int_{0}^{L} dx \, xN(x,t) \int_{L-x}^{+\infty} dy \, K(x,y)N(y,t), \quad (7)$$

Now, in the absence of an infinite cluster, the mass conservation through collisions requires that $J_L(t) \to 0$ when $L \to \infty$. If gelation occurs at t_g , the infinite cluster collects some mass through exogenous growth and collisions, and $J_L(t > t_g)$ has a finite $L \to \infty$ limit, which implies that N(s,t) has a slowly vanishing large s tail. Inserting the ansatz $N(s,t) \sim A(t)s^{-\tau}$ for $t \ge t_g$, we find $\tau = \max(1 + \beta, (3 + \lambda)/2)$. One must have $\tau > 2$, since the total mass contained in finite clusters must be finite. This shows that gelation can occur only if $\lambda > 1$ or $\beta > 1$.

Exact solutions - For Eq. (1), exact solution are known only for K = 1, K = x + y, K=xy, but the existence of even few exact time-dependent solutions is important to check the scaling theories. Here we shall provide two exact solutions for Eq. (5), corresponding to K = 1, with $\beta = 0$ and $\beta = 1$. To do this, we consider the Fourier-Laplace transform of N(s,t), $Z(z,t) = \int_0^{+\infty} e^{-zs} N(s,t) \, ds$. Z(0,t) is the total density of clusters n(t).

For $\beta = 0$, the Laplace transform of Eq. (5) with K = 1 reads,

$$\partial_t Z + zZ = Z^2/2 - Z(0, t)Z,$$
 (8)

With n(0) = 1 and $Z(z, 0) = Z_0(z)$, we find Z(0, t) = n(t) = 2/(t+2) and,

$$Z(z,t) = \frac{e^{-zt}}{(t+2)^2 \left(\frac{1}{4Z_0(z)} - \frac{1}{2} \int_0^t \frac{e^{zt'}}{(t'+2)^2} dt'\right)}.$$
 (9)

Which leads in the scaling regime $t \to \infty$, and for a monodispersed initial condition $N(s,t) = \delta(s-1)$ to,

$$N(s,t) \sim \frac{2}{t^2 \ln t} e^{-\frac{s}{t \ln t}}.$$
 (10)

The total mass in the system is, $M_1(t) = 1 + 2\ln(t+1/2)$, from $M_1(t) = -\partial_z Z(0,t)$.

For $\beta = 1$, the equation for Z is,

$$\partial_t Z - z \partial_z Z = Z^2 / 2 - Z(0, t) Z.$$
(11)

Once again, n(t) = Z(0,t) = 2/(t+2), and the solution is,

$$Z(z,t) = \frac{2}{t+2} \frac{2Z_0(ze^t)}{(1-Z_0(ze^t))(t+2) + 2Z_0(ze^t)}.$$
 (12)

For a monodispersed initial distribution, $Z_0(z) = e^{-z}$, Z(z,t) has a pole at $z_0(t) = -e^{-t} \ln(1+2/t)$, and we can explicitly compute N(s,t),

$$N(s,t) = \frac{4}{(t+2)^2 e^t} \exp\left(-se^{-t}\ln(1+\frac{2}{t})\right), \quad (13)$$

which leads in the large time limit to $N(s,t) \sim \frac{4}{t^2 e^t} e^{-\frac{2s}{te^t}}$.

These two exact solutions support the fact that Eq. (5) exhibits dynamic scaling, but they also show some unusual time dependence of S(t), and a more general scaling form,

$$N(s,t) \sim Y(t)^{-1} f(s/S(t)),$$
 (14)

where Y(t) does not have the hyperscaling form $Y(t) \propto S(t)^{\theta}$.

Scaling theory - Now we want to study the scaling solutions of Eq. (5), with the scaling form of Eq. (14), starting from a monodispersed initial distribution $\delta(s-1)$. A first remark is that the distribution N(s,t) is at any time strictly zero below $s_0(t) = (1 + (1 - \beta)t)^{1/(1-\beta)}$. In physical terms, this is due to the fact that the smallest clusters are the one which have never collided, and thus have grown since t = 0 with $\dot{s} = s^{\beta}$. Consequently, the scaling function is strictly zero below $x_0 = \lim(s_0(t)/S(t))$. x_0 can be zero if $S(t) \gg t^{1/(1-\beta)}$, but cannot be infinite.

For kernels with $\lambda < 1$ and $\beta < 1$, we can have a qualitative understanding of the scaling results to be expected. Let us examine Eq. (5). On the one hand, if we switch off the collision term (*i.e.* the right hand side of the equation), the equation describes a set of particles which grow in time with $\dot{s} = s^{\beta}$, and is associated with the size scale $S_g(t) \propto t^{\frac{1}{1-\beta}}$. On the other hand, if we forget the growth term in the left hand side, we are back with a standard Smoluchowski equation describing clustering with mass conservation. The typical size in the scaling regime is $S_c(t) \propto t^{1/(1-\lambda)}$, and $\theta = 2$ [8,9].

Thus, if we switch on both growth and collision, we shall observe a "competition" between the dynamic size scales corresponding to both processes. If $\beta < \lambda$, $S_g(t) \gg S_c(t)$, and in the scaling regime we expect $S(t) \propto S_g(t)$ and $z = 1/(1 - \beta)$. If $\beta > \lambda$, on the contrary, the typical size of particles increases essentially due to collisions, hence $S(t) \propto S_c(t)$ and $z = 1/(1 - \lambda)$. For the marginal case $\lambda = \beta$, logarithmic corrections may be observed, and are indeed present for the exact solution K = 1, $\beta = 0$.

A detailed demonstration of the scaling results is quite intricate, since it involves the treatment of many subcases, and will be published elsewhere [23]. However the main idea is quite simple. Let us define the α -th moment, $M_{\alpha}(t) = \int_{0}^{+\infty} s^{\alpha} N(s, t) ds$. With the scaling form of Eq. (14), the different terms of Eq. (5) read,

$$\partial_t N(s,t) \sim -\frac{1}{Y} \left(\frac{\dot{Y}}{Y} f(x) + \frac{\dot{S}}{S} x f'(x) \right),$$
 (15)

$$\partial_s(s^\beta N)(s,t) \sim \frac{S^{\beta-1}}{Y}(x^\beta f)'(x),\tag{16}$$

Collision term
$$\sim \frac{S^{1+\lambda}}{Y^2}(...).$$
 (17)

We look for an asymptotically time independent equation consistent with the evolution equation for the total mass, obtained by multiplying Eq. (5) by s and integrating over all s,

$$\dot{M}_1 = M_\beta. \tag{18}$$

We also use the fact that S(t) cannot be much smaller than $s_0(t)$. Results for Y(t) and S(t), are in agreement with the qualitative picture above. For $\beta > \lambda$, it is found that $S(t) \propto s_0(t)$, thus $z = 1/(1-\beta)$, and $Y(t) \propto S(t)^{\theta}$, with $\theta = 2 + \lambda - \beta$. The three terms in the kinetic equation are of the same order at large time, and, from the remark above, the scaling function is zero below $x_0 = \lim(s_0(t)/S(t)) > 0$. For the droplets nucleation kernel with d < D, we have $\beta > \lambda$ and these results yield $z = D/(1-\omega)$ and $\theta = 1 + d/D$, in agreement with the results for the model of Family and Meakin [17]. This scaling theory also shows that f(x) cannot diverge at small x, as observed for droplets models without nucleation [19], and the fact that the scaling function is even zero below a *finite* $x_0 > 0$ is well supported by numerical simulations [23].

For $\beta \leq \lambda$, $S(t) \gg s_0(t)$, and the exogenous growth term (16) is negligible at large time. The scaling equation is found to be the same as for standard Smoluchowski's equation with the same kernel,

$$xf'(x) + 2f(x) = f(x) \int_0^{+\infty} f(x_1) K(x, x_1) dx_1 - \frac{1}{2} \int_0^x f(x_1) f(x - x_1) K(x_1, x - x_1) dx_1,$$
(19)

and the results of [8,9] can be applied. The nontrivial polydispersity exponents occurring for $\mu = 0$ kernels can be computed using the variational method introduced recently by the present authors [10].

For $\lambda > \beta$, we find $z = 1/(1 - \lambda)$, and $\theta = 2$, the total mass in the system being asymptotically conserved at large time.

For the marginal case $\lambda = \beta$, we find different results for $\mu \leq 0$ kernels and $\mu > 0$ kernels. For $\mu \leq 0$ kernels, corresponding to a bell-shaped scaling functions $(\mu < 0)$, or a nontrivial $\tau < 1 + \lambda$ $(\mu = 0)$, the total mass in the system grows logarithmically in the scaling regime, $M_1(t) \propto \ln t$, we have $S(t) \propto (t \ln t)^z$, with $z = 1/(1 - \beta)$, and $Y(t) \propto S^2(t)/M_1(t)$. For $\mu > 0$ kernels, with $\tau = 1 + \lambda$, there is an additional sublogarithmic correction, leading to $M_1(t) \propto (\ln t) \ln(\ln t)$, $S(t) \propto (tM_1(t))^z$, with z unchanged, and $Y \propto S^2/M_1$ as before.

These results are in full agreement with the exact solution for K = 1 and $\beta = 0$, both for the asymptotics of S, Y, and M_1 , and for the scaling function itself. Indeed, the scaling function is a pure exponential, just as for the exact solution of Eq. (1) with K = 1 [6].

Results for systems on the gelling boundary ($\lambda = 1$ or $\beta = 1$) can also be found and will be reported elsewhere [23]. Here, we would like to show how this Smoluchowski equation approach can be applied to the nucleation of discs on a plane. More generally, we consider the nucleation model described above, but with d = D. Family and Meakin [17] have simulated growth and coalescence of droplets for d = D = 2 and $\omega = 0.5$, and found that in contrast with the d < D case, the scaling function is polydispersed with a small x divergence $f(x) \propto x^{-\tau}$ [19]. Can we understand this feature ?

For this model, the collision kernel is given by Eq. (6), and we have $\beta = \lambda + (D - d)/D$. Thus, d < D corresponds to $\lambda < \beta$, while d = D corresponds to $\lambda = \beta$, and consequently our scaling theory based on a Smoluchowski equation states that the scaling function vanishes below a finite x_0 , for d < D, while f is bell-shaped or polydispersed for d = D, depending on μ . For this kernel, we have $\mu = 0$ for $\omega \ge 0$, and $\mu = \omega/D$ for $\omega < 0$, which leads to the cross-over from a bell-shaped scaling function for $\omega < 0$, to a small x power law for $\omega > 0$. To check this prediction, we performed simulations of the droplets growth and coalescence model for d = D = 2and various values of ω . These simulations are quite difficult since the number of droplets decreases very quickly leading to poor statistics and time range limitation. Results for the small x behavior of the scaling function for different values of ω shown on Fig. 1 are in good agreement with the expected results. For $\omega = -3$, f(x) has a maximum and vanishes faster than any power law as $x \to 0$. When $\omega \to 0^-$, the position of the maximum of the scaling function tends to 0 very quickly, and the scaling function crosses over to a power-law at $\omega = 0$.



FIG. 1. Small x = s/S behavior of the size distributions obtained in numerical simulations for different values of the growth exponent ω .

Therefore, the mean-field scaling theory qualitatively describes the behavior of the scaling function for d = D, and makes for the difference between d < D and d = D. Besides, for $\omega = 0$, the mean-field polydispersity exponent can be computed by the method described in [10]. We find $\tau = 1.109$ [23], which compares well with the exponent extracted from the numerics $\tau \approx 1.2$.

Despite this qualitative success of the Smoluchowski equation approach, it should be noticed that the upper critical dimension of the model with d = D is probably infinite, since we have in mean-field $M_1(t) \sim \ln t$, whereas $M_1(t)$ is bounded for the actual model, being proportional to the surface coverage.

In conclusion, we have generalized Smoluchowski's equation to coagulating systems for which clusters grow between collisions with $\dot{s} = s^{\beta}$. This equation appears to have interesting scaling properties, as found both from exact solutions and a general study. This approach recovers scaling results for droplets nucleation models with d < D, and qualitatively predicts a transition in the shape of the scaling function when varying β , in agreement with numerical simulations, despite mean-field limitations. It would be interesting to see if this kind of approach could also be used to describe dropwise condensation with renucleation in empty spaces [17,19], for which nontrivial polydispersity exponents appear.

We are very grateful to M. Albrecht for a critical reading of the manuscript.

- S. K. Friedlander, Smoke, dust and haze, Wiley Interscience, New York, 1977.
- [2] F. Family and D. Landau, editors, *Kinetics of Aggrega*tion and Gelation, North Holland, Amsterdam, 1984.
- [3] P. Meakin, Phys. Rev. Lett. 51, 1119 (1983).
- [4] M. Kolb, R. Botet, and R. Jullien, Phys. Rev. Lett. 51, 1123 (1983).
- [5] P. Meakin, Physica Scripta 46, 46 (1992).
- [6] M. von Smoluchowski, Z. Phys. Chem 92, 129 (1918).
- [7] T. Vicsek, P. Meakin, and F. Family, Phys. Rev. A 32, 1122 (1985).
- [8] P. G. J. van Dongen and M. H. Ernst, Phys. Rev. Lett. 54, 1396 (1985).
- [9] P. G. J. van Dongen and M. H. Ernst, J. Stat. Phys. 50, 295 (1987).
- [10] S. Cueille and C. Sire, Phys. Rev. E 55, 5465 (1997).
- [11] F. Family, P. Meakin, and J. Deutch, Phys. Rev. Lett. 57, 727 (1986).
- [12] W. H. White, J. Colloid Interface Sci. 87, 204 (1982).
- [13] Z. Rácz, Phys. Rev. A 32, 1129 (1985).
- [14] H. Hayakawa, J. Phys. A 20, L801 (1987).
- [15] D. Beysens and C. Knobler, Phys. Rev. Lett. 57, 1433 (1986).
- [16] F. Family and P. Meakin, Phys. Rev. Lett. 61, 428 (1988).
- [17] F. Family and P. Meakin, Phys. Rev. A 40, 3836 (1989).
- [18] D. Fritter, C. Knobler, and D. Beysens, Phys. Rev. A 43, 2858 (1991).
- [19] P. Meakin, Rep. Prog. Phys. 55, 157 (1992).
- [20] B. Derrida, C. Godrèche, and I. Yekutieli, Phys. Rev. A 44, 6241 (1991).
- [21] M. Marcos-Martin, D. Beysens, J.-P. Bouchaud, C. Godrèche, and I. Yekutieli, Physica A 214, 396 (1995).
- [22] R. Vincent, Proc. Roy. Soc. A **321**, 53 (1971).
- [23] S. Cueille and C. Sire, in preparation, 1997.