

# Hidden Duality and associated instabilities of Tomonaga-Luttinger Liquid on Lattice

Z. N. C. Ha\*

*National High Magnetic Field Laboratory, Florida State University, Tallahassee, FL 32306,  
U.S.A.*

(July 24, 1997)

## Abstract

Hidden duality and associated instabilities of the spinless Luttinger liquid on lattice are reported. The local quantum fluctuations and the long-distance chiral modes compete and as a result produce a hierarchy of exotic charge/density instabilities. Explicit bosonic quantum operators for the local density fluctuations are constructed and are used to make identification of the Luttinger liquid with the classical 2D Coulomb gas with  $\theta$ -term and with the rich hidden duality.

*Keywords:* duality, Luttinger liquid, low dimension, phase transition.

---

\*zha@magnet.fsu.edu (Email); 904-644-5038 (Fax).

I report novel instabilities and hidden duality structure of the spinless Luttinger liquid on lattice with anomalous *local* quantum fractional statistical fluctuations which, I argue, are induced by the possible multi-particle umklapp and pairing processes at rational densities. The local parameter is, in general, different from the usual charge stiffness  $K$  which characterizes the *long-distance* physics. It is shown that the interplay between the *short-* and *long-distance* physics gives rise to new exotic charge/density instabilities which expose the hidden duality of the Luttinger liquid on lattice.

First, I need to consider two conjugate phase fields  $\theta(x)$  and  $\phi(x)$ . The phase  $\theta(x)$  essentially defines a field that measures the density modulations, and it acquires phase  $\pm 2\pi$  going from one to a neighboring particle. The canonically conjugate phase field  $\phi(x)$  is associated with the  $U(1)$  charge degrees of freedom such that  $[\phi(x), \theta(x')] = i\pi \text{sgn}(x - x')$ . Explicit constructions of the two operators are straightforward via Fourier transform. Usually, the renormalization coupling constant  $e^{-2\varphi}$  ( $= 2\pi K$ ) is introduced to code the effects of quantum fluctuation [1].

In the Luttinger liquid universality class it is always possible to find the right and left eigenmodes which carry, in general, fractional statistics. In order to show this more explicitly I use the following right and left Mandelstam modes [2]

$$\Psi_R^\dagger(x) = e^{i\phi(x)} e^{i\beta\theta(x)}; \quad \Psi_L^\dagger(x) = e^{i\phi(x)} e^{-i\beta\theta(x)}. \quad (1)$$

The time-dependent correlation function for large  $x$  and  $t$  is given by

$$\langle \Psi_R^\dagger(x, t) \Psi_R(0, 0) \rangle \propto \frac{1}{(x - v_s t)^{2x_R} (x + v_s t)^{2x_L}}, \quad (2)$$

where  $x_{R,L} = (2\beta \exp(\varphi) \pm \exp(-\varphi))^2/4$ . If  $2\beta = \exp(-2\varphi)$  then the correlation function involves either the right- or the left-movers only. Therefore, the Mandelstam modes with  $2\beta = \exp(-2\varphi)$  can be regarded as the FQS-carrying *long-distance* chiral eigenmodes of the Luttinger liquid.

I conjecture that there are two parameters for describing the Luttinger liquid on lattice. One is the long-distance charge stiffness  $K (= \exp(-2\varphi)/2\pi)$  previously discussed, and the other the local FQHE-like parameter which is presumably determined by the filling fraction and *allowed* local interactions such as the umklapp processes.

Now, consider perturbing the Luttinger liquid with the density fluctuations containing the following

$$\Psi_{\lambda,n}^{\dagger m} = e^{in\theta(x)} e^{im\phi(x)} e^{im\lambda\theta(x)}, \quad (3)$$

where  $m$  is integer content of the  $U(1)$  charge and  $n$  integer index for the low-energy sectors. The parameters  $m$ ,  $n$ , and  $\lambda$  are chosen such that the overall operator be bosonic and the resulting damping term equal to constant (allowed umklapp condition). The scaling dimension,

$$x_{n,m}^\lambda = \frac{1}{4\pi K} (n + \lambda m)^2 + \pi K m^2, \quad (4)$$

is invariant under the following duality  $\hat{D}$  and periodicity  $\hat{T}$  transformations [3]

$$\hat{D} : \eta \rightarrow 1/\eta; \quad (n, m) \rightarrow (-m, n), \quad (5)$$

$$\hat{T} : \eta \rightarrow \eta + i; \quad (n, m) \rightarrow (n - m, m), \quad (6)$$

where  $\hat{D}^2 = 1$  and  $\eta$  is a complex parameter defined as  $2\pi K + i\lambda$ . This duality generalizes the well-known duality for  $\lambda = 0$ .

This work is supported by DOE grant 5024-528-23.

## REFERENCES

1. F. D. M. Haldane, Phys. Rev. **B25**, 4925 (1982).
2. S. Mandelstam, Phys. Rev. **D11**, 3026 (1975).
3. J. L. Cardy, Nucl. Phys. **B205**, 17 (1982).