

Percolation-like phase transition in a non-equilibrium steady state

Indrani Bose and Indranath Chaudhuri
 Department of Physics,
 Bose Institute,
 93/1, Acharya Prafulla Chandra Road,
 Calcutta-700 009, India.

Abstract

We study the Gierer-Meinhardt model of reaction-diffusion on a site-disordered square lattice. Let p be the site occupation probability of the square lattice. For p greater than a critical value p_c , the steady state consists of stripe-like patterns with long-range connectivity. For $p < p_c$, the connectivity is lost. The value of p_c is found to be much greater than that of the site percolation threshold for the square lattice. In the vicinity of p_c , the cluster-related quantities exhibit power-law scaling behaviour. The method of finite-size scaling is used to determine the values of the fractal dimension d_f , the ratio, $\frac{2}{\nu}$, of the average cluster size exponent γ and the correlation length exponent ν and also ν itself. The values appear to indicate that the disordered GM model belongs to the universality class of ordinary percolation.

PACS number(s): 05.70. Ln

I. Introduction

Phase transitions in the non-equilibrium are not as well understood as in the case of equilibrium systems. The percolation model provides an example of phase transition in disordered systems [1, 2]. The model is simply defined. Consider a square lattice of sites. The lattice can be made disordered by assuming that each site is present with probability p and absent with probability $1 - p$. There is a critical value p_c of p , called the percolation threshold, below which long-range connectivity in the system under study is missing and above which the connectivity is present with probability one. The percolation transition is analogous to equilibrium thermodynamic phase transitions and exhibits “critical” phenomena in the vicinity of the

percolation threshold. Percolation-like transitions have been observed in the non-equilibrium in a surface-reaction model [3], an autocatalytic reaction model [4] and chemical wave propagation on a lattice of excitable and non-excitable clusters [5]. Ziff, Gulari and Barshad(ZGB) [3] have studied a kinetic reaction model describing the chemical reaction $CO+O \rightarrow CO_2$ on a catalytic surface. The surface is represented by a square lattice, the sites of which can be singly occupied by a CO molecule or an O atom. let X_{CO} and $1-X_{CO}$ be the molecular concentrations of CO and O_2 molecules in the gas above the surface. CO and O_2 molecules are added randomly to the lattice in the relative ratio of their molecular concentrations. Two adjacent empty sites are necessary for the adsorption of an oxygen molecule. The reaction consists of $CO - O$ pairs being randomly selected from nearest-neighbour (n.n) lattice sites occupied by O atoms and CO molecules. The reaction produces CO_2 molecules which escape from the surface. ZGB found that when X_{CO} is less than a certain value X_1 , the lattice is completely covered by oxygen atoms in the steady state and no further reaction takes place. For X_{CO} greater than X_2 , the lattice is filled with CO . The system exists in a catalytically active mixed-phase only in the range $X_1 < X_{CO} < X_2$. The transitions at $X_1(X_2)$ have been found to be of second (first) order. Later studies [6, 7] have shown that the second-order phase transition belongs to the universality class of Reggeon-field theory (RFT) to which directed percolation (DP) also belongs. Aukrust et al [4] have studied an irreversible kinetic reaction model for a one-component autocatalytic reaction $A + A \rightarrow A_2$. In this model, an atom adsorbing on a lattice site reacts, with probability $1-p$, with one of the n.n. atoms, if present. After the reaction, the two atoms leave the lattice. Otherwise the atom occupies the site. As p is varied, the model undergoes a second-order kinetic phase transition from a partial occupation (chemically-reactive state) of the lattice to a completely covered inert state. Again, the phase transition has been shown to belong to the universality class of Reggeon-field theory-directed percolation. Sendiña-Nadal et al [5] have studied the propagation of chemical waves in a disordered excitable medium in terms of percolation theory. They performed an experiment with Belousov-Zhabotinsky reagent catalyzed by $Ru(bpy)$ complex which is sensitive to visible light thus making it possible to control the excitability of the system experimentally. The system consists of excitable (black) and nonexcitable (white) clusters. The effective wave front velocity has been observed to jump from 0 finite values at a threshold $p = p_c$, where p is the proportion

of black sites. At p_c , a cluster of black sites spans the medium. The data obtained appear to indicate that the percolation process is similar to ordinary percolation. In this paper, we study the Gierer-Meinhardt (GM) model [8] of reaction-diffusion (RD) in two dimensions (2d) and show that a percolation-like transition occurs in the non-equilibrium steady state as the underlying lattice is made disordered. In the vicinity of the percolation threshold p_c , evidence of power-law scaling behaviour of cluster-related quantities is obtained. The percolation threshold appears to be in the universality class of ordinary percolation. The study is based on the finite-size scaling analysis of the percolation problem. In Section II, we define the GM model of RD and describe the various results obtained by us. Section III contains concluding remarks.

II. Reaction-diffusion in the presence of disorder

Turing [9] made a remarkable suggestion, several years back, that diffusion need not always act to smooth out concentration differences in a chemical system. Two interacting chemicals can generate a stable, inhomogeneous pattern if one of the substances (the inhibitor) diffuses much faster than the other (the activator). The activator is autocatalytic, i.e, a small increase in its concentration ‘a’ over its homogeneous steady-state concentration leads to a further increase of ‘a’. The activator besides promoting its own production also promotes the production of the inhibitor. The inhibitor, as the name implies, is antagonistic to the activator and inhibits its production. Suppose, originally the system is in a homogeneous steady state. A local increase in the activator concentration leads to a further increase in the concentration of the activator due to autocatalysis. The concentration of the inhibitor is also increased locally. The inhibitor, having a diffusion coefficient much larger than that of the activator, diffuses faster to the surrounding region and prevents the activator from coming there. This process of autocatalysis and long-rang inhibition finally lead to a stationary state consisting of islands of high activator concentration within a region of high inhibitor concentration. The islands constitute what is known as the Turing pattern. In some Turing systems, there is a possibility of saturation of the autocatalytic process. In

this case, the inhibitor production is limited. Locally activated regions now have activated neighbours. Nonactivated areas are, however, close by into which the inhibitor can diffuse. This results in the emergence of stripe-like patterns which have long-range connectivity. We study the effect of disorder on the connectivity. The Turing patterns are regions of chemical concentration gradients. Turing's original idea was that the gradients are responsible for the patterns seen in biological systems. Later, several physical, chemical and biological systems have been identified [10, 11, 12, 13, 14, 15] which exhibit Turing patterns.

The GM model of RD is described by the following two partial differential equations :

$$\frac{\partial a}{\partial t} = D_a \Delta a + \rho_a \frac{a^2}{(1 + k_a a^2)h} - \mu_a a \quad (1a)$$

$$\frac{\partial h}{\partial t} = D_h \Delta h + \rho_h a^2 - \mu_h h \quad (1b)$$

where Δ is the Laplacian given by $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$, 'a' and 'h' denote the concentrations of the activator and the inhibitor, D_a , D_h , are the respective diffusion coefficients, μ_a , μ_h are the removal rates and ρ_a , ρ_h are the cross-reaction coefficients. The conditions for the formation of stable Turing patterns are $D_h \gg D_a$ and $\mu_h > \mu_a$ [8]. We also assume that $\frac{\rho_a}{1+k_a} = \mu_a$ and $\rho_h = \mu_h$. In this case, the steady state solution of equations (1a) and (1b) is given by (a,h)=(1,1), i.e., the steady state is homogeneous. We use the parameter values $D_a = 0.005$, $D_h = 0.2$, $\rho_h = \mu_h = 0.02$, $\rho_a = 0.0125$ and $k_a = 0.25$ for our study. Bose and Chaudhuri [16] have studied the effect of disorder on the formation of Turing islands in the GM model with $k_a = 0$. We extend this study to the case of $k_a \neq 0$. As in Ref.[16], a very simple discretization scheme is used. The lattice chosen is of size $L \times L$, with L ranging from 20 to 70 in steps of 10. Also, periodic boundary conditions are used. Disorder is introduced into the underlying substrate (lattice) by stipulating that the probability of a particular site being part of the RD network is p . The disordered lattice configuration is generated with the help of a random number generator [16]. For the disordered lattice, the Laplacian is written as

$$\Delta a(x_{ij}, t) = iocc(i + 1, j) \times (a(x_{i+1j}, t) - a(x_{ij}, t))$$

$$\begin{aligned}
& + iocc(i, j + 1) \times (a(x_{ij+1}, t) - a(x_{ij}, t)) \\
& + iocc(i - 1, j) \times (a(x_{i-1j}, t) - a(x_{ij}, t)) \\
& + iocc(i, j - 1) \times (a(x_{ij-1}, t) - a(x_{ij}, t))
\end{aligned} \tag{2}$$

x_{ij} denotes a lattice site (i,j). The array $iocc$ keeps track of the occupation status of the sites of the square lattice. If the site x_{ij} is occupied then $iocc(i, j)=1$, otherwise it is equal to zero. Eqn.(2) expresses the fact that diffusion to a site from a neighbouring site takes place only if the neighbouring site belongs to the RD network. One can easily check that the discretized differential equations (with $\frac{\rho_a}{1+k_a} = \mu_a$ and $\rho_h = \mu_h$) have a steady state solution given by $a=1$ and $h=1$ for all the lattice sites. Random fluctuations of magnitude less than 0.1 are created in the steady state with the help of the random number generator. This fixes the values of a and h at all the cluster sites at time $t=0$. The values of a and h at time $t+1$ are determined at a site x_{ij} belonging to a cluster with the help of the discrete equations for a and h . This process is repeated till the steady state is reached, i.e., the values of a and h at all the cluster sites do not change within a specified accuracy. We define an ‘activated’ site as one at which the activator concentration has a value greater than 1, which is the homogeneous steady-state value. Figs. 1(a)-1(c) show the patterns of activated sites on a lattice of size 50×50 and for site occupation probability $p = 1.0$, $p = 0.9$, and $p = 0.59$. Fig. 1(a) shows a fully-connected stripe pattern with long-range connectivity. For p greater than a critical value p_c , a cluster of n.n. activated sites still spans the lattice. Fig. 1(b) shows a pattern in which long-range connectivity still exists. Fig. 1(c) shows a pattern without long-range connectivity ($p < p_c$). The spanning of the largest cluster is checked as in the percolation problem and the value of p_c is measured by the method of binary chopping [17]. Table I shows the average p_c values determined for lattices of various sizes. For each lattice size, p_c is calculated by taking the average of 100 values. Because of the finite size of the lattice, the value of p_c has a small spread. As in the percolation problem, one expects that p_c has a sharp, unique value as the lattice size $L \rightarrow \infty$. The critical value p_c is greater than the site percolation threshold $p_c^{site} = 0.59$ for the square lattice.

We now describe some cluster-related properties. Fig. 2 shows the spanning cluster for RD on a 50×50 lattice and at $p=p_c$ (Table I). Let M be the size of the spanning cluster, size denotes the number of sites in the cluster. Size of the largest cluster is determined for lattice size $L \times L$, L ranging from

20 to 70 in steps 10. To calculate M for each lattice size, the average of 200 values is taken. Table II shows the data for average M for various lattice sizes. The fractal dimension d_f of the largest cluster is defined as

$$M \sim L^{d_f} \quad (3)$$

Fig. 3 shows a plot of $\log(M)$ vs. $\log(L)$. From the slope, the value of d_f is obtained as $d_f=1.86 \pm 0.01$. The error quoted is the least-square fitting error. The value of d_f is close to that of the spanning cluster ($d_f=1.89$) of ordinary percolation. At any value of $p \leq p_c$, the non-equilibrium steady-state pattern consists of clusters of various sizes. Let n_s be the number of clusters of size s per site of the lattice. The average cluster size S_{AV} is then defined as

$$S_{AV} \sim \sum'_s s^2 n_s \quad (4)$$

The prime indicates that the largest cluster is not included in the sum. As $p \rightarrow p_c$, S_{AV} exhibits a power-law scaling of the form

$$S_{AV} \sim |p - p_c|^{-\gamma} \quad (5)$$

The critical exponent γ can be determined by various numerical methods. We use the method of finite-size scaling to obtain an estimate of γ in our model. In this method, the average cluster size as a function of p and L is given by

$$S_{AV}(p, L) \sim L^{\frac{\gamma}{\nu}} f((p - p_c)L^{\frac{1}{\nu}}) \quad (6)$$

The exponent ν is the correlation length exponent. As $p \rightarrow p_c$, the correlation length ξ diverges as

$$\xi \sim |p - p_c|^{-\nu} \quad (7)$$

The function f in Eqn.(6) is a suitable scaling function and at $p=p_c$, $f(0)=\text{constant}$. The average cluster size is determined for lattice size L ranging from 20 to 70 in steps of 10. For each lattice size, the average of 200 values is taken. Table III shows the data obtained for $p=p_c$. Fig. 4 shows a plot of $\log(S_{AV})$ versus $\log(L)$ for $p=p_c$. From the slope of the straight line, the exponent $\frac{\gamma}{\nu}$ is estimated as $\frac{\gamma}{\nu} = 1.72 \pm 0.01$. Again, the error quoted is the least-square fitting error. The value of $\frac{\gamma}{\nu} = 1.79$. The average cluster size S_{AV} is further

determined for $(p - p_c)$ in the range 0.03 to 0.33. Fig. 5 shows a plot of $\frac{S_{AV}(p,L)}{L^{\frac{1}{\nu}}}$ versus $z=(p - p_c) L^{\frac{1}{\nu}}$. Data for three lattice sizes (L=50, 60 and 70) are shown using the symbols: solid circle (L=50), open circle (L=60) and solid square (L=70). According to Eqn. (6), all the data should fall on a single curve with the functional form $f(z)$. The value of the correlation length exponent ν for which the best collapse of data is obtained is $\nu = 1.66$. The value of ν is $\nu = 1.75$ in the case of ordinary percolation.

III. Conclusion

We have studied a RD model on the square lattice. The non-equilibrium steady state of the model consists of stripe-like patterns with long-range connectivity. A stripe is a region of activator concentration greater than the homogeneous, steady state value of 1. The sites of the underlying square lattice are occupied with probability p . As p falls below a critical value p_c , the long-range connectivity of the steady state pattern disappears. As $p \rightarrow p_c$, the average cluster size diverges. The spanning cluster at $p = p_c$ is a fractal object. The values of the exponent $\frac{2}{\nu}$ and the fractal dimension d_f appears to indicate that the percolation model belongs to the universality class of ordinary percolation. The value 1.66 of the correlation length exponent ν is different from the value 1.75 in the case of ordinary percolation. However, the difference may be due to finite-size effects and also due to the fact that the number of trials (~ 200) over which an average is taken is not large. This is because each trial takes a considerable amount of time. First, a steady state has to be generated which is reached only after several thousands of time steps. It is for this very reason that the lattice size could not be made large. The surface reaction models, we have discussed before, belong to the universality class of directed percolation whereas the GM model of RD appears to belong to the universality class of ordinary percolation. In the ordinary percolation problem, one looks at the connectivity of sites which are randomly occupied with probability p . The occupation status of a particular site is independent of the occupation status of neighbouring sites. In our study, we focus on the connectivity of a RD steady state pattern formed on a disordered lattice. The sites of the pattern are not randomly occupied, as in the case of ordinary percolation, but there is some degree of correlation in the occupancy of sites. The distribution of sites in the steady-state depends on

the parameters of the RD model, the diffusion coefficients D_a , D_h , the saturation parameter k_a etc. The cluster-related properties of the steady state pattern, say, the average cluster size, have a power-law scaling behaviour when occupation probability p of the underlying lattice is close to a critical value p_c . As already mentioned in the Introduction, there are many physical, chemical and biological systems which exhibit Turing patterns. Some of these patterns consist of stripes which may sometimes form in a disordered environment. The present study is of relevance to such systems. In the percolation picture, one studies a transition from localised to extended states as a function of disorder. In the GM model, instead of disorder, one can vary the parameter k_a to bring about the transition. For $k_a=0$, the steady state pattern consists of isolated Turing islands. For $k_a \neq 0$, the steady state consists of connected stripes. An interesting question to ask is whether the transition from localised to extended states occurs as soon as k_a is made different from zero. Preliminary studies [18] shows that the transition occurs at a finite value of k_a . The connection of this transition with the one in the presence of disorder will be explored in future. The effect of changing the parameters of the RD model and also changing the form of the nonlinearity in the RD model (provided it still gives rise to connected patterns in the steady state) will also be investigated.

Acknowledgment

One of the author I.C. is supported by the Council of Scientific and Industrial Research, India under Sanction No. 9/15(173)/96-EMR-I.

Table I. The average value of p_c for lattices of various sizes.

Lattice size	Average p_c
20×20	0.81089
30×30	0.80735
40×40	0.80992
50×50	0.80714
60×60	0.80873
70×70	0.80843

Table II. The average size M of the spanning cluster at $p = p_c$ for lattices of various sizes

Lattice size	Average M
20×20	96.54
30×30	195.77
40×40	350.65
50×50	508.77
60×60	754.81
70×70	974.22

Table III. The average cluster size S_{AV} at $p = p_c$ for lattices of various sizes

Lattice size	Average S_{AV}
20×20	4.01
30×30	8.22
40×40	14.11
50×50	19.65
60×60	27.82
70×70	33.96

Figure Captions

Fig.1 Non-equilibrium steady state patterns in the GM model on a disordered square lattice of size 50×50 . The probability that a site of the original square lattice exists is p . The patterns are shown for (a) $p=1$, (b) $p=0.9$, (c) $p=0.59$.

Fig.2 The spanning cluster on a 50×50 lattice for $p = p_c$ (see Table I.) .

Fig.3 $\log(M)$ versus $\log(L)$ for $p = p_c$ (Eqn.(3)). The slope of the straight line gives the fractal dimension $d_f=1.86\pm 0.01$.

Fig.4 $\log(S_{AV})$ versus $\log(L)$ for $p = p_c$ (Eqn.(6)). The slope of the straight line gives the exponent $\frac{\gamma}{\nu}=1.72\pm 0.01$.

Fig.5 The collapse of data for three different lattice sizes: solid circles, open circles and solid squares for $L=50, 60$ and 70 respectively.

References

- [1] D. Stauffer and A. Aharony, Introduction to Percolation Theory (Taylor and Francis, London, 1994)
- [2] M. Sahimi, Applications of Percolation Theory (Taylor and Francis, London, 1994)
- [3] R. M. Ziff, E. Gulari, and Y. Barshad, Phys.Rev.Lett 56, 2553 (1986)
- [4] T. Aukrust, D. A. Browne and I. Webman, Phys.Rev.A 41, 5294 (1990)
- [5] I. Sendiña-Nadal, D. Vives, V. Pérez-Muñuzuri, M. Gómez-Gesteira, V. Pérez-Villar, J. Echave, J. Casademunt, L. Ramirez-Piscina, and F. Sagueś, Phys.Rev.E 58, 1183 (1998)
- [6] R. Dickman and M. A. Burschka, Phys.Lett.A 127, 132 (1988).
- [7] G. Grinstein, Z. W. Lai and D. A. Browne, Phys.Rev.A 40, 4820, (1989)
- [8] A. J. Koch and H. Meinhardt, Rev.Mod.Phys. 66, 1481 (1994)
- [9] A.M.Turing, Phil.Trans.R.Soc.B 327, 37 (1952)
- [10] V.Castets, E.Dulos, J.Boissonade and P.De Kepper, Phys.Rev.Lett. 64, 2953 (1990)
- [11] Q. Ouyang and H.L. Swinney, Nature 352, 610 (1991)
- [12] K.J.Lee, W.D.McCormick, Z.Noszticzius and H.L.Swinney, J.Chem.Phys. 96, 4048 (1992)
- [13] M.C.Cross and P.C.Hohenberg, Rev.Mod.Phys. 65, 851 (1993)
- [14] Y.Astrov, E.Ammelt, S.Teperick and H.G.Purwins, Phys.Lett. A 211, 184 (1996)
- [15] S. Kondo and R. Asai, Nature 376, 765 (1995)
- [16] Indrani Bose and Indranath Chaudhuri, Phys.Rev.E 55, 5291 (1997)

- [17] D. Stauffer, in *Disordered System and Localization*, ed. by C. Castellani, C. Di Castro and L. Peliti (Lecture Notes in Physics, Vol.149, Springer-Verlag, Berlin 1981)
- [18] Indranath Chaudhuri and Indrani Bose, unpublished.

Ÿ





