

# Phase diagram of the $\frac{1}{2}$ - $\frac{1}{2}$ -1-1 spin chain by the nonlinear $\sigma$ model

Ken'ichi Takano<sup>1</sup>

*Laboratory of Theoretical Condensed Matter Physics and Research Center for Advanced Photon Technology,  
Toyota Technological Institute, Nagoya 468-8511, Japan*

---

## Abstract

We examine a periodic mixed spin chain with spin magnitudes  $\frac{1}{2}$  and 1 which are arrayed as  $\frac{1}{2}$ - $\frac{1}{2}$ -1-1. The three independent parameters are ratios of the four exchange couplings. We determine phase boundaries in the parameter space by using the gapless condition which was previously derived by mapping a general inhomogeneous spin chain to the nonlinear  $\sigma$  model. We find two gapless boundaries separating three disordered phases. The features of the phases are explained in terms of singlet clusters.

*Keywords:* spin gap; mixed spin chain; phase diagram; nonlinear  $\sigma$  model

---

Since the prediction of the Haldane gap for spin chains [1], the nonlinear  $\sigma$  model (NLSM) is considered to be a useful tool to investigate spin systems. After then an NLSM for a spin chain with bond alternation is derived and examined [2]. The application of the NLSM to spin chains with more than one spin species is an interesting problem. In a previous paper [3] we derived an NLSM for a general mixed quantum spin chain with arbitrary finite period. However its various applications have not been examined in detail [4].

In this paper we examine the case that the spin magnitudes in a unit cell are arrayed as  $\frac{1}{2}$ - $\frac{1}{2}$ -1-1 with spacing  $a$ . The Hamiltonian is written as

$$H = \sum_j (J \mathbf{S}_{4j+1} \cdot \mathbf{S}_{4j+2} + J_+ \mathbf{S}_{4j+2} \cdot \mathbf{S}_{4j+3} + \mathbf{S}_{4j+3} \cdot \mathbf{S}_{4j+4} + J_- \mathbf{S}_{4j+4} \cdot \mathbf{S}_{4j+5}), \quad (1)$$

where the magnitudes of  $\mathbf{S}_{4j+1}$  and  $\mathbf{S}_{4j+2}$  are  $\frac{1}{2}$  and those of  $\mathbf{S}_{4j+3}$  and  $\mathbf{S}_{4j+4}$  are 1. We reparametrize the coupling constants as

$$J_+ = J'/(1 + \delta), \quad J_- = J'/(1 - \delta). \quad (2)$$

Following Ref. [3], we map the spin chain described by Eq. (1) to an NLSM. The resultant NLSM action is given by

$$S_{\text{eff}} = \int d\tau \int dx \left\{ -i \frac{J^{(0)}}{J^{(1)}} \mathbf{n} \cdot (\partial_\tau \mathbf{n} \times \partial_x \mathbf{n}) + \frac{1}{2aJ^{(1)}} \left( \frac{J^{(1)}}{J^{(2)}} - \frac{J^{(0)}}{J^{(1)}} \right) (\partial_\tau \mathbf{n})^2 + \frac{a}{2} J^{(0)} (\partial_x \mathbf{n})^2 \right\},$$

$$\frac{4}{J^{(0)}} = \frac{4}{J} + \frac{4}{J'} + 1, \quad \frac{4}{J^{(1)}} = \frac{2}{J} + 1, \quad \frac{4}{J^{(2)}} = \frac{1}{J} + 1.$$

The NLSM has gapless excitations if the topological angle  $4\pi J^{(0)}/J^{(1)}$  is a half-odd-integer multiple of  $2\pi$ . This condition is written as

$$\frac{1}{J'} = -\frac{2l-3}{2l-1} \frac{1}{J} - \frac{1}{4} \frac{2l-5}{2l-1} \quad (l = 1, 2). \quad (3)$$

---

<sup>1</sup> E-mail: takano@toyota-ti.ac.jp




Fig. 1. The phase diagram in the  $J$ - $J'$  plane for arbitrarily fixed  $\delta$ . Three disordered phases are separated by two gapless lines.

This equation for each  $l$  determines a phase boundary between gapful disordered phases. Since the equation is independent of  $\delta$ , the phase boundaries in the  $J$ - $J'$ - $\delta$  space is uniform in the  $\delta$  direction. We present the phase diagram in a  $J$ - $J'$  plane perpendicular to the  $\delta$  axis in Fig. 1. In the special case of  $J_+ = J_- (=J')$ , Tonegawa et al. performed a quantum Monte Carlo simulation and obtained a phase diagram [5]. For example, their phase boundary between phases A and B passes the point  $(J, J') = (1, 0.77)$  instead of  $(1, 4/7)$  in Fig. 1. However the overall feature of Fig. 1 well agrees with the numerical phase diagram.

We explain the phases in Fig. 1 in terms of singlet clusters (SC's). An SC means a local singlet state formed by more than one  $\frac{1}{2}$ -spins. We call a singlet state consisting of SC's the singlet cluster solid (SCS). The SCS is an extension of the valence bond solid [6] and special versions are used to explain phases for other spin chains [7,8]. We show the SCS pictures for phases A, B and C in Fig. 2. Here a spin with magnitude 1 is decomposed into two  $\frac{1}{2}$ -spins (circles) and the symmetrization retrieves the original spin. A closed loop means an SC which is formed by  $\frac{1}{2}$ -spins in it. In phase A, adjacent spins of magnitude 1 are strongly connected




Fig. 2. The SCS pictures for phases A, B and C.

to form two valence bonds, since the coupling 1 between them is larger than  $J'$ . A pair of remnant  $\frac{1}{2}$ -spins form a valence bond almost irrespective of the value of  $J$ . In phase C, both of  $J$  and  $J'$  are larger than 1 and hence four  $\frac{1}{2}$ -spins connected by them form an SC. Phase B is an intermediate; one of two valence bonds remains for a pair of spins of magnitude 1 and a 4-spin cluster is formed in a unit cell.

The SCS picture will be more generally explained [9].

## References

- [1] F. D. M. Haldane, Phys. Lett. **93A**, 464 (1983); Phys. Rev. Lett. **50**, 1153 (1983).
- [2] I. Affleck, Nucl. Phys. B **257**, 397 (1985); **265**, 409 (1986); I. Affleck and F. D. M. Haldane, Phys. Rev. B **36**, 5291 (1987).
- [3] K. Takano, to be published in Phys. Rev. Lett.
- [4] Special cases have been examined by a different NLSM method: T. Fukui and N. Kawakami, Phys. Rev. **B56**, 8799 (1997).
- [5] T. Tonegawa, private communication; T. Tonegawa, T. Hikihara, M. Kaburagi, T. Nishino, S. Miyashita and H.-J. Mikeska, J. Phys. Soc. Jpn. **67**, 1000 (1998).
- [6] I. Affleck, T. Kennedy, E. H. Lieb and H. Tasaki, Phys. Rev. Lett. **59**, 799 (1987).
- [7] K. Takano, K. Kubo and H. Sakamoto, J. Phys.: Condens. Matter **8**, 6405 (1996).
- [8] W. Chen and K. Hida, J. Phys. Soc. Jpn. **67**, 2910 (1998).
- [9] K. Takano, in preparation.