Fast Multipoint Evaluation of Bivariate Polynomials

Michael Nüsken and Martin Ziegler*

University of Paderborn, 33095 Paderborn, GERMANY {nuesken, ziegler}@upb.de

Abstract. We generalize univariate multipoint evaluation of polynomials of degree n at sublinear amortized cost per point. More precisely, it is shown how to evaluate a bivariate polynomial p of maximum degree less than n, specified by its n^2 coefficients, simultaneously at n^2 given points using a total of $\mathcal{O}(n^{2.667})$ arithmetic operations. In terms of the input size N being quadratic in n, this amounts to an amortized cost of $\mathcal{O}(N^{0.334})$ per point.

1 Introduction

By Horner's Rule, any polynomial p of degree less than n can be evaluated at a given argument x in $\mathcal{O}(n)$ arithmetic operations which is optimal for a generic polynomial as proved by Pan (1966), see for example Theorem 6.5 in Bürgisser, Clausen & Shokrollahi (1997).

In order to evaluate p at several points, we might sequentially compute $p(x_k)$ for $0 \le k < n$. However, regarding that both the input consisting of n coefficients of p and n points x_k and the output consisting of the n values $p(x_k)$ have only linear size, information theory provides no justification for this quadratic total running time. In fact, a more sophisticated algorithm permits to compute all $p(x_k)$ simultaneously using only $\mathcal{O}(n \cdot \log^2 n \cdot \log \log n)$ operations. Based on the Fast Fourier Transform, the mentioned algorithms and others realize what is known as Fast Polynomial Arithmetic. For ease of notation, we use the 'soft-Oh' notation, namely $\mathcal{O}^{\sim}(f(n)) := \mathcal{O}\left(f(n)(\log f(n))^{\mathcal{O}(1)}\right)$. This variant of the usual asymptotic 'big-Oh' notation ignores poly-logarithmic factors like $\log^2 n \cdot \log \log n$.

Fact 1. Let R be a commutative ring with one.

(i) Multiplication of univariate polynomials: Suppose we are given polynomials $p, q \in R[X]$ of degree less than n, specified by their coefficients. Then we can compute the coefficients of the product polynomial $p \cdot q \in R[X]$ using $\mathcal{O}^{\sim}(n)$ arithmetic operations in R.

^{*} Supported by the DFG Research Training Group GK-693 of the Paderborn Institute for Scientific Computation (PaSCo)

- (ii) Multipoint evaluation of a univariate polynomial: Suppose we are given a polynomial $p \in R[X]$ of degree less than n, again specified by its coefficients, and points $x_0, \ldots, x_{n-1} \in R$. Then we can compute the values $p(x_0), \ldots, p(x_{n-1}) \in R$ using $\mathcal{O}^{\sim}(n)$ arithmetic operations in R.
- (iii) Univariate interpolation: Conversely, suppose we are given points $(x_k, y_k) \in \mathbb{R}^2$ for $0 \le k < n$ such that $x_k x_\ell$ is invertible in R for all $k \ne \ell$. Then we can compute the coefficients of a polynomial $p \in R[X]$ of degree less than n such that $p(x_k) = y_j$, $0 \le k < n$, that is, determine the interpolation polynomial to data (x_k, y_k) using $\mathcal{O}^{\sim}(n)$ arithmetic operations in R.

Proof. These results can be found for example in von zur Gathen & Gerhard (2003) including small constants:

(i) can be done using at most $63.427 \cdot n \cdot \log_2 n \cdot \log_2 \log_2 n + \mathcal{O}(n \log n)$ arithmetic operations in R by Theorem 8.23. The essential ingredient is the Fast Fourier Transform. If $R = \mathbb{C}$ then even $\frac{9}{2}n \log_2 n + \mathcal{O}(n)$ arithmetic operations suffice. This goes back to Schönhage & Strassen (1971) and Schönhage (1977).

In the following M(n) denotes the cost of one multiplication of univariate polynomials over R of degree less then n.

- (ii) can be done using at most $\frac{11}{2}M(n)\log_2 n + \mathcal{O}(n\log n)$ operations in R according to Corollary 10.8. Here, Divide & Conquer provides the final building block. This goes back to Fiduccia (1972).
- (iii) can be done using at most $\frac{13}{2}M(n)\log_2 n + \mathcal{O}(n\log n)$ operations in R according to Corollary 10.12. This, too, is completed by Divide & Conquer. The result goes back to Horowitz (1972).

You also find an excellent account of all these in Borodin & Munro (1975).

Fast polynomial arithmetic and in particular multipoint evaluation has found many applications in algorithmic number theory (see for example Odlyzko & Schönhage 1988), computer aided geometric design (see for example Lodha & Goldman 1997), and computational physics (see for example Ziegler 2003b).

Observe that the above claims apply to the univariate case. What about multivariate analogues? Let us for a start consider the bivariate case: A bivariate polynomial $p \in R[X,Y]$ of maximum degree max $\deg p := \max \{\deg_X p, \deg_Y p\}$ less than n has up to n^2 coefficients, one for each monomial X^iY^j with $0 \le i,j < n$. Now corresponding to Fact 1, the following questions emerge:

- **Question 2.** (i) Multiplication of bivariate polynomials: Can two given bivariate polynomials of maximum degree less than n be multiplied within time $\mathcal{O}^{\sim}(n^2)$?
- (ii) Multipoint evaluation of a bivariate polynomial: Can a given bivariate polynomial of maximum degree less than n be evaluated simultaneously at n^2 arguments in time $\mathcal{O}^{\sim}(n^2)$?
- (iii) Bivariate interpolation: Given n^2 points $(x_k, y_k, z_k) \in \mathbb{R}^3$, is there a polynomial $p \in \mathbb{R}[X,Y]$ of maximum degree less than n such that $p(x_k, y_k) = z_k$ for all $0 \le k < n^2$? And, if yes, can we compute it in time $\mathcal{O}^{\sim}(n^2)$?

64 generic points.

Such issues also arise for instance in connection with fast arithmetic for polynomials over the skew-field of hypercomplex numbers (Ziegler 2003a, Section 3.1).

A positive answer to Question 2(i) is achieved by embedding p and q into univariate polynomials of degree $\mathcal{O}(n^2)$ using the Kronecker substitution $Y = X^{2n-1}$, applying Fact 1(i) to them, and then re-substituting the result to a bivariate polynomial; see for example Corollary 8.28 in von zur Gathen & Gerhard (2003) or Section 1.8 in Bini & Pan (1994).

Note that the first part of (iii) has negative answer for instance whenever the points (x_k, y_k) are co-linear or, more generally, lie on a curve of small degree: Here, a bivariate polynomial of maximum degree less than n does not even exist in general.

Addressing (ii), observe that Kronecker substitution is not compatible with evaluation and thus of no direct use for reducing to the univariate case. The methods that yield Fact 1(ii) are not applicable either as they rely on fast polynomial division with remainder which looses many of its nice mathematical and computational properties when passing from the univariate to the bivariate case.

Nevertheless, (ii) does admit a rather immediate positive answer provided the arguments (x_k, y_k) , $0 \le k < n^2$ form a Cartesian $n \times n$ -grid (also called tensor product grid). Indeed, consider $p(X,Y) = \sum_{0 \le j < n} q_j(X)Y^j$ as a polynomial in Y with coefficients q_j being univariate polynomials in X. Then multi-evaluate q_j at the n distinct values x_k : as q_j has degree less than n, this takes time $\mathcal{O}^{\sim}(n)$ for each j, adding to a total of $\mathcal{O}^{\sim}(n^2)$. Finally take the n different univariate polynomials $p(x_k, Y)$ in Y of degree less than n and multi-evaluate each at the n distinct values y_ℓ : this takes another $\mathcal{O}^{\sim}(n^2)$.

The presumption on the arguments to form a Cartesian grid allows for a slight relaxation in that this grid may be rotated and sheared: Such kind of



Fig. 1. Cartesian 8×8 -grid, same rotated and sheared;

affine distortion is easy to detect, reverted to the arguments, and then instead applied to the polynomial p by transforming its coefficients within time $\mathcal{O}^{\sim}(n^2)$, see Lemma 14 below. The obtained polynomial \hat{p} can then be evaluated on the now strictly Cartesian grid as described above. However, $n \times n$ grids, even rotated and sheared ones, form only a zero-set within the $2n^2$ -dimensional space of all possible configurations of n^2 points. Thus this is a severe restriction.

2 Goal and Idea

The big open question and goal of the present work is concerned with fast multipoint evaluation of a multivariate polynomial. As a first step in this direction we consider the bivariate case.

The naïve approach to this problem, namely of sequentially calculating all $p(x_k, y_k)$, takes quadratic time each, thus inferring total cost of order n^4 . A first improvement to $\mathcal{O}^{\sim}(n^3)$ is based on the simple observation that any n points in the plane can easily be extended to an $n \times n$ grid on which, by the above considerations, multipoint evaluation of p is feasible in time $\mathcal{O}^{\sim}(n^2)$. So we may partition the n^2 arguments into n blocks of n points and multi-evaluate p sequentially on each of them to obtain the following

Theorem 3. Let R be a commutative ring with one. A bivariate polynomial $p \in R[X,Y]$ of $\deg_X(p) < n$ and $\deg_Y(p) < n$, given by its coefficients, can be evaluated simultaneously at n^2 given arguments (x_k, y_k) using at most $\mathcal{O}(n^3 \cdot \log^2 n \cdot \log \log n)$ arithmetic operations in R.

We reduce this softly cubic upper complexity bound to $\mathcal{O}(n^{2.667})$. More precisely, by combining fast univariate polynomial arithmetic with fast matrix multiplication we will prove:

Result 4. Let \mathbb{K} denote an arbitrary field. A bivariate polynomial $p \in \mathbb{K}[X,Y]$ of $\deg_X(p) < n$ and $\deg_Y(p) < m$, specified by its coefficients, can be evaluated simultaneously at N given arguments $(x_k, y_k) \in \mathbb{K}^2$ with pairwise different first coordinates using $\mathcal{O}\left((N+nm)m^{\omega_2/2-1+\varepsilon}\right)$ arithmetic operations in \mathbb{K} for any fixed $\varepsilon > 0$.

Here, ω_2 denotes the exponent of the multiplication of $n \times n$ - by rectangular $n \times n^2$ -matrices, see Section 3. In fact this problem is well-known to admit a much faster solution than naïve $\mathcal{O}(n^4)$, the current world record $\omega_2 < 3.334$ being due to Huang & Pan (1998). By choosing m = n and $N = n^2$, this yields the running time claimed in the abstract.

The general idea underlying Result 4, illustrated for the case of n=m, is to reduce the bivariate to the univariate case by substituting Y in p(X,Y) with the interpolation polynomial g(X) of degree less than n^2 to data (x_k,y_k) . It then suffices to multi-evaluate the univariate result p(X,g(X)) at the n^2 arguments x_k . Obviously, this can only work if such an interpolation polynomial g is available, that is any two evaluation points $(x_k,y_k)\neq (x_{k'},y_{k'})$ differ in their first coordinates, $x_k\neq x_{k'}$. However, this condition can be asserted easily later on, see Section 6, so for now assume it is fulfilled.

This naïve substitution leads to a polynomial of degree up to $\mathcal{O}(n^3)$. On the other hand, it obviously suffices to obtain p(X, g(X)) modulo the polynomial $f(X) := \prod_{0 \le k < n^2} (X - x_k)$ which has degree less than n^2 . The key to efficient bivariate multipoint evaluation is thus an efficient algorithm for this modular bi-to-univariate composition problem, presented in Theorem 9.

As we make heavy use of fast matrix multiplication, Section 3 recalls some basic facts, observations, and the state of the art in that field of research. Section 4 formally states the main result of the present work together with two tools (affine substitution and modular composition) which might be interesting on their own, their proofs being postponed to Section 5. Section 6 describes three ways to deal with arguments that do have coinciding first coordinates. Section 7 gives some final remarks.

3 Basics on Fast Matrix Multiplication

Recall that, for a field \mathbb{K} , $\omega = \omega(\mathbb{K}) \geq 2$ denotes the *exponent of matrix multiplication*, that is, the least real such that $m \times m$ matrix multiplication is feasible in asymptotic time $\mathcal{O}(m^{\omega+\varepsilon})$ for any $\varepsilon > 0$; see for example Chapter 15 in Bürgisser *et al.* (1997). The current world-record due to Coppersmith & Winograd (1990) achieves $\omega < 2.376$ independent of the ground field \mathbb{K} . The Notes 12.1 in von zur Gathen & Gerhard (2003) contain a short historical account.

Clearly, a rectangular matrix multiplication of, say, $m \times m$ -matrices by $m \times m^t$ -matrices can always be done partitioning into $m \times m$ square matrices. Yet, in some cases there are better known algorithms than this. We use the notation introduced by Huang & Pan (1998): $\omega(r, s, t)$ denotes the exponent of the multiplication of $\lceil m^r \rceil \times \lceil m^s \rceil$ - by $\lceil m^s \rceil \times \lceil m^t \rceil$ -matrices, that is

$$\omega(r,s,t) = \inf \left\{ \tau \in \mathbb{R} \left| \begin{array}{l} \text{Multiplication of } \lceil m^r \rceil \times \lceil m^s \rceil \text{- by} \\ \lceil m^s \rceil \times \lceil m^t \rceil \text{-matrices can be done} \\ \text{with } \mathcal{O}(m^\tau) \text{ arithmetic operations} \end{array} \right\}.$$

Clearly, $\omega = \omega(1, 1, 1)$. We always have

$$\max\{r+s,r+t,s+t\} \leq \omega(r,s,t) \leq r+s+t. \tag{5}$$

Note that $\omega(r, s, t)$ is in fact invariant under permutation of its arguments. We collect some known bounds on fast matrix multiplication algorithms.

Fact 6. (i) $\omega = \omega(1,1,1) \le \log_2(7) < 2.8073549221$ (Strassen 1969). (ii) $\omega = \omega(1,1,1) < 2.3754769128$ (Coppersmith & Winograd 1990). (iii) $\omega_2 := \omega(1,1,2) < 3.3339532438$ (Huang & Pan 1998).

Partitioning into square matrices only yields $\omega_2 \leq \omega + 1 < 3.3754769128$. Bounds for further rectangular matrix multiplications can be also be found in Huang & Pan (1998). It is conjectured that $\omega = 2$. Then by partitioning into square blocks also $\omega(r,s,t)$ touches its lower bound in (5), that is $\omega(r,s,t) = \max\{r+s,r+t,s+t\}$. In particular, $\omega_2 = 3$ then.

We point out that the definition of ω and $\omega(r, s, t)$ refers to arbitrary algebraic computations which furthermore may be non-uniform, that is, use for each matrix size m a different algorithm. However, closer inspection of Section 15.1 in Bürgisser *et al.* (1997) reveals the following

Observation 7. Rectangular matrix multiplication of $\lceil m^r \rceil \times \lceil m^s \rceil$ - by $\lceil m^s \rceil \times \lceil m^t \rceil$ -matrices over \mathbb{K} can be done with $\mathcal{O}(m^{\omega(r,s,t)+\varepsilon})$ arithmetic operations in \mathbb{K} by a uniform, bilinear algorithm for any fixed ε .

A bilinear computation is a very special kind of algorithm where apart from additions and scalar multiplications only bilinear multiplications occur; see for example Definition 14.7 in Bürgisser *et al.* (1997) for more details. In particular, no divisions are allowed.

4 Main results

Our major contribution concerns bivariate multi-evaluation at arguments (x_k, y_k) under the condition that their first coordinates x_k are pairwise distinct. This amounts to a weakened *general position* presumption as is common for instance in Computational Geometry.

For notational convenience, we define ' \mathcal{O}^{\approx} ' (smooth-Oh) which, in addition to polylogarithmic factors in n, also ignores factors n^{ε} as long as $\varepsilon > 0$ can be chosen arbitrarily small. Formally, $\mathcal{O}^{\approx}(f(n)) := \bigcap_{\varepsilon > 0} \mathcal{O}(f(n)^{1+\varepsilon})$. Note that $\mathcal{O}^{\sim}(f(n)) \subset \mathcal{O}^{\approx}(f(n))$.

Theorem 8. Let \mathbb{K} denote a field. Suppose $n, m \in \mathbb{N}$. Given the nm coefficients of a bivariate polynomial p with $\deg_X(p) < n$ and $\deg_Y(p) < m$ and given nm points $(x_k, y_k) \in \mathbb{K}^2$, $0 \le k < nm$ such that the first coordinates x_k are pairwise different, we can calculate the n values $p(x_k, y_k)$ using $\mathcal{O}^{\approx}(nm^{\omega_2/2})$ arithmetic operations over \mathbb{K} . The algorithm is uniform.

Observe that this yields the first part of Result 4 by performing $\lceil N/(nm) \rceil$ separate multipoint evaluations at nm points each. Let us also remark that any further progress in matrix multiplication immediately carries over to our problem. As it is conjectured that $\omega=2$ holds, this would lead to bivariate multipoint evaluation within time $\mathcal{O}^{\approx}(nm^{1.5})$.

Our proof of Theorem 8 is based on the following generalization of Brent & Kungs efficient *univariate* modular composition, see for example Section 12.2 in von zur Gathen & Gerhard (2003), to a certain 'bi-to-univariate' variant:

Theorem 9. Fix a field \mathbb{K} . Given $n, m \in \mathbb{N}$, a bivariate polynomial $p \in \mathbb{K}[X, Y]$ with $\deg_X(p) < n$ and $\deg_Y(p) < m$ and univariate polynomials $g, f \in \mathbb{K}[X]$ of degree less than nm, specified by their coefficients. Then p(X, g(X)) rem f(X) can be computed with $\mathcal{O}^{\approx}(nm^{\omega_2/2})$ arithmetic operations in \mathbb{K} .

We remark that true bivariate modular computation requires Gröbner basis methods which for complexity reasons are beyond our interest here.

5 Proofs

Now we come to the proofs.

Lemma 10. Let \mathbb{K} denote a field and fix t > 0.

- (i) Let both A be an $m \times m$ -matrix and B an $m \times m^t$ -matrix whose entries consist of polynomials $a_{ij}(X), b_{ij}(X) \in \mathbb{K}[X]$ of degree less than n. Given m and the $n \cdot (m^2 + m^t)$ coefficients, we can compute the coefficients of the polynomial entries $c_{ij}(X)$ of $C := A \cdot B$ within $\mathcal{O}^{\approx}(nm^{\omega(1,1,t)})$ arithmetic operations.
- (ii) If A denotes an $m \times m$ square matrix with polynomial entries of degree less than n and b denotes an m-component vector of polynomials of degree less than nm^t , then $(A,b) \mapsto A \cdot b$ is computable within $\mathcal{O}^{\approx}(nm^{\omega(1,1,t)})$.
- (iii) Let $p_0, \ldots, p_{m-1} \in \mathbb{K}[X, Y]$ denote bivariate polynomials with $\deg_X(p_i) < n$ and $\deg_Y(p_i) < m$, given their nm^2 coefficients, and let furthermore univariate polynomials $g, f \in \mathbb{K}[X]$ of degree less than nm^t be given by their coefficients. Then the coefficients of the m univariate polynomials

$$p_i(X, g(X)) \operatorname{rem} f(X)$$

can be computed with $\mathcal{O}^{\approx}(nm^{\omega(1,1,t)})$ arithmetic operations.

In particular, for t=1 we have $cost\ \mathcal{O}^{\approx}(nm^{\omega}) \subset \mathcal{O}^{\sim}(nm^{2.376})$ and for t=2 we have $cost\ \mathcal{O}^{\approx}(nm^{\omega_2}) \subset \mathcal{O}^{\sim}(nm^{3.334})$.

- *Proof.* (i) By scalar extension to $R = \mathbb{K}[X]$ we obtain an algorithm with cost $\mathcal{O}^{\approx}(m^{\omega(1,1,t)})$ arithmetic operations in R using Observation 7. For the algorithm scalar extension simply means that we perform any multiplication in R instead of \mathbb{K} , multiplications with constants become scalar multiplications. And the cost for one operation in R is $\mathcal{O}^{\sim}(n)$ as only polynomials of degree n have to be multiplied.
- (ii) For each j, $0 \le j < m$, decompose the polynomial b_j of degree less than nm^t into m^t polynomials of degree less than n, that is, write $b_j(X) = \sum_{0 \le k \le m^t} b_{jk}(X) \cdot X^{kn}$. The desired polynomial vector is then given by

$$(A \cdot b)_{i}(X) = \sum_{1 \leq j \leq m} a_{ij}(X) \cdot \left(\sum_{0 \leq k < m^{t}} b_{jk}(X) \cdot X^{kn}\right)$$
$$= \sum_{0 \leq k < m^{t}} (A \cdot B)_{ik}(X) \cdot X^{kn}$$
(*)

where $0 \le i < m$ and $B := (b_{jk})$ denotes an $m \times m^t$ matrix of polynomials of degree less than n. The product $A \cdot B$ can be computed according to (i) in the claimed running time. Multiplication by X^{kn} amounts to mere coefficient shifts rather than arithmetic operations. And observing that $\deg ((A \cdot B)_{ik}) < 2n$, only two consecutive terms in the right hand side of (*) can overlap. So evaluating this sum amounts to m^t -fold addition of pairs of polynomials of degree less than n. Since $\omega(1, 1, t) \ge 1 + t$ by virtue of (5), this last cost of nm^{1+t} is also covered by the claimed complexity bound.

(iii) Write each p_i as a polynomial in Y with coefficients from $\mathbb{K}[X]$, that is

$$p_i(X,Y) = \sum_{0 \le j \le m} q_{ij}(X) \cdot Y^j$$

with all $q_{ij}(X)$ of degree less than n. Iteratively compute the m polynomials $g_j(X) := g^j(X)$ rem f(X), each of degree less than nm^t , within time $\mathcal{O}^{\sim}(nm^{1+t})$ by fast division with remainder (see for example Theorem 9.6 in von zur Gathen & Gerhard 2003).

By multiplying the matrix $A := (q_{ij})$ to the vector $b := (g_j)$ according to (ii), determine the m polynomials

$$\tilde{p}_i(X) := \sum_{0 \le j < m} q_{ij}(X) \cdot g_j(X), \qquad 0 \le i < m$$

of degree less than $n + nm^t$. For each i reduce again modulo f(X) and obtain $p_i(X, g(X))$ rem f(X) using another $\mathcal{O}^{\sim}(nm^{1+t})$ operations. Since $\omega(1, 1, t) \geq 1 + t$ according to (5), both parts are covered by the claimed running time $\mathcal{O}^{\approx}(nm^{\omega(1, 1, t)})$.

Lemma 10 puts us in position to prove Theorem 9.

Proof (Theorem 9). Without loss of generality we assume that m is a square. We use a baby step, giant step strategy: Partition p into \sqrt{m} polynomials p_i of $\deg_V(p_i) < \sqrt{m}$, that is

$$p(X,Y) = \sum_{0 \le i < \sqrt{m}} p_i(X,Y) \cdot Y^{i\sqrt{m}} .$$

Then apply Lemma 10(iii) with t=2 and m replaced by \sqrt{m} to obtain the \sqrt{m} polynomials $\tilde{p}_i(X) := p_i(X, g(X))$ rem f(X) within $\mathcal{O}^{\approx}(nm^{\omega_2/2})$ operations. Iteratively determine the \sqrt{m} polynomials $\tilde{g}_i(X) := (g(X)^{\sqrt{m}})^i$ rem f(X) for $0 \le i < \sqrt{m}$ within $\mathcal{O}^{\sim}(nm^{3/2})$. Again, $\omega_2 \ge 3$ asserts this to remain in the claimed bound. Finally compute

$$p(X, g(X)) \operatorname{rem} f(X) = \sum_{0 \le i < \sqrt{m}} (\tilde{p}_i(X) \cdot \tilde{g}_i(X)) \operatorname{rem} f(X)$$

using another time $\mathcal{O}^{\sim}(nm^{3/2})$.

Based on Theorem 9, the following algorithm realizes the idea expressed in Section 2.

Algorithm 11. Generic multipoint evaluation of a bivariate polynomial.

Input: Coefficients of a polynomial $p \in \mathbb{K}[X,Y]$ of $\deg_X(p) < n$, $\deg_Y(p) < m$ and points (x_k,y_k) for $0 \le k < nm$ with pairwise different first coordinates x_k .

Output: The values $p(x_k, y_k)$ for $0 \le k < nm$.

- 1. Compute the univariate polynomial $f(X) := \prod_{0 \le k \le nm} (X x_k) \in \mathbb{K}[X]$.
- 2. Compute an interpolation polynomial $g \in \mathbb{K}[X]$ of degree less than nm satisfying $g(x_k) = y_k$ for all $0 \le k < nm$.
- 3. Apply Theorem 9 to obtain $\tilde{p}(X) := p(X, g(X))$ rem f(X).
- 4. Multi-evaluate this univariate polynomial $\tilde{p} \in \mathbb{K}[X]$ of degree less than nm at the nm arguments x_k .
- 5. Return $(\tilde{p}(x_k))_{0 \leq k < nm}$.

Proof (Theorem 8). The algorithm is correct by construction.

Step 1 in Algorithm 11 can be done in $\mathcal{O}^{\sim}(nm)$ arithmetic operations. As the points (x_k, y_k) have pairwise different first coordinates, the interpolation problem in Step 2 is solvable and, by virtue of Fact 1(iii), in running time $\mathcal{O}^{\sim}(nm)$. For Step 3 Theorem 9 guarantees running time $\mathcal{O}^{\approx}(nm^{\omega_2/2})$. According to Fact 1(ii), Step 4 is possible within time $\mathcal{O}^{\sim}(nm)$. Summing up, we obtain the claimed running time.

6 Evaluating at degenerate points

Here we indicate how certain fields \mathbb{K} permit to remove the condition on the evaluation point set imposed in Theorem 8. The idea is to rotate or shear the situation slightly, so that afterwards the point set has pairwise different first coordinates. To this end choose $\theta \in \mathbb{K}$ arbitrary such that

$$\#\{x_k + \theta y_k \mid 0 \le k < N\} = N \tag{12}$$

where N := nm denotes the number of points. Then replace each (x_k, y_k) by $(x'_k, y'_k) := (x_k + \theta y_k, y_k)$ and the polynomial p by $\hat{p}(X, Y) := p(X - \theta Y, Y)$. This can be done with $\mathcal{O}^{\sim}(n^2 + m^2)$ arithmetic operations, see the more general Lemma 14 below. In any case a perturbation like this might even be a good idea if there are points whose first coordinates are 'almost equal' for reasons of numerical stability.

Lemma 13. Let \mathbb{K} denote a field and $P = \{(x_k, y_k) \in \mathbb{K}^2 \mid 0 \le k < N\}$ a collection of N planar points.

- (i) If $\#\mathbb{K} \geq N^2$, then $\theta \in \mathbb{K}$ chosen uniformly at random satisfies (12) with probability at least $\frac{1}{2}$. Using $\mathcal{O}(\log N)$ guesses and a total of $\mathcal{O}(N \cdot \log^2 N)$ operations, we can thus find an appropriate θ with high probability. If \mathbb{K} is even infinite, a single guess almost certainly suffices.
- (ii) In case $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$, we can deterministically find an appropriate θ in time $\mathcal{O}(N \cdot \log N)$.
- (iii) For a fixed proper extension field \mathbb{L} of \mathbb{K} , any $\theta \in \mathbb{L} \setminus \mathbb{K}$ will do.

Applying (i) or (ii) together with Lemma 14 affects the running time of Theorem 8 only by the possible change in the Y-degree. Using (iii) means that all subsequent computations must be performed in \mathbb{L} . This increases all further costs by no more than an additional constant factor depending on the degree $[\mathbb{L} : \mathbb{K}]$ only.

- Proof. (i) Observe that an undesirable θ with $x_k + \theta y_k = x_{k'} + \theta y_{k'}$ implies $y_k = y_{k'}$ or $\theta = \frac{x_k x_{k'}}{y_{k'} y_k}$. In the latter case, θ is thus uniquely determined by $\{k, k'\}$. Since there are at most $\binom{N}{2} < N^2/2$ such choices $\{k, k'\}$, no more than half of the $\#\mathbb{K} \geq N^2$ possible values of θ can be undesirable.
- (ii) If $\mathbb{K} = \mathbb{R}$ choose $\theta > 0$ such that $\theta \cdot (y_{\max} y_{\min}) < \min\{x_k x_{k'} \mid x_k > x_{k'}\}$. Such a value θ can be found in linear time after sorting the points with respect to their x-coordinate.
 - In case $\mathbb{K} = \mathbb{C}$, we can do the same with respect to the real parts.
- (iii) Simply observe that 1 and θ are linearly independent.

We now state the already announced

Lemma 14. Let R be a commutative ring with one. Given $n \in \mathbb{N}$ and the n^2 coefficients of a polynomial $p(X,Y) \in R[X,Y]$ of degree less than n in both X and Y. Given furthermore a matrix $A \in R^{2\times 2}$ and a vector $b \in R^2$. From this, we can compute the coefficients of the affinely transformed polynomial $p(a_{11}X + a_{12}Y + b_1, a_{21}X + a_{22}Y + b_2)$ using $\mathcal{O}(n^2 \cdot \log^2 n \cdot \log \log n)$ or $\mathcal{O}^{\sim}(n^2)$ arithmetic operations over R.

In the special case $R = \mathbb{C}$ we can decrease the running time to $\mathcal{O}(n^2 \log n)$.

Lemma 14 straight-forwardly generalizes to d-variate polynomials and d-dimensional affine transformations being applicable within time $\mathcal{O}^{\sim}(n^d)$ for fixed d.

Proof. We prove this in several steps.

- First we note that, over any commutative ring S with one, we can compute the Taylor shift p(X+a) of a polynomial $p \in S[X]$ of degree less than n by an element $a \in S$ using $\mathcal{O}(n \cdot \log^2 n \cdot \log \log n)$ arithmetic operations in S. There are many solutions for computing the Taylor shift of a polynomial. We would like to sketch the divide and conquer solution from Fact 2.1(iv) in von zur Gathen (1990) that works over any ring S: Precompute all powers $(X+a)^{2^i}$ for $0 \le i \le \nu := \lfloor \log_2 n \rfloor$. Then recursively split $p(X) = p_0(X) + X^{2^\nu}p_1(X)$ with $\deg p_0 < 2^\nu$ and calculate $p(X+a) = p_0(X+a) + (X+a)^{2^\nu}p_1(X+a)$. This amounts to $\mathcal{O}(n \cdot \log^2 n \cdot \log \log n)$ multiplications in S and $\mathcal{O}(n \log n)$ other operations. So we achieve this over any ring S with $\mathcal{O}(n \cdot \log^2 n \cdot \log \log n)$ operations.
- Next let S = R[Y]. Then we can use the previous to compute p(X + a, Y) or p(X + aY, Y) for a polynomial $p \in R[X, Y] = S[X]$ of maximum degree less than n and an element $a \in R$. Using Kronecker substitution for the multiplications in R[X, Y] this can be done with $\mathcal{O}(n^2 \cdot \log^2 n \cdot \log \log n)$ arithmetic operations in R.

- Now we prove the assertion. Scaling is easy: $p(x,y) \mapsto p(\alpha x,y)$ obviously works within $\mathcal{O}(n^2)$ steps. Use this and the discussed shifts once or twice.

The solution to Problem 2.6 in Bini & Pan (1994) allows to save a factor $\log n \cdot \log \log n$ when $R = S = \mathbb{C}$.

7 Conclusion and Further Questions

We lowered the upper complexity bound for multi-evaluating dense bivariate polynomials of degree less than n with n^2 coefficients at n^2 points with pairwise different first coordinates from naïve $\mathcal{O}(n^4)$ and $\mathcal{O}^{\sim}(n^3)$ to $\mathcal{O}(n^{2.667})$. The algorithm is based on fast univariate polynomial arithmetic together with fast matrix multiplication and will immediately benefit from any future improvement of the latter.

With the same technique, evaluation of a trivariate polynomial of maximum degree less than n at n^3 points can be accelerated from naïve $\mathcal{O}(n^6)$ to $\mathcal{O}(n^{4.334})$.

Regarding that the matrix multiplication method of Huang & Pan (1998) has huge constants hidden in the big-Oh notation, it might in practice be preferable to use either the naïve $2m^3$ or Strassen's $4.7m^{2.81}$ algorithm (with some tricks). Applying them to our approach still yields bivariate multipoint evaluation within time $\mathcal{O}(n^3)$ or $\mathcal{O}(n^{2.91})$, respectively, with small big-Oh constants and no hidden factors $\log n$ in the leading term, that is, faster than Theorem 3.

Further questions to consider are:

- Is it possible to remove even the divisions? This would give a much more stable algorithm and it would also work over many rings.
- As $\omega \geq 2$, the above techniques will never get below running times of order $n^{2.5}$. Can we achieve an upper complexity bound as close as $\mathcal{O}^{\sim}(n^2)$ to the information theoretic lower bound?
- Can multipoint evaluation of trivariate polynomials $p(X_1, X_2, X_3)$ be performed in time o(n^4)?

Acknowledgements The authors wish to thank David Eppstein (2004) for an inspiring suggestion that finally led to Lemma 13(ii).

References

- DARIO BINI & VICTOR Y. PAN (1994). Polynomial and matrix computations, volume 1 of Progress in theoretical computer science. Birkhäuser Verlag, Boston, Basel, Berlin. ISBN 0-8176-3786-9, 3-7643-3786-9.
- A. Borodin & I. Munro (1975). The Computational Complexity of Algebraic and Numeric Problems. Number 1 in Theory of computation series. American Elsevier Publishing Company, New York.
- R. P. Brent & H. T. Kung (1978). Fast Algorithms for Manipulating Formal Power Series. *Journal of the ACM* **25**(4), 581–595.

- Peter Bürgisser, Michael Clausen & Mohammed Amin Shokrollahi (1997). Algebraic Complexity Theory. Number 315 in Grundlehren der mathematischen Wissenschaften. Springer-Verlag.
- Don Coppersmith & Shmuel Winograd (1990). Matrix Multiplication via Arithmetic Progressions. *Journal of Symbolic Computation* 9, 251–280.
- DAVID EPPSTEIN (2004). Re: Geometry problem: Optimal direction. Known results? Usenet news article. URL http://mathforum.org/epigone/sci.math.research/slexyaxsle.
- Charles M. Fiduccia (1972). Polynomial evaluation via the division algorithm: the fast Fourier transform revisited. In *Proceedings of the Fourth Annual ACM Symposium on the Theory of Computing*, Denver CO, 88-93. ACM Press.
- Joachim von zur Gathen (1990). Functional Decomposition of Polynomials: the Tame Case. Journal of Symbolic Computation 9, 281–299.
- JOACHIM VON ZUR GATHEN & JÜRGEN GERHARD (2003). Modern Computer Algebra. Cambridge University Press, Cambridge, UK, 2nd edition. ISBN 0-521-82646-2. URL http://www-math.upb.de/~aggathen/mca/. First edition 1999.
- ELLIS HOROWITZ (1972). A fast method for interpolation using preconditioning. *Information Processing Letters* 1, 157–163.
- XIAOHAN HUANG & VICTOR Y. PAN (1998). Fast Rectangular Matrix Multiplication and Applications. *Journal of Complexity* 14, 257–299. ISSN 0885-064X.
- SURESH K. LODHA & RON GOLDMAN (1997). A unified approach to evaluation algorithms for multivariate polynomials. *Mathematics of Computation* **66**(220), 1521-1559. ISSN 0025-5718. URL http://www.ams.org/mcom/1997-66-220/S0025-5718-97-00862-4.
- A. M. Odlyzko & Arnold Schönhage (1988). Fast algorithms for multiple evaluations of the Riemann zeta function. *Transactions of the American Mathematical Society* 309(2), 797–809.
- V. Ya. Pan (1966). О способах вычисления значении многочленов. Успехи Математических Наук **21**(1(127)), 103–134. V. Ya. Pan, Methods of computing values of polynomials, *Russian Mathematical Surveys* **21** (1966), 105–136.
- Arnold Schönhage (1977). Schnelle Multiplikation von Polynomen über Körpern der Charakteristik 2. Acta Informatica 7, 395–398.
- Arnold Schönhage & Volker Strassen (1971). Schnelle Multiplikation großer Zahlen. Computing 7, 281–292.
- Volker Strassen (1969). Gaussian Elimination is not Optimal. Numerische Mathematik 13, 354-356.
- MARTIN ZIEGLER (2003a). Fast Relative Approximation of Potential Fields. In Algorithms and Data Structures, Frank Dehne, Jörg-Rüdiger Sack & Michiel Smid, editors, number 2748 in Lecture Notes in Computer Science, 140-149. Springer-Verlag, Berlin, Heidelberg. ISBN 3-540-40545-3. ISSN 0302-9743. URL http://www.springerlink.com/openurl.asp?genre=article&issn=0302-9743&volume=2748&spage=140.
- MARTIN ZIEGLER (2003b). Quasi-optimal Arithmetic for Quaternion Polynomials. In Proceedings of 14th International Symposium on Algorithms and Computation ISAAC 2003, Kyoto, Japan, Toshihide Ibariki, Naoki Katoh & Hirotaka Ono, editors, number 2906 in Lecture Notes in Computer Science, 705-715. Springer-Verlag, Berlin, Heidelberg. ISBN 3-540-20695-7. ISSN 0302-9743. URL http://www.springerlink.com/openurl.asp?genre=article&issn=0302-9743&volume=2906&spage=705.