

Isolated Horizons and their Applications

Abhay Ashtekar¹, Christopher Beetle¹, Olaf Dreyer¹,
Stephen Fairhurst¹, Badri Krishnan¹, Jerzy Lewandowski^{2,1} and Jacek Wisniewski¹

1. Physics Department, 104 Davey, Penn State, University Park, PA 16802, USA

2. Institute of Theoretical Physics, University of Warsaw, ul. Hoza 69, 00-681 Warsaw, Poland

A set of boundary conditions defining isolated horizons (possibly with distortion and rotation) is introduced. Space-times containing a black hole, itself in equilibrium but possibly surrounded by radiation, satisfy these conditions. In spite of this generality, the conditions have rich consequences. They lead to a framework, somewhat analogous to null infinity, for extracting physical information, but now in the strong field regions. The framework also generalizes the zeroth and first laws of black hole mechanics to more realistic situations and sheds new light on the 'origin' of the first law. Finally, it provides a point of departure for black hole entropy calculations in non-perturbative quantum gravity.

A great deal of analytical work on black holes in general relativity centers around event horizons in globally stationary space-times (see, e.g., [1,2]). While it is a natural starting point, this idealization seems overly restrictive from a physical point of view. In a realistic gravitational collapse, or a black hole merger, the final black hole is expected to rapidly reach equilibrium. However, the exterior space-time region will not be stationary. Indeed, a primary goal of many numerical simulations is to study radiation emitted in the process. Similarly, since event horizons can only be determined retroactively after knowing the entire space-time evolution, they are not directly useful in many situations. For example, when one speaks of black holes in centers of galaxies, one does not refer to event horizons. The idealization seems unsuitable also for black hole mechanics and statistical mechanical calculations of entropy. Firstly, in ordinary equilibrium statistical mechanics, one only assumes that the system under consideration is stationary, not the whole universe. Secondly, from quantum field theory in curved space-times, thermodynamic considerations are known to apply also to cosmological horizons [3]. Thus, it seems desirable to replace event horizons by a quasi-local notion and develop a detailed framework tailored to diverse applications, from numerical relativity to quantum gravity, without the assumption of global stationarity. The purpose of this letter is to present such a framework.

Specifically, we will provide a set of quasi-local boundary conditions which define an isolated horizon representing, for example, the last stages of a collapse or a merger, and focus on space-time regions admitting such horizons as an inner boundary. Although the boundary conditions are motivated purely by geometric considerations, they lead to a well-defined action principle and Hamiltonian framework. This, in turn, leads to a definition of the horizon mass M and angular momentum J . These quantities refer only to structures intrinsically available on Σ , without any reference to infinity, and yet lead to a generalization of the familiar laws of black hole mechanics. We will also introduce invariantly

defined coordinates near Σ and a Bondi-type expansion of the metric.

For brevity, in the main discussion we will restrict ourselves to the Einstein-Maxwell theory in four space-time dimensions. Throughout, $\equiv$ will stand for equality restricted to Σ ; an arrow under an index will denote pull-back of that index to Σ ; V^a will be a generic vector field tangential to Σ and ∇^a any of its extensions to space-time. The electromagnetic potential and fields will be denoted by bold-faced letters. All fields are assumed to be smooth, and bundles, trivial. For details, generalizations and subtleties, see [4,7].

Definition: A sub-manifold Σ of a space-time $(M; g_{ab})$ is said to be an isolated horizon if:

- i) It is topologically $S^2 \times \mathbb{R}$, null, with zero shear and expansion. This condition implies, in particular, that the space-time Σ induces a unique derivative operator D on Σ via $D_a V^b \equiv \nabla_a V^b$.
- ii) $(L \cdot D_a - D_a L \cdot)V^b \equiv 0$ and $L \cdot A_a \equiv 0$ for some null normal L to Σ ; and,
- iii) Field equations hold at Σ .

These conditions are local to Σ . The first two imply that the intrinsic metric and connection on Σ are 'time-independent' and spell out the precise sense in which Σ is 'isolated'. Every Killing horizon which is topologically $S^2 \times \mathbb{R}$ is an isolated horizon. However, in general, space-times with isolated horizons need not admit any Killing field even in a neighborhood of Σ . The local existence of such space-times was shown in [8]. A global example is provided by Robinson-Trautman space-times which admit an isolated horizon but have radiation in every neighborhood of it [9]. Finally, on a general Σ , the null normal L of ii) plays a role analogous to that of the Killing field on a Killing horizon. Generically, L satisfying ii) is unique up to a constant rescaling $L \rightarrow cL$. (In particular, this is true of the Kerr family.) We will denote by $[L]$ the equivalence class of null normals satisfying ii). One can not hope to eliminate this constant rescaling freedom because, without reference to infinity, it exists already on Killing horizons.

Geometry of isolated horizons: Although the boundary conditions are rather weak, they have surprisingly rich consequences. We now summarize the most important ones.

1. Intrinsic geometry: γ is a symmetry of the degenerate, intrinsic metric $q_{ab} := g_{ab}|_{\gamma}$; $L \cdot q_{ab} \neq 0$. γ is naturally equipped with a 2-form ω_{ab} , the pullback to γ of the volume 2-form on the 2-sphere of integral curves of χ , satisfying $\omega_{ab} \chi^b \neq 0$ and $L \cdot \omega_{ab} \neq 0$. The area of any cross-section S is given by $\int_S \omega$ and is the same for all cross-sections. We will denote it by A .

2. Connection coefficients: γ is geodesic and free of divergence, shear and twist. Hence there exists a 1-form λ on γ such that $r_a \chi^b = \lambda_a \chi^b$. The surface gravity κ defined by χ is given by $\kappa = \lambda_a \chi^a$. The boundary conditions imply κ is constant on γ [6]. Thus, the zeroth law holds. Similarly, the electromagnetic potential $\phi = A_a \chi^a$ is constant on γ . Note, however, that other connection components or the scalar curvature of the intrinsic metric q_{ab} need not be constant; the horizon may be distorted arbitrarily.

3. Weyl curvature: Let us pick an ℓ in $[\gamma]$ and construct a null tetrad $\{\ell, n, m, \bar{m}\}$ on γ . Thus, $m, \bar{m}$ are tangential to γ and n is transverse. Then, the Weyl components $\psi_0 = C_{abcd} \ell^a m^b \ell^c m^d$ and $\psi_1 = C_{abcd} \ell^a m^b \chi^c n^d$ vanish, implying that there is no flux of gravitational radiation across γ and the Weyl tensor at γ is of Petrov type II [6]. Hence $\psi_2 := C_{abcd} \ell^a m^b \bar{m}^c n^d$ is gauge invariant. Its imaginary part is given by: $dI = 2\text{Im} \psi_2$. We will see that it contains the angular momentum information. While ψ_2 is time independent on the horizon, in general, $\psi_3 = C_{abcd} \chi^a m^b \bar{m}^c n^d$ and $\psi_4 = C_{abcd} n^a \bar{m}^b n^c \bar{m}^d$ are not [7].

4. A natural foliation: Let us consider the non-extremal case when κ is non-zero. Then, γ admits a natural foliation, thereby providing a natural horizon rest frame e' [7]. The 2-sphere cross-sections of the horizon defined by this foliation are completely analogous to the 'good cuts' that null infinity admits in absence of Bondi news. Therefore, we will refer to them as good cuts of the horizon. On a non-rotating horizon, since $\text{Im} \psi_2$ vanishes, so does dI . Hence, there exists a function ϕ on γ with $\chi \cdot \phi = dI$. Since $L \cdot \phi = 0$, ϕ is constant on γ , the constant surfaces foliate γ . In the rotating case, the argument is more involved but the foliation is again determined invariantly by the geometrical structure of γ . This foliation turns out to be very useful (see below).

5. Symmetries of γ : In view of our main Definition, it is natural to define the symmetry group G of a given isolated horizon to be the sub-group of $D(\gamma)$ which preserves $[\chi]$; $q_{ab}; D; A_a$. Since $q_{ab}; D; A_a$ can vary from one isolated horizon to another, G is not canonical. For simplicity, let us again restrict ourselves to the non-extremal case $\kappa \neq 0$. Then, isolated horizons fall into three universality classes [7]: I. $\dim G = 4$: in this case,

q_{ab} is spherically symmetric, good cuts are invariant under the natural $SO(3)$ action and G is the direct product of $SO(3)$ with translations along χ ; II. $\dim G = 2$: in this case, q_{ab} is axisymmetric, the general infinitesimal symmetry ξ^a has the form $\xi^a = c^a + \chi^a$, where c^a are constants on γ and χ^a is a rotational vector field tangential to good cuts; and, III. $\dim G = 1$: in this case, the infinitesimal horizon symmetry has the form $\xi^a = c^a$. In case I, as one might expect, $\text{Im} \psi_2 \neq 0$ and the horizon is non-rotating. Case III corresponds to general distortion.

Extracting physics: The isolated horizon framework can be used to extract invariant physical information in the strong field region near black holes, formed by gravitational collapse or merger of compact objects. At a sufficiently late time, the space-time would contain an (approximate) isolated horizon. In the most interesting case, γ would be of universality class II above. One can ask for its angular momentum and mass. Recall that, for asymptotically flat space-times without internal boundaries, one obtains expressions of the ADM mass M_1 and angular momentum J_1 using a Hamiltonian framework. This strategy can be extended to the present case (see below). When constraints are satisfied, the total Hamiltonian is now a sum of two surface terms, one at infinity and the other at γ . The terms at infinity again yield M_1 and J_1 . General arguments lead one to interpret the surface terms at γ as the horizon mass M and angular momentum J . We have [7]:

$$J = \frac{1}{8\pi G} \int_S (\lambda_a + 2GA_a) \chi^a$$

$$= \frac{1}{4\pi G} \int_S f (\text{Im} \psi_2 + 2G \text{Im} \psi_1) \quad (1)$$

where S is any 2-sphere cross-section of γ , f is related to χ by $D_a f = \chi_b \chi'^b$ and $\text{Im} \psi_1 = (i=2) F_{ab} m^a \bar{m}^b$ is a Newman-Penrose component of the Maxwell field. In a vacuum, axisymmetric space-time, $J = J_1$. However, in general, the two differ by the angular momentum in the gravitational radiation and the Maxwell field in the region between γ and infinity. Even in presence of such radiation, the horizon mass is given by [7]

$$M = \frac{1}{2\pi G R} (R^2 + GQ^2)^2 + 4G^2 J^2 \quad (2)$$

where R is the horizon radius, given by $a = 4R^2$, and $Q = \frac{1}{4\pi G} \int_S F$ is the horizon charge. Somewhat surprisingly, M has the same dependence on area, angular momentum and charge as in the Kerr-Newman family (provided J is defined via (1)). However, this is a result of the calculation, not an assumption. In a Kerr-Newman space-time, we have $M_1 = M$ for all values of Q . (Thus, if $Q \neq 0$, M does not agree with any of the known quasi-local expressions of mass.) However, in general M is different from M_1 . Under certain physically reasonable assumptions on the behavior of fields

near future time-like infinity i^+ , one can show that the difference is the energy radiated across I^+ by gravitational and electromagnetic waves.

If $\gamma \neq 0$, irrespective of the universality class, one can introduce (essentially) invariant coordinates and tetrads in a neighborhood of Σ . Fix an ℓ in $[\gamma]$. Let $v; \bar{v}$ be coordinates on Σ such that $L \cdot v \neq 0$ and good cuts are given by $v = \text{const}$. Let n^a be the unique future-directed null vector field which is orthogonal to the good cuts and normalized so that $\ell \cdot n = 1$. Consider past null geodesics emanating from the good cuts, with n^a as their tangent at Σ . Finally, define r via $L \cdot n r = 1$ and $r = r^0$ on Σ , and Lie drag $v; \bar{v}$ along n^a . We now have a natural set of coordinates, $(r; v; \bar{v})$, the only arbitrariness being in the initial choice of $(v; \bar{v})$ and adding constants to $r; v$. Next, let us parallel transport $\ell; m; \bar{m}$ along n to obtain this neighborhood. The tetrad is unique up to local $m; \bar{m}$ rotations at Σ . Now, assuming the vacuum equations hold in this neighborhood, one can give a Bondi-type expansion for the metric components in powers of $(r-r^0)$ to any desired order. For example, retaining terms to second order, we have [7]:

$$\begin{aligned} g_{ab} = & 2m^0_{(a} \bar{m}^0_{b)} + 2r_{; (a} v_{; b)} (r - r^0)^4 + 2m^0_{(a} \bar{m}^0_{b)} + 2v_{; (a} (2!_{b)} (v; \bar{v})_{b)} \\ & + (1=2) (r - r^0)^2 4((v)^2 + (v^0)^2) m^0_{(a} \bar{m}^0_{b)} \\ & + (4(v^0 v^0 - 2(v^0 v^0)) m^0_{(a} \bar{m}^0_{b)} + (4(v^0 v^0 - 2(v^0 v^0)) \bar{m}^0_{(a} m^0_{b)} \\ & + 4(v^0 v^0 + v^0 v^0 - v^0 v^0) v_{; (a} \bar{m}^0_{b)} \\ & + 4(v^0 v^0 + v^0 v^0 - v^0 v^0) v_{; (a} m^0_{b)} \\ & + (2(v^0 v^0 - 2(v^0 v^0 - 2(v^0 v^0)) v_{; (a} v_{; b)} + O((r - r^0)^3); \end{aligned}$$

where quantities with the a superscript 0 are evaluated on Σ , and the Newman-Penrose spin coefficients are defined as: $\rho = m^a \bar{m}^b r_{; a} n_{; b}$, $\sigma = \bar{m}^a m^b r_{; a} n_{; b}$ and $\tau = v^a \bar{m}^b r_{; a} n_{; b}$. Using the boundary conditions and field equations, at the horizon these spin coefficients as well as the Weyl components can be expressed in terms of the dyad $m^0; \bar{m}^0$ defining the intrinsic horizon geometry, 1-form ℓ_a and the value of ρ_a on any one good cut [7].

The null surfaces $v = \text{const}$: are invariantly defined. Therefore (modulo the small freedom mentioned above) the tetrad components of the Weyl tensors on these surfaces are gauge invariant. This property will be useful in physically interpreting the outcomes of numerical simulations of mergers of compact objects. For example, it will enable a gauge invariant comparison between the radiation fields $j_4 j$ created in two simulations, say with somewhat different initial conditions. Finally, one can give a systematic procedure to extend any infinitesimal symmetry $\ell^a \in \mathfrak{c}^a + \ell^a$ on Σ to a potential Killing field ℓ^a in a neighborhood [7]. If the space-time does admit a Killing field ℓ^a which coincides with ℓ^a on Σ , then ℓ^a must equal ℓ^a in the neighborhood. Again, since they are defined invariantly, the vector fields ℓ^a can be useful

to extract physical information coded in the strong field geometry.

Finally, note that all this structure — particularly the definitions of M and J — is defined intrinsically, using local geometry of the physical space-time under consideration. To extract physical information, one does not have to embed this space-time in a Kerr solution which, in the light of the no-hair theorems, presumably approximates the physical, near horizon geometry at late times. In practice this is a significant advantage because the embedding problem can be very difficult: typically, one knows little about the form of the desired Kerr metric in the coordinate system in which the numerical simulation is carried out. More importantly, a priori, one does not know which Kerr parameters to use in the embedding, nor does one have a quantitative control on precisely how the physical near-horizon geometry is to approach Kerr.

Isolated Horizon Mechanics: We already saw that the zeroth law holds on all isolated horizons. Let us consider the first law: $M = (8G) a + J + Q$. In the stationary context the law is somewhat hybrid in that M and J are defined at infinity, a at the horizon and Q and ℓ are evaluated at the horizon but refer to the normalization of the Killing field carried out at infinity. In the non-stationary context now under consideration, there are two additional problems: due to the presence of radiation, M_1 and J_1 have little to do with the horizon mass and since we no longer have a global Killing field, there is an ambiguity in the normalization of ℓ and ℓ .

As in [10], our strategy is to arrive at the first law through a Hamiltonian framework, but now adapted to the isolated horizon boundary conditions. For brevity, we will again focus on the physically most interesting universality class II. Let us fix on the (abstract) isolated horizon boundary a rotational vector field ℓ^a . Consider the space of asymptotically flat solutions to the Einstein-Maxwell equations for which ℓ as an isolated horizon inner-boundary with symmetry ℓ^a . $\mathcal{H}$ will be our covariant phase-space [6,7]. Denote by ℓ^a any extension of ℓ^a which is an asymptotic rotational Killing field at spatial infinity. Then, one can show that the vector field ℓ on $\mathcal{H}$ is defined by the Lie derivative of basic fields along ℓ^a is a phase space symmetry, i.e., Lie drags the symplectic structure. Its generator is given by [7]

$$H_\ell = J_1 - J$$

where J is given by (1). Hence, it is natural to interpret (1) as the horizon angular momentum.

To define the horizon energy, one needs to select a 'time translation'. On $\mathcal{H}$, it should coincide with a horizon symmetry $\ell^a \in \mathfrak{c}^a + \ell^a$. While $\mathfrak{c}; \ell$ are constants on Σ , in the phase space we must allow them to vary from one solution to another. (In the numerical relativity language, we must allow ℓ^a — or, the lapse and shift at Σ — to be live.) For, unlike at infinity, the 4-geometries under

consideration do not approach a fixed 4-geometry at ∞ , whence it is not a priori obvious how to pick the same time-translation for all geometries in the phase space. Let t^a be any extension of t^a to the whole space-time which approaches a fixed time translation at infinity. We can ask if the corresponding ξ_t is a phase space symmetry. The answer is rather surprising: yes, if and only if there exists a function E^t on the phase space, involving only the horizon fields, such that the first law,

$$E^t = \frac{(t)}{8\pi G} \int_a + \int_t J + \int_{(t)} Q; \quad (3)$$

holds [6,7]. Thus, not only does the isolated horizon framework enable one to extend the first law beyond the stationary context, but it also brings out its deeper role: it is a necessary and sufficient condition for a consistent Hamiltonian evolution.

However, there are many choices of t^a on the horizon for which this condition can be met, each with a corresponding time-evolution, horizon energy function and first law. Can we make a canonical choice of t^a ? In the Einstein-Maxwell theory, the answer is in the affirmative. The requirement that the (live) vector field t^a coincide, on each Kerr-Newman solution, with that stationary Killing field which is unit at infinity uniquely fixes t^a on the isolated horizon of every space-time in the phase space. With this canonical choice, say $t = t_0$, in Einstein-Maxwell theory we can define the horizon mass to be

$$M = E^{t_0}.$$

Then, M is given by (2).

We will conclude with three remarks.

1. We focused our discussion on the physically most interesting universality class II. Class I was treated in detail in [4,5] and is a special case of non-rotating, class III horizons discussed in [6]. All these cases have been analyzed in detail. However, the current understanding of class III with $\int_{\mathcal{H}} \epsilon \neq 0$ is rather sketchy.
2. The framework that led us to the zeroth and first laws can be easily extended to other space-time dimensions. The 2+1-dimensional case has already been analyzed in detail [11] and has some special interesting features in the context of a negative cosmological constant. In the non-rotating, class III case, dilaton and Yang-Mills fields have also been incorporated [4,6]. In the Yang-Mills case, although the zeroth and first laws can be proved, the analog of the mass formula (2) is not known because one does not have as much control on the space of all stationary solutions. Nonetheless, the framework has been used to derive new relations between masses of static black holes with hair and their solitonic analogs in Einstein-Yang-Mills theory [5,6]. More importantly, as is well-known, the standard no-hair theorems fail in this case and the framework has been used to conjecture new no-hair theorems tailored to isolated horizons rather than infinity [5].

3. In the non-rotating case, the framework has been used to carry out a systematic and detailed entropy calculation using non-perturbative quantum gravity [12]. The analysis encompasses all black holes without any restriction of near-extremality made in string theory calculations. Furthermore, it also naturally incorporates the cosmological horizons to which thermodynamic considerations are known to apply [3]. Recently, sub-leading corrections to entropy have also been calculated [13]. However, the non-perturbative quantization scheme faces a quantization ambiguity — analogous to the θ -ambiguity in QCD — which permeates all these calculations. Its role is not fully understood. Carlip [14] and others have suggested the use of horizon symmetries in entropy calculations and this approach could shed light on the quantization ambiguity and relate the analysis of [12] to conformal field theories. Conversely, the isolated horizon framework may offer a more systematic avenue for implementing Carlip's ideas. Finally, since rotation has now been incorporated in the classical theory [7], one can hope to extend the entropy calculation to this case.

Acknowledgements: We would like to thank A. Corichi, S. Hayward, J. Pullin, D. Sudarsky and R. Wald for discussions. This work was supported in part by the NSF grants PHY 95-14240, INT 97-22514, the Polish CSR grant 2 P 03B 060 17, the Albert Einstein Institute and the Eberly research funds of Penn State.

-
- [1] M. Heusler, Black Hole Uniqueness Theorems (Cambridge University Press: Cambridge) (1996).
 - [2] R. M. Wald, Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics (University of Chicago Press) (1994).
 - [3] G. Gibbons and S.W. Hawking, Phys. Rev. D 15 2738 (1977).
 - [4] A. Ashtekar, C. Beetle and S. Fairhurst, Class. Quantum Grav. 17 253 (2000); A. Ashtekar and A. Corichi, Class. Quantum Grav. 17 1317 (2000).
 - [5] A. Corichi, U. Nucamendi and D. Sudarsky, gr-qc 0002078.
 - [6] A. Ashtekar, S. Fairhurst and B. Krishnan, gr-qc 0005083
 - [7] A. Ashtekar, C. Beetle and J. Lewandowski, Mechanics of rotating isolated horizons (in preparation); Geometry of general isolated horizons (in preparation).
 - [8] J. Lewandowski, Class. Quantum Grav. 17 L53-L59 (2000).
 - [9] P.T. Chruściel, Proc. Roy. Soc. London A 436 299 (1992).
 - [10] D. Brown and J.W. York, Phys. Rev. D 47 1407 (1993).
 - [11] A. Ashtekar, O. Dreyer and J. Wisniewski, (in preparation).
 - [12] A. Ashtekar, J. Baez, A. Corichi and K. Kasnov, Phys. Rev. Lett. 80 904-907 (1998); A. Ashtekar, J. Baez, and K. Kasnov, gr-qc 0005126.
 - [13] R. Kaul and P. Majumdar, Phys. Rev. Lett. (in press).
 - [14] S. Carlip, Phys. Rev. Lett. 82 2828-2831 (1999).