

Comment on negative squares of the Weyl tensor

Kayll Lake [*]

Department of Physics and Department of Mathematics and Statistics,
Queen's University, Kingston, Ontario, Canada, K7L 3N6

(Dated: April 11, 2019)

Recently Schmidt has given an example which shows that the square of the Weyl tensor can be negative. This phenomenon occurs in the Kerr metric and it is demonstrated here in a parameter independent way.

PACS numbers: 04.20.Cv, 04.20.Jb, 04.40.Dg

Recently Schmidt [1] has pointed out that the metric

$$ds^2 = -dt^2 + 2yzdtdx + dx^2 + dy^2 + dz^2 \quad (1)$$

has $W < 0$ in an open neighborhood of $t = x = y = z = 0$ where

$$W \equiv C_{ijkl}C^{ijkl} \quad (2)$$

and C_{ijkl} is the Weyl tensor. There is no need for a restriction on t since, for example, the metric

$$ds^2 = -dt^2 + 2(\alpha xy + \beta xz + \gamma yz + \delta(y^2 + z^2))dtdx + dx^2 + dy^2 + dz^2, \quad (3)$$

where α, β, γ and δ are constants, is not flat but regular at $x = y = z = 0$ and gives

$$W = -(\alpha^2 + \beta^2 + 4\gamma^2) \quad (4)$$

there. The purpose of this comment is not to elaborate on the metric (3), the physical meaning of which is unclear, but rather to emphasize the fact that $W < 0$ in an open region in the Kerr metric.

In terms of the familiar Boyer-Lindquist coordinates we find

$$\mathcal{W} \equiv \frac{W a^6}{48 m^2} = \frac{(x - \cos(\theta))(x + \cos(\theta)) \left(x^2 + 4x \cos(\theta) + (\cos(\theta))^2 \right) \left(x^2 - 4x \cos(\theta) + (\cos(\theta))^2 \right)}{\left(x^2 + (\cos(\theta))^2 \right)^6} \quad (5)$$

where $x \equiv r/a, a \neq 0$. Along $x = 0$ then

$$\mathcal{W} = -\frac{1}{\cos(\theta)^6} \quad (6)$$

and so the Kerr metric is singular at $x = 0$ only in the equatorial plane $\theta = \pi/2$, as is well known. Along the rotation axis ($\theta = 0$ or π) \mathcal{W} oscillates between negative and positive values in the range $0 < x < 2 + \sqrt{3}$ while crossing the axis $\mathcal{W} = 0$ at $x = 2 - \sqrt{3}, 1$ and $2 + \sqrt{3}$. This is shown in Figure 1. A three dimensional picture, showing \mathcal{W} for a range in both x and θ , is shown in Figure 2. These figures are independent of a and m . For related, but parameter dependent discussions, see [2] and [3].

Acknowledgments

This work was supported by a grant from the Natural Sciences and Engineering Research Council of Canada. Portions of this work were made possible by use of *GRTensorII* [4].

[*] Electronic Address: lake@astro.queensu.ca

[1] H. -J. Schmidt, Gen. Relat. Grav (2003) (gr-qc/0302078)

[2] C. Cherubini, D. Bini, S. Capozziello, R. Ruffini, Int. J. Mod. Phys. **D 11** (2002) 827.

[3] R. C. Henry, Ap. J. **535**, 350 (2000).

[4] This is a package which runs within Maple. It is entirely distinct from packages distributed with Maple and must be obtained independently. The GRTensorII software and documentation is distributed freely on the World-Wide-Web from the address <http://grtensor.org>




FIG. 1: Plot of $\mathcal{W}(\equiv \frac{C_{ijkl}C^{ijkl}a^6}{48m^2})$ along the rotation axis ($\theta = 0$ or π). $\mathcal{W} > 0$ for $x(\equiv \frac{r}{a}) > 2 + \sqrt{3}$ and asymptotically approaches zero as $x \rightarrow \infty$.




FIG. 2: Plot of $\mathcal{W}(\equiv \frac{C_{ijkl}C^{ijkl}a^6}{48m^2})$ showing a variation in both $x \equiv \frac{r}{a}$ and θ .