

Refined algebraic quantisation with the triangular subgroup of $SL(2, \mathbb{R})$

Jorma Louko^{*} and Alberto Molgado[†]

*School of Mathematical Sciences, University of Nottingham,
Nottingham NG7 2RD, UK*

(April 2004)

$\langle \text{arXiv:gr-qc/0404100} \rangle$

Abstract

We investigate refined algebraic quantisation with group averaging in a constrained Hamiltonian system whose gauge group is the connected component of the lower triangular subgroup of $SL(2, \mathbb{R})$. The unreduced phase space is $T^*\mathbb{R}^{p+q}$ with $p \geq 1$ and $q \geq 1$, and the system has a distinguished classical $\mathfrak{o}(p, q)$ observable algebra. Group averaging with the geometric average of the right and left invariant measures, invariant under the group inverse, yields a Hilbert space that carries a maximally degenerate principal unitary series representation of $O(p, q)$. The representation is nontrivial iff $(p, q) \neq (1, 1)$, which is also the condition for the classical reduced phase space to contain a symplectic manifold. We present a detailed comparison to an algebraic quantisation that imposes the constraints in the sense $\hat{H}_a \Psi = 0$ and requires self-adjointness of the $\mathfrak{o}(p, q)$ observables. Under certain technical assumptions that parallel those of the group averaging theory, this algebraic quantisation gives no quantum theory when $(p, q) = (1, 2)$ or $(2, 1)$, or when $p \geq 2$, $q \geq 2$ and $p + q \equiv 1 \pmod{2}$.

^{*}jorma.louko@nottingham.ac.uk

[†]pmxam@nottingham.ac.uk

1 Introduction

In quantisation of constrained systems, an elegant idea for constructing the physical Hilbert space is to average states in an auxiliary Hilbert space \mathcal{H}_{aux} over a suitable action of the gauge group [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14]. For a noncompact gauge group the averaging need not converge on all of \mathcal{H}_{aux} , but when the averaging is formulated within refined algebraic quantisation [4, 8, 12], convergence on a suitable linear subspace will suffice, and such convergence has been found to occur in concrete examples [9, 10, 14]. Results on the equivalence of refined algebraic quantisation with other methods [13, 15] further show that group averaging provides considerable control over the quantisation.

In this paper we study group averaging in a system with a nonunimodular gauge group. The interest of this situation arises from the rather different senses in which group averaging satisfies the Dirac constraints for unimodular and nonunimodular groups [8]. To guarantee that the would-be inner product provided by group averaging is real, the averaging measure needs to be invariant under the group inverse. For a unimodular group, the left and right invariant Haar measure has this property. For a nonunimodular group, the left and right invariant Haar measures do not coincide, and neither is invariant under the group inverse, but their geometric average d_0g is. Suppose \hat{H}_a are the constraint operators that generate the unitary gauge group action on the auxiliary Hilbert space, with the commutators

$$[\hat{H}_a, \hat{H}_b] = iC^c_{ab}\hat{H}_c \quad , \quad (1.1)$$

where C^c_{ab} are the structure constants of the Lie algebra. It was shown in [8] that group averaging with d_0g gives physical states Ψ that satisfy

$$\hat{H}_a\Psi = -\frac{i}{2}C^b_{ab}\Psi \quad , \quad (1.2)$$

which agrees with the naïve Dirac quantum constraints, $\hat{H}_a\Psi = 0$, if and only if the group is unimodular. The term on the right-hand side of (1.2) leads to no known inconsistencies: For systems amenable to geometric quantisation in both reduced and unreduced phase space, (1.2) is in fact the form of Dirac constraints equivalent to reduced phase space quantisation [16, 17, 18].¹ Related observations for first class constrained

¹A comment on the notation is in order. In [8], the physical states $\langle\phi|$ live in the dual of the test space and satisfy $\langle\phi|U(g) = \langle\phi|[\Delta(g)]^{1/2}$, where $U(g)$ is a unitary representation of the gauge group generated by the constraints and acts on the bra vector $\langle\phi|$ in the dual sense from the right: see the discussion after equation (2.6) in [8]. To compare with a picture in which $U(g)$ acts on a ket vector $|\phi\rangle$ from the left, we map $\langle\phi|$ to $|\phi\rangle$ by the antilinear isomorphism obtained by complex conjugating the group averaging sesquilinear form. As $U(g)$ is unitary in the auxiliary Hilbert space, this map sends the right action of $U(g)$ to the left action of $U(g^{-1})$, which implies $U(g^{-1})|\phi\rangle = [\Delta(g)]^{1/2}|\phi\rangle$, or $U(g)|\phi\rangle = [\Delta(g)]^{-1/2}|\phi\rangle$. Equation (4.2) in [8] is therefore to be understood in this sense. Consequently, equation (B6) in [8] is to be understood as $(\mathcal{J}_a)|\phi\rangle = -(i/2)\text{tr}(\text{ad}_a)|\phi\rangle$, which agrees with equation (6.4) in [18] and with our (1.2). We thank Nico Giulini and Don Marolf for correspondence on this point.

systems with one constraint quadratic in the momenta and several constraints linear in the momenta were made in [19].

We shall consider a system obtained by replacing a unimodular gauge group G_u by its nonunimodular subgroup G . The effect of this replacement on the constraints (1.2) is not just that some of the constraint equations are dropped: In the constraints that remain, new terms appear on the right-hand side. An observable algebra that commutes strongly with the old constraints is still represented on the solution space to the new constraints, but the new representation need not be isomorphic to the old representation. In group averaging, where no observables need be explicitly constructed, these changes are encoded in the integration measures on G_u and G . The integrals over G_u and G may also have differing convergence properties, and it may hence be necessary to choose the test spaces differently even when \mathcal{H}_{aux} is unchanged and the representation of G is obtained from that of G_u by restriction. Our system will indeed exemplify all these phenomena.

Our unimodular gauge system [14] has gauge group $G_u \simeq \text{SL}(2, \mathbb{R})$, unreduced phase space $T^*\mathbb{R}^{p+q} \simeq \mathbb{R}^{2(p+q)}$ with $p \geq 1$ and $q \geq 1$ and a distinguished classical $\mathfrak{o}(p, q)$ observable algebra. The reduced phase space contains a symplectic manifold if and only if $p \geq 2$ and $q \geq 2$: This manifold has dimension $2(p + q - 3)$ and is separated by the $\mathfrak{o}(p, q)$ observables. The system was quantised with group averaging in [10, 14], recovering a quantum theory with a nontrivial representation of the quantum $\mathfrak{o}(p, q)$ observables when $p \geq 2$, $q \geq 2$ and $p + q \equiv 0 \pmod{2}$. The quadratic $\mathfrak{o}(p, q)$ Casimir was found to take the value $-\frac{1}{4}(p+q)(p+q-4)$. Quantisations of this system for special values of p and q by a variety of other methods can be found in [10, 14, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31].

Our nonunimodular gauge group $G \subset G_u$ is the connected component of the lower triangular subgroup of $\text{SL}(2, \mathbb{R})$. G is two-dimensional and nonabelian, and hence isomorphic to every two-dimensional connected nonabelian group. The reduced phase space contains a symplectic manifold if and only if $(p, q) \neq (1, 1)$: This manifold is symplectomorphic to $T^*(S^{p-1} \times S^{q-1})$, and it is separated by the $\mathfrak{o}(p, q)$ observables up to a set of measure zero. Quantisation with group averaging gives a quantum theory with a nontrivial representation of the quantum $\mathfrak{o}(p, q)$ observables for all $(p, q) \neq (1, 1)$. This representation is the end-point of the maximally degenerate principal unitary series [32], and the quadratic Casimir takes the value $-\frac{1}{4}(p + q - 2)^2$.

For comparison, we quantise the system also with the constraints $\hat{H}_a \Psi = 0$, adopting the algebraic quantisation framework of [33, 34] and requiring the $\mathfrak{o}(p, q)$ observables to become self-adjoint operators. For $(p, q) = (1, 3)$, $(3, 1)$ and $(2, 2)$, the group averaging theory and the algebraic quantisation theory are qualitatively fairly similar, with minor differences in the spectra of certain operators. Our algebraic quantisation results for general (p, q) remain incomplete, but we show that under certain technical assumptions that parallel those of the group averaging theory, the algebraic quantisation gives no quantum theory for $(p, q) = (1, 2)$ and $(2, 1)$, or for $p \geq 2$, $q \geq 2$ and $p + q \equiv 1 \pmod{2}$.

The rest of the paper is as follows. Section 2 introduces and analyses the classical

system. Sections 3 and 4 discuss algebraic quantisation, with respectively the constraints (1.2) and $\hat{H}_a \Psi = 0$. Refined algebraic quantisation with group averaging is carried out in section 5 for $(p, q) \neq (1, 1)$ and in section 6 for $p = q = 1$. Section 7 presents a summary and concluding remarks.

Appendix A collects some basic properties of the group G . The separation properties of the $\mathfrak{o}(p, q)$ observables on the reduced phase space are proved in appendix B. Certain technical results for refined algebraic quantisation are proved in appendix C.

2 Classical system

In this section we introduce a classical constrained system with the unreduced phase space $T^*\mathbb{R}^{p+q}$, where $p \geq 1$ and $q \geq 1$. The system is obtained from the $\text{SL}(2, \mathbb{R})$ system of [14] by dropping the constraint H_1 therein and relabelling one of the canonical pairs by $(\mathbf{v}, \boldsymbol{\pi}) = (-\boldsymbol{\varpi}, \mathbf{w})$. The gauge transformations and the $\mathfrak{o}(p, q)$ observables are therefore obtained directly from [14]. We shall however see that the structure of the reduced phase space differs markedly from that in [14].

2.1 The system

The system is defined by the action

$$S = \int dt (\mathbf{p} \cdot \dot{\mathbf{u}} + \boldsymbol{\varpi} \cdot \dot{\mathbf{w}} - MH - \lambda D) , \quad (2.1)$$

where \mathbf{u} and \mathbf{p} are real vectors of dimension $p \geq 1$, \mathbf{w} and $\boldsymbol{\varpi}$ are real vectors of dimension $q \geq 1$, and the overdot denotes differentiation with respect to t . The symplectic structure is

$$\Omega = \sum_{i=1}^p dp_i \wedge du_i + \sum_{j=1}^q d\varpi_j \wedge dw_j , \quad (2.2)$$

and the phase space is $\Gamma := T^*\mathbb{R}^{p+q} \simeq \mathbb{R}^{2(p+q)}$. M and λ are Lagrange multipliers associated with the constraints

$$\begin{aligned} H &:= \frac{1}{2}(\mathbf{w}^2 - \mathbf{u}^2) , \\ D &:= \mathbf{u} \cdot \mathbf{p} + \mathbf{w} \cdot \boldsymbol{\varpi} . \end{aligned} \quad (2.3)$$

The Poisson algebra of the constraints is

$$\{H, D\} = 2H , \quad (2.4)$$

and the system is a first class constrained system [35, 36]. The finite gauge transformations on Γ generated by the constraints are

$$\begin{pmatrix} \mathbf{u} \\ \mathbf{p} \end{pmatrix} \mapsto g \begin{pmatrix} \mathbf{u} \\ \mathbf{p} \end{pmatrix}, \quad \begin{pmatrix} \mathbf{w} \\ \boldsymbol{\varpi} \end{pmatrix} \mapsto \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} g \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} \mathbf{w} \\ \boldsymbol{\varpi} \end{pmatrix}, \quad (2.5)$$

where $g \in \text{SL}(2, \mathbb{R})$ is a lower diagonal matrix with positive diagonal elements. Relevant properties of the gauge group G and its Lie algebra are collected in appendix A.

2.2 Classical observables

Recall that an observable is a function on Γ whose Poisson brackets with the first class constraints vanish when the first class constraints hold [36]. As discussed in [14], the system has the observables

$$\begin{aligned} A_{ij} &:= u_i p_j - u_j p_i, & 1 \leq i \leq p, \quad 1 \leq j \leq p; \\ B_{ij} &:= w_i \varpi_j - w_j \varpi_i, & 1 \leq i \leq q, \quad 1 \leq j \leq q; \\ C_{ij} &:= -u_i \varpi_j - p_i w_j, & 1 \leq i \leq p, \quad 1 \leq j \leq q, \end{aligned} \quad (2.6)$$

whose Poisson brackets close as the $\mathfrak{o}(p, q)$ Lie algebra, $\{A_{ij}\}$ and $\{B_{ij}\}$ spanning respectively the $\mathfrak{o}(p)$ and $\mathfrak{o}(q)$ Lie subalgebras. We denote the algebra generated by the observables (2.6) by $\mathcal{A}_{\text{class}}$. The finite transformations that $\mathcal{A}_{\text{class}}$ generates on Γ are

$$\begin{pmatrix} \mathbf{u} \\ \mathbf{w} \end{pmatrix} \mapsto R \begin{pmatrix} \mathbf{u} \\ \mathbf{w} \end{pmatrix}, \quad \begin{pmatrix} \mathbf{p} \\ \boldsymbol{\varpi} \end{pmatrix} \mapsto (R^T)^{-1} \begin{pmatrix} \mathbf{p} \\ \boldsymbol{\varpi} \end{pmatrix}, \quad (2.7)$$

where R is an $O(p, q)$ matrix, in the connected component $O_c(p, q)$.

The quadratic Casimir element in $\mathcal{A}_{\text{class}}$ is [37]

$$\mathcal{C} := \sum_{i < j} (A_{ij})^2 + \sum_{i < j} (B_{ij})^2 - \sum_{i, j} (C_{ij})^2. \quad (2.8)$$

When the constraints hold, \mathcal{C} vanishes [14].

2.3 Reduced phase space

Let $\bar{\Gamma}$ be the subset of Γ where the constraints hold. The reduced phase space, denoted by \mathcal{M} , is the quotient of $\bar{\Gamma}$ under the gauge action (2.5). We now discuss the structure of \mathcal{M} .

Note first that as the Hamiltonian is a linear combination of the constraints, there is no dynamics on \mathcal{M} , and \mathcal{M} can be identified with the space of classical solutions. As the functions in $\mathcal{A}_{\text{class}}$ are gauge invariant, they project to functions on \mathcal{M} .

$\bar{\Gamma}$ is clearly connected. Hence also \mathcal{M} is connected.

Let $\bar{\Gamma}_0 = \{\mathbf{q}_0\}$, where \mathbf{q}_0 is the origin of Γ , $\mathbf{u} = \mathbf{p} = \mathbf{0} = \mathbf{w} = \boldsymbol{\varpi}$. Let $\bar{\Gamma}_{\text{ex}}$ contain all other points on $\bar{\Gamma}$ at which $\mathbf{u} = \mathbf{0}$. Finally, let $\bar{\Gamma}_{\text{reg}}$ contain the points on $\bar{\Gamma}$ at which $\mathbf{u} \neq \mathbf{0}$. We refer to $\bar{\Gamma}_{\text{ex}}$ and $\bar{\Gamma}_{\text{reg}}$ as respectively the “exceptional” and “regular” parts of $\bar{\Gamma}$. As $\bar{\Gamma}_0$, $\bar{\Gamma}_{\text{ex}}$ and $\bar{\Gamma}_{\text{reg}}$ are preserved by the gauge transformations, they project onto disjoint subsets of \mathcal{M} . We denote these sets respectively by \mathcal{M}_0 , \mathcal{M}_{ex} and \mathcal{M}_{reg} and analyse each in turn.

2.3.1 \mathcal{M}_0

\mathcal{M}_0 contains only one point, the projection of \mathbf{q}_0 . All observables in $\mathcal{A}_{\text{class}}$ vanish on \mathcal{M}_0 .

2.3.2 \mathcal{M}_{ex}

Each point in $\bar{\Gamma}_{\text{ex}}$ is gauge-equivalent to a unique point that satisfies

$$\mathbf{u} = \mathbf{0} = \mathbf{w} \quad , \quad \mathbf{p}^2 + \boldsymbol{\varpi}^2 = 1 \quad . \quad (2.9)$$

\mathcal{M}_{ex} has thus topology S^{p+q-1} . By (2.2), the projection of Ω vanishes on \mathcal{M}_{ex} . All observables in $\mathcal{A}_{\text{class}}$ vanish on \mathcal{M}_{ex} .

2.3.3 \mathcal{M}_{reg}

Each point in $\bar{\Gamma}_{\text{reg}}$ is gauge-equivalent to a unique point that satisfies

$$\mathbf{u}^2 = \mathbf{w}^2 = 1 \quad , \quad \mathbf{u} \cdot \mathbf{p} = \mathbf{w} \cdot \boldsymbol{\varpi} = 0 \quad . \quad (2.10)$$

\mathcal{M}_{reg} can therefore be represented as the set (2.10), which is the cotangent bundle over $S^{p-1} \times S^{q-1}$, with (\mathbf{u}, \mathbf{w}) forming the base space and $(\mathbf{p}, \boldsymbol{\varpi})$ the fibres. By (2.2), the symplectic structure on \mathcal{M}_{reg} induced from Γ is precisely the symplectic structure of this cotangent bundle description. \mathcal{M}_{reg} is connected when $p > 1$ and $q > 1$, and it has two connected components when exactly one of p and q equals 1. When $p = q = 1$, \mathcal{M}_{reg} contains just four points.

$\mathcal{A}_{\text{class}}$ does not separate all of \mathcal{M}_{reg} . However, we show in appendix B that when $p > 1$ and $q > 1$, $\mathcal{A}_{\text{class}}$ separates the subset of \mathcal{M}_{reg} in which $0 \neq \mathbf{p}^2 \neq \boldsymbol{\varpi}^2 \neq 0$, in the gauge (2.10), up to the twofold degeneracy

$$(\mathbf{u}, \mathbf{w}, \mathbf{p}, \boldsymbol{\varpi}) \mapsto (-\mathbf{u}, -\mathbf{w}, -\mathbf{p}, -\boldsymbol{\varpi}) \quad . \quad (2.11)$$

We also show that when $p = 1$ and $q > 1$ (respectively $p > 1$ and $q = 1$), $\mathcal{A}_{\text{class}}$ separates the subset of \mathcal{M}_{reg} in which $\boldsymbol{\varpi}^2 \neq 0$ ($\mathbf{p}^2 \neq 0$), again up to the degeneracy (2.11). When $p = q = 1$, all observables in $\mathcal{A}_{\text{class}}$ vanish on \mathcal{M}_{reg} .

3 Algebraic quantisation with the constraints (1.2)

In this section we quantise the system by the algebraic techniques of [33, 34], imposing the constraints in the form (1.2) and requiring the classical $\mathfrak{o}(p, q)$ observables to become self-adjoint operators. Subsection 3.1 shows that this procedure leads to a quantum theory for arbitrary (p, q) : It will be seen in sections 5 and 6 that these quantum theories are isomorphic to those emerging from refined algebraic quantisation with group averaging. The remaining subsections analyse in detail the representations of the $\mathfrak{o}(p, q)$ observables for $p + q \leq 4$.

3.1 General (p, q)

Treating \mathbf{u} and \mathbf{w} as the ‘configuration’ variables, we represent the elementary operators as

$$\begin{aligned}\hat{\mathbf{u}}\Psi &= \mathbf{u}\Psi, & \hat{\mathbf{p}}\Psi &= -i\nabla_{\mathbf{u}}\Psi, \\ \hat{\mathbf{w}}\Psi &= \mathbf{w}\Psi, & \hat{\boldsymbol{\omega}}\Psi &= -i\nabla_{\mathbf{w}}\Psi,\end{aligned}\tag{3.1}$$

where the class of ‘functions’ $\Psi(\mathbf{u}, \mathbf{w})$ will be specified shortly. For the quantum constraint operators, we take

$$\hat{H} := \frac{1}{2}(\mathbf{w}^2 - \mathbf{u}^2), \tag{3.2a}$$

$$\hat{D} := -i\left(\mathbf{u} \cdot \nabla_{\mathbf{u}} + \mathbf{w} \cdot \nabla_{\mathbf{w}} + \frac{p+q}{2}\right), \tag{3.2b}$$

so that the commutator algebra reads

$$[\hat{H}, \hat{D}] = 2i\hat{H}. \tag{3.3}$$

The choice of the additive constant in (3.2b) is inherited from the $\text{SL}(2, \mathbb{R})$ constraint algebra in [14] and will lead to a transparent comparison with refined algebraic quantisation in sections 5 and 6.

The quantum constraints (1.2) read

$$\hat{H}\Psi = 0, \tag{3.4a}$$

$$(\hat{D} - i)\Psi = 0. \tag{3.4b}$$

As the only continuous solution to (3.4a) is $\Psi = 0$, we seek solutions in the distributional sense. Equation (3.4b) is equivalent to the homogeneity condition $\Psi(r\mathbf{u}, r\mathbf{w}) = r^{-(p+q+2)/2}\Psi(\mathbf{u}, \mathbf{w})$ for $r > 0$. The set (3.4) is thus satisfied by

$$\Psi(\mathbf{u}, \mathbf{w}) = \delta(u^2 - w^2)f(\mathbf{u}, \mathbf{w}), \tag{3.5a}$$

where $u := \sqrt{\mathbf{u}^2}$ and $w := \sqrt{\mathbf{w}^2}$, δ is the Dirac delta-distribution, and $f(\mathbf{u}, \mathbf{w})$ is smooth for $(\mathbf{u}, \mathbf{w}) \neq (\mathbf{0}, \mathbf{0})$ and homogeneous of degree $-(p+q-2)/2$,

$$f(r\mathbf{u}, r\mathbf{w}) = r^{-(p+q-2)/2}f(\mathbf{u}, \mathbf{w}) \text{ for } r > 0. \tag{3.5b}$$

We denote the vector space of the solutions (3.5) by $\mathcal{V}_{\text{phys}}$.

We define the quantum counterparts of the classical observables (2.6) as

$$\begin{aligned}\hat{A}_{ij} &:= -i(u_i\partial_{u_j} - u_j\partial_{u_i}), & 1 \leq i \leq p, \quad 1 \leq j \leq p; \\ \hat{B}_{ij} &:= -i(w_i\partial_{w_j} - w_j\partial_{w_i}), & 1 \leq i \leq q, \quad 1 \leq j \leq q; \\ \hat{C}_{ij} &:= i(u_i\partial_{w_j} + w_j\partial_{u_i}), & 1 \leq i \leq p, \quad 1 \leq j \leq q,\end{aligned}\tag{3.6}$$

These operators commute with the quantum constraints (3.2) and are thus quantum observables, and their commutator algebra closes as (i times) the $\mathfrak{o}(p, q)$ Lie algebra.

As in [14], we define the full star-algebra $\mathcal{A}_{\text{phys}}^{(*)}$ as the algebra generated by (3.6), the antilinear star-relation being defined so that it leaves the observables (3.6) invariant. The quantum quadratic Casimir observable is [37]

$$\hat{\mathcal{C}} := \sum_{i < j} (\hat{A}_{ij})^2 + \sum_{i < j} (\hat{B}_{ij})^2 - \sum_{i,j} (\hat{C}_{ij})^2 . \quad (3.7)$$

On states satisfying (3.4), $\hat{\mathcal{C}}$ takes the value $-\frac{1}{4}(p+q-2)^2$ [14].

By (2.6) and (2.7), the observables (3.6) generate on $\mathcal{V}_{\text{phys}}$ the representation T of $O_c(p, q)$ given by

$$[T(R)\Psi](\mathbf{u}, \mathbf{w}) = \Psi(R^{-1}(\mathbf{u}, \mathbf{w})) , \quad (3.8)$$

where $R \in O_c(p, q)$ acts on the configuration space as in (2.7). Writing $\Psi = \delta f$ as in (3.5), and noting that $u^2 - w^2$ is invariant under $O_c(p, q)$, we find

$$[T(R)\delta f](\mathbf{u}, \mathbf{w}) = \delta(u^2 - w^2)f(R^{-1}(\mathbf{u}, \mathbf{w})) . \quad (3.9)$$

T is thus isomorphic to the representation of $O_c(p, q)$ on homogeneous functions of degree $-(p+q-2)/2$ on the light cone of $\mathbb{R}^{p,q}$. This representation was investigated in [32], Section 9.2.9, and shown to be unitary with respect to the inner product

$$(\Psi_1, \Psi_2) = \int_{S^{p-1} \times S^{q-1}} d\mathbf{u} d\mathbf{w} \overline{f_1(\mathbf{u}, \mathbf{w})} f_2(\mathbf{u}, \mathbf{w}) , \quad (3.10)$$

where the overline denotes complex conjugation and the integration is over the product of the unit spheres in \mathbf{u} and \mathbf{w} , $\mathbf{u}^2 = \mathbf{w}^2 = 1$. Completion of $\mathcal{V}_{\text{phys}}$ in the inner product (3.10) therefore gives a physical Hilbert space on which T is a unitary representation of $O_c(p, q)$ and the infinitesimal generators (3.6) of T are densely-defined self-adjoint operators.

For any p and q , T reduces into subrepresentations T^ϵ on functions of parity $\epsilon \in \{0, 1\}$, $f(-\mathbf{u}, -\mathbf{w}) = (-1)^\epsilon f(\mathbf{u}, \mathbf{w})$. In the terminology of [32], T^ϵ is the end-point of the maximally degenerate principal unitary series of representations of $O_c(p, q)$, denoted by $T_{pq}^{(p+q-2)/2, \epsilon}$. $T_{pq}^{(p+q-2)/2, \epsilon}$ is nontrivial iff $(p, q) \neq (1, 1)$. $T_{pq}^{(p+q-2)/2, \epsilon}$ is known to be irreducible when p and q are of opposite parity, and also when p and q are both even and $(p+q)/2 + \epsilon \equiv 0 \pmod{2}$.

In the following subsections we decompose T into irreducible representations for $p+q \leq 4$. By interchange of p and q , it suffices to consider $p \geq q$.

3.2 $(p, q) = (1, 1)$

When $(p, q) = (1, 1)$, $\mathcal{V}_{\text{phys}}$ has dimension four, corresponding to the four branches of the 1+1 light cone. The representation of $\mathcal{A}_{\text{phys}}^{(*)}$ on $\mathcal{V}_{\text{phys}}$ is trivial and does not restrict the choice of an inner product on $\mathcal{V}_{\text{phys}}$.

3.3 $(p, q) = (2, 1)$

When $(p, q) = (2, 1)$, we consider the subspaces $\mathcal{V}^\kappa := \text{span}\{\psi_m^\kappa\} \subset \mathcal{V}_{\text{phys}}$, where $\kappa \in \{1, -1\}$,

$$\psi_m^\kappa := (-i\kappa)^m \delta(u^2 - w^2) \theta(\kappa w_1) u^{-1/2} e^{im\alpha} , \quad (3.11)$$

$m \in \mathbb{Z}$, θ is the Heaviside function, and we have written $u_1 + iu_2 = ue^{i\alpha}$. Writing $\hat{C}_\pm := \hat{C}_{11} \pm i\hat{C}_{21}$, the set $\{\hat{A}_{12}, \hat{C}_+, \hat{C}_-\}$ forms a standard raising and lowering operator basis for $\mathfrak{o}(2, 1)$ [38, 39], and we find

$$\hat{A}_{12}\psi_m^\kappa = m\psi_m^\kappa , \quad (3.12a)$$

$$\hat{C}_\pm\psi_m^\kappa = (m \pm \frac{1}{2})\psi_{m\pm 1}^\kappa . \quad (3.12b)$$

Each \mathcal{V}^κ hence carries an irreducible representation of $\mathcal{A}_{\text{phys}}^{(*)}$, and requiring \hat{A}_{12} , \hat{C}_{11} and \hat{C}_{21} to be self-adjoint fixes $\{\psi_m^\kappa\}$ for each κ to be an orthonormal set up to an overall scale [38]. Note that this agrees with the inner product (3.10). The Hilbert space is obtained by Cauchy completion. In the terminology of [38], the representation of $\text{O}_c(2, 1)$ on the Hilbert space is the principal series irreducible unitary representation $C_{1/4}^0$.

The representations on \mathcal{V}^1 and \mathcal{V}^{-1} are isomorphic. The representations $T_{2,1}^{\frac{1}{2},\epsilon}$ arise by starting from the parity ϵ subspaces $\text{span}\{\psi_m^1 + (-1)^\epsilon \psi_m^{-1}\} \subset \mathcal{V}^1 \oplus \mathcal{V}^{-1}$.

The quantum theories appear reasonable in view of the properties of the reduced phase space. In particular, both the classical observable A_{12} and the spectrum of the quantum observable \hat{A}_{12} are unbounded above and below.

3.4 $(p, q) = (2, 2)$

When $(p, q) = (2, 2)$, we write [10]

$$\begin{aligned} \hat{\tau}_0^\eta &:= \frac{1}{2}(\hat{A}_{12} + \eta\hat{B}_{12}) , \\ \hat{\tau}_1^\eta &:= \frac{1}{2}(\hat{C}_{11} - \eta\hat{C}_{22}) , \\ \hat{\tau}_2^\eta &:= \frac{1}{2}(\hat{C}_{21} + \eta\hat{C}_{12}) , \end{aligned} \quad (3.13)$$

where $\eta \in \{1, -1\}$ and

$$\hat{\tau}_\pm^\eta := \hat{\tau}_1^\eta \pm i\hat{\tau}_2^\eta . \quad (3.14)$$

The commutators of the $\hat{\tau}$ s are

$$[\hat{\tau}_0^\eta, \hat{\tau}_\pm^{\eta'}] = \pm\delta^{\eta\eta'} \hat{\tau}_\pm^\eta , \quad (3.15a)$$

$$[\hat{\tau}_+^\eta, \hat{\tau}_-^{\eta'}] = -2\delta^{\eta\eta'} \hat{\tau}_0^\eta , \quad (3.15b)$$

which shows that the $\hat{\tau}$ s provide standard $\mathfrak{sl}(2, \mathbb{R})$ raising and lowering operator bases in the decomposition $\mathfrak{o}(2, 2) \simeq \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{sl}(2, \mathbb{R})$. In the polar coordinates $u_1 + iu_2 = ue^{i\alpha}$,

$w_1 + iw_2 = we^{i\beta}$, we find

$$\hat{\tau}_0^\eta = -\frac{1}{2}i(\partial_\alpha + \eta\partial_\beta) \quad , \quad (3.16a)$$

$$\hat{\tau}_\pm^\eta = \frac{1}{2}ie^{\pm i(\alpha+\eta\beta)} \left\{ u[\partial_w \pm \eta(i/w)\partial_\beta] + w[\partial_u \pm (i/u)\partial_\alpha] \right\} \quad . \quad (3.16b)$$

Let $\mathcal{V} \subset \mathcal{V}_{\text{phys}}$ be the subspace spanned by the vectors $\delta(u^2 - w^2)u^{-1}e^{i(m\alpha+n\beta)}$, where m and n are integers. We label these vectors by $\mu := \frac{1}{2}(m+n)$ and $\nu := \frac{1}{2}(m-n)$, defining

$$\psi_{\mu\nu} := (-i)^{\mu+\nu} \delta(u^2 - w^2)u^{-1}e^{i[(\mu+\nu)\alpha+(\mu-\nu)\beta]} \quad , \quad (3.17)$$

where μ and ν are either both integers or both half-integers. A direct computation gives

$$\begin{aligned} \tau_0^+ \psi_{\mu\nu} &= \mu \psi_{\mu\nu} \quad , \\ \tau_\pm^+ \psi_{\mu\nu} &= \left(\mu \pm \frac{1}{2}\right) \psi_{\mu\pm 1, \nu} \quad , \\ \tau_0^- \psi_{\mu\nu} &= \nu \psi_{\mu\nu} \quad , \\ \tau_\pm^- \psi_{\mu\nu} &= \left(\nu \pm \frac{1}{2}\right) \psi_{\mu, \nu\pm 1} \quad , \end{aligned} \quad (3.18)$$

which shows that \mathcal{V} carries a representation of $\mathcal{A}_{\text{phys}}^{(\star)}$.

We decompose \mathcal{V} as $\mathcal{V} = \mathcal{V}^e \oplus \bigoplus_{\epsilon_1 \epsilon_2} \mathcal{V}_{\epsilon_1 \epsilon_2}^o$, where

$$\begin{aligned} \mathcal{V}^e &:= \text{span}\{\psi_{\mu\nu} \mid \mu, \nu \in \mathbb{Z}\} \quad , \\ \mathcal{V}_{\epsilon_1 \epsilon_2}^o &:= \text{span}\{\psi_{\mu\nu} \mid \mu, \nu \in \mathbb{Z} + \tfrac{1}{2}; \text{sgn}(\mu) = \epsilon_1; \text{sgn}(\nu) = \epsilon_2\} \quad , \end{aligned} \quad (3.19)$$

and $\epsilon_i \in \{1, -1\}$. Equations (3.18) show that each of the five spaces in (3.19) carries a representation of $\mathcal{A}_{\text{phys}}^{(\star)}$. These representations are irreducible: Given a nonzero vector in one of the spaces, acting on this vector repeatedly with τ_0^η and taking suitable linear combinations generates some $\psi_{\mu_0\nu_0}$, and repeatedly acting on this $\psi_{\mu_0\nu_0}$ by τ_\pm^η and taking linear combinations generates all of the space. Note that the difference between integer and half-integer indices in (3.19) arises because in the latter case the numerical coefficients in the raising and lowering operator action in (3.18) may vanish.

On each of the five spaces in (3.19), the $\mathfrak{sl}(2, \mathbb{R})$ analysis of [38] in each index shows that the adjoint relations

$$(\hat{\tau}_0^\eta)^\dagger = (\hat{\tau}_0^\eta)^\dagger \quad , \quad (\hat{\tau}_\pm^\eta)^\dagger = (\hat{\tau}_\mp^\eta)^\dagger \quad , \quad (3.20)$$

determine uniquely an inner product in which $\{\psi_{\mu\nu}\}$ is an orthonormal set up to an overall scale. This agrees with the inner product (3.10). The Hilbert spaces \mathcal{H}^e and $\mathcal{H}_{\epsilon_1 \epsilon_2}^o$ are obtained by Cauchy completion. In the terminology of [38], the representations of $\text{O}_c(2, 2) \simeq (\text{SL}(2, \mathbb{R}) \times \text{SL}(2, \mathbb{R}))/\mathbb{Z}_2$ are respectively $C_{1/4}^0 \times C_{1/4}^0$ and $D_{1/2}^{\epsilon_1} \times D_{1/2}^{\epsilon_2}$. The first of these is $T_{2,2}^{1,0}$ and the other four constitute $T_{2,2}^{1,1}$.

The quantum theory on \mathcal{H}^e appears reasonable in view of the properties of the reduced phase space. In particular, the classical observables $A_{12} \pm B_{12}$ and the spectra of $\hat{A}_{12} \pm \hat{B}_{12}$ are all unbounded above and below. By contrast, on each $\mathcal{H}_{\epsilon_1 \epsilon_2}^o$ the spectra of $\hat{A}_{12} \pm \hat{B}_{12}$ have a definite sign. This classically unexpected property may be related to the failure of $\mathcal{A}_{\text{class}}$ to separate the subsets of \mathcal{M}_{reg} where one of $A_{12} \pm B_{12}$ vanishes.

3.5 $(p, q) = (3, 1)$

When $(p, q) = (3, 1)$, the value of the quadratic Casimir operator (3.7) in the representations $T_{3,1}^{2,\epsilon}$ is -1 , and a direct computation shows that the Casimir operator $\hat{A}_{12}\hat{C}_{31} + \hat{A}_{23}\hat{C}_{11} + \hat{A}_{31}\hat{C}_{21}$ has value zero. It follows that the representations $T_{3,1}^{2,\epsilon}$ are each isomorphic to the principal series unitary irreducible representation $\mathfrak{S}_{0,0}$ of $O_c(3, 1)$ [40].

4 Algebraic quantisation with $\hat{H}_a\Psi = 0$

In this section we discuss how the algebraic quantisation of section 3 is modified when the constraints (1.2) are replaced by $\hat{H}_a\Psi = 0$. We give a complete analysis for $p+q \leq 4$ and partial results for other values of p and q .

4.1 General (p, q)

The quantum constraints (3.4) are replaced by

$$\hat{H}\Psi = 0 \quad , \quad (4.1a)$$

$$\hat{D}\Psi = 0 \quad . \quad (4.1b)$$

Proceeding as in subsection 3.1, the exponent of r in (3.5b) is replaced by $-(p+q-4)/2$, and the value of the quadratic Casimir (3.7) is $-\frac{1}{4}(p+q)(p+q-4)$ [14].

The representation of $O_c(p, q)$ generated by the quantum observables is now isomorphic to the representation on homogeneous functions of degree $-(p+q-4)/2$ on the light cone of $\mathbb{R}^{p,q}$. The outstanding question is whether this representation or some subrepresentation thereof is unitary in some inner product.

We analyse different ranges of (p, q) in the following subsections. By interchange of p and q , it suffices to consider $p \geq q$.

4.2 $(p, q) = (1, 1)$

When $(p, q) = (1, 1)$, the discussion follows subsection 3.2. The representation of $\mathcal{A}_{\text{phys}}^{(*)}$ is trivial.

4.3 $(p, q) = (2, 1)$

When $(p, q) = (2, 1)$, the factor $u^{-1/2}$ in (3.11) is replaced by $u^{1/2}$, and the numerical factor on the right-hand side of (3.12b) is replaced by $(m \mp \frac{1}{2})$. The representations of $\mathcal{A}_{\text{phys}}^{(*)}$ on the counterparts of \mathcal{V}^κ are irreducible, but there is no inner product in which \hat{A}_{12} , \hat{C}_{11} and \hat{C}_{21} would be self-adjoint [38]. No quantum theory is recovered.

4.4 $(p, q) = (2, 2)$

When $(p, q) = (2, 2)$, (3.17) is replaced by

$$\tilde{\psi}_{\mu\nu} := (-i)^{\mu+\nu} \delta(u^2 - w^2) e^{i[(\mu+\nu)\alpha + (\mu-\nu)\beta]} , \quad (4.2)$$

where μ and ν are again either both integers or both half-integers. We write $\tilde{\mathcal{V}} := \text{span}\{\tilde{\psi}_{\mu\nu}\}$. A direct computation gives

$$\begin{aligned} \tau_0^+ \tilde{\psi}_{\mu\nu} &= \mu \tilde{\psi}_{\mu\nu} , \\ \tau_{\pm}^+ \tilde{\psi}_{\mu\nu} &= \mu \tilde{\psi}_{\mu\pm 1, \nu} , \\ \tau_0^- \tilde{\psi}_{\mu\nu} &= \nu \tilde{\psi}_{\mu\nu} , \\ \tau_{\pm}^- \tilde{\psi}_{\mu\nu} &= \nu \tilde{\psi}_{\mu, \nu\pm 1} . \end{aligned} \quad (4.3)$$

Hence $\tilde{\mathcal{V}}$ carries a representation of $\mathcal{A}_{\text{phys}}^{(*)}$.

We decompose $\tilde{\mathcal{V}}$ as $\tilde{\mathcal{V}} = \tilde{\mathcal{V}}^o \oplus \bigoplus_{\epsilon_1 \epsilon_2} \tilde{\mathcal{V}}_{\epsilon_1 \epsilon_2}^e$, where

$$\begin{aligned} \tilde{\mathcal{V}}^o &:= \text{span}\{\tilde{\psi}_{\mu\nu} \mid \mu, \nu \in \mathbb{Z} + \tfrac{1}{2}\} , \\ \tilde{\mathcal{V}}_{\epsilon_1 \epsilon_2}^e &:= \text{span}\{\tilde{\psi}_{\mu\nu} \mid \mu, \nu \in \mathbb{Z}; \text{sgn}(\mu) = \epsilon_1; \text{sgn}(\nu) = \epsilon_2\} , \end{aligned} \quad (4.4)$$

and $\epsilon_i \in \{1, 0, -1\}$. Equations (4.3) show that each of the ten spaces in (4.4) carries a representation of $\mathcal{A}_{\text{phys}}^{(*)}$, given by (4.3) except that whenever a raising (respectively lowering) operator raises (lowers) the index -1 ($+1$) to zero, the vector on the right-hand side is replaced by the zero vector. It can be verified as in subsection 3.4 that these representations are irreducible. The $\mathfrak{sl}(2, \mathbb{R})$ analysis of [38] in each index then shows that there is no inner product on $\tilde{\mathcal{V}}^o$ compatible with the adjoint relations (3.20), whereas on each $\tilde{\mathcal{V}}_{\epsilon_1 \epsilon_2}^e$ these relations determine an inner product that is unique up to an overall scale. For $\epsilon_1 \neq 0 \neq \epsilon_2$, this inner product reads $(\tilde{\psi}_{\mu'\nu'}, \tilde{\psi}_{\mu\nu}) = |\mu\nu| \delta_{\mu\mu'} \delta_{\nu\nu'}$, while the formulas for $\epsilon_1 \neq 0 = \epsilon_2$ and $\epsilon_1 = 0 \neq \epsilon_2$ are respectively $(\tilde{\psi}_{\mu'0}, \tilde{\psi}_{\mu 0}) = |\mu| \delta_{\mu\mu'}$ and $(\tilde{\psi}_{0\nu'}, \tilde{\psi}_{0\nu}) = |\nu| \delta_{\nu\nu'}$. On the one-dimensional space $\tilde{\mathcal{V}}_{00}^e$ the representation of $\mathcal{A}_{\text{phys}}^{(*)}$ is trivial, and we have $(\tilde{\psi}_{00}, \tilde{\psi}_{00}) = 1$.

The Hilbert spaces $\tilde{\mathcal{H}}_{\epsilon_1 \epsilon_2}^e$ are obtained by Cauchy completion. In the terminology of [38], the representation of $\text{O}_c(2, 2) \simeq (\text{SL}(2, \mathbb{R}) \times \text{SL}(2, \mathbb{R}))/\mathbb{Z}_2$ on $\tilde{\mathcal{H}}_{\epsilon_1 \epsilon_2}^e$ is $D_1^{\epsilon_1} \times D_1^{\epsilon_2}$ for $\epsilon_1 \neq 0 \neq \epsilon_2$, and for $\epsilon_i = 0$, $D_1^{\epsilon_i}$ is replaced by the trivial representation.

The quantum theories on $\tilde{\mathcal{H}}_{\epsilon_1 \epsilon_2}^e$ with $\epsilon_1 \neq 0 \neq \epsilon_2$ are qualitatively similar to the theories on $\mathcal{H}_{\epsilon_1 \epsilon_2}^o$ obtained in subsection 3.4, with the roles of integer and half-integer eigenvalues of $\hat{\tau}_0^\eta$ interchanged. The remaining five theories are degenerate in that at least one $\hat{\tau}_0^\eta$ annihilates the whole space. Note that in contrast to subsection 3.4, we obtained no theory in which the representation of $\mathcal{A}_{\text{phys}}^{(*)}$ would be irreducible and the spectra of $\hat{A}_{12} \pm \hat{B}_{12}$ would be unbounded both above and below.

$$\begin{aligned}
\hat{A}_{12}\tilde{\psi}_{lm} &= m\tilde{\psi}_{lm} \\
(\hat{A}_{23} \pm i\hat{A}_{31})\tilde{\psi}_{lm} &= \sqrt{(l \pm m + 1)(l \mp m)} \tilde{\psi}_{l,m \pm 1} \\
\hat{C}_{31}\tilde{\psi}_{lm} &= -il \sqrt{\frac{(l+m+1)(l-m+1)}{(2l+1)(2l+3)}} \tilde{\psi}_{l+1,m} \\
&\quad + i(l+1) \sqrt{\frac{(l+m)(l-m)}{(2l+1)(2l-1)}} \tilde{\psi}_{l-1,m} \\
(\hat{C}_{11} \pm i\hat{C}_{21})\tilde{\psi}_{lm} &= \pm il \sqrt{\frac{(l \pm m + 1)(l \pm m + 2)}{(2l+1)(2l+3)}} \tilde{\psi}_{l+1,m \pm 1} \\
&\quad \pm i(l+1) \sqrt{\frac{(l \mp m)(l \mp m - 1)}{(2l+1)(2l-1)}} \tilde{\psi}_{l-1,m \pm 1}
\end{aligned}$$

Table 1: The action of $\mathcal{A}_{\text{phys}}^{(*)}$ on $\tilde{\mathcal{V}}$ for $(p, q) = (3, 1)$.

4.5 $(p, q) = (3, 1)$

When $(p, q) = (3, 1)$, the branches $w_1 > 0$ and $w_1 < 0$ of the light cone decouple and give isomorphic quantum theories. We consider the branch $w_1 > 0$.

We set $\tilde{\mathcal{V}} := \text{span}\{\tilde{\psi}_{lm}\}$, where

$$\tilde{\psi}_{lm} := \delta(u^2 - w^2)\theta(w_1)Y_{lm} \quad , \quad (4.5)$$

and Y_{lm} are the usual spherical harmonics on unit S^2 in \mathbf{u} [41]. The action of the operators (3.6) on $\tilde{\mathcal{V}}$ can be computed from standard properties of the spherical harmonics [41, 42] and is displayed in Table 1. This shows that $\tilde{\mathcal{V}}$ carries a representation of $\mathcal{A}_{\text{phys}}^{(*)}$.

We decompose $\tilde{\mathcal{V}}$ as $\tilde{\mathcal{V}} = \tilde{\mathcal{V}}_0 \oplus \tilde{\mathcal{V}}_+$, where

$$\begin{aligned}
\tilde{\mathcal{V}}_0 &:= \text{span}\{\tilde{\psi}_{00}\} \quad , \\
\tilde{\mathcal{V}}_+ &:= \text{span}\{\tilde{\psi}_{lm} \mid l > 0\} \quad .
\end{aligned} \quad (4.6)$$

Table 1 shows that $\mathcal{A}_{\text{phys}}^{(*)}$ is represented trivially on $\tilde{\mathcal{V}}_0$, while $\tilde{\mathcal{V}}_+$ carries a representation that is as in the Table except that any term on the right-hand side with the first index taking the value zero is replaced by the zero vector. Comparison with the infinitesimal representations of $\text{O}_c(3, 1)$ ([40], section 8.3) shows that the representation on $\tilde{\mathcal{V}}_+$ is isomorphic to the principal series irreducible representation $\mathfrak{S}_{2,0}$, which is unitary precisely when the inner product is

$$(\tilde{\psi}_{l'm'}, \tilde{\psi}_{lm}) = l(l+1) \delta_{ll'} \delta_{mm'} \quad , \quad (4.7)$$

up to an overall multiple. The Casimir operator $\hat{A}_{12}\hat{C}_{31} + \hat{A}_{23}\hat{C}_{11} + \hat{A}_{31}\hat{C}_{21}$ takes value zero.

4.6 $2 \leq q \leq p$, $(p, q) \neq (2, 2)$

When $2 \leq q \leq p$ and $(p, q) \neq (2, 2)$ we set $\tilde{\mathcal{V}} := \text{span}\{\tilde{\psi}_{lj k_u k_w}\}$, where

$$\tilde{\psi}_{lj k_u k_w} := \delta(u^2 - w^2) u^{-(p+q-4)/2} Y_{l k_u}(\theta^{(u)}) Y_{j k_w}(\theta^{(w)}) \quad , \quad (4.8)$$

where l and j are non-negative integers and $Y_{l k_u}$ (respectively $Y_{j k_w}$) are the spherical harmonics on unit S^{p-1} in \mathbf{u} (S^{q-1} in \mathbf{w}) [32, 41]. The notation for the spherical harmonics follows [14]. The construction of the spherical harmonics implies that $\tilde{\mathcal{V}}$ carries a representation of $\mathcal{A}_{\text{phys}}^{(*)}$.

We seek a linear subspace $\tilde{\mathcal{V}}_0 \subset \tilde{\mathcal{V}}$ with the following properties:

1. $\tilde{\mathcal{V}}_0 := \text{span}\{\tilde{\psi}_{lj k_u k_w} \mid (l, j) \in I\}$, where I is some nonempty index set.
2. $\tilde{\mathcal{V}}_0$ carries a representation of $\mathcal{A}_{\text{phys}}^{(*)}$.
3. The generators (3.6) of $\mathcal{A}_{\text{phys}}^{(*)}$ are self-adjoint in an inner product of the form

$$(\tilde{\psi}_{l' j' k'_u k'_w}, \tilde{\psi}_{l j k_u k_w}) = K_{lj} \delta_{l' l} \delta_{j' j} \delta_{k'_u k_u} \delta_{k'_w k_w} \quad , \quad (4.9)$$

where the positive numbers K_{lj} depend only on l and j .

By the properties of the spherical harmonics, the rotation generators \hat{A}_{ij} and \hat{B}_{ij} in (3.6) leave $\tilde{\mathcal{V}}_0$ invariant and are self-adjoint in the inner product (4.9). What remains is to examine the boost generators \hat{C}_{ij} .

Let Y_{l0} (respectively Y_{j0}) denote the zonal spherical harmonics, which can be expressed in terms of Gegenbauer polynomials of argument u_p/u (w_q/w) [41]. The recursion relations of the Gegenbauer polynomials [43] allow an explicit computation of the action of \hat{C}_{pq} on $\tilde{\psi}_{lj00}$. Suppressing the indices $k_u = k_w = 0$, we find

$$\begin{aligned} 4i\hat{C}_{pq}\tilde{\psi}_{lj} &= [l+j+\tfrac{1}{2}(p+q-4)] \left(W_{p,l+1} W_{q,j+1} \tilde{\psi}_{l+1,j+1} - W_{pl} W_{qj} \tilde{\psi}_{l-1,j-1} \right) \\ &\quad + [l-j+\tfrac{1}{2}(p-q)] \left(W_{p,l+1} W_{qj} \tilde{\psi}_{l+1,j-1} - W_{pl} W_{q,j+1} \tilde{\psi}_{l-1,j+1} \right) \quad , \end{aligned} \quad (4.10)$$

where

$$\begin{aligned} W_{qj} &:= 2 \left[\frac{j(j+q-3)}{(2j+q-2)(2j+q-4)} \right]^{1/2} \quad \text{for } j > 0, (q, j) \neq (2, 1) \quad , \\ W_{21} &:= \sqrt{2} \quad , \\ W_{q0} &:= 0 \quad . \end{aligned} \quad (4.11)$$

By (4.9) and (4.10), self-adjointness of \hat{C}_{pq} implies the recursion relations

$$[l+j+\tfrac{1}{2}(p+q-4)] K_{l+1,j+1} = [l+j+\tfrac{1}{2}(p+q)] K_{lj} \quad , \quad (4.12a)$$

$$[l-j-1+\tfrac{1}{2}(p-q)] K_{l+1,j} = [l-j+1+\tfrac{1}{2}(p-q)] K_{l,j+1} \quad . \quad (4.12b)$$

Note that the coefficients in (4.12a) are always positive.

Suppose first that $p + q$ is odd. From (4.10) it follows that the index set I must contain all pairs (l, j) where $l + j$ is odd or all pairs where $l + j$ is even. The coefficients in (4.12b) are always nonzero, but the coefficients on the two sides have opposite sign for $j - l = \frac{1}{2}(p - q \pm 1)$. Hence there are both positive and negative K_{lj} , and the inner product does not exist. We have proved:

Theorem 4.1 *Let $p \geq 2$, $q \geq 2$ and $p + q \equiv 1 \pmod{2}$. Then there is no $\tilde{\mathcal{V}}_0$ satisfying 1–3.*

Suppose then that $p + q$ is even. If $\tilde{\mathcal{V}}_0$ contains a vector for which $l - j + \frac{1}{2}(p - q)$ is odd, (4.10) shows that it must contain all such vectors, and examination of the signs in (4.12b) shows that the inner product does not exist. Hence $\tilde{\mathcal{V}}_0$ can contain only vectors for which $l - j + \frac{1}{2}(p - q)$ is even. From (4.10) and (4.12) we obtain:

Theorem 4.2 *Let $p \geq 2$, $q \geq 2$, $(p, q) \neq (2, 2)$ and $p + q \equiv 0 \pmod{2}$. For a $\tilde{\mathcal{V}}_0$ satisfying 1–3, I is either $I_0 := \{(l, j) \mid l - j + \frac{1}{2}(p - q) = 0\}$ or one of $I_{\pm} := \{(l, j) \mid l - j + \frac{1}{2}(p - q) = 2k, k \in \mathbb{Z}_{\pm}\}$ or a union of two or all of these. For indices within each of I_0 , I_+ and I_- , (4.12) determines the positive numbers K_{lj} uniquely up to an overall scale.*

For $p + q$ even, Theorem 4.2 severely restricts the possible candidates for $\tilde{\mathcal{V}}_0$. A more complete understanding of the \hat{C}_{ij} action would be required to determine whether the candidate spaces of Theorem 4.2 indeed have properties 2 and 3. It is known ([32], Section 9.2.10) that there exist subspaces of $\tilde{\mathcal{V}}$ carrying an irreducible representation of $\mathcal{A}_{\text{phys}}^{(*)}$ such that the generators (3.6) are self-adjoint in an appropriate inner product, and these representations are equivalent to certain discrete series representations of $O_c(p, q)$. We shall however not examine correspondences between these subspaces and the spaces $\tilde{\mathcal{V}}_0$ of Theorem 4.2 further.

4.7 $p \geq 4$, $q = 1$

When $p \geq 4$ and $q = 1$, the branches $w_1 > 0$ and $w_1 < 0$ of the light cone decouple and lead to isomorphic situations. We consider the branch $w_1 > 0$ and set $\tilde{\mathcal{V}} := \text{span}\{\tilde{\psi}_{lk_u}\}$, where

$$\tilde{\psi}_{lk_u} := \delta(u^2 - w^2)\theta(w_1)u^{-(p-3)/2}Y_{lk_u} \quad , \quad (4.13)$$

and Y_{lk_u} are the spherical harmonics in on unit S^{p-1} in \mathbf{u} as in subsection 4.6. By construction of the spherical harmonics, $\tilde{\mathcal{V}}$ carries a representation of $\mathcal{A}_{\text{phys}}^{(*)}$. Assuming an inner product of the form

$$(\tilde{\psi}_{l'k'_u}, \tilde{\psi}_{lk_u}) = K_l \delta_{ll'} \delta_{k_u k'_u} \quad , \quad (4.14)$$

where the positive numbers K_l depend only on l , the rotation generators \hat{A}_{ij} are self-adjoint, and an analysis of the action of \hat{C}_{pq} on the $k_u = 0$ states as in subsection 4.6

implies $K_l = [l + \frac{1}{2}(p-1)][l + \frac{1}{2}(p-3)]$, up to an overall scale. However, further analysis would be required to determine whether \hat{C}_{i1} are indeed self-adjoint in this inner product.

5 Refined algebraic quantisation for $(p, q) \neq (1, 1)$

In this section we perform refined algebraic quantisation for $(p, q) \neq (1, 1)$. The special case $p = q = 1$ will be treated in section 6.

We employ refined algebraic quantisation with group averaging as formulated in [8]. We do not find a test space to which the uniqueness theorem of [8] would apply, but we find a test space on which the convergence of the averaging and the physical observable algebra can be controlled.

5.1 Auxiliary Hilbert space and representation of the gauge group

We use the auxiliary Hilbert space $\mathcal{H}_{\text{aux}} \simeq L^2(\mathbb{R}^{p+q})$ of square integrable functions $\Psi(\mathbf{u}, \mathbf{v})$ in the inner product

$$(\Psi_1, \Psi_2)_{\text{aux}} := \int d^p \mathbf{u} d^q \mathbf{v} \overline{\Psi_1} \Psi_2 \quad . \quad (5.1)$$

The algebra of the quantum constraints (3.2) exponentiates to a unitary representation U of G on \mathcal{H}_{aux} . The group elements in the decomposition (A.3) are represented by

$$U(\exp(\mu e^-)) = \exp(-i\mu \hat{H}) \quad , \quad (5.2a)$$

$$U(\exp(\lambda h)) = \exp(-i\lambda \hat{D}) \quad , \quad (5.2b)$$

where

$$[\exp(-i\mu \hat{H})\Psi](\mathbf{u}, \mathbf{w}) = \exp\left[\frac{i\mu(\mathbf{u}^2 - \mathbf{w}^2)}{2}\right] \Psi(\mathbf{u}, \mathbf{w}) \quad , \quad (5.3a)$$

$$[\exp(-i\lambda \hat{D})\Psi](\mathbf{u}, \mathbf{w}) = \exp\left[-\frac{\lambda(p+q)}{2}\right] \Psi(e^{-\lambda}\mathbf{u}, e^{-\lambda}\mathbf{w}) \quad . \quad (5.3b)$$

5.2 Test space

Let

$$\Psi_{l j m n k_u k_w}(\mathbf{u}, \mathbf{w}) := u^{l+2m} w^{j+2n} e^{-\frac{1}{2}(u^2 + w^2)} Y_{l k_u}(\theta^{(u)}) Y_{j k_w}(\theta^{(w)}) \quad , \quad (5.4)$$

where l, j, m and n are non-negative integers and the spherical harmonics are as in subsection 4.6. $p = 1$ is covered as the special case in which $\theta^{(u)} := u_1/u \in \{1, -1\}$, $l \in \{0, 1\}$, the index k_u takes only a single value and can be dropped and $Y_l(\theta^{(u)}) := (\theta^{(u)})^l / \sqrt{2}$, and similarly for $q = 1$.

We set $\Phi_0 := \text{span}\{\Psi_{ljmnk_uk_w}\} = \{P(\mathbf{u}, \mathbf{w}) \exp[-\frac{1}{2}(u^2 + w^2)]\}$, where $P(\mathbf{u}, \mathbf{w})$ is an arbitrary polynomial in $\{u_i\}$ and $\{w_i\}$. Φ_0 is dense in \mathcal{H}_{aux} and mapped to itself by $\mathcal{A}_{\text{phys}}^{(*)}$. We adopt as our test space Φ the closure of Φ_0 under the algebra generated by $\{U(g) \mid g \in G\}$. Note that although Φ satisfies the test space axioms in [8], it is not invariant under the group algebra defined in [8]. We shall return to this point in section 7.

5.3 Rigging map

We construct a rigging map by averaging states in Φ over G .

We define on Φ the sesquilinear form

$$(\phi_2, \phi_1)_{\text{ga}} := \int_G d_0g (\phi_2, U(g)\phi_1)_{\text{aux}} \quad , \quad (5.5)$$

where d_0g is the symmetric measure (see appendix A), normalised so that $d_0g = e^\lambda d\lambda d\mu$ in the parametrisation (A.3). We shall show that $(\cdot, \cdot)_{\text{ga}}$ is well defined and evaluate it explicitly.

From (A.3), (5.2) and (5.3) we find

$$\begin{aligned} [U(g)\Psi_{ljmnk_uk_w}](\mathbf{u}, \mathbf{w}) &= z^{-[\frac{1}{4}(p+q)+\frac{1}{2}(l+j)+m+n]} u^{l+2m} w^{j+2n} Y_{lk_u}(\theta^{(u)}) Y_{jk_w}(\theta^{(w)}) \\ &\quad \times \exp \left[-\frac{1}{2} \left(\frac{1}{z} - i\mu \right) u^2 - \frac{1}{2} \left(\frac{1}{z} + i\mu \right) w^2 \right] \quad , \quad (5.6) \end{aligned}$$

where $z := e^{2\lambda}$. In $(\Psi_{l'j'm'n'k'_uk'_w}, U(g)\Psi_{ljmnk_uk_w})_{\text{aux}}$, the angular integrals give the factor $\delta_{ll'}\delta_{jj'}\delta_{k_uk'_u}\delta_{k_wk'_w}$ and the integrals over u and w give Gamma-functions [44], with the result

$$\begin{aligned} &(\Psi_{l'j'm'n'k'_uk'_w}, U(g)\Psi_{ljmnk_uk_w})_{\text{aux}} \\ &= \delta_{ll'}\delta_{jj'}\delta_{k_uk'_u}\delta_{k_wk'_w} 2^{\frac{1}{2}(p+q)+l+j+m+m'+n+n'-2} \Gamma\left(\frac{1}{2}p+l+m+m'\right) \Gamma\left(\frac{1}{2}q+j+n+n'\right) \\ &\quad \times \frac{z^{\frac{1}{4}(p+q)+\frac{1}{2}(l+j)+m'+n'}}{(1-i\omega)^{\frac{1}{2}p+l+m+m'}(1+i\omega)^{\frac{1}{2}q+j+n+n'}(1+z)^{\frac{1}{2}(p+q)+l+j+m+m'+n+n'}} \quad , \quad (5.7) \end{aligned}$$

where we have written $\mu = \omega(1+z)/z$. An elementary analysis shows that necessary and sufficient conditions for (5.7) to be integrable in absolute value in the measure $d_0g = \frac{1}{2}z^{-3/2}(1+z)d\omega dz$ are

$$\begin{aligned} \frac{1}{4}(p+q-2) + \frac{1}{2}(l+j) + m+n &> 0 \quad , \\ \frac{1}{4}(p+q-2) + \frac{1}{2}(l+j) + m'+n' &> 0 \quad . \end{aligned} \quad (5.8)$$

As $p+q > 2$ by assumption, (5.8) is satisfied, and the integral of (5.7) in the measure d_0g is well defined. To evaluate this integral, we perform first the dz integral by 3.194.3 in [44]. If one of the exponents in the remaining $d\omega$ integral is less than unity, a contour

deformation brings the integral to a form to which 3.194.3 in [44] applies, and an analytic continuation in the exponents shows that the result is valid also for larger exponents. Collecting, we find

$$\begin{aligned} (\Psi_{l'j'm'n'k'_uk'_w}, \Psi_{lmnk_uk_w})_{\text{ga}} &= \frac{\pi}{2} \delta_{ll'} \delta_{jj'} \delta_{k_uk'_u} \delta_{k_wk'_w} \\ &\times \Gamma\left(\frac{1}{4}(p+q-2) + \frac{1}{2}(l+j) + m+n\right) \Gamma\left(\frac{1}{4}(p+q-2) + \frac{1}{2}(l+j) + m'+n'\right) . \end{aligned} \quad (5.9)$$

$(\cdot, \cdot)_{\text{ga}}$ is hence well defined on Φ_0 and given by (5.9). The relations [8] $d_0(g_1g) = \Delta^{1/2}(g_1)d_0g$, $d_0(gg_1) = \Delta^{-1/2}(g_1)d_0g$ show that $(\cdot, \cdot)_{\text{ga}}$ is well defined on all of Φ and given by (5.9) and

$$(\phi_1, U(g)\phi_2)_{\text{ga}} = (U(g)\phi_1, \phi_2)_{\text{ga}} = \Delta^{1/2}(g)(\phi_1, \phi_2)_{\text{ga}} . \quad (5.10)$$

Let Φ^* be the algebraic dual of Φ and let $f[\phi]$ denote the dual action of $f \in \Phi^*$ on $\phi \in \Phi$. Refined algebraic quantisation uses $(\cdot, \cdot)_{\text{ga}}$ to define the antilinear map $\eta : \Phi \rightarrow \Phi^*$ by

$$\eta(\phi_1)[\phi_2] := (\phi_1, \phi_2)_{\text{ga}} , \quad (5.11)$$

and then defines on the image of η the sesquilinear form $(\cdot, \cdot)_{\text{RAQ}}$ by

$$(\eta(\phi_1), \eta(\phi_2))_{\text{RAQ}} := \eta(\phi_2)[\phi_1] . \quad (5.12)$$

From (5.9) it follows that the image of η is nontrivial. We now give an explicit characterisation of the image of η and evaluate $(\cdot, \cdot)_{\text{RAQ}}$.

By (5.10) we have $\eta(\phi_1)[U(g)\phi_2] = \eta(U(g)\phi_1)[\phi_2] = [\Delta(g)]^{1/2}\eta(\phi_1)[\phi_2]$, and hence $\eta(U(g)\phi) = [\Delta(g)]^{1/2}\eta(\phi)$. It therefore suffices to evaluate $\eta(\phi_1)[\phi_2]$ for $\phi_1, \phi_2 \in \Phi_0$.

Let

$$\chi_{lj k_uk_w}(\mathbf{u}, \mathbf{w}) := \delta(u^2 - w^2) u^{-(p+q-2)/2} Y_{lk_u}(\theta^{(u)}) Y_{jk_w}(\theta^{(w)}) . \quad (5.13)$$

We interpret $\overline{\chi_{lj k_uk_w}}$ as an element of Φ^* , acting on states $\phi \in \Phi$ by

$$\overline{\chi_{lj k_uk_w}}[\phi] = \int d^p \mathbf{u} d^q \mathbf{w} \overline{\chi_{lj k_uk_w}(\mathbf{u}, \mathbf{w})} \phi(\mathbf{u}, \mathbf{w}) . \quad (5.14)$$

An explicit computation gives

$$\overline{\chi_{lj k_uk_w}}[\Psi_{lmnk_uk_w}] = \frac{1}{4} \delta_{ll'} \delta_{jj'} \delta_{k_uk'_u} \delta_{k_wk'_w} \Gamma\left(\frac{1}{4}(p+q-2) + \frac{1}{2}(l+j) + m+n\right) . \quad (5.15)$$

From (5.9), (5.11) and (5.15) we find

$$\eta(\Psi_{lmnk_uk_w}) = 2\pi \Gamma\left(\frac{1}{4}(p+q-2) + \frac{1}{2}(l+j) + m+n\right) \overline{\chi_{lj k_uk_w}} . \quad (5.16)$$

Hence the image of η is spanned by $\{\overline{\chi_{lj k_u k_w}}\}$. From (5.9), (5.12), (5.15), and (5.16) we obtain

$$(\overline{\chi_{l'j'k'_u k'_w}}, \overline{\chi_{lj k_u k_w}})_{\text{RAQ}} = (8\pi)^{-1} \delta_{l'l'} \delta_{jj'} \delta_{k_u k'_u} \delta_{k_w k'_w} . \quad (5.17)$$

We see that $(\cdot, \cdot)_{\text{RAQ}}$ is positive definite. It follows that η is a rigging map, and the physical Hilbert space \mathcal{H}_{RAQ} is the Cauchy completion of the image of η in $(\cdot, \cdot)_{\text{RAQ}}$.

The representation (3.6) of $\mathcal{A}_{\text{phys}}^{(*)}$ on \mathcal{H}_{aux} leaves Φ invariant and commutes with $U(g)$, and the star-relation in this representation coincides with the adjoint map on \mathcal{H}_{aux} . It follows, as in [14], that \mathcal{H}_{RAQ} carries an antilinear representation ρ of $\mathcal{A}_{\text{phys}}^{(*)}$, such that the star-relation coincides with the adjoint map on \mathcal{H}_{RAQ} . Comparing (5.13) and (5.14) to (3.5) shows that ρ is antilinearly isomorphic to the representation of $\mathcal{A}_{\text{phys}}^{(*)}$ obtained in the algebraic quantisation in subsection 3.1

6 Refined algebraic quantisation for $p = q = 1$

In this section we perform refined algebraic quantisation for $p = q = 1$.

When $p = q = 1$, the convergence conditions (5.8) fail to hold when $l = j = 0$ and at least one of the pairs (m, n) and (m', n') equals $(0, 0)$. Further, the integral of (5.7) is unambiguously divergent for $l = l' = j = j' = m = m' = n = n' = 0$. We shall remedy this problem by modifying the test space.

Dropping the redundant indices k_u and k_w , we write

$$\phi_{ljmn}(u_1, w_1) := \Psi_{ljmn00} = u^{l+2m} w^{j+2n} e^{-\frac{1}{2}(u^2+w^2)} Y_l(\theta^{(u)}) Y_j(\theta^{(w)}) , \quad (6.1)$$

where $l, j \in \{0, 1\}$. We also define

$$\psi_{mn} := -2\phi_{00,m+1,n+1} + (2m+1)\phi_{00,m,n+1} + (2n+1)\phi_{00,m+1,n} . \quad (6.2)$$

$$\Phi_0^{\text{mod}} := \text{span}(\{\phi_{ljmn} \mid l+j > 0\} \cup \{\psi_{mn}\}) . \quad (6.3)$$

An explicit computation shows

$$\hat{C}_{11}\phi_{00mn} = 2i(-\phi_{11mn} + m\phi_{11,m-1,n} + n\phi_{11,m,n-1}) , \quad (6.4a)$$

$$\hat{C}_{11}\phi_{01mn} = i[-2\phi_{10,m,n+1} + 2m\phi_{10,m-1,n+1} + (2n+1)\phi_{10mn}] , \quad (6.4b)$$

$$\hat{C}_{11}\phi_{10mn} = i[-2\phi_{01,m+1,n} + 2n\phi_{01,m+1,n-1} + (2m+1)\phi_{01mn}] , \quad (6.4c)$$

$$\hat{C}_{11}\phi_{11mn} = i\psi_{mn} , \quad (6.4d)$$

where \hat{C}_{11} (3.6) is the single generator of $\mathcal{A}_{\text{phys}}^{(*)}$. Hence Φ_0^{mod} is invariant under $\mathcal{A}_{\text{phys}}^{(*)}$. We show in appendix C that Φ_0^{mod} is dense in \mathcal{H}_{aux} .

We denote by Φ^{mod} the closure of Φ_0^{mod} under the algebra generated by $\{U(g) \mid g \in G\}$ and adopt Φ^{mod} as the test space. By construction, Φ^{mod} is invariant under $\mathcal{A}_{\text{phys}}^{(*)}$. From (5.8), (6.2) and (6.3) it follows that the integral in (5.5) converges in absolute value, and $(\cdot, \cdot)_{\text{ga}}$ is hence well defined. The evaluation of $(\cdot, \cdot)_{\text{ga}}$ for $l+j > 0$ proceeds

as in section 5, while $(\psi_{m'n'}, \psi_{mn})_{\text{ga}} = 0$ by an explicit computation using (5.9) and (6.2), in the index range where (5.9) is valid. Hence the formulas of section 5 hold for $l+j > 0$, while η sends the whole $l = j = 0$ sector of Φ^{mod} to zero.

\mathcal{H}_{RAQ} has dimension three. The representation of $\mathcal{A}_{\text{phys}}^{(*)}$ on \mathcal{H}_{RAQ} is trivial.

7 Discussion

In this paper we have discussed refined algebraic quantisation with group averaging in a constrained Hamiltonian system with unreduced phase space $\mathbb{R}^{2(p+q)}$, where $p \geq 1$ and $q \geq 1$, and a nonunimodular gauge group. The system was constructed by restricting the gauge transformations in a system with the unimodular gauge group $\text{SL}(2, \mathbb{R})$, previously analysed in [10, 14], to a two-dimensional nonunimodular subgroup G . We obtained a Hilbert space with a nontrivial representation of the distinguished $\mathfrak{o}(p, q)$ observable algebra for $(p, q) \neq (1, 1)$, which is precisely the condition for the classical reduced phase space to contain a symplectic manifold. The representation was found to be the end-point of the maximally degenerate principal unitary series [32], with the quadratic Casimir taking the value $-\frac{1}{4}(p+q-2)^2$.

By contrast, the reduced phase space of the $\text{SL}(2, \mathbb{R})$ system contains a symplectic manifold for $\min(p, q) \geq 2$, but group averaging in this system gives a Hilbert space with a nontrivial representation of the $\mathfrak{o}(p, q)$ observables only when $\min(p, q) \geq 2$ and $p+q \equiv 0 \pmod{2}$ [10, 14]. The quadratic Casimir in this representation takes the value $-\frac{1}{4}(p+q)(p+q-4)$. The difference in the Casimirs in the G system and the $\text{SL}(2, \mathbb{R})$ system arises from the different senses in which the physical states produced by group averaging satisfy the constraints for unimodular and nonunimodular gauge groups [8]. The physical states in the $\text{SL}(2, \mathbb{R})$ system are invariant under $\text{SL}(2, \mathbb{R})$, in a representation induced from the unitary representation on the auxiliary Hilbert space, while the physical states in the G system are not invariant under the corresponding representation of G .

As neither $\text{SL}(2, \mathbb{R})$ nor G is compact, convergence of the group averaging was an issue for both systems. Choosing the test space to be a suitable extension of the linear span of the harmonic oscillator eigenstates on \mathbb{R}^{p+q} guaranteed absolute convergence for $p+q > 4$ in the $\text{SL}(2, \mathbb{R})$ system and for $(p, q) \neq (1, 1)$ in the G system. The remaining values of p and q could be handled with minor modifications in the low angular momentum sectors of the test space, taking care to preserve the inclusion of the $\mathfrak{o}(p, q)$ observables in the physical observable algebra. In the $\text{SL}(2, \mathbb{R})$ system, it was straightforward to choose the extension so that the theorem of Giulini and Marolf [8] implied the uniqueness of the group averaging rigging map as the only rigging map admitted by the auxiliary Hilbert space, the representation of the gauge group and the test space. In our G system it was straightforward to choose the extension so that refined algebraic quantisation in the formulation of [8] could be applied and the $\mathfrak{o}(p, q)$ observables could be controlled, but we have not found a test space to which the uniqueness theorem of Giulini and Marolf would apply. For a nonunimodular gauge

group, the theorem assumes absolute convergence not just in the measure d_0g in which the actual averaging is performed, but in the whole family of measures $\{\Delta^{n/2}(g) d_0g \mid n \in \mathbb{Z}\}$, and for our test space this convergence fails for large $|n|$. Uniqueness of the rigging map for the G system remains thus an open question.

Finally, we discussed quantisation of the nonunimodular gauge system by algebraic methods, requiring the physical states to be strictly annihilated by the generators of the G -action and the classical $\mathfrak{o}(p, q)$ observables to become self-adjoint operators. The group averaging theory and the algebraic quantisation theory were found to be qualitatively fairly similar for $(p, q) = (1, 3)$, $(3, 1)$ and $(2, 2)$, but for $(p, q) = (1, 2)$ and $(2, 1)$, and for $p \geq 2$, $q \geq 2$ and $p + q \equiv 1 \pmod{2}$, algebraic quantisation produced no quantum theory, under certain technical assumptions that parallel those of the group averaging theory. It would be interesting to understand these phenomena from the $O(p, q)$ representation theory viewpoint. From a broader perspective, it would be interesting to explore whether similar phenomena arise when techniques akin to group averaging are extended to systems where the constraint algebra closes with nonconstant structure functions [13].

Acknowledgements

We thank Abhay Ashtekar, John Barrett, Nico Giulini, Eli Hawkins, Don Marolf and Jacek Wiśniewski for discussions and correspondence, and in particular Abhay Ashtekar for raising the question addressed in section 4. J.L. thanks Nico Giulini for hospitality funded by DFG project Ro 864/7-2 at the University of Freiburg. A.M. was supported by a CONACYT (Mexico) Postgraduate Fellowship and an ORS Award to the University of Nottingham.

A Appendix: Gauge group

In this appendix we collect some relevant properties of the gauge group G . The notation follows the $\mathrm{SL}(2, \mathbb{R})$ notation of [45].

G is the lower triangular subgroup of $\mathrm{SL}(2, \mathbb{R})$ with positive diagonal elements. G is two-dimensional, nonabelian and connected, which properties characterise G uniquely up to isomorphisms.

The Lie algebra \mathfrak{g} is spanned by the matrices

$$h := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad , \quad e^- := \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \quad , \quad (\text{A.1})$$

whose commutator is

$$[h, e^-] = -2e^- \quad . \quad (\text{A.2})$$

Elements of G can be written uniquely as

$$\begin{aligned} g &= \exp(\mu e^-) \exp(\lambda h) \\ &= \begin{pmatrix} 1 & 0 \\ \mu & 1 \end{pmatrix} \begin{pmatrix} e^\lambda & 0 \\ 0 & e^{-\lambda} \end{pmatrix}, \end{aligned} \quad (\text{A.3})$$

where $\mu \in \mathbb{R}$ and $\lambda \in \mathbb{R}$.

From (A.3) we have $g^{-1}dg = h d\lambda + e^- e^{2\lambda} d\mu$ and $dg g^{-1} = h d\lambda + e^-(d\mu + 2\mu d\lambda)$, and the left and right invariant Haar measures are respectively $d_L g = e^{2\lambda} d\lambda d\mu$ and $d_R g = d\lambda d\mu$.

The adjoint action of G on \mathfrak{g} reads $\text{Ad}_g(h) = ghg^{-1} = h + 2\mu e^-$, $\text{Ad}_g(e^-) = ge^-g^{-1} = e^{-2\lambda}e^-$. Hence the modular function [8] is $\Delta(g) := \det(\text{Ad}_g) = e^{-2\lambda}$. The symmetric measure, invariant under $g \mapsto g^{-1}$, is $d_0 g = [\Delta(g)]^{1/2} d_L g = [\Delta(g)]^{-1/2} d_R g = e^\lambda d\lambda d\mu$.

B Appendix: Separation of \mathcal{M}_{reg} by $\mathcal{A}_{\text{class}}$

In this appendix we verify the separation properties of $\mathcal{A}_{\text{class}}$ on \mathcal{M}_{reg} stated in subsection 2.3. We represent \mathcal{M}_{reg} as the set \mathcal{N}_{reg} defined by (2.10).

Theorem B.1 *Let $p > 1$ and $q > 1$, and let $\mathcal{N}_{\text{reg}}^+$ be the subset of \mathcal{N}_{reg} where $0 \neq \mathbf{p}^2 \neq \varpi^2 \neq 0$. Then $\mathcal{A}_{\text{class}}$ separates $\mathcal{N}_{\text{reg}}^+$ up to the twofold degeneracy (2.11).*

Proof. Let $a = (\mathbf{u}, \mathbf{p}, \mathbf{w}, \varpi) \in \mathcal{N}_{\text{reg}}^+$ and $b = (\mathbf{u}', \mathbf{p}', \mathbf{w}', \varpi') \in \mathcal{N}_{\text{reg}}^+$ such that $A(a) = A(b)$ for all $A \in \mathcal{A}_{\text{class}}$. We shall show that $a = \pm b$. We use the basis (2.6).

Using (2.10), the condition $A_{ij}(a) = A_{ij}(b)$ implies

$$\mathbf{u}' = \cos(\theta)\mathbf{u} + \sin(\theta)|\mathbf{p}|^{-1}\mathbf{p}, \quad (\text{B.1a})$$

$$\mathbf{p}' = \cos(\theta)\mathbf{p} - \sin(\theta)|\mathbf{p}|\mathbf{u}, \quad (\text{B.1b})$$

where $0 \leq \theta < 2\pi$. Similarly, the condition $B_{ij}(a) = B_{ij}(b)$ implies

$$\mathbf{w}' = \cos(\varphi)\mathbf{w} + \sin(\varphi)|\varpi|^{-1}\varpi, \quad (\text{B.2a})$$

$$\varpi' = \cos(\varphi)\varpi - \sin(\varphi)|\varpi|\mathbf{w}, \quad (\text{B.2b})$$

where $0 \leq \varphi < 2\pi$. With (B.1) and (B.2), the condition $C_{ij}(a) = C_{ij}(b)$ reads

$$\begin{aligned} 0 &= p_i w_j [\cos(\theta) \cos(\varphi) - 1 + \sin(\theta) \sin(\varphi) |\varpi| |\mathbf{p}|^{-1}] \\ &\quad - u_i \varpi_j [\cos(\theta) \cos(\varphi) - 1 + \sin(\theta) \sin(\varphi) |\mathbf{p}| |\varpi|^{-1}] \\ &\quad + u_i w_j [\cos(\theta) \sin(\varphi) |\varpi| - \sin(\theta) \cos(\varphi) |\mathbf{p}|] \\ &\quad + p_i \varpi_j [\cos(\theta) \sin(\varphi) |\varpi|^{-1} - \sin(\theta) \cos(\varphi) |\mathbf{p}|^{-1}]. \end{aligned} \quad (\text{B.3})$$

Contracting (B.3) with $u_i w_j$, $u_i \varpi_j$, $p_i w_j$ and $p_i \varpi_j$ shows that each of the four terms must vanish individually. Using $|\mathbf{p}| \neq |\varpi|$, elementary algebra shows that the only solutions are $\theta = \varphi = 0$ and $\theta = \varphi = \pi$. ■

Remark. The assumption $|\mathbf{p}| \neq |\varpi|$ is necessary, for otherwise (B.3) is satisfied by $\theta = \varphi$.

Theorem B.2 *Let $p = 1$ and $q > 1$ (respectively $p > 1$ and $q = 1$), and let $\mathcal{N}_{\text{reg}}^+$ be the subset of \mathcal{N}_{reg} where $\varpi^2 \neq 0$ ($\mathbf{p}^2 \neq 0$). Then $\mathcal{A}_{\text{class}}$ separates $\mathcal{N}_{\text{reg}}^+$ up to the twofold degeneracy (2.11).*

Proof. It suffices to consider $p = 1$ and $q > 1$.

Proceeding as in the proof of Theorem B.1, the condition $A_{ij}(a) = A_{ij}(b)$ is an identity, the condition $B_{ij}(a) = B_{ij}(b)$ leads to (B.2), and the condition $C_{ij}(a) = C_{ij}(b)$ reads $u_1 \varpi = u'_1 \varpi'$. As $(u'_1)^2 = (u_1)^2 = 1$, the result follows by contracting (B.2b) with ϖ . ■

C Appendix: Φ_0^{mod} is dense in \mathcal{H}_{aux}

Theorem C.1 *For $p = q = 1$, the test space Φ_0^{mod} defined in section 6 is dense in \mathcal{H}_{aux} .*

Proof. Let $\mathcal{H}_{\text{aux}}^{++} \subset \mathcal{H}_{\text{aux}}$ be the subspace in which the functions are even both in u_1 and in w_1 . By (6.3), it suffices to show that $\Phi_0^{\text{mod}++} := \text{span}\{\psi_{mn}\}$ is dense in $\mathcal{H}_{\text{aux}}^{++}$.

Let

$$\tilde{\phi}_{mn} := 2^{-(m+n)/2} (\pi m! n!)^{-1/2} H_m(u_1) H_n(w_1) \exp\left[-\frac{1}{2}(u_1^2 + w_1^2)\right] , \quad (\text{C.1})$$

where the H s are the Hermite polynomials [43]. $\{\tilde{\phi}_{mn}\}$ is an orthonormal basis of \mathcal{H}_{aux} , $\{\tilde{\phi}_{2m,2n}\}$ is an orthonormal basis of $\mathcal{H}_{\text{aux}}^{++}$, and the recursion relations of the Hermite polynomials imply

$$\hat{C}_{11} \tilde{\phi}_{mn} = i \left(-\sqrt{(m+1)(n+1)} \tilde{\phi}_{m+1,n+1} + \sqrt{mn} \tilde{\phi}_{m-1,n-1} \right) . \quad (\text{C.2})$$

As $\text{span}\{\phi_{11mn}\} = \text{span}\{\tilde{\phi}_{2m+1,2n+1}\}$, (6.4d) shows that $\Phi_0^{\text{mod}++} = \text{span}\{\hat{C}_{11} \tilde{\phi}_{2m+1,2n+1}\}$.

Suppose now that $\Phi_0^{\text{mod}++}$ is not dense in $\mathcal{H}_{\text{aux}}^{++}$. Then there exists a nonzero vector $y = \sum_{mn} a_{mn} \tilde{\phi}_{2m,2n} \in \mathcal{H}_{\text{aux}}^{++}$ that is orthogonal to each $\hat{C}_{11} \tilde{\phi}_{2m'+1,2n'+1}$. By (C.2), this implies

$$a_{m+1,n+1} = \sqrt{\frac{(m+\frac{1}{2})(n+\frac{1}{2})}{(m+1)(n+1)}} a_{mn} , \quad (\text{C.3})$$

from which it follows by elementary analysis that y has finite norm only if it is the zero vector. Hence $\Phi_0^{\text{mod}++}$ is dense in $\mathcal{H}_{\text{aux}}^{++}$, and Φ_0^{mod} is dense in \mathcal{H}_{aux} . ■

References

- [1] A. Higuchi, *Quantum linearization instabilities of de Sitter space-time 2*, Class. Quantum Grav. **8**, 1983 (1991).
- [2] N. P. Landsman, *Rieffel induction as generalized quantum Marsden-Weinstein reduction*, J. Geom. Phys. **15**, 285 (1995). [⟨arXiv:hep-th/9305088⟩](#)
- [3] D. Marolf, *Quantum observables and recollapsing dynamics*, Class. Quantum Grav. **12**, 1199 (1995). [⟨arXiv:gr-qc/9404053⟩](#)
- [4] A. Ashtekar, J. Lewandowski, D. Marolf, J. Mourão, and T. Thiemann, *Quantization of diffeomorphism invariant theories of connections with local degrees of freedom*, J. Math. Phys. **36**, 6456 (1995). [⟨arXiv:gr-qc/9504018⟩](#)
- [5] D. Marolf, *Refined algebraic quantization: Systems with a single constraint*, in: *Symplectic Singularities and Geometry of Gauge Fields* (Banach Center Publications, Polish Academy of Sciences, Institute of Mathematics, Warsaw, 1997). [⟨arXiv:gr-qc/9508015⟩](#)
- [6] N. P. Landsman, *Against the Wheeler-DeWitt equation*, Class. Quantum Grav. **12**, L119 (1995). [⟨arXiv:gr-qc/9510033⟩](#)
- [7] N. P. Landsman and K. K. Wren, *Constrained quantization and θ -angles*, Nucl. Phys. **B502**, 537 (1997). [⟨arXiv:hep-th/9706178⟩](#)
- [8] D. Giulini and D. Marolf, *A uniqueness theorem for constraint quantization*, Class. Quantum Grav. **16**, 2489 (1999). [⟨arXiv:gr-qc/9902045⟩](#)
- [9] A. Gomberoff and D. Marolf, *On group averaging for $SO(n, 1)$* , Int. J. Mod. Phys. D **8**, 519 (1999). [⟨arXiv:gr-qc/9902069⟩](#)
- [10] J. Louko and C. Rovelli, *Refined algebraic quantization in the oscillator representation of $SL(2, \mathbb{R})$* , J. Math. Phys. **41**, 132 (2000). [⟨arXiv:gr-qc/9907004⟩](#)
- [11] D. Giulini, *Group averaging and refined algebraic quantization*, Nucl. Phys. Proc. Suppl. **88**, 385 (2000). [⟨arXiv:gr-qc/0003040⟩](#)
- [12] D. Marolf, *Group averaging and refined algebraic quantization: Where are we now?*, in: *The Ninth Marcel Grossmann meeting : Proceedings*, edited by V. G. Gurzadyan, R. T. Jantzen and R. Ruffini (World Scientific, Singapore, 2002). [⟨arXiv:gr-qc/0011112⟩](#)
- [13] O. Yu. Shvedov, *On correspondence of BRST-BFV, Dirac and refined algebraic quantisations of constrained systems*, Ann. Phys. (N.Y.) **302**, 2 (2002). [⟨arXiv:hep-th/0111270⟩](#)

- [14] J. Louko and A. Molgado, *Group averaging in the (p, q) oscillator representation of $SL(2, \mathbb{R})$* , [⟨arXiv:gr-qc/0312014⟩](#).
- [15] D. Giulini and D. Marolf, *On the generality of refined algebraic quantization*, Class. Quantum Grav. **16**, 2479 (1999). [⟨arXiv:gr-qc/9812024⟩](#)
- [16] G. M. Tuynman, *Reduction, quantization, and nonunimodular groups*, J. Math. Phys. **31**, 83 (1990).
- [17] C. Duval, J. Elhadad, M. G. Gotay and G. M. Tuynman, *Nonunimodularity and the quantization of the pseudorigid body*, in: *Hamiltonian Systems, Transformation Groups and Spectral Transform Methods*, edited by J. Harnad and J. Marsden (Les Publications du CRM, 1990), pp 149–160.
- [18] C. Duval, J. Elhadad, M. G. Gotay, J. Śniatycki and G. M. Tuynman, *Quantization and bosonic BRST theory*, Ann. Phys. (N.Y.) **206**, 1 (1991).
- [19] P. Hájíček and K. V. Kuchař, *Constraint quantization of parametrized relativistic gauge systems in curved spacetimes*, Phys. Rev. D **41**, 1091 (1990).
- [20] M. Montesinos, C. Rovelli and T. Thiemann, *$SL(2, \mathbb{R})$ model with two Hamiltonian constraints*, Phys. Rev. D **60**, 044009 (1999). [⟨arXiv:gr-qc/9901073⟩](#)
- [21] M. Montesinos, *Relational evolution of the degrees of freedom of generally covariant quantum theories*, Gen. Rel. Grav. **33**, 1 (2001). [⟨arXiv:gr-qc/0002023⟩](#)
- [22] R. Gambini and R. A. Porto, *Relational time in generally covariant quantum systems: Four models*, Phys. Rev. D **63**, 105014 (2001). [⟨arXiv:gr-qc/0101057⟩](#)
- [23] M. Trunk, *An $SL(2, \mathbb{R})$ model of constrained systems: Algebraic constraint quantization*, University of Freiburg preprint THEP 99/3. [⟨arXiv:hep-th/9907056⟩](#)
- [24] M. Bojowald, H. A. Kastrup, F. Schramm, and T. Strobl, *Group theoretical quantization of a phase space $S^1 \times \mathbb{R}^+$ and the mass spectrum of Schwarzschild black holes in D space-time dimensions*, Phys. Rev. D **62**, 044026 (2000). [⟨arXiv:gr-qc/9906105⟩](#)
- [25] M. Bojowald and T. Strobl, *Group theoretical quantization and the example of a phase space $S^1 \times \mathbb{R}^+$* , J. Math. Phys. **41**, 2537 (2000). [⟨arXiv:quant-ph/9908079⟩](#)
- [26] H. A. Kastrup, *Quantization of the optical phase space $\mathcal{S}^2 = \{\varphi \bmod 2\pi, I > 0\}$ in terms of the group $SO^\uparrow(1, 2)$* , Fortsch. Phys. **51**, 975 (2003). [⟨arXiv:quant-ph/0307069⟩](#)
- [27] I. Bars and C. Kounnas, *Theories with Two Times*, Phys. Lett. **B402**, 25 (1997). [⟨arXiv:hep-th/9703060⟩](#)

- [28] I. Bars, *Two-Time Physics*, in: *Proceedings of the 22nd International Colloquium On Group Theoretical Methods In Physics*, edited by S. P. Corney, R. Delbourgo and P. D. Jarvis (International Press, Cambridge, Massachusetts, 1999), 2–17. [⟨arXiv:hep-th/9809034⟩](#).
- [29] I. Bars, *Survey of Two-Time Physics*, *Class. Quantum Grav.* **18**, 3113 (2001). [⟨arXiv:hep-th/0008164⟩](#)
- [30] I. Bars, *2T-Physics 2001*, in: *New developments in fundamental interaction theories: Proceedings of the 37th Karpacz Winter School Of Theoretical Physics* (AIP Conference Proceedings, Vol. 589), edited by J. Lukierski and J. Rembielinski (American Institute of Physics, Melville, NY, 2001), 18–30. [⟨arXiv:hep-th/0106021⟩](#).
- [31] B. Dittrich, private communication (2003).
- [32] N. Ja. Vilenkin and A. V. Klimyk, *Representation of Lie Groups and Special Functions*, Vol. 2 (Kluwer Academic Publishers, 1993).
- [33] A. Ashtekar, *Lectures on Non-Perturbative Canonical Gravity* (World Scientific, Singapore, 1991).
- [34] A. Ashtekar and R. S. Tate, *An algebraic extension of Dirac quantization: Examples*, *J. Math. Phys.* **35**, 6434 (1994). [⟨arXiv:gr-qc/9405073⟩](#)
- [35] P. A. M. Dirac, *Lectures on Quantum Mechanics* (Belfer Graduate School of Science, New York, 1964).
- [36] M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems* (Princeton University Press, Princeton, 1992).
- [37] J. A. Azcárraga, A. J. Macfarlane, A. J. Mountain and J. C. Pérez Bueno, *Invariant tensors for simple groups*, *Nucl. Phys. B* **510**, 657 (1998). [⟨arXiv:physics/9706006⟩](#)
- [38] V. Bargmann, *Irreducible unitary representations of the Lorentz group*, *Ann. Math.* **38**, 568 (1947).
- [39] S. Lang, *$SL_2(\mathbf{R})$* (Addison-Wesley, Reading, Massachusetts, 1975).
- [40] M. A. Naimark, *Linear Representations of the Lorentz Group* (Pergamon, Oxford, 1964).
- [41] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, *Higher Transcendental Functions, Vol. 2* (Bateman Manuscript project, McGraw-Hill, New York, 1953).

- [42] G. Arfken, *Mathematical Methods for Physicists*, second edition (Academic, New York, 1970).
- [43] W. Magnus, F. Oberhettinger, and R. P. Soni, *Formulas and Theorems for the Special Functions of Mathematical Physics*, 3rd ed. (Springer, Berlin, 1966).
- [44] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series and Products*, 4th edition (Academic, New York, 1980).
- [45] R. Howe and E. C. Tan, *Non-Abelian Harmonic Analysis: Applications of $SL(2, \mathbb{R})$* (Springer, New York, 1992).