Quantum Effect and Curvature Strength of Naked Singularities

Umpei MIYAMOTO,^{1,*)} Hideki MAEDA^{2,**)} and Tomohiro HARADA^{3,***)}

¹Department of Physics, Waseda University, Okubo 3-4-1, Shinjuku, Tokyo 169-8555, Japan

²Advanced Research Institute for Science and Engineering, Waseda University, Okubo 3-4-1, Shinjuku-ku, Tokyo 169-8555, Japan

³Astronomy Unit, School of Mathematical Sciences, Queen Mary, University of London, Mile End Road, London E1 4NS, UK

There are many solutions to Einstein field equations, which demonstrate naked singularity (NS) formation after regular evolution. If the cosmic censorship is to be recovered, one of the possibilities is that these NSs may disappear or be hidden behind horizons if quantum effects are taken into account. We investigate the relation between the curvature strength and the quantum effects of NSs in a very wide class of spherical dust collapse. Through a perturbative calculation, we find that if the NS is very strong, the quantum particle creation diverges as the Cauchy horizon is approached, while if the NS is very weak, the creation should be finite. In the context of the cosmic censorship, the strong NSs will be subjected by the backreaction of the quantum effects, while the weak ones will not.

§1. Introduction

The cosmic censorship hypothesis (CCH) presents one of the most important unsolved problems in general relativity.¹⁾ Its validity is often assumed in the analysis of physical phenomena in strong gravitational fields. There are two versions of this hypothesis. The weak hypothesis states that all singularities in gravitational collapse are hidden within black holes. This version implies the future predictability of the spacetime outside the event horizon. The strong one asserts that no singularities visible to any observer can exist. This version states that all physically reasonable spacetimes are globally hyperbolic. Despite several attempts neither general proof nor precise mathematical formulation of the hypothesis has been available so far. On the contrary, some solutions of the Einstein field equation with regular initial conditions evolving into spacetimes containing naked singularities (NSs) have been found.^{2),3)}

For a naked-singular spacetime to be a counterexample against the CCH, it is at least necessary that the Cauchy horizon (CH) be stable. Although the CCH was originally stated in the classical context, CHs may be unstable due to the backreaction of quantum effects such as particle creation, i.e., the particle creation may prevent NSs from forming. Research on such a possibility can be traced back to the pioneering works of Ford and Parker⁴⁾ and Hiscock, Williams, and Eardley.⁵⁾ Ford and Parker considered the particle creation during the formation of a shell-crossing

^{*)} Electronic address:umpei@gravity.phys.waseda.ac.jp

^{**)} Electronic address:hideki@gravity.phys.waseda.ac.jp

^{***)} Electronic address:T.Harada@qmul.ac.uk

NS to obtain a finite amount of flux.⁴⁾ On the other hand, Hiscock *et al.* considered the formation of a shell-focusing NS in the collapse of a null-dust fluid to obtain a diverging amount of flux.⁵⁾ Subsequently, such quantum phenomena have been studied in the models of a self-similar dust,⁶⁾⁻⁸⁾ a self-similar null dust,^{8),9)} and an analytic dust.¹⁰⁾ It is sometimes argued that the quantum radiation from a strong NS such as a shell-focusing one must diverge as the CH is approached,^{6),9)} although there is not enough evidence. See Ref. 11) for a recent review of the quantum and classical radiation processes during NS formation.

Recently, two of the present authors computed the quantum radiation in spherically symmetric self-similar collapse without specifying the collapsing matter.⁸⁾ It was found that in the generic class of self-similar spacetimes resulting in a NS, the luminosity of particle creation diverges as the inverse square of the remaining time until the CH. There was an another interesting result, leading us to the present study. In a class of self-similar spacetimes, which seem to be non-generic, the luminosity remains finite as the CH is approached. Indeed, it was shown that in the self-similar collapse of a massless scalar field, described by the Roberts solution,¹²⁾ the luminosity remains finite at the CH. We recognized that the results in Ref. 8) can be interpreted in terms of the curvature strength of the NS along the CH. Although NSs forming in generic spherically symmetric self-similar spacetimes are known to be strong¹³⁾ along the CH in the sense of Tipler,¹⁴⁾ the NS appearing in the Roberts solution is weak^{*)} even in the sense of Królak.¹⁶⁾ The relation between the curvature strength and quantum effect of NSs has already been suggested by Ref. 10). They showed the divergence of the quantum radiation during NS formation in the so-called analytic dust model, in which the forming NS is known to be weak,^{17),18)} weaker than that in the self-similar dust collapse. However, comprehensive understanding of the relation between the curvature strength and quantum effect of NSs is not available yet. The purpose here is to clarify how the amount of quantum radiation during the formation of NSs depends on such a nature of singularities as curvature strength. This analysis will help us obtain knowledge about the instability of CH, which would be predicted by a full semiclassical theory, taking into account the backreaction of quantum fields to gravity. In addition, it is clarified here how the coupling manner of quantized scalar fields to gravity changes the amount of quantum radiation. The dependence on the coupling manner is important because the CHs will suffer from the semiclassical instability, caused by all fundamental quantum fields.

The organization of this paper is as follows. In Sec. 2.1 we introduce a class of the Lemaître-Tolman-Bondi (LTB) solutions¹⁹⁾ that result in the formation of a NS. Then, the class of LTB solutions is divided into three sub-classes, depending on the curvature strength of NSs in Sec. 2.2. In Sec. 3 we calculate a map of null rays passing near the NS, which plays crucial roles in the estimate of quantum emission in the geometrical-optics approximation. In Sec. 4 the luminosity and total energy of emitted particles are computed. Section 5 is devoted to discussion. Throughout this

^{*)} Indeed, in the Roberts solution, the quantity of $R_{\mu\nu}k^{\mu}k^{\nu}$ vanishes along the CH, where $R_{\mu\nu}$ is the Ricci curvature tensor and k^{μ} is the tangent vector of the CH. This fact implies that the NS is not strong in the sense of Królak or Tipler.¹⁵

paper, the units of $c = G = \hbar = 1$ and the signature of (-, +, +, +) for spacetime metrics are used.

§2. Naked singularity in spherically symmetric dust collapse

2.1. Lemaître-Tolman-Bondi solutions admitting a naked singularity

The LTB solution,¹⁹⁾ which describes the collapse of a dust ball, in a comoving coordinate system is written as

$$ds^{2} = -dt^{2} + \frac{R'^{2}}{1 + f(r)}dr^{2} + R^{2}(t, r)d\Omega^{2},$$

$$\dot{R}^{2} = \frac{F(r)}{R} + f(r),$$

$$\rho = \frac{F'}{8\pi R^{2}R'},$$

(2.1)

where ρ is the energy density, $d\Omega^2$ is the line element of a unit two-dimensional sphere, and the prime and dot denote the partial derivatives with respect to r and t, respectively. Since we are concerned with the collapse of a dust fluid, we require $\dot{R} < 0$. Arbitrary functions F(r) and 1 + f(r) > 0 are twice the conserved Misner-Sharp mass and the specific energy, respectively. Hereafter we concentrate on the marginally bound case for simplicity, which is defined by f = 0. In such a case, Eq. (2.1) is integrated to give

$$R^{3} = \frac{9}{4} F(r) \left[t - t_{s}(r) \right]^{2}, \qquad (2.2)$$

where $t_s(r)$ is an arbitrary function of r. The time $t = t_s(r)$ corresponds to one when a dust shell at r meets the shell-focusing singularity, which is defined by R = 0. It is possible to choose as $t_s(r) = r$ by the scaling of r. LTB solutions can describe the formation of a shell-focusing NS from regular spacetimes. It has been shown that the shell-focusing singularity occurring at R = 0 with r > 0 is totally spacelike,²⁰⁾ and therefore our discussion will be confined to the singularity at (t, r) = (0, 0).

Here, we introduce a class of marginally bound LTB solutions in which the leading term of mass function F near the regular center takes the form of

$$F(r) = \frac{4\lambda^3}{9(\mu+1)^3} r^{3\mu+1} + o(r^{3\mu+1}), \qquad (2.3)$$

where $\mu \ (\geq 0)$ and $\lambda \ (> 0)$ are constants. The nontrivial form of factor and power in Eq. (2·3) is just for later convenience. In Appendix A, it is shown that this class of LTB spacetimes results in the formation of a shell-focusing NS, which could be globally naked and could violate the weak version of CCH. Radial null geodesics are described near the center by

$$\frac{dt}{dr} = \pm R' \simeq \pm \lambda r^{\mu} f(t/r), \qquad (2.4)$$

where

$$f(z) \equiv \left[1 - \frac{3\mu + 1}{3(\mu + 1)}z\right](1 - z)^{-1/3}.$$

Here, the upper and lower signs correspond to outgoing and ingoing null geodesics, respectively.

It will be helpful for later discussion to look into the initial regular density profile of the dust fluid near the regular center. The initial density profile at an initial time slice of $t = t_{in} < 0$ is written as,

$$\rho(t_{in}, r) = \frac{1}{6\pi t_{in}^2} \left[1 + 2\left(1 + \frac{F}{rF'}\right) \frac{r}{t_{in}} + O\left((r/t_{in})^2\right) \right].$$

Therefore, in the present case, the initial density profile in terms of physical radius $R\propto r^{\mu+1/3}$ takes the form of

$$\rho_{\rm in}(R) = \rho_0 + \rho_1 R^{\gamma} + \cdots, \qquad (2.5)$$

where

$$\gamma \equiv \frac{3}{3\mu+1}, \ \ \rho_0 \equiv \frac{1}{6\pi t_{in}^2}, \ \ \rho_1 \equiv -\frac{(3\mu+2)(\mu+1)^{3/(3\mu+1)}}{3(3\mu+1)\pi\lambda^{3/(3\mu+1)}(-t_{in})^{(9\mu+5)/(3\mu+1)}}. \ (2.6)$$

Note that the parameter γ is in the region of $0 < \gamma \leq 3$ for $\mu \geq 0$. In particular, the analytic and self-similar LTB models are the cases with $\gamma = 2$ ($\mu = 1/6$) and $\gamma = 3$ ($\mu = 0$), respectively.

2.2. Curvature strength of the naked singularities

The curvature strength of spacetime singularities is defined in a hope that weak convergence would imply the extendibility of the spacetime in a distributional sense. In this context, Tipler defined the *strong curvature condition* (SCC),¹⁴⁾ while Królak defined a weaker condition, which we call the *limiting focusing condition* (LFC).¹⁶⁾ The sufficient and necessary conditions for the singularities in spherically symmetric spacetimes with vanishing radial pressure satisfying the LFC or SCC are given in simple forms.¹⁸⁾ Here, we explain the results obtained in Ref. 18). If a singularity is naked, the relation between the circumferential radius R and the Misner-Sharp mass m is given by

$$R \simeq 2y_0 m^\beta \tag{2.7}$$

near the singularity along the null geodesics terminating at or emanating from the NS. The constant β is in the region of

$$1/3 < \beta \leqslant 1. \tag{2.8}$$

The constants y_0 and β are determined by requiring that there exist a positive finite limit of $y_0 \equiv \lim_{m\to 0} R/(2m^{\beta})$ along the null geodesics terminating at or emanating from the NS. If the geodesic satisfies the "gravity-dominance condition",¹⁸⁾ which

the null geodesic terminating at or emanating from the shell-focusing NS in LTB spacetime satisfies, the sufficient and necessary conditions are summarized as the following theorem:¹⁸⁾ if and only if $1/3 < \beta < 1/2$ is satisfied, neither the SCC nor the LFC holds; if and only if $1/2 \leq \beta < 1$ is satisfied, not the SCC but only the LFC holds; and if and only if $\beta = 1$, both the SCC and the LFC hold, for the radial null geodesic terminating at or emanating from the NS.

In the case of the LTB spacetimes we consider, from Eq. (A·2), $R \propto r^{\mu+1}$ holds along the outgoing null geodesic emanating from the NS, i.e., along the CH.^{*)} On the other hand, $m = F(r)/2 \propto r^{3\mu+1}$ holds along the outgoing null geodesic near the singularity. Therefore, the constant β is determined as

$$\beta = \frac{\mu + 1}{3\mu + 1} = \frac{2\gamma + 3}{9}.$$
(2.9)

With the above theorem by Harada *et al.*, the NS for $0 < \gamma < 3/4$ satisfy neither the LFC nor the SCC along the outgoing null geodesic emanating from the NS. The NS for $3/4 \leq \gamma < 3$ does not satisfy the SCC but does the LFC. The NS for $\gamma = 3$ satisfies both the LFC and the SCC. See also the Table I for the relation between γ and the curvature strength of NSs.

§3. Map of null rays passing near the naked singularity

3.1. Local map

The global map, $v = \mathcal{G}(u)$, is defined as the relation between the moments when a null ray leaves \mathcal{I}^- and when it terminates at \mathcal{I}^+ after passing through the regular center (see Fig. 1). The global map plays crucial roles in the estimate of quantum radiation in the geometrical-optics approximation. The global map cannot be obtained without solving the null geodesic equation from \mathcal{I}^- to \mathcal{I}^+ globally.

However, the main properties of the global map for NS formation will be determined by the behavior of null rays near the singularities, since the particle creation will be caused by the curvature around singularities. From this point of view, Tanaka and Singh proposed an alternative map, which we call the *local map*.²¹⁾ They considered an observer on a comoving shell who sends ingoing null rays. These null rays are reflected at the regular center, and come back to the same comoving observer. A radial null geodesic crosses a comoving shell located at a fixed comoving radius r before and after the reflection at the center. Thus these null rays define a map between the sending time and the receiving time measured by the proper time for the comoving observer. See Fig. 2 for a schematic illustration of the local map. The local map is expected to have the same structure as the global map because there are no singular features in the map between the proper time on a comoving shell at a finite distance and those measured by the null coordinates naturally defined at infinity.

^{*)} Although we assume that Eq. (A·2) is the asymptotic form of the CH, it is not trivial. Strictly speaking, it is necessary to show that Eq. (A·2) is the asymptotic form of the *first* outgoing null geodesic emanating from the NS. The analysis in the rest of this paper, however, does not depend on this assumption crucially.

This expectation has been confirmed for the self-similar dust^{8), 21)} and the analytic dust²¹⁾ models. Two of the present authors generalized the local map to a general class of self-similar spacetimes without specifying the collapsing matter, and the validity of such a local map was confirmed for the self-similar Vaidya model.⁸⁾ Therefore, one can safely assume that the local map has the same structure as the global map in the LTB spacetimes dealt with in this paper.

In this section, we calculate the local map by solving the radial null geodesic equation for the non-self-similar LTB spacetimes ($\mu > 0$). The local map for the self-similar LTB spacetime ($\mu = 0$) was obtained in Refs. 21) and 8).

3.2. Outline of the calculation of local map

To make the notation simple, we change the coordinates as

$$r' = \lambda^{1/\mu} r,$$

$$t' = \lambda^{1/\mu} t,$$

and we abbreviate r' and t' as r and t, respectively, hereafter in this section. Then, the null geodesic equation (2.4) becomes

$$\frac{dt}{dr} = \pm r^{\mu} f(t/r). \tag{3.1}$$

Before getting into a substantial calculation, it is convenient to summarize the calculation scheme for obtaining the local map, which we will implement in the following subsections. To obtain the local map, we have to obtain solutions of Eq. (3.1) near r = 0, which correspond to radial null rays passing through the center at $t = t(0) = -t_0$ ($t_0 > 0$) and then take the limit $t_0 \to 0$. We cannot expect, however, to have general exact solutions to this equation. Hence, here we adopt the following scheme to obtain the local map. First, we apply three different approximation regimes A, B and C, and find three kinds of expressions, $t = t^{A}(r)$, $t = t^{B}(r)$ and $t = t^{C}(r)$, respectively. Next we show that these three regimes have an overlapping region, where all three approximations are valid and the obtained approximate solutions can be matched with each other. Finally, we calculate the local map, that is, we calculate the relation between the sending time and the receiving time of the null ray at a comoving observer near the center.

In Sec. 3.3, we apply the approximation regime A, where $0 \leq r \ll t_0^{1/(1+\mu)}$ is satisfied. In this regime, we can deal with the center and then we can relate the ingoing null ray and the outgoing null ray both of which reach the center at the same time $t = t_0$. In Sec. 3.4, we apply the approximation regime B, where $t_0 \ll r \ll t_0^{1/(1+\mu)}$ is satisfied. This regime is possible only when $t_0 \ll 1$. Although regime B is completely included in regime A, regime B enables us to have an explicit expression for solutions and is therefore essential to obtain the local map. In Sec. 3.5, we apply the approximation regime C, where $t/r \ll 1$ is assumed. When we put $t/r = O(\eta)$ ($\eta \ll 1$), it turns out that the approximation is valid for $t_0/\eta \lesssim r \lesssim \eta^{1/\mu}$. Regimes A and B trivially have an overlapping region. When we take the limit $t_0 \to 0$, regime B and regime C have an overlapping region and regime C is still valid at a small but finite radius. In Sec. 3.6, we implement the matching between the approximate solutions $t = t^B(r)$ and $t = t^C(r)$ in the overlapping region. In Sec. 3.7, the local map is finally obtained for a comoving observer at a small but finite radius in the region for regime C. This is the generalization of Tanaka and Singh²¹'s calculation scheme.

3.3. Regime $A: 0 \leq r \ll t_0^{1/(1+\mu)}$

To find a null geodesic for $0\leqslant r\ll t_0^{1/(1+\mu)},$ we put

$$\frac{r}{t_0^{1/(\mu+1)}} = \epsilon \zeta, \tag{3.2}$$

$$\frac{r}{t_0} = \frac{\zeta}{\delta},\tag{3.3}$$

where $\epsilon \ll 1$ and $\zeta = O(1)$. δ is introduced for later convenience and can be arbitrarily large in regime A. From Eqs. (3.2) and (3.3), the following relations hold:

$$t_0 = \epsilon^{(\mu+1)/\mu} \delta^{(\mu+1)/\mu}, \tag{3.4}$$

$$r = \epsilon^{(\mu+1)/\mu} \delta^{1/\mu} \zeta. \tag{3.5}$$

The null geodesic $t = t^{A}(r)$ can be expanded by ϵ as follows:

$$t^{\mathcal{A}}(r) = -t_0 + \sum_{n=1}^{\infty} \epsilon^{(\mu+1)(n\mu+1)/\mu} t_n^{\mathcal{A}}(\zeta), \qquad (3.6)$$

where $t_n^{A}(\zeta)$ $(n = 1, 2, \dots)$ are the functions of ζ of order unity. Substituting Eq. (3.6) into Eq. (3.1), the following differential equations are obtained,

$$\frac{dt_1^{\mathcal{A}}}{d\zeta}(\zeta) = \pm \delta^{(\mu+1)/\mu} \zeta^{\mu} f(-\delta/\zeta), \qquad (3.7)$$

$$\frac{dt_2^{\rm A}}{d\zeta}(\zeta) = \pm \delta \zeta^{\mu-1} f'(-\delta/\zeta) t_1^{\rm A}(\zeta), \qquad (3.8)$$

$$\frac{dt_3^{\rm A}}{d\zeta}(\zeta) = \pm \frac{1}{2} \delta^{(\mu-1)/\mu} \zeta^{\mu-2} f''(-\delta/\zeta) (t_1^{\rm A})^2(\zeta) \pm \delta \zeta^{\mu-1} f'(-\delta/\zeta) t_2^{\rm A}(\zeta), \quad (3.9)$$

and so on, where f' and f'' denote the derivatives of f with respect to its argument. Since $t^{A}(0) = -t_0$, $t_n^{A}(0) = 0$ must be satisfied for $n \ge 1$. Equation (3.7) can be integrated immediately to give

$$t_1^{\mathcal{A}}(\zeta) = \pm \frac{1}{\mu+1} \delta^{(\mu+1)/\mu} \zeta^{\mu+1} (1+\delta/\zeta)^{2/3}.$$
 (3.10)

With $t_1^{\mathcal{A}}(\zeta)$ obtained above, Eq. (3.8) is integrated to give

$$t_2^{\rm A}(\zeta) = \frac{1}{\mu+1} \delta^{(2\mu+1)/\mu} \int_0^{\zeta} \hat{\zeta}^{2\mu} f'(-\delta/\hat{\zeta}) (1+\delta/\hat{\zeta})^{2/3} d\hat{\zeta}$$
(3.11)

$$= \delta^{(\mu+1)(2\mu+1)/\mu} I_2(\zeta/\delta), \tag{3.12}$$

where

$$I_2(y) \equiv \frac{1}{\mu+1} \int_0^y x^{2\mu} f'(-1/x)(1+1/x)^{2/3} dx.$$
 (3.13)

This integration cannot be expressed in terms of elementary functions. In a similar way, one can write $t_3^{\rm A}(\zeta)$ in an integral form,

$$t_{3}^{A}(\zeta) = \pm \int_{0}^{\zeta} \left[\frac{1}{2(\mu+1)^{2}} \delta^{(3\mu+1)/\mu} \hat{\zeta}^{3\mu} f''(-\delta/\hat{\zeta}) (1+\delta/\hat{\zeta})^{4/3} + \delta^{(2\mu^{2}+4\mu+1)/\mu} \hat{\zeta}^{\mu-1} f'(-\delta/\hat{\zeta}) I_{2}(\hat{\zeta}/\delta) \right] d\hat{\zeta}$$

= $\pm \delta^{(\mu+1)(3\mu+1)/\mu} I_{3}(\zeta/\delta),$

where

$$I_3(y) \equiv \int_0^y \left[\frac{1}{2(\mu+1)^2} x^{3\mu} f''(-1/x)(1+1/x)^{4/3} + x^{\mu-1} f'(-1/x) I_2(x) \right] dx. \quad (3.14)$$

It should be noted that we can safely take the limit $r \to 0$ in this regime because we do not assume δ is small.

3.4. Regime B: $t_0 \ll r \ll t_0^{1/(1+\mu)}$

For the approximation regime B, we will additionally assume that $\delta \ll 1$, which means $t_0 \ll r$. This also requires that $t_0 \ll 1$ from Eq. (3.4). The approximate solution $t = t^{\rm B}(r)$ can be obtained by approximating $t = t^{\rm A}(r)$ with $\delta \ll 1$. Hence, we define $t_n^{\rm B}(\zeta)$ as the function which is obtained by approximating $t_n^{\rm A}(\zeta)$ with $\delta \ll 1$. Thus, we have

$$t_1^{\rm B} = \pm \frac{1}{\mu+1} \delta^{(\mu+1)/\mu} \zeta^{\mu+1} \left[1 + \frac{2}{3} \zeta^{-1} \delta + \mathcal{O}(\delta^2) \right].$$
(3.15)

The approximate form of t_2^A for $\delta \ll 1$ can be obtained by using the asymptotic form of $I_2(y)$ for large y in Eq. (3.13) given by

$$I_2(y) = C_2 - \frac{2\mu}{3(\mu+1)^2(2\mu+1)} y^{2\mu+1} \left[1 + O(1/y)\right],$$

where C_2 is a constant. Except for the term of C_2 , all other terms are completely determined by integrating the expanded integrand. Therefore, $t_2^{\rm B}(\zeta)$ is obtained as

$$t_2^{\rm B}(\zeta) = C_2 \delta^{(\mu+1)(2\mu+1)/\mu} - \frac{2\mu}{3(\mu+1)^2(2\mu+1)} \delta^{(2\mu+1)/\mu} \zeta^{2\mu+1} \left[1 + \mathcal{O}(\delta)\right].$$
(3.16)

Similarly, the asymptotic form of I_3 is given by

$$I_3(y) = C_3 + \left[\frac{2\mu^2 + \mu + 1}{9(\mu + 1)^3(2\mu + 1)(3\mu + 1)}y^{3\mu + 1} - \frac{2C_2}{3(\mu + 1)}y^{\mu}\right] \left[1 + O(1/y)\right].$$

and hence $t_3^{\rm B}(\zeta)$ is

$$t_{3}^{\mathrm{B}}(\zeta) = \pm C_{3} \delta^{(\mu+1)(3\mu+1)/\mu} \pm \frac{2\mu^{2} + \mu + 1}{9(\mu+1)^{3}(2\mu+1)(3\mu+1)} \delta^{(3\mu+1)/\mu} \zeta^{3\mu+1} [1 + \mathrm{O}(\delta)] \\ \mp \frac{2C_{2}}{3(\mu+1)} \delta^{(2\mu^{2}+4\mu+1)/\mu} \zeta^{\mu} [1 + \mathrm{O}(\delta)] . \quad (3.17)$$

Then, an approximate solution is obtained by Eqs. $(3\cdot4)$ - $(3\cdot6)$ and $(3\cdot15)$ - $(3\cdot17)$ as

$$t^{\mathrm{B}}(r) = -t_{0} + C_{2}t_{0}^{2\mu+1} \pm C_{3}t_{0}^{3\mu+1} + \left[\pm\frac{1}{\mu+1}r^{\mu+1} - \frac{2\mu}{3(\mu+1)^{2}(2\mu+1)}r^{2\mu+1} \pm \frac{2\mu^{2}+\mu+1}{9(\mu+1)^{3}(2\mu+1)(3\mu+1)}r^{3\mu+1}\right] \times [1 + \mathcal{O}(\delta)]. \quad (3.18)$$

For later use, we have obtained the explicit expression up to the third order. It is straightforward to compute higher orders. It should be noted that we cannot take the limit $r \to 0$ in this expression (3.18).

3.5. Regime C: $t/r \ll 1$

Suppose $t/r \ll 1$. Here, we introduce $\eta \ll 1$, which controls the order of t/r, i.e., $t/r = O(\eta)$. In this approximation regime, we can expand f by t/r on the right hand side of Eq. (3.1) and obtain the expanded form of solutions.

Let us consider the critical outgoing and ingoing null geodesics $t = t_{\pm}^{\text{crit}}(r)$, which emanate from and terminate at the NS, i.e., t = r = 0. $t = t_{\pm}^{\text{crit}}(r)$ gives CH by definition. If we assume $t/r = O(\eta)$, they take the following form:

$$t_{\pm}^{\text{crit}}(r) = \pm \frac{1}{\mu+1} r^{\mu+1} - \frac{2\mu}{3(\mu+1)^2(2\mu+1)} r^{2\mu+1} \\ \pm \frac{2\mu^2 + \mu + 1}{9(\mu+1)^3(2\mu+1)(3\mu+1)} r^{3\mu+1} + \mathcal{O}(\eta^4), \quad (3.19)$$

where the upper (lower) sign corresponds to emanating (terminating) null ray. For $0 \leq r \leq \eta^{1/\mu}$, $t/r = O(\eta)$ is satisfied on the critical null rays and this approximation is justified. That is, the approximation regime C is valid for $0 \leq r \leq \eta^{1/\mu}$.

However, we are interested in null rays which are slightly earlier than these critical null rays. We expand the solution as follows:

$$t = t^{\rm C}(r) = \sum_{n=1}^{\infty} t_n^{\rm C}(r),$$
 (3.20)

where we assume $t_n^{\rm C}(r)/r = O(\eta^n)$. Substituting Eq. (3.20) into Eq. (3.1) and expanding the right hand side, the following differential equations are obtained:

$$\frac{dt_1^{\rm C}}{dr} = \pm r^{\mu},\tag{3.21}$$

$$\frac{dt_2^{\rm C}}{dr} = \mp \frac{2\mu}{3(\mu+1)} r^{\mu-1} t_1^{\rm C}(r), \qquad (3.22)$$

U. Miyamoto, H. Maeda and T. Harada

$$\frac{dt_3^{\rm C}}{dr} = \mp \frac{2\mu}{3(\mu+1)} r^{\mu-1} t_2^{\rm C}(r) \mp \frac{\mu-1}{9(\mu+1)} r^{\mu-2} (t_1^{\rm C})^2(r).$$
(3.23)

These equations are integrated to yield

$$t_1^{\mathcal{C}}(r) = D_{\pm} \pm \frac{1}{\mu+1} r^{\mu+1}, \qquad (3.24)$$

$$t_2^{\rm C}(r) = \mp \frac{2}{3(\mu+1)} D_{\pm} r^{\mu} - \frac{2\mu}{3(\mu+1)^2(2\mu+1)} r^{2\mu+1}, \qquad (3.25)$$

$$t_{3}^{C}(r) = \mp \frac{1}{9(\mu+1)} D_{\pm}^{2} r^{\mu-1} - \frac{3\mu-1}{9\mu(\mu+1)^{2}} D_{\pm} r^{2\mu} \\ \pm \frac{2\mu^{2}+\mu+1}{9(\mu+1)^{3}(2\mu+1)(3\mu+1)} r^{3\mu+1}, \quad (3.26)$$

where D_{\pm} is an integration constant which appears in the integration of Eq. (3.21). Integration constants for Eqs. (3.22) and (3.23) are set to be zero. From Eqs. (3.20), (3.24)-(3.26), the solution takes the following form:

$$t^{C}(r) = D_{\pm}$$

$$\pm \frac{1}{\mu+1}r^{\mu+1}$$

$$\mp \frac{2}{3(\mu+1)}D_{\pm}r^{\mu} - \frac{2\mu}{3(\mu+1)^{2}(2\mu+1)}r^{2\mu+1}$$

$$\mp \frac{1}{9(\mu+1)}D_{\pm}^{2}r^{\mu-1} - \frac{3\mu-1}{9\mu(\mu+1)^{2}}D_{\pm}r^{2\mu} \pm \frac{2\mu^{2}+\mu+1}{9(\mu+1)^{3}(2\mu+1)(3\mu+1)}r^{3\mu+1}$$

$$+ O(\eta^{4}).$$
(3.27)

Now, we can examine in what region the solution (3.27) is valid. To justify the expansion (3.20), $t_n^{\rm C}(r)/r = O(\eta^n)$ must be satisfied. This is the case if

$$\max\left(r^{\mu}, \frac{D_{\pm}}{r}\right) = \mathcal{O}(\eta).$$

If D_{\pm} is sufficiently small, the above condition implies

$$D_{\pm}/\eta \lesssim r \lesssim \eta^{1/\mu}.$$

3.6. Matching the approximation regimes

Since we are interested in the null rays which are close to the critical null geodesics, we take the limit $t_0 \to 0$. The region for the approximation regime B is completely included in that for regime A and the matching is trivially implemented. If we assume $D_{\pm} = O(t_0)$, the region where regime C is valid must have an overlap with the region for regime B if $t_0 \ll \eta^{(1+\mu)/\mu}$. This means that we can relate the integration constants D_{\pm} which appear in the regime C solution $t = t^{\rm C}(r)$ to t_0 which appear in the regime B solution $t = t^{\rm B}(r)$ by matching these two solutions in the overlapping region.

10

Different expressions $t = t^{B}(r)$ and $t = t^{C}(r)$ for the null geodesic are obtained independently in Secs. 3.4 and 3.5. We can see the solution $t = t^{B}(r)$ given by Eq. (3.18) coincide with the solution $t = t^{C}(r)$ given by Eq. (3.27) in several lowest orders if the constant terms satisfy the following relation:

$$D_{\pm} \simeq -t_0 + C_2 t_0^{2\mu+1} \pm C_3 t_0^{3\mu+1}. \tag{3.28}$$

This justifies the assumption that D_{\pm} is sufficiently small and $D_{\pm} = O(t_0)$ in the limit $t_0 \to 0$. This also implies that the approximation regime C is valid for $t_0/\eta \lesssim r \lesssim \eta^{1/\mu}$.

3.7. Obtaining the local map

To obtain the local map for this spacetime, we consider a comoving observer at $r = r_0$, where r_0 satisfies $r_0 \leq \eta^{1/\mu}$. As time proceeds, this observer approaches the ingoing critical null ray $t = t_{-}^{\text{crit}}(r)$ and therefore enters the region $t_0/\eta \leq r \leq \eta^{1/\mu}$, where the approximation regime C is valid. Then, $t_{\pm}(r_0) = t_{\pm}^{C}(r_0)$ are regarded as the sending time and receiving time of the null geodesic, where the sign \pm is introduced to distinguish the outgoing and ingoing null rays. From Eq. (3.27), the constant D_{\pm} is specified as

$$-D_{\pm} = t_{\pm}^{\text{crit}}(r_0) - t_{\pm}(r_0) + \mathcal{O}(t_0\eta),$$

where $t_{\pm}^{\text{crit}}(r_0)$ is the moment when the observer crosses the null geodesic (3.19). Therefore, if we take η to be sufficiently small, D_- (D_+) is interpreted as the time difference between the moments when the observer emits (receives) the null ray and crosses the null ray terminating at (emanating from) the NS. See also Fig. 2.

Since we consider a set of an ingoing null ray which reaches the center at $t = t_0$ and an outgoing null ray which can be regarded as a reflected ray of the former at $t = t_0$, we pick up both the ingoing and outgoing null rays with the common value for t_0 . When we eliminate t_0 from Eq. (3.28) for both signs, we have the relation between D_+ and D_- as

$$D_{-} \simeq D_{+} - 2C_3(-D_{+})^{3\mu+1},$$

which can be rewritten in terms of the sending and receiving times $t_{\pm}(r_0)$ at $r = r_0$ for sufficiently small r_0 as

$$t_{-}^{\rm C}(r_0) \simeq t_{-}^{\rm crit}(r_0) - \left[t_{+}^{\rm crit}(r_0) - t_{+}^{\rm C}(r_0)\right] - 2C_3 \left[t_{+}^{\rm crit}(r_0) - t_{+}^{\rm C}(r_0)\right]^{3\mu+1}.$$
 (3·29)

Equation (3.29) is the very local map, relating the moments when the comoving observer locating at $r = r_0$ sends the ingoing null ray and receives the reflected outgoing null rays. If we revive λ , the final result becomes

$$t_{-}^{\rm C}(r_0) \simeq t_{-}^{\rm crit}(r_0) - \left[t_{+}^{\rm crit}(r_0) - t_{+}^{\rm C}(r_0)\right] - 2C_3\lambda^3 \left[t_{+}^{\rm crit}(r_0) - t_{+}^{\rm C}(r_0)\right]^{3\mu+1}.$$
 (3·30)

§4. Luminosity and energy of particle creation

We consider the massless scalar fields coupled to the scalar curvature as

$$\left(\Box - \xi R\right)\phi = 0,\tag{4.1}$$

where R is the Ricci scalar curvature and ξ is an arbitrary real constant. In particular, the scalar fields with $\xi = 0$ and $\xi = 1/6$ are minimally and conformally coupled ones, respectively. The stress-energy tensor of the scalar field in an asymptotically flat region is given by

$$T^{(\xi)}_{\mu\nu} \simeq \nabla_{\mu}\phi\nabla_{\nu}\phi - \frac{1}{2}g_{\mu\nu}\nabla^{\alpha}\phi\nabla_{\alpha}\phi - \xi\nabla_{\mu}\nabla_{\nu}\phi^{2} + \xi g_{\mu\nu}\Box\phi^{2}.$$

To calculate the luminosity of emitted particles we need an estimate of the vacuumexpectation value of the above stress-energy tensor. A suitable regularization is required in the calculation of vacuum-expectation value because the stress-energy tensor is quadratic in fields at the same point. The regularization for minimally and conformally coupled scalar fields was given in Ref. 4) via the point-splitting regularization scheme. One can easily generalize the point-splitting scheme in Ref. 4) to the above more general scalar fields to give

$$L_{lm}^{(\xi)}(u) = \frac{1}{4\pi} \left[\left(\frac{1}{4} - \xi \right) \left(\frac{\mathcal{G}''}{\mathcal{G}'} \right)^2 + \left(\xi - \frac{1}{6} \right) \frac{\mathcal{G}'''}{\mathcal{G}'} \right], \tag{4.2}$$

where the prime denotes the differentiation with respect to u, and $l (= 0, 1, \dots)$ and $m (= 0, \pm 1, \dots, \pm l)$ are angular momentum and magnetic quantum numbers, respectively. The luminosity of quantum emission is the sum of all these modes, but it diverges because the above luminosity (4·2) is independent of (l, m). Such divergence is due to the neglect of the back-scattering by the potential barrier, which will reduce the emission for highly rotational modes. Hereafter, we shall omit the quantum numbers (l, m) and one should keep in mind that the above expression holds only for small l. The total energy of emitted particles is estimated by integrating the luminosity with respect to u,

$$E^{(\xi)}(u) \equiv \int_{-\infty}^{u} L^{(\xi)}(u') du'.$$

4.1. Quantum radiation in the non-self-similar LTB spacetimes: $0 < \gamma < 3$

As described in Sec. 3, we assume that the local map and the global map have the same structure. It means that from Eq. $(3\cdot30)$, the asymptotic form of the global map will take the form of

$$\mathcal{G}(u) \simeq v_0 - A(u_0 - u) - Ag\lambda^3 (u_0 - u)^{3\mu + 1},$$

where $u = u_0$ and $v = v_0$ are the CH and the ingoing null ray that terminates at the NS, respectively. A and g are constants. Noting that $\mu = (3 - \gamma)/(3\gamma)$, one can calculate the luminosity of particle creation as

$$L^{(\xi)} \simeq \left(\xi - \frac{1}{6}\right) \frac{3(3-\gamma)(3-2\gamma)}{4\pi\gamma^3} g\omega_s^{(3-\gamma)/\gamma} (u_0 - u)^{-3+3/\gamma} \\ + \left[\xi - \frac{7\gamma - 15}{36(\gamma - 2)}\right] \frac{27(\gamma - 2)(3-\gamma)}{4\pi\gamma^4} g^2 \omega_s^{2(3-\gamma)/\gamma} (u_0 - u)^{-4+6/\gamma}, \quad (4.3)$$

where we have defined the "frequency" of the NSs as

$$\omega_s \equiv \lambda^{1/\mu} = \lambda^{3\gamma/(3-\gamma)}. \tag{4.4}$$

In Appendix B, we show the frequency (4·4) is equivalent to the one defined in Ref. 10). One can easily recognize that depending on whether the scalar field couples to the scalar curvature conformally ($\xi = 1/6$) or not ($\xi \neq 1/6$), the leading term of the luminosity changes. Let us examine the time dependence of the luminosity and total energy of emitted particles into detail for the cases of $\xi \neq 1/6$ and $\xi = 1/6$ in order.

In the case of $\xi \neq 1/6$, the first term in Eq. (4.3) dominates except for the special case of $\gamma = 3/2$. In the case of $0 < \gamma \leq 1$, the leading term vanishes as $u \to u_0$ as the CH is approached. On the other hand, in the case of $1 < \gamma < 3$ and $\gamma \neq 3/2$, the luminosity diverges as the CH is approached as a negative power of the remaining time until the CH. The special case of $\gamma = 3/2$, in which the factor of the first term in Eq. (4.3) vanishes, is divided into two sub-cases depending on whether $\xi \neq 1/4$ or $\xi = 1/4$. If $\xi \neq 1/4$, the second term in Eq. (4.3) survives to be a finite constant. On the other hand, if $\xi = 1/4$, the factor of the second term also vanishes. Therefore, higher order terms contribute to the luminosity, which turn out to be finite. Therefore, in the both sub-cases of $\gamma = 3/2$, the luminosity remains finite at the CH. Now, let us examine the total energy of emitted particles. In the case of $3/2 < \gamma < 3$, the leading term in the energy is

$$E^{(\xi)} \simeq -\left(\xi - \frac{1}{6}\right) \frac{3(3-\gamma)}{4\pi\gamma^2} g\omega_s^{(3-\gamma)/\gamma} (u_0 - u)^{-2+3/\gamma}, \qquad (4.5)$$

so that the total energy of emitted particles diverges as the CH is approached. In particular, in the case of analytic LTB solution ($\gamma = 2$), the energy diverges as $(u_0 - u)^{-1/2}$, which coincides with the result in Ref. 10). In the case of $0 < \gamma < 3/2$, the total energy remains finite as the CH is approached. In the special case of $\gamma = 3/2$, the energy also remains finite. See also Table I(a).

In the case of the conformally coupled scalar field, which is defined by Eq. (4.1) with $\xi = 1/6$, the second term in Eq. (4.3) dominates. It is found that in the case of $0 < \gamma \leq 3/2$, the luminosity remains finite at most. While, in the case of $3/2 < \gamma < 3$, the luminosity diverges due to the negative power of the remaining time until the CH. Let us examine the total energy of the emitted particles. In the case of $2 < \gamma < 3$,

$$E^{(1/6)} \simeq \frac{(3-\gamma)^2}{16\pi\gamma^3(\gamma-2)} g^2 \omega_s^{2(3-\gamma)/\gamma} (u_0 - u)^{-3+6/\gamma}, \qquad (4.6)$$

which diverges as the CH is approached. In the case of $\gamma = 2$,

$$E^{(1/6)} \simeq \frac{3}{256\pi} g^2 \omega_s \ln[\omega_s(u_0 - u)]^{-1}, \qquad (4.7)$$

so that the energy diverges logarithmically, which coincides with the result in Ref. 10) again. On the other hand, if $0 < \gamma < 2$ the energy remains finite at most. See also Table I(b).

It follows from what has been seen that the radiation by the conformally coupled scalar field is milder than that of the non-conformally coupled scalar fields for a given value of γ . Such a consequence would result from the fact that the coupling of the conformal scalar field to gravity is weaker than that of the other scalar fields. It has been known that a conformally coupled scalar field must have non-zero and finite mass to be created in the early universe, while non-conformally coupled scalar particles are created regardless of their mass.²²

4.2. Quantum radiation in the self-similar LTB spacetime: $\gamma = 3$

The global map for the self-similar LTB spacetime ending in NS formation was calculated analytically in Ref. 6). Then, its main property was re-produced with the local-map method.^{8,21} The global map for the null rays passing near the CH is given by

$$\mathcal{G}(u) \simeq v_0 - B(u_0 - u)^{\alpha} \left[1 + q(u_0 - u) + O\left((u_0 - u)^2 \right) \right], \tag{4.8}$$

where α , B, and q are constants. The terms in the square brackets in Eq. (4.8) is an analytic function of $(u_0 - u)$,^{*)} The constant α depends only on the parameter λ in Eq. (2.3) and is shown to be greater than unity for the region of λ in which the singularity is naked.⁶⁾ Using the global map (4.8), we can compute the luminosity and energy of the particle creation as

$$L^{(\xi)} \simeq \frac{(\alpha - 1)(\alpha + 1 - 12\xi)}{48\pi} (u_0 - u)^{-2} + \frac{\alpha^2 - 1}{24\pi\alpha} q(u_0 - u)^{-1}, \qquad (4.9)$$

$$E^{(\xi)} \simeq \frac{(\alpha - 1)(\alpha + 1 - 12\xi)}{48\pi} (u_0 - u)^{-1} + \frac{\alpha^2 - 1}{24\pi\alpha} q \ln q^{-1} (u_0 - u)^{-1}.$$
 (4.10)

Each first term in Eqs. (4.9) and (4.10) dominates except for the special case of $\xi = (\alpha + 1)/12$. Therefore, the luminosity and energy generically diverge like the inverse square and the inverse of the remaining time to the CH, respectively. On the other hand, in the special case of $\xi = (\alpha + 1)/12$, each second term in Eqs. (4.9) and (4.10) dominates so that the power and energy diverge inversely and logarithmically, respectively. This case, however, should be regarded as an accidental cancellation in the sense that the coupling constant ξ happens to be a special value, which is determined by the detail of the collapse, α . See also Tables I(a) and I(b).

§5. Discussion

In this paper we have considered particle creation during the formation of shellfocusing NSs in the wide class of spherical dust collapse, which is described by the

^{*)} In previous works on particle creation during NS formation in the self-similar LTB spacetime, only the constant term in the square bracket in Eq. (4.8) was considered,^{6,8,21} which was sufficient to obtaining results. It is easy to calculate the higher order terms by the local-map method and to show that they constitute an analytic function near the CH. It is possible, however, that the emergence of the scale, q, in Eq. (4.8) from such a scale invariant spacetime as self-similar LTB solution indicates the breakdown of the local-map method. What extent the local-map method is valid leaves room for discussion.

marginally bound LTB solutions. Each solution has different initial density profile, and the resulting NSs have a variety of curvature strength along the CHs. The luminosity and energy of particle creation were estimated for each LTB solution and each scalar field that couples to scalar curvature in the linear form.

From the results, which are summarized in Tables I(a) and I(b), it is found that following statements hold for the generic naked-singular LTB spacetimes: the SCC along the CH is a sufficient condition for the luminosity and energy of the created scalar particles to diverge as the CH is approached; while, not to satisfy the LFC is a sufficient condition for the luminosity and energy to be finite at the CH; if the NS does not satisfy the SCC but does the LFC, the luminosity and energy can be either divergent or finite. It should be stressed that we only considered the dust collapse for simplicity; however, the above statements as to the curvature strength and the quantum radiation would be independent of the collapsing matter because the particle creation is a purely kinematic phenomenon and not directly related to the Einstein field equation. Therefore, we conjecture that the above statements hold for spherically symmetric collapsing spacetimes with any kinds of collapsing matter. Here, we have to mention also the coupling manner of scalar fields to gravity. It is found that the quantum radiation due to the conformally coupled scalar field is less than that of other scalar fields, including the minimally coupled scalar field. It is found, however, that the dependence of the amount of radiation on the coupling manner is not so drastic as to modify the above statements. Having observed the results and noticed the neglect of the backreaction in the analysis, one can then go on to consider their implication for the CCH. In general, it is difficult to assert rigidly that diverging quantum radiation prevents NSs from forming and recovers the cosmic censor because no semiclassical formalism taking into account the backreaction completely is available. It is possible, however, that the strong quantum field causes the instability of NS formation. If we interpret the results from this point of view, the formation of the NSs of which curvature strength is strong (weak) tends to (not to) be affected by the quantum effect. In particular, the result that the quantum radiation during the formation of weak NSs remains finite is novel and striking from the viewpoint of the CCH. It suggests that the weak NSs need another mechanism if they are to hide behind the event horizons.

The analysis was based on three assumptions: the validity of the local-map method, geometrical-optics approximation, and quantum field theory in curved spacetime. The validity of each approximation seems to leave room for discussion. In particular, quantum field theory in curved spacetime should be crucial.^{7),8),23)} Since the curvature radius in the collapsing dust ball decreases arbitrarily even below the scale of quantum gravity before the appearance of a CH, the radiation will be cut off by quantum gravitational effects. It had been shown that the total energy of emitted particles within this phase will only amount to the Planck energy in the dust model²³⁾ and that the backreaction would not affect the appearance of NS in the two-dimensional self-similar dust model.⁷⁾ Recently, the possibility was discussed in Ref. 8) that the radiated energy within the semiclassical phase in self-similar spacetimes can be arbitrarily large when the CH and the event horizon are very close. However, it remains to be shown whether or not such a mechanism is effective in non-self-similar spacetimes.

We do not find a necessary and sufficient condition on the curvature strength of NSs for the quantum radiation to be divergent or finite. We believe that a new definition of strength of (naked) singularities should be proposed from the viewpoint of the behavior of quantum fields on spacetimes rather than the viewpoint of the behavior of classical particles. Such a philosophy can be seen also in the wave-probe approach to NSs,²⁴ which is based on the theory of dynamics in non-globally hyperbolic spacetimes developed first by Wald.²⁵ On this point, there is room for further investigation.

Acknowledgements

We are grateful to Kei-ichi Maeda for his continuous encouragement. U.M. thanks L.H. Ford for a helpful comment on the coupling of scalar fields and thanks also T.P. Singh and D.A. Konkowski for useful comments on the Cauchy horizon instability. Thanks are due to A. Ishibashi for thoughtful comments on the implication and application of the results. H.M. was supported by a Grant for The 21st Century COE Program (Holistic Research and Education Center for Physics Self-Organization Systems) at Waseda University. T.H. was supported from JSPS.

Appendix A

— Nakedness of the singularity —

In order to determine whether the singularity is naked or not, we investigate the future-directed outgoing null geodesics emanating from the singularity at (t, r) = (0, 0). We find the asymptotic solutions that obey a power law near the center²⁶ as

$$t \simeq X_0 r^p, \tag{A.1}$$

where $X_0 > 0$ and $p \ge 1$ are constants. The latter condition is due to the fact that the orbit of the shell-focusing singularity is $t = t_s(r) = r$. After some straightforward calculations, one can find an asymptotic solution for $\mu > 0$ as

$$t \simeq \frac{\lambda}{\mu+1} r^{\mu+1}.\tag{A.2}$$

With Eq. (A·2) and the fact that the apparent horizon, which is defined by F = R, behaves as $t = t_{ah}(r) = r - 2F(r)/3 \simeq r$ for $\mu > 0$ near the center, the singularity is at least locally naked. In the case of the self-similar case ($\mu = 0$), similar discussion is possible and the singularity is known to be naked for small values of λ .²⁶⁾ We consider the situation in which the collapsing dust ball is attached to an outer vacuum region at a comoving radius r = constant, within which the null ray (A·2) is outside the apparent horizon. Then, the singularity is globally naked and the weak version of CCH is violated.

Appendix B

— Frequency of the singularity —

It will be helpful to compare the gauge used in this paper with one used in much literature. Such a comparison shows that ω_s defined in Sec. 4 coincides with the characteristic frequency of singularity introduced in Ref. 10) except for a numerical factor.

Let us denote the comoving coordinates by (\tilde{t}, \tilde{r}) , in which \tilde{r} is chosen to coincide with the physical radius R at the initial regular epoch of $\tilde{t} = 0$, i.e., $R(0, \tilde{r}) = \tilde{r}$. We assume that the mass function $F(\tilde{r})$ can be expanded near the regular center as

$$F(\tilde{r}) = F_1 \tilde{r}^a + F_2 \tilde{r}^b + \cdots$$

where a and b are constants satisfying a < b. Then the initial density profile is written as

$$\rho(0,\tilde{r}) = \frac{aF_1}{8\pi}R^{a-3} + \frac{bF_2}{8\pi}R^{b-3} + \cdots .$$
 (B·1)

Comparing Eq. (B·1) with Eqs. (2·5) and (2·6), one obtains the powers and coefficients of $F(\tilde{r})$ as,

$$a = 3, b = \frac{3(3\mu + 2)}{3\mu + 1},$$

$$F_1 = \frac{4}{9t_{in}^2}, F_2 = -\frac{8(\mu + 1)^{3/(3\mu + 1)}}{9\lambda^{3/(3\mu + 1)}(-t_{in})^{(9\mu + 5)/(3\mu + 1)}}.$$

It is found that the power b is in the region of $3 < b \leq 6$ for $\mu \geq 0$. In Ref. 10), Harada *et al.* determined the characteristic frequency of the naked singularity in the analytic model ($\mu = 1/6$) through physical discussion. It is easy to repeat their discussion for the general value of $\mu > 0$. One possible quantity, which is composed only of F_1 and F_2 , independent of the choice of initial time slice, and has the dimension of frequency is

$$F_1^{(9\mu+5)/(6\mu)}(-F_2)^{-(3\mu+1)/(3\mu)} = (\mu+1)^{-1/\mu} \left(\frac{2}{3}\right)^{(9\mu+5)/(3\mu)} \left(\frac{9}{8}\right)^{(3\mu+1)/(3\mu)} \omega_s, \qquad (B.2)$$

where we used Eq. (4.4). In terms of γ , the quantity of (B.2) is written as follows:

$$\Omega_{\gamma}(\lambda) \equiv F_1^{(2\gamma+9)/(2(3-\gamma))} (-F_2)^{-3/(3-\gamma)}.$$

In the case of the analytic LTB model ($\gamma = 2$),

$$\Omega_2(\lambda) = F_1^{13/2} (-F_2)^{-3},$$

which coincides with the frequency defined in Ref. 10) except for a numerical factor. This shows that it is valid to define $\omega_s \equiv \lambda^{1/\mu}$ as the frequency of singularity. In the self-similar LTB solution ($\mu = 0$), such a quantity does not exist because of the scale-invariant nature of self-similar spacetimes.

References

- R. Penrose, Riv. Nuovo Cim. 1 (1969), 252, reprinted in Gen. Relativ. Gravit. 34 (2002), 1141; in *General Relativity, An Einstein Century Survey*, edited by S.W. Hawking and W. Israel (Cambridge University Press, Cambridge, England, 1979), p. 581.
- P.S. Joshi, Global Aspects in Gravitation and Cosmology (Oxford University Press, New York, 1993).
- 3) T. Harada, Pramana 63(4) (2004), 741, Proceedings of the 5th International Conference on Gravitation and Cosmology, edited by B.R. Iyer, V. Kuriakose, and C.V. Vishveshwara, gr-qc/0407109.
- 4) L.H. Ford and L. Parker, Phys. Rev. D 17 (1978), 1485.
- 5) W.A. Hiscock, L.G. Williams, and D.M. Eardley, Phys. Rev. D 26 (1982), 751.
- S. Barve, T.P. Singh, C. Vaz, and L. Witten, Nucl. Phys. B **532** (1998), 361; Phys. Rev. D **58** (1998), 104018; C. Vaz and L. Witten, Phys. Lett. B **442** (1998), 90.
- 7) H. Iguchi and T. Harada, Class. Quantum Gravit. 18 (2001), 3681.
- 8) U. Miyamoto and T. Harada, Phys. Rev. D 69 (2004), 104005.
- 9) T.P. Singh and C. Vaz, Phys. Lett. B 481 (2000), 74.
- 10) T. Harada, H. Iguchi, K. Nakao, Phys. Rev. D 61 (2000), 101502; *ibid.* 62 (2000), 084037.
- 11) T. Harada, H. Iguchi and K. Nakao, Prog. Theor. Phys. 107 (2002), 449.
- 12) M.D. Roberts, Gen. Relativ. Gravit. 21 (1989), 907.
- 13) B. Waugh and K. Lake, Phys. Rev. D 38 (1988), 1315.
- 14) F.J. Tipler, Phys. Rev. Lett. A 64 (1977), 8.
- 15) C.J.S. Clarke and K. Królak, J. Geom. Phys. 2 (1985), 127.
- 16) A. Królak, J. Math. Phys. 28 (1987), 138.
- 17) S.S. Deshingkar, P.S. Joshi, and I.H. Dwivedi, Phys. Rev. D 59 (1999), 044018.
- 18) T. Harada, K. Nakao, and H. Iguchi, Class. Quantum Gravit. 16 (1999), 2785.
- G. Lemaître, Ann. Soc. Sci. Bruxelles I A 53 (1933), 51; R.C. Tolman, Proc. Nat. Acad. Sci. 20 (1934), 169; H. Bondi, Mon. Not. R. Astron. Soc. 107 (1947), 410.
- 20) D. Christodoulou, Commun. Math. Phys. 93 (1984), 171.
- 21) T. Tanaka and T.P. Singh, Phys. Rev. D 63 (2001), 124021.
- 22) N.D. Birrell and P.C.W. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, England, 1982); L.H. Ford, Phys. Rev. D 15 (1987), 2955.
- 23) T. Harada, H. Iguchi, K. Nakao, T. Tanaka, T.P. Singh, and C. Vaz, Phys. Rev. D 64 (2001), 041501.
- 24) G.T. Horowitz and D. Marolf, Phys. Rev. D 52 (1995), 5670; A. Ishibashi and A. Hosoya, Phys. Rev. D 60 (1999), 104028; D.A. Konkowski and T.M. Helliwell, Gen. Relativ. Gravit. 33 (2001), 1131.
- 25) R.M. Wald, J. Math. Phys. 21 (1980), 2802; A. Ishibashi and R.M. Wald, Class. Quantum Gravit. 20 (2003), 3815.
- 26) P.S. Joshi and I.H. Dwivedi, Phys. Rev. D 47 (1993), 5357.




Fig. 1. A possible causal structure of the LTB spacetimes considered in this article. A null singularity occurs at the spacetime point (u_0, v_0) and is visible from \mathcal{I}^+ , where (u, v) is a suitable double null coordinate system. An outgoing null ray u = constant can be traced backward in time from \mathcal{I}^+ to \mathcal{I}^- , which turns out to be an ingoing null ray $v = \mathcal{G}(u)$ after passing the regular center, locating at r = 0 with t < 0. The outgoing null ray $u = u_0$ and ingoing null ray $v = v_0$ represent the CH and the null ray that terminates at the NS, respectively.




Fig. 2. A schematic spacetime diagram of the dust-filled region with the illustration of the local map defined in Sec. 3. A pair of an ingoing null ray $t_{-}(r)$ and an outgoing null ray $t_{+}(r)$ is depicted (solid line), which passes near the NS, locating at (t, r) = (0, 0). Null rays terminating at and emanating from the NS, $t_{-}^{\text{crit}}(r)$ and $t_{+}^{\text{crit}}(r)$, are also depicted (dashed lines), where the latter is the CH. A comoving observer is locating at $r = r_0 = \text{constant}$. The local map is defined as the relation between $t_{-}(r_0)$ and $t_{+}(r_0)$.

Table I. The relation among the curvature strength of naked singularities, the luminosity, and the energy of scalar fields near the Cauchy horizons (a) for the non-conformally coupled scalar fields ($\xi \neq 1/6$) and (b) for the conformally coupled scalar field ($\xi = 1/6$). The constant γ parameterizes the initial density profile of a dust fluid. SCC implies LFC.

				(a)				
γ	0		3/4	1		3/2		3
Strength	—	Weak	LFC				SCC	
Luminosity	-	Finite			Divergent	Finite	Divergent	
Energy	—	Finite Divergent						

_	(D)									
ſ	γ	0		3/4		3/2		2		3
	Strength	_	Weak	Veak LFC					SCC	
	Luminosity	-	Finite			Divergent				
	Energy	_	Finite Divergent							

~)