

Quantum Cosmological Models

D.H. Coule

Institute of Cosmology and Gravitation,
University of Portsmouth, Mercantile House, Hampshire Terrace,
Portsmouth PO1 2EG.

Abstract

We contrast the initial condition requirements of various contemporary cosmological models including inflationary and bouncing cosmologies.

Quantum cosmology is used, as a first approximation to full quantum gravity, to determine whether suitable initial conditions are present.

Various proposals such as Hartle-Hawking's "no boundary", or Tunnelling boundary conditions are assessed on grounds of naturalness and fine tuning. Alternatively a quiescent initial state or an initial closed time-like curve "time machine" is considered. Possible extensions to brane models are also addressed. Further ideas about universe creation from a meta-universe are outlined.

We compare the recent loop quantum cosmology of Bojowald and co-workers with these earlier proposals. A number of possible difficulties and limitations are outlined.

PACS numbers: 98.80.Qc

Contents

- 1.0 Introduction
- 2.0 Classical Initial Conditions
 - 2.1 Initial conditions for Inflation
 - 2.2 Finite domain size for Inflation
 - 2.3 Inflation from previously non-inflationary conditions
 - 2.4 Variable constant models
- 3.0 Quantum Cosmology
 - 3.1 Cosmological constant Λ case
 - 3.2 Massless scalar field case
 - 3.3 Scalar potential $V(\phi)$ case
 - 3.4 Further Topological and Geometric aspects
 - 3.5 Brane quantum cosmology
 - 3.6 Universe from a quiescent or static state
- 4.0 Loop Quantum Cosmology
 - 4.1 Bouncing and Inflationary model?
 - 4.2 Modified Wheeler-De Witt equation
 - 4.3 Loop cosmology and super-inflation
 - 4.4 Is loop quantum cosmology unstable?
- 5.0 Conclusions

1.0 Introduction

In this review we intend to outline and compare various ideas about the origin of the universe as suggested by quantum cosmological reasoning. A number of earlier reviews are already available especially concerning technical issues [1-12]. We also consider the implications of possible discreteness of space, as suggested by loop quantum gravity - for extensive reviews see [13,14]. For other approaches and more general issues of quantum gravity see e.g. [15-24].

The recent history of cosmology is one of gradually more extreme matter sources being considered. Originally in standard big bang (SBB) cosmology only ordinary matter like radiation or dust, that satisfies the strong energy condition, was believed suitable. This resulted in singularities being present and other so-called *puzzles* of SBB cosmology.

A significant step was made when false vacuum states were considered

that violate the strong energy condition: so driving an antigravity phase, now generally known as inflationary expansion - see e.g. [25,26], while for definition of various energy conditions see [27-29]. Further cosmological issues and alternative models are given in [30-34].

Recent evidence suggests the universe is again undergoing inflationary expansion although with perhaps a very small effective cosmological constant, this and other possibilities are reviewed in [35]. However, in the early universe many of the original puzzles for inflation have since been shown to be either avoidable, solvable by other means or anyway unsatisfactory treated by inflation. For example, the *horizon problem* [25,26] is something that is actually generated during the Planck time [36], so a quantum gravity solution is really required: perhaps by the presence of wormholes in the quantum foam at that time e.g.[37]. Since thermodynamical quantities become ill-defined at the Planck epoch cf.[38] it might be expected anyway that initial homogeneity or smoothness could occur [39]. Another drawback is that a secondary *horizon problem* caused by phase transitions proceeding after inflation could occur and so spoil the earlier smoothness [23].

The *flatness problem* [25,26] might also be solved by quantum consideration during the Planck time [40], (see p.501 in ref.[32] for a related solution) or by choosing natural values for arbitrary constants [41].¹ This choice either given by quantum gravity arguments or from purely classical reasoning.

Neither is the presence of singularities in SBB cosmology alleviated by inflation. Recent proofs suggest inflation also cannot be extended indefinitely into the past [44]: essentially because the flat de Sitter metric is then geodesically incomplete [27] (to null rays) unless one makes topological identifications that remove time orientability [45]. Earlier incompleteness theorems required the weak energy condition be satisfied [44]. One unorientable example of patching together two de Sitter universes with time running oppositely is given in ref.[46], although this idea has long been known [47]. The result can also be questioned on the grounds that actual finite particle lifetimes never allows timelike paths to actually become null cf.[48]. Recycling behaviour might also help circumscribe the geodesic incompleteness results [49].

One major advantage of inflation is that it can give a large universe even starting with natural Planck values for various quantities: so an initial “small

¹There is a Bayesian argument [42] that also apparently solves the flatness problem, but it assumes an invalid scale invariance property of General Relativity [43].

bang” can be a sufficient starting point. Without inflation the energy density when extrapolated back vastly exceed the Planck energy density when the universe was the Planck time old: the *Planck problem* of SBB cosmology [50,25,26]. Another use of inflation is that Hawking radiation can provide a mechanism for providing scale-invariant perturbations for galaxy formation e.g. [51].

For a simple FRW universe the Friedmann equation is given by [25-28]²

$$H^2 + \frac{k}{a^2} = \frac{8\pi G}{3}\rho \quad (1)$$

For a perfect fluid with equation of state $p = (\gamma - 1)\rho$ the energy density ρ scales as $a^{-3\gamma}$. The strong energy condition is violated for $0 \leq \gamma < 2/3$. Such an example is that of a scalar field with potential $V(\phi)$ dominating over the kinetic energy $\dot{\phi}^2$. In general, while keeping ρ positive, and since

$$\gamma = \frac{2\dot{\phi}^2}{\dot{\phi}^2 + V(\phi)} \quad (2)$$

one can obtain more extreme violation of the weak e.g. ($\gamma < 0$) or dominant energy conditions ($\gamma > 2$) by choosing, in turn, either a negative kinetic term or negative potential. The negative γ example, from a “wrong sign” kinetic term, causes super or pole-law inflation: the scale factor a goes as $a \sim t^n$ with $n < 0$. If the present expansion is of this form it has been dubbed a phantom matter source proceeding towards a “big-rip” singularity [52]. Although such models were earlier simply described as a “whimper expanding to a big bang” e.g.[30]. Numerous examples of super-inflation are known such as in Brans-Dicke gravity e.g[31], with parameter $\omega < -3/2$ or with coupled dilaton-axion scalar fields [53]. Super-inflation suffers from a number of problems stemming from a growing Hubble parameter giving a blue spectrum of perturbations and the presence of a future singularity - so was generally discounted as a suitable form of inflation for the early universe [54].

Also if the present universe is undergoing such a rapid expansion it is difficult to see how a phantom scalar field is compatible with an earlier inflationary phase in the universe. The required “wrong sign” kinetic term should have been diluted or else it would have dominated over the earlier

²We use notations and units of the quoted references, generally Planck units.

inflationary source. Having “wrong sign” spatial gradient terms is also likely to cause instability [55]. Another possibility is to add various new terms to the gravitational action and so allow super-inflation today e.g.[56], although extra non-analytic Ricci scalar terms generally only give power-law inflation i.e. $a \sim t^n$ with now $\infty > n > 1$ [57].

The ekpyrotic model [58] is a recent attempt at producing a cyclic universe inspired by the supergravity limit of string theory. During the contraction phase a negative potential can dominate to produce an effective “super-stiff” equation of state $\gamma \gg 2$ which is said to produce an ultralocal phase.

Another extreme case is to take ρ negative, which is a more drastic violation of the weak energy condition. One can now produce a bounce for arbitrary curvature k unlike for the closed case which can bounce by violation of the strong energy condition alone [59]. In general a FRW bouncing model is described by an equation of the form

$$H^2 = \frac{A}{a^n} - \frac{B}{a^m} \quad (3)$$

A, B arbitrary positive constants and provided $m > n$, a bounce proceeds. Note that the stiffer (larger γ) matter component requires the minus sign. For the closed universe the curvature plays this role i.e. $m = 2$, so only the strong energy condition has to be violated i.e. $0 \leq n < 2$ for a bounce to proceed.

Such bouncing models tend to suffer from rapidly growing classical perturbations since the usual Jeans instability is no longer damped by rapid expansion [60,61]. Further examples of bounces are reviewed in ref.[62], while the eventual dominance of any (small) cosmological constant was shown in ref.[63]. The related quasi-steady-state model [64] has an underlying de Sitter expansion with superimposed oscillations: the model now has a “no hair” property that inhomogeneity is suppressed [65]. Although the bouncing models can avoid a possible singularity they still require suitable initial conditions, albeit at an arbitrary distant past. If such models collapse as much as they expand it will tend to negate any advantage of having an inflationary phase. We later will use this to restrict the usefulness of combining bouncing behaviour with subsequent inflationary expansion.

Alternatively the universe evolved from a high curvature region where quantum gravity would have dominated. As a first approximation one can attempt to quantize Einstein’s field equations and obtain quantum cosmology using predominantly the Wheeler De Witt (WDW) equation [66,67].

Ultimately, the true theory for the forces of nature will automatically include quantum aspects and such an approximation would be superseded: it is sometimes said that quantum mechanics will have to forgo some of its principles for this unification, see e.g. [21]. However, quantum mechanics alone does not necessarily regularize singularities. Recall the Schrödinger equation for the hydrogen atom in N spatial dimensions. The total energy is given by

$$E \sim \frac{h^2}{2mr^2} - \frac{e^2}{r^{N-2}} \quad (4)$$

where the Heisenberg uncertainty principle is used to estimate the kinetic energy resisting localization. Following the presentation in ref.[32], when $N > 5$ the potential is too divergent and there is no minimum energy level [68]. Using more rigorous arguments that include special relativistic effects this result can be strengthened such that no stable bound exists for $N > 3$ [69]. So the quantization alone does not prevent the ($N > 3$) hydrogen atom being singular. Similar problems are expected to occur in quantum gravity if the potential term is initially too divergent.

Even if the singularity becomes regularized it is still perhaps a mixed blessing. If there is no longer any breakdown in the evolution one then requires an explanation as to what instead happens. If there is some automatic limiting curvature then perhaps a de Sitter phase occurs near the singularity [70] or simply some entire new universe results [71]. If one simply reenters the same universe or passes into a parallel one then we no longer have any natural starting point to the universe. Or perhaps the singularity then matches to some ubiquitous natural state.

One widely used notion in quantum cosmology is to assume such a state of “nothing” or zero size from which universes can initially tunnel [1,2]. More recently a possible meta-universe has instead been suggested as a potential source of universes [72-74]. The idea is that a, usually disfavoured, low entropy initial state is then possible by tunnelling like behaviour from the meta-universe. This unfortunately then pushes problems of naturalness into this pre-existing state, which then itself requires further explanation. If this meta-universe is already perhaps at a maximum entropy state then fluctuations could just as well surpass the holography bound [75] causing singularities. If the meta-universe is continually expanding then again it will likely suffer from related geodesic incompleteness theorems cf.[74,44].

Alternatively a continual cyclic evolution of a single universe, through possible regulated singularities, as in the ekpyrotic example, might be a less

drastic solution if one wishes to entertain the notion of an eternal state. But there are still issues as to whether a “heat death” or maximum entropy state should rather be expected [76].

If the present universe is undergoing super-inflation with a growing Hubble parameter it might eventually reach a quantum gravity singular state: perhaps eventually matching to a more moderate high energy de Sitter expansion. This state might then spawn new universes in a cyclic like manner. Whether such a scheme is possible is presently a topic of some interest but again would appear profligate in terms of growing space and overall entropy production. The geodesic incompleteness results might again be alleviated by giving all matter only a finite lifetime.

We can consider different examples of how quantum effects might play a role in the quantum gravity, or early, phase of the universe.

- *Different matter sources.*

Quantization will only have significant differences when the classical evolution reaches regions of small action. This is the most studied case where different matter sources are investigated and quantum tunnelling of space-time itself considered when the total action is small. One problem is that action is rather more ambiguous when gravity is included.

- *Alter time.*

Time might have more unusual properties during an quantum gravitational epoch. One possible example is that closed timelike curves [27] might occur during a quantum gravitational epoch. Adapting quantum mechanics to this example of a “time machine” is rather involved before one even considers including high curvatures.

- *Alter space.*

This approach stems from the idea that space itself should be quantized so that it occurs in discrete packets or “cells” [14]. Eventually at larger distances the cells merge together to approximate a continuum. Recently loop quantum cosmology [13] has been developed and is said to allow matter to take different properties when this granularity of space is apparent. One difficulty we will address is that absolute distance can be a rather nebulous quantity especially during rapid cosmological expansion.

Just as the hydrogen atom is dependent upon the form of the potential we first need to consider a suitable cosmological model before any quantum aspects might further be introduced.

2.0 Classical Initial conditions

There are essentially two approaches one can take for the initial condi-

tions: chaotic or orderly. In the earlier chaotic cosmology program dissipation or quantum particle effects were supposed to smooth the universe - see e.g.[32]. In practice such processes can only work if the initial anisotropy is constrained. One could choose unwanted conditions in the universe today and evolve backwards with the deterministic Einstein's equations to obtain initial conditions that are not-consistent with our universe [77]. The celebrated “no go” theorem of Collins and Hawking [78] also shows that only a measure zero of universes become isotropic at late time.

Inflationary cosmology tried essentially the same program but with a cosmological constant Λ or scalar potential $V(\phi)$ playing the role of the smoothing process. The hope that “no hair” of the chaotic initial conditions would remain can be contradicted by choosing large initial inhomogeneities - see [79] for a review. Likewise one could evolve backwards unwanted conditions through a finite amount of inflationary expansion [77,80,81]. The initial conditions therefore require some restrictions in order for a smooth FRW universe to be eventually produced. One can obtain a similar Collins-Hawking result for an inflationary convex potential $V(\phi)$ with zero minimum [82] or (flat) exponential potential provided some inhomogeneity is initially present [83]. In some general sense the scalar field is less robust at removing hair than the pure cosmological constant, e.g. [84].

It would therefore seem that inflation alone does not entirely explain the obedience of the *cosmological principle* e.g.[33]. Although the Collins-Hawking requirements are probably unnecessarily strict, since smoothing for finite future time and shear remaining bounded $\sigma/H \rightarrow \text{constant}$ [32,77], would suffice in practice to create a temporary observance of the *principle*. However counter examples are known so some restrictions on initial conditions are required if inflation is to actually produce a FRW universe from arbitrary initial conditions. One question is whether a (quantum) principle alone can provide this restriction so allowing inflation some chance of success?

The ekpyrotic universe also attempts to smooth space by using both a low-energy driven inflationary phase and a “super-stiff” $\gamma > 2$ phase during a subsequent contraction [85]. This combination might then have a more robust “no hair” property than inflation alone. But again this could also be overwhelmed by choosing irregular starting conditions: massive black holes for example that are not ripped apart by inflation or have time to evaporate [76]. The argument now for reducing anisotropy is that as the bounce proceeds, the super-stiff matter scales as $\sim a^{-3\gamma}$ while the shear $\sigma^2 \sim a^{-6}$. So for $\gamma > 2$ the shear remains subdominant and the universe is forced isotropic

[85,86]. However it is also important that the individual components should remain below Planck values if the model is to be described by the low energy supergravity limit of string theory. Without inflation around the conventional big bang point one requires energy densities vastly bigger than Planck values; the *Planck problem* of conventional cosmology [50]. A similar problem occurs in another recent attempt at avoiding inflation during the conventional big bang type evolution [87].

Inflation must also restrict the nature of space at the Planck scale: length scales are being blown up from below the Planck length to macroscopic size. If space had a fractal like structure this complexity would simply be extended to large size by inflationary expansion [88,77]. This problem is particularly severe for perpetually expanding models such as, from the higher dimensional perspective, the ekpyrotic scenario. Since it uses low energy string theory as a starting point it is eventually inconsistent that all length scales within the universe have come from below the Planck or string length of previous cycles [76].

The present string paradigm is that branes are present in higher dimensional bulk spaces - see e.g. [89] for a review. Our usual matter is confined to 4 dimensional spacetime while gravity, and some types of exotic matter, extends freely over the extra dimensions [90]. The branes are believed to undergo cosmological evolution presumably from some finite time in the past. But so far these models have been done with highly symmetric bulk spaces particularly Anti-de Sitter (AdS) [27] space with maximal symmetry. This is a strict departure from the idea of random initial conditions to spaces that obey the unchanging *perfect cosmological principle* e.g.[33]. Some attempts have been made to first have inflation within the bulk space but this is rather contrived if a negative cosmological constant is the true ground state [91]. Also inflation, driven by matter that introduces a preferred time cf.[92], does not provide such idealized symmetry, as previously discussed, at best it can only achieve the *cosmological principle* for finite times. Apart from simply passing to a higher dimensional cosmology, that requires explanation, one requires the initial conditions or processes that can produced the original smooth bulk space.

To more closely match the bulk space one can attempt to make the brane eternal by achieving a bounce. One example was by taking a Reissner-

Nordström AdS bulk which can modify the Friedmann equation to [93,94]

$$H^2 + \frac{1}{a^2} = \frac{8\pi G}{3} \left(\rho + \frac{\rho^2}{2\lambda} \right) + \frac{M}{a^4} - \frac{Q^2}{a^6} \quad (5)$$

where λ is the brane tension and M and Q represent the mass and charge of the bulk space. However this charged black hole appears unstable to perturbations since the bounce occurs within the outer event horizon [95,96]. Related work has suggested only weak singularities occur because of related “blue sheet” effects [97], but this probably required restricted initial conditions. The difficulty is that one is effectively trying to violate the weak energy condition and this often leads to instabilities. Note also that by choosing the wrong sign of the brane tension in eq.(5) you can also get a bounce without the bulk being charged [98] cf. eq. (3). Being on a negative tension brane is probably not consistent with other solar system constraints, and would also require more than a single brane to stabilize [99].

We can contrast this with the earlier pre-big bang model [100] that used the low energy limit of string theory with a dilaton term - reviewed in [101]. The extra dimensions are now assumed tightly compactified. This model is taken to start at time $t = -\infty$ before approaching a singularity at, say, time $t = 0$. However this model is not inflationary in the Einstein frame, so still having a *Planck problem* [102]. Also having no “no hair” like suppression requires fine tuning of initial conditions. A suggested initial state of plane gravitational and dilatonic waves is justified as having zero gravitational entropy “asymptotic past triviality” [103]. These waves after gravitationally interacting are expected to form a space-like singularity into the future which plays the role of the usual big bang singularity

During any collapsing phase quantum fluctuations can be produced by Hawking radiation although one has to have the correct contraction rate to get a scale invariant spectrum: dust $\gamma = 1$ [104,105] or super-stiff equation of state $\gamma \gg 2$ as in the ekpyrotic case [58,85]. Also quantum fluctuations require the presence of an event horizon, this coming from the time reversal of a particle horizon [33] which means the strong energy condition now has to be satisfied for quantum fluctuations to be present: so being now the opposite to the usual inflationary case. However, because if the impending singularity is to be avoided, the energy conditions have eventually to be violated. This, strictly speaking, contradicts the requirements for the production of Hawking radiation during the collapsing phase. There are similar uncertainties when

considering the Unruh radiation within a Rindler wedge (see e.g.[106]) when acceleration proceeds for only a finite time cf.[107].

Obtaining the weak energy violating source that turns around the pre-big bang has proved elusive, although certain higher order gravitational action corrections terms from string theory are possible. But requiring, for example, the wrong signs before the Gauss-Bonnet term can cause other instabilities [108].

In summary most emphasis in cosmology has been on assuming random initial conditions and then hoping some process will smooth things gradually towards a FRW like universe. In brane cosmology highly symmetric models have been assumed *ab initio*. Can quantum considerations suggest which of these complementary cases is most reasonable?

2.1 Initial conditions for Inflation

We can consider a simpler example of a scalar field with energy density $\rho = \dot{\phi}^2 + V(\phi)$ within a FRW universe to demonstrate some of the issues involved. Such an example can display chaotic inflation [109,110]. Unlike a simple cosmological constant which only has a de Sitter solution the scalar field has both inflationary and non inflationary solutions. In order to classify which is most probable Belinsky and coworkers [111-114] considered a simple equipartition at the Planck boundary $\rho \sim M_{pl}^4$. This was motivated by the, admittedly rough, argument that the uncertainty principle $\Delta E \Delta t \geq 1$ applied during the Planck time would give an uncertainty in the energy $E \sim$ Planck value [110,26]. This argument also assumes that the universe be closed so that the total energy is zero: positive matter energy density being cancelled by negative gravitational energy. Otherwise the uncertainty in energy would have to be paid back before the scalar field could take large classical values: or the non-zero total energy would have to be justified further.

This sort of analysis for a massive scalar field $V(\phi) = 1/2m^2\phi^2$ gives the probability of sufficient inflation ~ 70 e-foldings around $\sim (1 - m/M_{pl})$ [111] so almost certain for mass $m \ll 1$. For the correct size of fluctuations to be produced requires $m \sim 10^{-5}$ [25,26]. The slight chance of non-inflation is when the initial kinetic energy $\dot{\phi}^2$ is large ~ 1 in Planck units [111].

A more rigorous approach is to use a canonical measure [115-118]

$$\omega = -dp_a \wedge da + dp_\phi \wedge d\phi \quad (6)$$

where the canonical momenta are

$$p_\phi = a^3 \dot{\phi} \quad p_a = -3a\dot{a} \quad (7)$$

at fixed scale factor $a = a_0$ this measure simplifies to [41,117]

$$\omega = a_0^3 d\dot{\phi} \wedge d\phi \quad (8)$$

Because this measure is peaked at large energy densities it can solve the flatness problem i.e. that $\Omega = 1$ regardless of inflation [41]. Although if the potential is bounded above, Ω can again become arbitrary [116,117]. Such a bounded potential might be expected from a conformal anomaly or higher derivative R^2 like terms in the action [119].

The canonical measure can be shown to give an ambiguous probability of whether inflation occurs, there are infinite numbers of both inflationary and non-inflationary solutions [41]. Although this sort of ambiguity might be rectified if some input from quantum cosmology could, for example, determine if the universe started small or with a certain energy density [117]. One can also investigate the probability of bounces occurring. In simple models most collapsing universes have only finite measure of bouncing compared to infinite measure of singular solutions [116,120]. We are therefore not sure that one can say that deflation is as likely as inflation with this measure cf.[118]. There is also an extremely small measure for perpetually oscillating solutions: both periodic and non-periodic [121,122].

Once the scale factor is fixed one can obtain a finite total measure and obtain the probability of inflation [41,117,118]. It depends as expected with the chosen energy density. For the massive scalar field case the fraction of inflationary solutions f_I , for at least z e-foldings of inflation is [41]

$$f_I \sim 1 - \frac{m \ln(1+z)}{\rho^{1/2}} \quad (9)$$

Taking $m \sim 10^{-5}$ and requiring 100 e-foldings of inflation gives $f_I \sim 1 - 10^{-5}$ for an assumed energy density $\rho \sim$ Planck value. The above expression gives that as the initial value of ρ is reduced $f_I \rightarrow 0$. Although a more detailed calculation shows f_I actually tends to a very small finite value as ρ is taken to zero [41]. However, if anisotropy or other inhomogeneous degrees of freedom are included then f_I would also approach zero as $\rho \rightarrow 0$ [123]. In order to give a high probability of inflation therefore requires that the energy density should be large with this measure once the initial scale factor is fixed. We later will consider how loop quantum effects attempts to remedy this prediction and obtain inflation even for small initial values of the energy density.

2.2 Finite domain size for Inflation

So far this analysis assumes the scalar fields are homogeneous throughout the universe. But if only some finite domain is to inflate then this homogeneous region must be of sufficient size. Recall that the scalar field behaves as a negative pressure and any outside positive or zero pressure will wish to equalize the disparity by “rushing in” to cancel any pressure differences. Assuming this equalization can proceed at the speed of light one finds that the homogeneous domain must be of a size greater than $\sim 3H^{-1}$ [81]. This can be alleviated in some topologically non-trivial universes cf. [124]. Also topological driven inflation [125] using defects, such as monopoles, can possibly weaken this requirement due to topological charge conservation [126]; but we leave aside these possible complications. Once such a suitable domains occur then inflation will proceed.

Because of a combination of the fluctuations that are generated during inflation and the finite horizon size $\sim H^{-1}$ in de Sitter space inflation never stops entirely once started, provided a requirement on the size of the scalar field $\phi > \phi_*$ is satisfied [110]. In general the potential must satisfy $V(\phi)^{3/2} > dV(\phi)/d\phi$ for eternal inflation [26].

If the initial domain does not have such a sufficiently large scalar field value one can hope that a quantum “instanton” effect can produce a domain with sufficiently large scalar field provided there is a sufficient number $\sim \exp(1/V(\phi))$ of Hubble volumes: so that an especially large fluctuation will occur somewhere [127,128]. Note that this would require an astronomical $\sim \exp(10^{120})$ Hubble volumes for the apparently present inflationary state of the universe to actually produce a suitable eternal domain.

However, there are a number of uncertain aspects [129] about this eternal inflation mechanism. For example whether the fluctuations of matter transfer to the geometric left hand side of Einstein’s equation, or whether the black hole calculation entirely extends to the de Sitter case e.g.[130]. Extra dimensions, perhaps the bulk space in brane models might regulate such a mechanism cf. [131]. Other criticisms are given in ref.[132].

But even if the eternal mechanism proceeds you can argue that the total produced number of new inflationary domains is less than countable infinity $\aleph_0$ for a finite time of eternal universe production [129]. It does not turn an initially finite universe into an infinite one. It therefore is not clear that this alone can overcome, for example, the previous ambiguity in the classical measure that an infinity of both inflationary and non-inflationary solutions are possible. You might as well start with an initially infinite universe, like

the flat FRW one, and put down random conditions - but this seems a extravagant way of trying to counteract an infinity of unwanted solutions.

What is required is a definite prediction that inflation will occur - the idea that once inflation starts it will never finish does not supersede this requirement.

2.3 Inflation from previously non-inflationary conditions

What about the naturalness of inflation in models that do not inflate initially from their inception. Singularities seem a general consequence of producing conditions that give inflationary behaviour. The method of producing a “universe in the laboratory” was required to expand so rapidly to avoid re-collapse that a singularity would be present [133-135]. This can be seen clearer by noting that in a FRW universe, regions of size bigger than the so-called apparent horizon $\sim \rho^{-1/2}$, have a necessary singularity -see page 353 in ref.[27] where such a quantity is called the *Schwarzschild length* of matter density ρ . But this size, $\sim H^{-1}$ for the flat $k = 0$ case, is the minimum required to isolate an inflationary patch from its surroundings for sufficient time to start inflating, ignoring possible topological counterexamples. Now requiring this initial patch size to be larger than the apparent horizon size has been emphasized [136] as a problem in setting up inflationary conditions. However, depending on the matter source it need not strictly violate causality which is rather determined by the particle horizon: the distance light travels from the beginning of the universe. The large initial patch size does though imply that a singularity is present [7].

If we consider further the nature of the *horizon problem*. It occurs because the the particle horizon size, defined as

$$r = c \int_0^t \frac{dt}{a(t)} \quad (10)$$

is finite, see eg.[25,26]. The horizon proper distance R is this quantity r multiplied by the scale factor i.e. $R = a * r$. For any strong energy satisfying matter source this quantity R grows linearly with time. But in SSB cosmology the rate of change of the scale factor, given by $a \sim t^p$ and $0 < p < 1$, grows increasingly rapidly as $t \rightarrow 0$. The horizon cannot keep pace with the scale factor “velocity” $\dot{a} \sim 1/t^{1-p}$. But this only occurs for times below unity $0 < t < 1$. If the horizon problem was already solved at the Planck time t_{pl} it would remain permanently solved during the ensuing evolution [36]. Note also that in models that inflate from their inception the usual space-like sin-

gularity of the FRW universe becomes null like when $p > 1$ - so any horizon problem is absent, see eg.[27,137].

The main idea of inflation is to take an initial domain of size less than the corresponding particle horizon size and allow it to expand greatly to encompass our universe. Let us see how this requirement can be satisfied. Most of the relevant quantities have already been obtained in the context of the holography conjecture [138,139].

For this purpose a useful form of the FRW metric is

$$ds^2 = a^2(\eta) \left(-d\eta^2 + d\chi^2 + f^2(\chi)d\Omega^2 \right) \quad (11)$$

where $f(\chi) = \sinh \chi$, χ , $\sin \chi$, corresponding to open, flat and closed universes respectively. We can define a number of important quantities. The *Hubble horizon* is defined by

$$r_H = H^{-1} \quad (12)$$

The *particle horizon*, or the distance travelled by light from the initial moment of the universe, is simply,

$$\chi_{PH} = \eta \quad (13)$$

for this metric. The *apparent horizon* is given by [138]

$$\chi_{AH} = \frac{1}{\sqrt{H^2 + k/a^2}} \Rightarrow \frac{1}{\sqrt{\rho}} \quad (14)$$

Roughly speaking light rays beyond the apparent horizon are seen to move away from the origin, a so-called anti-trapped behaviour e.g. [27,28]. Note that in the flat case $k = 0$ the apparent horizon and Hubble horizon coincide.

In ref.[136] it is argued that the initial inflationary patch must have sufficient size x that it reaches the anti-trapped surface i.e. $x > \chi_{AH}$. Otherwise the weak energy condition is violated for light rays that could otherwise enter the inflating region from normal or trapped regions. For a perfect fluid $p = (\gamma - 1)\rho$, the apparent horizon has the following time dependence [138,139]

$$\chi_{AH} = \frac{d\gamma - 2}{2}\eta \quad (15)$$

with d the number of space dimension. However the causal particle horizon has a different time dependence simply $\chi_{PH} = \eta$ so the condition

$$\chi_{AH} < x < \chi_{PH} \quad (16)$$

can be satisfied for

$$\frac{d\gamma - 2}{2} < 1 \xrightarrow{d=3} \gamma < 4/3 \quad (17)$$

This does exclude the case of radiation ($\gamma = 4/3$) or stiffer equations of state. But if γ was gradually reducing before inflation occurred this causal constraint can be satisfied. The condition can be thought of as saying the effective value of γ cannot switch suddenly but rather must fall below $4/3$ for sufficient time to allow the causal or particle horizon to be larger than the apparent horizon. This result is independent of whether curvature is present.

In the closed case only during the expansion phase is an anti-trapped surface present - cf. Fig.(4) in ref.[139]. This means that producing inflation, within a larger domain, to avoid an impending big crunch singularity during a collapsing phase will violate also the weak energy condition. Now it is the case that needing $x > \chi_{AH}$ is difficult to justify in terms of particle physics processes, but if this patch could be smaller than χ_{AH} one could avoid the singularity in a FRW universe since the matter would be insufficient to converge the light rays into the past. See chapter 10 in ref.[27] for a proof of this argument. So allowing an initial domain of size $x < \chi_{AH}$ to inflate, would have allowed singularities to be expunged from this cosmology: the result that this cannot be done without violating the weak energy condition is therefore consistent with the studies of eternal inflation that singularities have to be present when the model is continued into the past [44]. Note that the more general geodesic incompleteness results show that even violation of the weak energy condition does not prevent past geodesics leaving the metric, although not necessarily possessing curvature singularities.

There are some alternative metrics with non-singular solutions, but like Minkowski space they do not have anti-trapped regions [137,140]. Achieving inflation in such spaces would likewise require the violation of the weak energy condition.

However such violations of the energy conditions are possible with quantum fluctuations of the vacuum. Universe creation in the laboratory by quantum tunnelling [141,142], or starting from Minkowski space or at high temperatures has been considered [143]. A black hole is produced in the laboratory that eventually evaporates allowing the new universe to disconnect from the original one: it does not simply supplant it [141-143,49]. The process generally is enhanced for larger coupling constants e.g. m or λ , and with temperatures approaching Planck values [143]. Topological inflation also produces charged black holes to observers outside the defect [125].

This black hole does not evaporate though, because of charge conservation, and any new inflationary universe remains connected by a wormhole to the original space.

A possible meta-universe could then take the place of the laboratory and cause, by Poincaré recurrence, universes to automatically branch off by quantum effects [72-74], see also Albrecht in ref.[22]. Although the probability is remote, if there is infinite space or time, then such events will eventually occur. This is not entirely a satisfactory explanation since the first cause is then simply switched into understanding the origin of the meta-universe.

2.4 Variable constant models

Another possible solution to the horizon problem is to postulate that the various constants, particularly the speed of light c , could take different values during the early universe [144], for a review see [145]. This alone is not too helpful since a space-like singularity cannot be crossed by any finite value of c and a higher c just means one has to go further back in time to see an equivalent horizon problem [146]. There is also a causality problem, of sorts, as to why c can change simultaneously over the whole universe and constantly stay equal throughout, once the value of c has started to reduce and causal contact lost. This behaviour for c really has to be pre-programmed in the universe from its conception: so the horizon problem appears simply switched into how constants “know how to vary”.

Changing such constants also tends to suppress any quantum gravitational epoch at the beginning of the universe [146]. The gravitational action at fixed time, for say a radiation dominated universe, scales as $S \propto c^3/G$ so increasing for bigger c . However, this quantum epoch can surface at a later time, which must be pushed sufficiently far into the future. In this regard these model have some similarity with the pre-big bang phase which also starts in a classical state and tend towards a quantum gravitational singular region - that becomes the start of the big bang phase. This makes such variable constant models more difficult to conceive of by quantum creation schemes if initially we expect a quantum gravity phase with some small action. However, quantum cosmology at present does not explain why the various constants take their actual values and so they must be imposed by fiat.

3.0 Quantum Cosmology

We now wish to consider how quantum considerations can suggest ways of providing initial conditions for classical evolution. We are here interested in conceptual and cosmological applications so we will avoid technical concerns

as much as possible.

In the general case there is the Wheeler-DeWitt equation $H\Psi$ and momentum constraint $H_i\Psi = 0$ equations [66,67]. Such an approach corresponds with Einstein's field equations [147]. A $3 + 1$ decomposition of spacetime is further assumed to define a Hamiltonian [1-3] although eventually this might be relaxed cf.[148]. In mini-superspace models the momentum constraint equations are trivially satisfied and the WDW can be written with kinetic and potential U pieces [1-3].

$$\left(-\frac{1}{2}\nabla^2 + U(a, \phi)\right)\Psi(a, \phi) = 0 \quad (18)$$

where we consider a FRW model with scale factor a and scalar field ϕ matter source.

Some important early developments are given in refs.[149-157,40]. A general idea is to start the universe with a small, in units of $\hbar$, action so that quantum effects can be expected to be relevant. As emphasized by Zeldovich [50] such a low action state necessarily requires an inflationary phase in order to drive a “small bang” universe to have correspondingly large amounts of energy and action: so eventually simulating the usual big bang quantities.

3.1 Cosmological constant Λ case

For the archetypal example consider a closed universe with a cosmological constant Λ . The WDW equation simplifies to, [158-162]

$$\left(\frac{d^2}{da^2} + \frac{p}{a}\frac{d}{da} - U\right)\Psi(a) = 0 \quad (19)$$

with p a factor ordering ambiguity and U the WDW potential is

$$U = a^2 - \Lambda a^4 \quad (20)$$

see Fig.(1). For $U > 0$ there is a forbidden region: recall the classical de Sitter solution $a = \Lambda^{-1/2} \cosh(\Lambda^{-1/2}t)$ that bounces before reaching the origin. But instead of taking the classical collapsing evolution from time $t = -\infty$ to $t = 0$ it is envisioned that the universe can tunnel from $a = 0$ or “nothing” to the smallest classically allowed value $a = \Lambda^{-1/2}$.

For the case $p = -1$ the WDW equation can be solved in terms of Airy functions - see e.g.[159]. The presence of a forbidden region allows both exponentially growing and decaying solutions so there can be large differences in behaviour e.g.[2,3]. The solutions are sketched in Fig.(1).




Figure 1: The WDW potential (solid line) for the de Sitter example. In the forbidden region ($U > 0$) exponentially growing and decaying modes are present. Quantum uncertainty might allow a small classical universe to develop around $a = 0$.

One can also work with the gravitational action [28,163].

$$S = \int d^4x \sqrt{g} (R - \mathcal{L}_{\mathcal{M}}) - 2 \int d^3\sqrt{h} K \quad (21)$$

where $\mathcal{L}_{\mathcal{M}}$ is the matter component. One generally needs to work with compact spaces to create a finite action or else add further boundary counter-terms [164]. The wavefunction is given by

$$\Psi(h, \Lambda) = \int [g] \exp(iS) \quad (22)$$

Formally the integration is over all geometries and topologies. Hartle and Hawking consider a rotation to Euclidean space $t \rightarrow i\tau$ to improve the regularization properties [153]. They further integrate over Euclidean manifolds with one boundary, specified by the 3-metric h .³To semi-classical order the Hartle-Hawking wavefunction is

$$\Psi_{HH} \sim \exp(-S) \quad (23)$$

We assume a number of technical difficulties can be resolved such as dealing with conformal modes that can set $S \rightarrow -\infty$ [166]

³It has been suggested that one can obtain similar results working with purely Lorentzian metrics [165].

The Euclidean action for a simple cosmological constant model is given by [167,168]

$$S = -\Lambda V_4 \quad (24)$$

where V_4 is the 4-volume of the space. For positive Λ the four-sphere S^4 has the largest volume with $V_4 = \Lambda^{-2}$ so the action is negative $S = -1/\Lambda$.

⁴ For negative Λ the action is positive and infinite for non-compact cases [170,171]. One might expect to take the minimum volume of V_4 in this case to minimize the action. This in turn will restrict the 3 geometry of any created Lorentzian universe to also have small volume as in Weeks and Thurston spaces -see [172] for review of hyperbolic geometries in cosmology.

The wavefunction Ψ_{HH} for de Sitter is therefore $\sim \exp(1/\Lambda)$. If Λ is given by a distribution function then such expressions would suggest $\Lambda \rightarrow 0$: this was at one time part of the wormhole “big fix” approach of Coleman [173]. The inclusion of four form field strength $F_{\mu\nu\rho\lambda}$ can alter this prediction [174,175]. Recently a decoherence term has been invoked to also alter the prediction [176]. However, this term would correspond to adding an imaginary part to the scalar potential so causing absorption cf. e.g.[177]. Since we are presumably dealing with a single universe it is unclear how such absorption can be interpreted, although decoherence has been invoked to explain why different possible branches of the universe fail to interfere [178]. Back reaction effects from the presence of other matter sources could also affect semi-classical tunnelling predictions cf. [179]. We would also add that related earlier investigations using quantum cosmological reasoning suggested a small Λ could be expected so possibly coinciding with the present universe [180].

In the tunnelling approach the action is expected to be positive definite. This can be achieved in a path integral formulation by imposing “outgoing only” modes e.g. [161]. In simple models one can Wick rotate, the gravitational term in the action, in the opposite direction to that used on any matter component. In the simple cosmological model the wavefunction is now $\Psi = \exp(-|S|) = \exp(-1/\Lambda)$ [155]. This rotation has been justified more rigorously from path integral methods, in that the wavefunction should not be exponentially large [181]. Although wormhole solution could violate this requirement [182].

In analogy with the field emission of electrons problem, where the applied

⁴Note that for S^n and large n the action is more divergent $S \sim -1/\Lambda^{n/2}$ and it is suggested one should minimize the energy to help predict $n = 4$ [169].

electric field E gives the probability of emission $\sim \exp(-1/E)$, the tunnelling boundary condition suggests a large value of Λ is expected.

We have questioned whether this analogy of treating the universe like α decay or a Scanning Tunnelling microscope is entirely sensible [7]. In these atomic examples the various particles already exist. But now the “particle” is the universe itself coming into existence - this is claimed not to be possible with the Copenhagen interpretation of quantum mechanics [183]. Note also that the barrier to be tunnelled through is dependent on having closed $k = 1$ curvature. It is absent for the flat or open cases, which classically can tend towards zero scale factor in the distant past. Although such models can have apparently infinite size and so infinite action one can instead topologically compactify the space at arbitrary small size [156,184,125]. For such cases one can impose, for example, analogous outgoing like boundary conditions but difficulties are present for the Hartle-Hawking case [184].

3.2 Massless scalar field case

The previous barrier in the WDW potential is dependent on having a strong energy condition violating matter source. For the alternative example we next consider the massless scalar field in a closed FRW universe, given by the Wheeler-DeWitt (WDW) equation, e.g.[1-3]. We follow our earlier presentation of this example [185],

$$\left(\frac{\partial}{\partial a^2} + \frac{p}{a} \frac{\partial}{\partial a} - \frac{1}{a^2} \frac{\partial}{\partial \phi^2} - a^2 \right) \Psi(a, \phi) = 0 \quad (25)$$

where p again represents part of the factor ordering ambiguity. The WDW potential now has a forbidden region at large scale factors, where the classical closed universe starts re-collapsing - see Fig.(2).

The WDW equation can be separated to

$$\left(\frac{d^2}{da^2} + \frac{p}{a} \frac{d}{da} + \frac{\nu^2}{a^2} - a^2 \right) \Psi(a) = 0 \quad (26)$$

$$\left(\frac{d^2}{d\phi^2} + \nu^2 \right) \Psi(\phi) = 0 \quad (27)$$

with ν the separation constant.

The solution to these equations can be obtained using MAPLE [186],

$$\Psi(a) \sim a^{(1-p)/2} \left\{ \alpha J_{i\nu/2}(ia^2/2) + \beta Y_{i\nu/2}(ia^2/2) \right\} \quad (28)$$




Figure 2: The WDW potential (solid line) for the massless scalar field. Now a forbidden region occurs, due to curvature $k = 1$, at large scale factors with corresponding exponentially growing and decaying solutions. The oscillating solutions gain arbitrary short pitch as $a \rightarrow 0$ representing a singularity with diverging kinetic energy.

$$\Psi(\phi) \sim \exp(i\nu\phi) \quad (29)$$

where J and Y are Bessel functions (see e.g. [187]) and each term has an associated arbitrary constant α, β , which we can choose accordingly.

First consider the limit $a \rightarrow 0$. Using the asymptote $J_\mu(z) \sim z^\mu$ as $z \rightarrow 0$ enables the solution to be expressed as

$$\Psi(a) \sim a^{(1-p)/2} \exp(i\nu \ln a) \quad (30)$$

There is a divergence as $a \rightarrow 0$ producing an infinite oscillation representing the classical singularity as the kinetic energy of the scalar field diverges [188]. A similar divergence also occurs for $\phi \rightarrow \infty$.

But as it stands, the solution can also be regularized by integrating over the arbitrary separation constant. Now the integral

$$\Psi(a, \phi) \equiv \Psi(a)\Psi(\phi) \sim \int \exp(i\nu[\ln a + \phi]) d\nu \quad (31)$$

is of the form $\int \exp(ixt)dt$ which by means of the Riemann-Lebesgue Lemma tends to zero as $x \rightarrow \infty$ (see eg. ref.[189]). The wavefunction is now damped as $a \rightarrow 0$ or $\phi \rightarrow \infty$. There is another possible divergence for factor ordering $p > 1$ but we have assumed its coordinate invariant value of unity [190]. It has

also been suggested that in the context of wormhole solutions these milder divergences due to the factor ordering are not particularly serious [191]. They are also present for flat empty space, so they conceivably anyway should be renormalized away [7]. Further considerations of factor ordering on the various proposals are given in [192]. For large scale factor the wavefunction eq.(28), behaves as $\sim \exp(-a^2/2)$ so indicating asymptotically Euclidean space cf.[188].⁵

In general one can impose a boundary condition like De Witt's original $\Psi(a=0)=0$ suggestion [66]. As it is a bound state problem, square integrable wave functions can be obtained so giving a normalizable wavefunction cf.[193]. However, this boundary condition at short distance does not necessarily determine the behaviour at large distance where possible growing and decaying exponential solutions are present cf.[194].

For later use we can also mention, as an example of scalar-tensor gravity, the Brans-Dicke model which is derived from the following action e.g.[31]

$$S = \int d^4x \sqrt{g} \left(\phi R - \frac{\omega}{\phi} (\partial_\mu \phi)^2 \right) . \quad (32)$$

For stability in Lorentzian space one requires $\omega > -3/2$.

Using standard techniques, the corresponding WDW equation can be obtained [185]. Only the equation for $\Psi(\phi)$ differs from the previous case

$$\Psi(\phi) \sim \exp(i\sqrt{B\nu} \ln \phi) \quad (33)$$

for $q=1$ and $B=(3+2\omega)$. So the oscillatory divergence now occurs for $\phi \rightarrow 0$ as well. Again one can integrate over the separation constant to produce a regular wave function

Since in the limit $\phi \rightarrow 0$ the Planck length $l_p = G^{1/2} \rightarrow \infty$, you might have expected this divergence. There is also an increasing pitch or "wiggleness" of oscillation in the $\phi \rightarrow \infty$ limit: now at large distance beyond the Planck length. Although less severe than the previous divergence the pitch can still develop at arbitrarily short distance. We can expect these rapidly oscillatory wavefunctions more generally for scalar-tensor gravity models, including non-minimally coupled scalar fields cf. [195]. Also higher order correction to the gravitational action typically correspond to additional scalar fields in the Einstein frame, see e.g.[196].

⁵Provided the combination $J+iY$ which equals the first Hankel function $H^{(1)}$ [187,188] is chosen.

One can see a similar behaviour in the WDW solution for a pure cosmological constant. The pitch of the solution gets increasingly shorter at large distance a cf.(Fig.1). This is simply because the universe keeps accelerating and so the “velocity” $\dot{a} \rightarrow \infty$. In general although the $\Psi(a)$ part of the solution might be regularized at short distances the actual solution $\Psi(a, \phi)$ can oscillate at arbitrary short pitch due to the kinetic energy of the matter component. Only simple “on-shell” perfect fluid models allow the matter to be expressed in terms of the scale factor. For the FRW case the scalar field has an extra degree of freedom over a perfect fluid e.g.[197]. Note also that both signs for the kinetic term are automatically included without further restrictions on the separation constants e.g. [198].

3.3 Scalar potential $V(\phi)$ case

We now generalize to a inflationary scalar potential. In the common boundary conditions such as the no boundary or tunnelling ones the massless component matter is disfavoured or suppressed. This therefore allow any inflationary matter present to become dominant during the early stage of the universe. The separation constants are effectively forced to be zero by the imposition of the boundary condition which in turn prevents any kinetic energy component from causing an infinitely oscillatory wavefunction.

Approximately, the two boundary conditions give the same results for $V(\phi)$ as the earlier cosmological constant case but with $V(\phi)$ taking the place of the Λ . One therefore obtains $\Psi_{HH} \sim \exp(1/V(\phi))$ and $\Psi_T \sim \exp(-1/V(\phi))$ for the Hartle-Hawking and tunnelling cases respectively.⁶ One can obtain conditional probabilities for the initial value of field ϕ . The tunnelling boundary condition peaks at large values of ϕ so is more inflationary than the HH case [1-3]. It has been argued that if one allows arbitrary large energy densities then even the HH case can also give sufficient inflation [201]. Also other matter sources like the four-form or boundary effects can alter this prediction. Particle creation effects could also occur during the tunnelling process [202]. See also [203] for further debate on the relative merits of the various proposals.

It would seem preferable to try and work without a particular boundary condition in the absence of strong reasons for a particular choice. This is the usual “principle of indifference” in the absence of other knowledge. Grischuk and coworkers [204-206] tried to obtain the average solution for the WDW

⁶The HH case was extended to complex metrics by Hartle [199] and another tunnelling type proposal is given in [200].

equation with a massive scalar field source. They found that the typical solution was more like the tunnelling one, in that $\Psi(0) > \Psi(1/\sqrt{V(\phi)})$ so suggesting a tunnelling from the origin. The HH case corresponds to an exponentially increasing solution from the origin. Also with some arbitrariness in putting a measure on a “sphere of quantum states ” they obtained a similar measure at the quantum boundary as previously obtained by classical equipartition arguments: this strongly favours inflationary behaviour - see section (2.1) . One might hope that this sort of argument can be generalized to other matter sources and inhomogeneous degrees of freedom. Some more realistic supersymmetric matter sources are explored in [207].

One might still question whether this strong energy matter suppression is reasonable and indeed it has been suggested that “zero point” fluctuations alone could alter this picture [208]. We represent this in Fig.(1) by the solid arrows where quantum effects will cause the potential U to be ill defined and allowing a Lorentzian region to also develop around $a = 0$. A classically allowed region near the origin could therefore result. It is also the case that the presence of strong energy violating matter is a strict requirement for these common boundary conditions otherwise no natural Euclidean or forbidden region would be present. More generally it is suggested that the surface between Lorentzian and Euclidean regions should have zero extrinsic curvature $K_{ij} = 0$: so called *real tunnelling geometries* [209]. In simple geometries this requires being at a stationary point in the scale factor i.e. $\dot{a} = 0$. So the surface is either the bounce point (cf. de Sitter) or at the maximum in a closed re-collapsing model. In ref.[210] the HH approach was applied to start the universe at its maximum size before then collapsing.

Other predictions that distinguish various interpretations of quantum mechanics have been investigated Bohm “pilot wave” e.g. [211] or many-worlds [212]. A further extension to field theory “universe” creation operators or so-called 3rd quantization has also been pursued [213], but criticized in ref.[161].

3.4 Further Topological and Geometric aspects

There is also the issue of including different topologies and geometries for the tunnelling amplitude in the more general case. However, proving whether 4-manifolds differ and so knowing if they are being over-counted in the path integral is undecidable. It has been suggested that the wavefunction should not be such a turing non-computable number [214]. Although whether this would have serious limitations on using Ψ is disputed since we only require limited accuracy in our predictions [215]. One can also work with conifolds:

and other more general notion of manifold to help alleviate this problem [216].

If the number of manifolds for the hyperbolic case can approach infinity it can overwhelm the usual suppression factor for the creation of a single universe with a given set topology [217]. The “average” topology might be able to predict the spatial homogeneity of the universe [217,218]. But again is this really reasonable? It implies an initial state or reservoir of all infinite possible topologies that should be included in the amplitude. There are now infinitely many “particles” one for each possible topology and geometry. Neither is it clear why just a single universe with an average topology results and not that many universes each with different topology form together. Working with closed models, and so fewer possible topologies, see e.g. [219], could alleviate this problem but the notion of curvature itself will also become hazy at the Planck scale.

Indeed the way that curvature is treated as a constant is rather unsatisfactory. In FRW models the actually local characteristic k is taken to be globally constant. In more general metrics the curvature can become a function also of time and space $k(t, x)$ cf. Stephani models e.g.[34]. For the FRW model with perfect fluid $p = (\gamma - 1)\rho$ the WDW potential takes the form, e.g.[197]

$$U = ka^2 - Aa^{4-3\gamma} \quad (34)$$

where the constant A can be obtained from the relation $\rho = A/a^{3\gamma}$. For a forbidden or Euclidean region at small scale factor a requires $U > 0$ which requires $k = 1$ and violation of the strong-energy condition i.e. $0 \leq \gamma < 2/3$. However, in a more general inhomogeneous model this behaviour can be drastically altered. For example in a Stephani model the corresponding WDW potential becomes cf. [220]

$$U = \beta a^n - Aa^{4-3\gamma} \quad (35)$$

so for $n > 2$ the forbidden region can be either narrowed or absent entirely even for closed models $\beta > 0$ and when the strong energy condition is being violated. This example is symptomatic of what, more realistically, can be expected as the Planck epoch is approached. Non-minimally coupled scalar fields or the presence of large anisotropy can also remove the barrier [221]. The presence of forbidden regions that play such a prominent role might then actually be absent even for closed models that can display inflation. Another possible complication is that a negative energy density can also create a

forbidden region. For example in a toroidal model with compactification scale L , one typically obtains a Casimir term, e.g.[222]

$$\rho = \langle T_0^0 \rangle = -\frac{\alpha}{L^4 a^4} \quad (36)$$

with a the scale factor and the constant α depending on the nature and number of matter fields present. More elaborate twisted scalar fields can also be possible [223]. This Casimir term corresponds to a $\beta > 0$ and $n = 0$ term in eq. (35) [156].

Even without a forbidden region some boundary conditions might be adapted to purely Lorentzian metrics, although the underlying principle is then often less prescriptive cf.[184]- where the “outgoing only” aspect of the tunnelling boundary condition was implemented in such a case.

Further examples of the WDW equation with variable constants [224] or variable space dimensions [225] have also been done.

3.5 Brane quantum cosmology

We here just wish to address a number of complications that brane models appear to have when they are assumed to be created from nothing by quantum processes. Recall that usual matter is confined to a brane existing in a bulk space. In the Randall Sundrum II model the brane is embedded in 5 dimensional Anti-de Sitter (AdS) space [90].

i) One can consider either a pre-existing bulk with branes being created or the bulk and brane appearing simultaneously [226,227]. Some related schemes are given in [228]. A similar situation can occur with multidimensional models with tightly compactified extra dimensions where *external nothing* and *total nothing* were distinguished [229]. If there is already a pre-existing bulk it can help fix the relevant outgoing only tunnelling boundary condition [227]. However, if branes can be spontaneously produced within a pre-existing bulk there are questions as what production rate is possible? A similar concern was present in earlier ideas that the universe spontaneously occurred in Minkowski space cf.[149,150]. Without some removal mechanism an infinite number of branes would eventually be produced within a fixed bulk.

ii) If we also require creation of the AdS bulk space then one needs to use a rather convoluted “cut and paste” procedure to produce a compact space with a Cauchy surface [94]. Recall the geodesically complete closed AdS space is not globally hyperbolic [27] and boundary conditions would be required to regulate continuously the time-like infinity.

iii) For creation of say compact de Sitter branes in Euclidean AdS space we have a mixture of negative action (de Sitter) and positive action (AdS) parts. Can these be simply added or should the individual actions be first rotated to positive values as expected for tunnelling processes. One could “cut and paste” two manifolds with equal and opposite action so allowing unlimited creation possibilities. This suggests that action alone might not convey the true complexity of such processes.

iv) Usually we might expect Euclidean space to be confined to small size. But if a large AdS space is also produced the Euclidean nature is being allowed to arbitrary large scale. This is related to the previous point about cancelling different actions. However because the bulk is a static space it can obey the real tunnelling condition that the surface between Euclidean and Lorentzian has no extrinsic curvature cf.[209].

v) The brane can have an induced curvature from being embedded in the higher dimensional bulk. The full action can have extra terms that, although not contributing to the equations of motion of the brane itself, affect the quantum calculation cf.[230,231]. It is also important that the brane remain compact to prevent an extra surface term, something not automatically resulting from taking a compact bulk space.

A number of extra issues therefore have to be clarified before any calculation of brane creation can properly proceed. For this reason one can instead attempt to promote the branes to an eternal existence just as the static nature of the bulk space would suggest. This leads one to cyclic universe models with possible brane collisions. One example again is the ekpyrotic scenario [48]. We would just mention that we do not think that entropy growth has been adequately dealt with in this, and related scenarios, and that the models will generally suffer the same problem as the earlier Tolman oscillating model. Inflation is used as an entropy sink, but inflation does not actually reduce entropy which would contradict the generalized 2nd Law of Thermodynamics [130]. The problem is closely related to whether information is destroyed e.g.[232] or preserved by black holes [233]. If information is lost then the 2nd Law might be violated over large time scales cf.[226]. The problem is further compounded by using an already infinite sized universe *ab initio* to hide the problem of entropy production. This is also done in ref.[74] where continual expansion is apparently used to dilute the entropy in the universe to effectively flat space. For a finite sized universe continually increasing the entropy would eventually surpass the critical density and so produce re-collapse to a future big-crunch singularity. It therefore seems a

sleight of hand to claim that flat space is the maximum entropy state cf.[74].

3.6 Universe from a quiescent or static state

We have spoken of the universe starting from nothing or by bouncing from a previously collapsing phase. A third possibility is that originally the universe was initially stuck in some unchanging quiescent state. Perhaps involving the presence of closed timelike curves (CTCs), see e.g.[29] for a review. This is closely related to introducing a topological identification scale as in Misner space e.g. [29].

Starting with Misner space, Gott and Li [208,234] obtained a self-consistent adapted Rindler vacuum state for a conformally coupled scalar field that remains finite at the Cauchy horizon, unlike for the Minkowski case [235]. They then conformally transformed this state to give a suitable vacuum state for multiply connected de Sitter space. Such a de Sitter space with CTCs might then be an initial state for the universe. It only has retarded solutions so giving a possible “arrow of time” and is a state of low entropy, actually of zero temperature [208] .

However, in Misner space this state was only possible with identification scale $b = 2\pi$, or $b = 2\pi r_0$ for the multiple de Sitter case [208,234] . Such an exact value is itself inconsistent with notions of quantum uncertainty. We are therefore wary of claims that such a multiply connected de Sitter state is stable especially since the relevant time loop is approximately $\sim$ Planck time, only a plausibility argument has so far been made [236].

The actual procedure of balancing a negative starting vacuum with a Hawking radiation, due to the periodicity, to give an empty vacuum state has further possible difficulties. The calculation makes use of the periodicity producing a thermal state [237] . Such a state is required to be a many particle state with technically a suitably large Fock space, see e.g.[106]. But by being close to the Planck scale one starts reducing the number of allowed states due to holography type arguments [75]. This will start preventing an exact thermal state, as also is expected during the final stages of black hole evaporation [238] or in Planck scale de Sitter space [38,239]. This mismatch could then result in some fluctuations still being present in the vacuum instead of a pure empty state, so destabilizing the CTC.

Neither is it clear that the b value, or the corresponding de Sitter one, remain independent of different matter couplings ξ or potentials $V(\phi)$. A more realistic combination of matter sources still appears divergent at the Cauchy horizon [240], although an improved *self-consistent* renormalization procedure [241] in Euclidean space might help regulate some of these other

cases.

Creating this state in any case seems rather contrived. Recall that the Rindler vacuum of accelerating observers requires “mirrors and absorbing stray radiation”, before we then make any topological identification [242]. One would need some more general reason why such an initial state was actually present. The analogous zero temperature state for charged Black holes has proved difficult to obtain on grounds of stability [243].

Instead of requiring CTCs one might just allow a static state with time still evolving normally from say $-\infty$. There is a recent emergent model [244], an update of the Eddington-Lemaître model e.g. [30] that starts from an Einstein static universe. Because this model has no forbidden region, and requires a balance of ordinary matter and a cosmological constant, it again will be prevented by the usual boundary conditions that bias against the normal (e.g. radiation) matter component. Neither do we think that maximizing the entropy is a more suitable principle for determining the boundary condition since the entropy actually grows later during the inflationary stage cf.[238]. The emergent model is however geodesically complete to the past unlike the previous case of multiply connected de Sitter space. One might do a further identification of antipodal points in the multiply connected space but one then loses time orientability. This would make the problem even more involved cf. [245]

Also such a model also requires a mechanism to stabilize the Einstein static phase to homogeneous perturbations, although the model appears stable to inhomogeneous perturbations [246]. More general inhomogeneous models might allow this. For example, by altering the curvature dependence as in eq.(35) one could produce a stable static universe with a now flat $U = 0$ WDW potential; or perhaps, at least prevent collapse to the origin by means of a repelling potential $U \gg 0$ around the origin $a = 0$, cf. a Casimir term [156].

Such a repelling or Planck potential has been added previously in an *ad hoc* manner to produce a forbidden region around the origin [247]: a term $\propto a^{-2}$ is added to the WDW potential U . This can then allow a De Witt like boundary condition $\Psi(0) = 0$ [66]. Also a non-zero separation constant can display a similar effect [248] cf. eq. (26). Since, as previously mentioned, a massless scalar field automatically includes such separation constants it can alone be used to produce a bouncing model as in [249]. It is also claimed that a perfect fluid can give such behaviour [250], but this would require an asymmetric Wick rotation cf.[198].




Figure 3: The WDW potential (solid line) for the unstable Einstein static universe. Reversing the potential (dotted line) would produce a stable static state.

One might also try to stabilize the Einstein static universe more generally by surrounding the state entirely with forbidden or Euclidean regions. For example if the sign of the WDW potential U is flipped the corresponding Einstein static universe becomes stabilized. Such a model requires $k; \Lambda; \rho; \rightarrow -k; -\Lambda; -\rho$, so now this is an open AdS with an extra negative radiation component. So violation of the weak energy condition is now required for such stability, such as might occur with the Casimir effect.

A stabilized state could also be achieved without altering the matter component by use of a signature change, represented by the parameter ϵ : $\epsilon = 1$ for usual Lorentzian space and -1 for Euclidean space [251]. In the simplest case the corresponding WDW potential is altered $U \rightarrow \epsilon U$ [197]. So if for some reason $\epsilon = -1$ the previous static universe is again stabilized. Other possible examples of signature change, starting from different action principles, are also possible [252].

It has been suggested that oscillations of ϵ between the two cases are constantly occurring but that the “average” now favours a Lorentzian space-time [253]. One might imagine instead a preponderance of negative Euclidean

values for ϵ . This might help stabilize a static model before for some reason the sign changed and Lorentzian evolution then could proceed.

Despite these present difficulties the notion of finding a suitable quiescent state has some attraction. The difficulty, as in the examples given, is why the state should survive for semi-infinite times, but still have some slight instability that causes the expansionary or Lorentzian evolution to eventually begin. For example, a previous model [254] suggested eternal oscillations around the Planck size could exist but it did not then produce a large universe in the infinite future.

4.0 Loop Quantum Cosmology

Another trend to quantum gravity is the loop approach that treats gravity in a similar way to other gauge theories. One works with holonomies and Wilson loops, previously used in non-gravitational field theories [14]. A number of reviews have been done on technical issues involving this approach [14,15,21]. Earlier work obtained exact solutions of the WDW equation using Knot theory invariants: the so-called Kodama state [255]. So far these models have proved difficult to interpret or implement in cosmological model beyond a simple cosmological constant [256] or slow roll scalar potential case [257]. A number of related approaches to quantum gravity have emphasized the discrete nature of space as you approach the Planck scale [15,21]. This might alter the application of quantum mechanics, for example a Bohr quantization scheme was claimed to give a more effective avoidance of singularities [258].

We will mostly consider the Thiemann approach [14], within loop quantum gravity, that has most been developed to a level that can make contact with realistic cosmological models [13]. One claimed advantage of loop quantum gravity over strings is that a background independent formulism might easier be achieved [2]. However this might actually be a hindrance for realistic cosmologies cf.[259] where *restricted covariance* was considered more realistic. Another possible drawback is that in GUT theories the various forces of nature should eventually unify. Therefore the present weak force of gravity should increase with energy scale to eventually coincide with the other forces of nature. The Planck length $\sim G^{1/2}$ will correspondingly grow as the unification scale is approached. But in loop gravity this aspect of “running” Planck length does not appear incorporated at present. A different initial Planck scale would correspondingly alter the various cosmological *puzzles* and so possibly alter the amount of inflation required.

At first sight any possible discreteness around Planck scales does not ap-

pear too helpful in actual cosmology since, in particular, flat or open FRW models are infinite from the start, so the granularity is never actually apparent. One can make a cut to enclose the big bang within a finite volume as time $t \rightarrow 0$, but then the matter or energy momentum becomes infinite within such a domain [260]. By making a compactification, at a scale L , this infinity can be alleviated to some extent.⁷ But still the matter densities will diverge as the universe is evolved back towards the initial singularity: likewise for closed FRW models. Incidentally compactifying in this way introduces possible vacuum polarization and Casimir like effects e.g.[222]. Typically the energy density ρ would become negative so violating the weak energy condition, unless supersymmetry intervened to suppress such terms.

4.1 Bouncing and Inflationary model

Interestingly, loop quantum gravity is also said to alter, possibly every matter component such that the weak energy condition is effectively violated at short distances when the granularity of space becomes significant. Recall that usually matter is diluted for an expanding universe or remains constant for an exact de Sitter solution. Likewise for the Casimir type component above.

By allowing the energy density to grow during expansion there need not be a divergence in the energy density as the universe is evolved back in time. One possible advantage is that now a bounce from an earlier collapsing phase might be implemented close to the Planck length scale. Usually one needs to bounce before the Planck energy density is surpassed by ordinary matter.

However, if the previous phase of the universe already contained large amounts of matter and entropy it would not necessarily require inflation. But then the bounce would have to proceed long before Planck size lengths are reached in order for Planck energy densities not to be vastly exceeded. Alternatively if the previous phase was devoid of matter then it can collapse to the Planck size. Then a later inflationary phase could produce sufficient matter and entropy. The previous collapsing phase of the universe is effectively empty: a similar stage occurs in the pre-big bang model [100]. Another less likely scenario is that a deflationary phase (rapid collapse) could remove matter and radiation: but this would seem to violate standard notions of quantum unitarity together with the generalized 2nd Law of thermodynamics e.g.[28,130]. In this case all matter component energy densities scale as $\rho \propto a^n$ with $n > 1$ as the bounce approaches and a rapid collapsing or

⁷This scale L seems a further arbitrary constant cf.[257].




Figure 4: (i) Bouncing universe - matter content symmetrical before and after bounce; (ii) Bouncing universe with extra inflationary expansion - presently favoured in loop quantum approach ; (iii) Bouncing universe with deflationary phase and inflationary phase - allows universe to remove matter and so bounce at smaller size without violating Planck energy densities.

deflationary behaviour is possible.

It is therefore not entirely clear whether an inflationary phase, following a previous collapsing epoch of the universe, should be implemented. Many of the so-called *puzzles* will be different to that of the usual big bang model and need a more careful consideration. Recall that inflation can take a “small bang” universe to that of a big bang: so effectively behaving as a universe amplifier. So only if there is a bounce away from a “small crunch” does inflation have to intervene to give a later “big” universe. But then there is a mismatch with the earlier preceding universe that it was not collapsing towards a big crunch before evading the singularity. This can be contrasted with the general point that producing inflation requires low entropy conditions [23,24]. Whereas a collapsing universe would be expected to have increasingly growing entropy for the usual direction for the arrow of time.

4.2 Modified Wheeler - De Witt equation

We can now consider a few specific issues for the loop quantum cosmology approach. Again we stick to the cosmological issues as much as possible. The scale factor is defined in terms of the triad component p where $a^2 = |p|$ [13,14]. At large distance the usual WDW equation should be produced. But

this continuum limit is given by, for a closed model and ignoring the matter Hamiltonian, [261-266]

$$\left(\frac{d^2}{dp^2} - 1\right) \sqrt{|p|} \Psi(p) = 0 \quad (37)$$

Writing this in terms of a gives

$$\left(\frac{d^2}{da^2} + \frac{1}{a} \frac{d}{da} - \frac{1}{a^2} - a^2\right) \Psi(a) = 0 \quad (38)$$

But this now automatically includes a repulsive Planck potential around the origin cf. eq.(26). It is unclear whether this operator ordering is actually imposed by the formulism or whether an arbitrary choice has been made cf.[263]. Partly this is imposed by “pre-classicality” that the pitch should not fall below the Planck length. However a constant inflationary solution develops arbitrarily short pitch and if such solutions are prevented it could, for example, prevent eternal inflation to the future. It is claimed [263] that this de Sitter example is only an “infrared problem” and can be ignored since the local curvature is still small. But this distinction seems arbitrary and so using the pitch alone would not impose an unambiguous choice of factor ordering.

If the factor ordering that gives an effective Planck potential is totally justified it can be used to produce a $\Psi(0) = 0$ type boundary condition [267]. By introducing a forbidden region around the origin it also can be responsible for causing a collapsing universe to rebound cf. [248]. This Planck potential suppresses any decaying mode corresponding to $D > 1$ in section (3.3) so disallowing any tunnelling possibility from the origin. Only an exponentially increasing mode is present in the forbidden region so the boundary condition is closer to the Hartle-Hawking one cf.[267].

With this reasoning loop quantum cosmology has given a justification for earlier ideas about modifying the WDW potential at small a . Incidentally, such a correction to the WDW potential could alone prevent a Mixmaster phase in Bianchi IX models: since it is known that the presence of an extra stiff fluid prevents chaos cf.[268,269]. Equation (38) corresponds to a stiff fluid but with the wrong sign. Chaos is also removed by reducing the number of degrees of freedom, if matter ϕ simply becomes a function of a , chaos cannot occur anyway e.g. [270].

At still shorter distances the loop effects cause the WDW equation to be replaced by a difference equation. One can apparently iterate to regions

with negative p , a so-called orientation change during the previous collapsing universe phase. However, if a forbidden region occurs at large scale as in a closed universe model the boundary conditions have to impose the relevant solution - see Fig.(2) where the growing and decaying modes are sketched. In order to obtain just the decaying mode one needed to chose a suitable Hankel function for the usual massless scalar field example. In ref.[271] they have fixed this solution correctly at the maximum size of the closed universe and iterated back to the corresponding point in the oppositely oriented universe. The solution there no longer just decreases into the forbidden region. This might have been anticipated since the Hankel function corresponds to a wave moving in, say, a right direction, so one would need it to switch to a left moving direction in the oriented epoch. In general it looks at though alterations at short distances does not resolve all the ambiguities of boundary conditions. Indeed it has previously been emphasized that quantum behaviour is not necessarily confined to small scale factors only cf. e.g.[272,254].

4.3 Loop cosmology and super-inflation

At intermediate scales there is a possible semiclassical effect that alters the behaviour of a scalar field matter source. This effect occurs up to distances a_* that depends on an arbitrary parameter j . Provided an inflationary phase is still necessary a new mechanism might now be possible. The idea is that a massless scalar field can itself become inflationary. The kinetic energy increasing as the universe expands, typically $\dot{\phi} \propto a^{12}$; this now corresponds to an effective $\gamma \sim -3$ so strongly violating the weak energy condition. This produces pole-law like inflation but alone is not sufficient to solve the usual puzzles of standard cosmology: for one thing the final value of the scale factor a_f should be $\sim cm$: so that the subsequent universe be large and produce mass density $\sim 10^{-30} gcm^{-3}$ today [50,54]. This would require the parameter j [13] to be extremely large so that loop effects are prevalent up to cm distances. Another problem is that pole-law inflation has a corresponding growing Hubble parameter that produces a blue spectrum of perturbations [273,54]. Because the size of gravitational waves is given by the Hubble parameter it should not be allowed to become larger than $\sim 10^{-5} m_{pl}$ [274].

It was then suggested that since the pole-law inflation was not alone sufficient, a second potential driven inflation could be produced [264,266,275]. This is because the friction term in the Klein-Gordon equation can change sign and possibly drive a scalar field up a potential. However, one can show that growth in ϕ is negligibly small while the kinetic term is growing at its maximum rate. Enormous changes in $\dot{\phi}$ are occurring for only small eventual

values of ϕ . In order to prevent kinetic energies beyond the Planck scale being produced, with allowance made for this large growth in $\dot{\phi}$, puts severe constraints on the initial values of $\dot{\phi}$. Typically for the chosen ($j \sim 100$) parameter the initial value is constrained such that $\dot{\phi}_i^2 < 10^{-12}$. This value alone is too small to drive the field significantly up the potential and so produce sufficient overall inflation. Later [276] other quantization schemes, with a second arbitrary parameter l , were suggested that can produce other functions of the scale factor that can also go in front of the potential term $V(\phi)$. But even then, contrary to the impression of ref. [276], very little phase space for sufficient inflation is actually present if the field is not initially displaced from its minimum. Also other, possibly massless scalar fields, would have to be suppressed and spatial gradient terms not allowed likewise to grow rapidly with scale factor. Incidentally, in Euclidean space the friction term is also switched in the corresponding Klein-Gordon equation. This “anti-friction” mechanism has been used heuristically in conventional quantum cosmology to explain a large initial field cf. [1].

The mechanism therefore has negligible effect on the overall measure for inflation. Recall earlier argument that equipartition for initial $\rho \sim 1$, or in term of the Planck mass $\rho \sim M_{pl}^4$, only failed to give inflation when $\dot{\phi}^2 \sim 1$. But these large values of kinetic energy are not suitable to improve further the probability of inflation f_i already strongly favoured around $\sim (1 - 10^{-5})$. However, the main uncertainty of quantum cosmology is to determine the initial value of ρ . Perhaps for small initial values of ρ the super-inflation can alleviate the decline in f_i as ρ is reduced. Working with the purely classical canonical measure does not resolve this question since the conjugate variable p_ϕ just changes its a dependence and a related measure is reproduced cf. eq.(8): an infinity of solution would surpass the Planck energy density and these must somehow be first excluded. An infinity of under-inflationary solutions is also present. Unless one can argue that “repeated tries” are allowed, which is closely related to an anthropic argument, or else suggest quantum gravity processes would intervene once the Planck energy density epoch is again reached. But the rough uncertainty principle in energy/time, which is here marginally more correct than a non-relativistic position/momentum uncertainty principle [275,276], already biases strongly towards inflation: at least for simple homogeneous scalar field models.

The Hartle-Hawking proposal seems to give a small ρ but the initial size of the universe is correspondingly large $a_i \sim \rho^{-1/2}$: this is vastly larger

than a_* so it would never be in the loop quantum regime. But if the model is actually bouncing not because of a small effective cosmological constant but rather because of a changing behaviour of the kinetic term this bounce could occur for size less than a_* . The size depends on the coefficient A in equation (3) and can be set $A \sim 1$ by scaling into the definition of a_* using the Planck length and Barbero-Immirzi parameter [14]. Incidentally looking at equation (3) we can see clearly why a bounce requiring $m > n$ proceeds in the closed ($k = 1$) case, since now $m = 2 > n \sim -12$; or by using a negative cosmological constant ($m = 0$) in place of the curvature cf.[277-279]. However, either a) any ordinary matter must be absent: which seemingly violates expected entropy increase, or b) all types of matter are altered to grow with a modified behaviour, as say $\rho \propto a^r$ with $r > 1$ during the loop gravity phase.

Normally the kinetic energy is the stiffest matter and $\rho \propto a^{-6}$ changes its a dependence by a factor a^{15} so that $r = 9$ [13,275]. It requires all other matter sources to have a smaller effective value for r less than 9, if such a mechanism is to be an attractor; or to prevent another source growing earlier to Planck densities before the kinetic energy itself can grow significantly. The spatial gradient terms should also not be allowed to have this different a dependence or else the universe is driven inhomogeneous and any subsequent scalar driven inflationary phase is unlikely. This dilemma awaits the consideration of going beyond simple scalar field examples. One might also worry that a purely gravitational vacuum terms like gravitational waves, or the Bianchi anisotropy parameters β_- , β_+ , could alter their behaviour during possible Kasner evolution. Although it is shown [268] that chaotic Mixmaster behaviour is prevented by means of the scalar curvature being bounded the shear term $\sigma^2 \propto a^{-6}$ itself might be altered. Especially since if the universe is contracting in one direction (while still expanding in the two others) this direction could be affected more by loop effects. Although still the total volume of the universe remains decreasing.

4.4 Is Loop quantum cosmology unstable?

There seem a number of dangers in promoting the scale factor into a true fundamental distance. For a massless scalar field the energy density ρ is now modified at short distance such that $\rho \propto a^n$. The energy density now disappears as $a \rightarrow 0$ and so is almost indistinguishable from flat space. But this suggests a danger that actually any Planck sized region is now potentially unstable to this inflationary expansion.

One might try and reason that for a Planck length region, within a pre-

existing universe, to inflate it requires a negative pressure that will be quickly equalized by the greater average pressure of the universe outside. This was one of the reasons that creating a universe in the lab is difficult because of the surrounding background metric [141]. But while such an equalization is taking place there is the possibility of a quantum tunnelling occurring to a new baby universe. This does not supplant the original universe but disconnects forming a new universe. In standard potential driven inflation such a scheme requires one to produce a high energy density false vacuum state that then has a minuscule chance of tunnelling to produce a new universe [141-143]. But now any Planck size region automatically could make such a transition providing topology changes are not forbidden on other grounds, see e.g.[29] for introduction to topology issues. In the lab it required huge effort to violate the strong energy condition, but now if the weak energy condition is continually being violated at short distances it seems easier to conceive of such tunnelling. One might try and quantitatively calculate this enhancement but there is another ambiguity: there is an arbitrary compactification scale for flat or open cosmological models. Usually, these cases have infinite action due to infinite size and are discounted, see e.g. [151]. But now with a finite volume V and energy density decreasing with size such universes are not apparently suppressed on action principles alone, $S = V \int a^n dt \rightarrow 0$ as $a \rightarrow 0$. In the closed case a forbidden region is present so that the created universe must start with at least a certain size cf.[266]. Placing this value beyond the weak energy violating region might help suppress the universe creation effect but this would introduce fine tuning.

We can also consider the creation of the original universe *ex nihilo*. Now even in standard quantum cosmology it is not entirely clear why universes are still not being created around us. You can try and argue that the forbidden region creates a barrier that to observers within the existing universe suppresses further universe creation e.g.[280]. But this barrier is either absent or reduced in loop quantum cosmology and also in some standard cosmological models. One can further distinguish between creating an isolated universe and one being formed within a pre-existing universe which requires gradient energy around the linkage [281]. If only new unconnected universes are easily created then you can argue that this is not a significant problem; although it might seem profligate if the mechanism causes constant new universe production.

There is second type of potential instability if the universe does starts inflating. Because we have particle or event horizons we only have causal

contact up to finite distance - see e.g.[27]. During high energy inflationary expansion this size could easily be below a_* . In other words, only from the perspective of being outside the universe is the scale factor really defined. To stop the inflationary expansion within a Hubble volume the scale factor needs to play another “non-local” messenger role. We seem in need of a sort of generalized Mach’s principle e.g.[30,33], telling the individual “granules of space” how big the total universe has become. A related concern [146] seems present in certain variable constant theories, that also use the scale factor to determine, for example, the actual speed of light value. Maybe this adapted Mach’s principle only needs to work up to some suitable size but it still occurs over distances not normally believed causally connected.

The idea of space being made of discrete quanta can introduce further conceptual problems. In an expanding model new cells have to be produced to fill in the gaps. But if we make analogy with cell division in living organisms, how are cells produced without error? Because presumably there is no analogy with DNA, there seems the need of providing “scaffolding” to force cells to have their correct form. Normally expanding space is simply stretched like an elastic band by the scale factor. Even then gravitationally bound systems e.g. galaxies can drop out of the expansionary global behaviour of the universe: so the scale factor never plays a universal messenger role to individual atoms. It seems difficult that loop quantum cosmology can therefore distinguish that normal matter should display weak energy condition violation only when the universe is small and not just at any small distance.

Bojowald has suggested to me that, in the above language, there now exists a scaffolding preventing such exotic behaviour due to the universe now obeying the “average” classical description. We are suspicious about how this is propagated and that the super-inflationary phase once started need never stop for $H^{-1} < a_*$.

We can also question how Einstein’s equations $G_{\mu\nu} \Leftrightarrow T_{\mu\nu}$ have been used to interchange between geometry and matter i.e. $a \leftrightarrow \phi$. It appears a rather “on-shell” restriction that matter can be expressed in terms of the scale factor cf.[197]. For one thing it reduces the number of independent degrees of freedom in the problem. Also, in more general scalar-tensor gravity, or with higher order corrections to the gravitational action, this distinction between the geometry and matter is even more involved. The total solution can have arbitrary high frequency oscillations that cannot easily be confined or excluded by discreteness in the scale factor alone.

Other standard models such as with a simple cosmological constant also get arbitrary short oscillation lengths corresponding to increasing kinetic energy. We have therefore suggested that this property of quantum gravity becoming important at short-distance is not sufficiently universal to resolve many problems. Requiring that also the energy-density be approaching large, or even Planck values, might be a more suitably and locally defined quantity. Future experiments might see whether a discrete structure does exist using gravitational waves, gamma-ray spectrometry or tabletop experiments e.g. [15,282-284].

5.0 Conclusions

A number of cosmological schemes have been considered that typically start from a region where quantum gravity effects become dominant. If the action becomes small then one might reason that quantum behaviour might be apparent. The corresponding universe is typically of Planck size and mass and so requires an inflationary phase in order to amplify the universe to a realistic size.

But such ideas are found to have certain *fragility* problems: like the forbidden region being strongly dependent on how the curvature behaves, or CTCs requiring extreme fine tuning. Analogies with atomic physics, such as tunnelling phenomena, are extrapolated to the universe as a whole. Usually the boundary conditions have been developed apparently with the sole aim of starting the universe in an inflationary state. One then at least must include matter sources in the starting mix that could produce inflation. Although, there is still some dispute whether this is achieved, it does help prevent ambiguities in purely classical measures for the probability of inflation. However, what preceded this inflationary state, and why and how it previously evolved, is far from clear.

Loop quantum gravity has a different approach in that any matter, because of discreteness, is promoted to be inflationary while the universe has a “small” scale factor. Although work has been so far confined to using massless scalar fields. Potentially this is a more robust, or definite prediction of inflation, in that possibly any matter source could initially be chosen. However, because distance is ill-defined such a super-inflationary phase is likely to suffer from a number of other problems: one is that inflation once started would never end if causal distance is fundamentally more valid than the scale factor *per se*.

If the laws of physics preceded the universe we might have some chance of describing the origin of the universe cf.[285]. But if there is only a single

universe then we also have fundamental problems in applying the notions of probability e.g.[286]. Sometimes a pre-existing meta-universe is introduced but its origin and laws are not explained further. Fluctuations within it, however remote, are taken as possibly allowing baby universes to branch off. These might have suitable low entropy so allowing dynamical evolution in line with the 2nd law of thermodynamics to then proceed. Whether the meta-universe itself has dynamical evolution is a further complication cf.[73,74].

We can finally consider some general principles that might be important for quantum gravity explanations of the universe.

- *Action Principle.*

If on evolving back the current universe to a state when the total action becomes small one can expect quantum effects to dominate. One complication is that the gravitational action, once suitably regulated, is not positive definite. One then might conceive of quantum tunnelling to this early minimal action regime. However, this assumes that notions gleaned from standard quantum mechanics can readily be extended to the universe as a whole. It also requires that a pre-existing state of nothing still obey the usual laws of physics: why such a state exists and whether it can spawn infinite other universes is also to be understood

In some brane type cosmologies, with infinite initial bulk space, the branes always have large action *ab initio*: they therefore seem less favourable to quantum type creation. But if no other structures with less action are actually allowed, perhaps on other grounds, they can also be produced. Since there is no notion of “lifetime” outside of the universe, arguments about likely and remote have less validity.

Quantum mechanics can also regulate possible singularities within cosmological models. But it still requires the various matter potentials to be constrained. Alternative quantization schemes like Bohr quantization could help further alleviate possible singularities although this is unlikely to prevent all such examples cf. hydrogen atom in higher dimensions. Large kinetic energies, corresponding to the pitch or “wiggleness” of the wavefunction, might be prevented because of possible discreteness of space. However, such rapid oscillatory behaviour is not readily confined to short distance in the scale factor *per se*. Care is also required if classical field equations are first used to simplify models, that although classically equivalent, can then alter the deduced quantum theory cf.[197].

The probabilistic nature of quantum arguments seems a mixed blessing: it might justify unusual events or prevent the imposition of a single deter-

mined history to the universe. Sometimes it is argued that determinism might still be present underlying quantum mechanics e.g. [287], or that our universe is again embedded within some vast or superior meta-universe. If quantum mechanics just essentially conserves energy but allows a Heisenberg like “borrowing before payback” it does not entirely help address why something is actually present. Energy is anyway ill defined in gravitational systems but in closed universes it can be formally defined to be zero: positive matter energy cancelled by negative gravitational potential energy. But this still does not explain why the universe should have the *potentiality* to occur, even if the universe has no overall apparent cost in energy to produce . This was referred to as the *irremovable problem* in ref.[286].

- *Entropy Arguments.*

One of the requirements of a cosmological scheme is to allow the generalized 2nd law of thermodynamics to occur. One might have expected the initial entropy to be maximized, or perhaps, at least the meta-universe to be in a state of high or maximum entropy.

One difficulty is that the laws of physics appear invariant under time reversal, although solutions do not necessarily have this property. For example, inflation is an attractor during expansion but the reverse deflation is an unstable repelling solution during the time-reversed contraction. Some have suggested that time asymmetry built into the laws themselves is required to explain the time-asymmetry of the actual universe e.g. [19,23]. Without this one needs to understand why the entropy within the observable universe is far below being maximized [23,24,88]. Alternatively you can simply point out that, for say a FRW universe, the entropy has to take the value it does [288] and this just comes from the various equation. Increasing the amount of entropy by adding radiation would simply increase the density parameter above unity i.e. $\Omega > 1$. It then just becomes then a question of why this model universe and not some other.

Quantum gravity could only help if entropy notions can be extended to general gravitational models and perhaps that the cosmological principle or simply inflationary expansion had some obvious preferred status.

The notion that a bounce could occur close to the Planck epoch also seems to have unforeseen consequences. There is the danger that if the entropy is correspondingly small at the bounce point then the arrow of time has run backwards during the collapsing phase. Or else the entropy must initially be set small at the start, presumably $t \rightarrow -\infty$. One then simply transfers the problem of initial conditions to this earlier state cf.[23]. Neither is a bounce

at small size, violating the holography bound, apparently consistent with quantum unitarity: that information should not readily be destroyed. Not only actual matter but the vacuum state itself must also be adjusted. Otherwise, vacuum polarization effects (like Casimir and Conformal anomaly) would be expected to dominate as the universe became small cf.[289].

- *Theory of Everything.*

Perhaps any final quantum gravity theory will be more prescriptive as to what values of various parameters are allowed. On recent example is that the topology might be restricted to only a few possible cases from consistency of string theory [290]. Eventually all arbitrary parameters, for example those that produce microwave background fluctuations, should be determined *a priori* from first principles leaving nothing to choice, see e.g [291] for this point of view. At present many variables are measured or imposed *a posteriori* i.e. after the fact, while others are given by distribution functions.

In such a closely regulated universe it would appear that randomness would play a rather subsidiary role. The final theory's laws might also be expected to display a Ockham's principle like razor conciseness, perhaps having incorporating notions like algorithmic complexity or Fisher information for their formulation [292]. These and related criteria might eventually play a role in explaining the universe's creation cf.[293].

However, the theory of everything might be limited by Gödel like incompleteness: our theories are developed within the universe and cannot take a vantage point independent of or preceding the universe [294]. Incidentally, if the universe evolved from a Planck sized nugget then the laws, vis-à-vis the holography principle, cannot even be represented within the universe at that time - this might be construed as a *reductio ad absurdum* for the principle. The incompleteness problem might be alleviated if, for example, the laws of physics used a restricted arithmetic and self reference did not produce paradoxical situations [295]. Or else the ultimate theory might actually be an infinite number of theories, but most having negligible effect, allows near certainty to be eventually attainable. In any case quantum mechanics alone seems dependent on observers making choices of possible measurements to take and so argues against a single objective history of the universe [296]. Randomness would then play a more significant part in understanding the properties and values of various constants in the actual universe.

Perhaps a single quanta of spacetime - a modern version of Lemaitre's "primeval atom"- is involved or an entirely new conceptual approach (from M theory?) is needed. The cosmological aspects of quantum gravity certainly

remains a fascinating topic for much future work.

Acknowledgement

This work developed from interesting email discussions with Martin Bjowald. I should also like to thank William Hiscock for remarks on the renormalization issue for CTCs.

References

1. J.J. Halliwell, in “Quantum Cosmology and Baby Universes” eds. S. Coleman, J.B. Hartle, T. Piran and S. Weinberg (World Scientific: Singapore) 1991.
2. D.N. Page, in “Proceedings of the Banff Summer institute on Gravitation”, eds: R.B. Mann and P.S. Wesson (World Scientific: Singapore) 1991.
3. D.L. Wiltshire, in “Cosmology: the Physics of the Universe” eds. B. Robson, N. Visvanathon and W.S. Woodcock (World Scientific: Singapore) 1996.
4. L.Z. Fang and Z.C. Wu, Int. J. of Mod. Phys. A 1 (1986) p.887.
5. D. Atkatz, Am. J. Phys. 62 (1994) p.619.
6. J.B. Hartle, in “Proceedings of the 11th Nishinomiya-Yukawa Symposium”, eds. K. Kikkawa, H. Kunitomo and H. Ohtsubo (World Scientific: Singapore) 1998.
7. D.H. Coule, Phys. Rev. D 62 (2000) p.124010.
8. Z.C. Zu, “No Boundary Universe” (Hunan Sci. Tech Press : Changsha) 1994.
9. A.O. Barvinsky, “Talk at 9th Marcel Grossmann Meeting”, gr-qc/0101046.
10. F. Englert, presented at Erice School on “Basics and Highlights in Fundamental Physics” 1999, hep-th/9911185.
11. T.P. Shestakova and C. Simeone, gr-qc/0409114 and gr-qc/0409119.
12. K.V. Kuchar, “Time and Interpretations of Quantum Gravity” in “Proceedings of the 4th Canadian Conference on General Relativity and relativistic astrophysics”, eds. G. Kunstatter, D. Vincent and J. Williams (World Scientific: Singapore) 1992.

13. M. Bojowald and H.A. Morales-Tecotl, Lect. Notes Phys. 646 (2004) p.421.
M. Bojowald, Gen. Rel. and Grav. 35 (2003) p.1877.
M. Bojowald, "Talk at IUFM Marseille , France" 2003 preprint astro-ph/0309478.
14. T. Thiemann, Lect. Notes. Phys. 631 (2003) p.41.
A. Ashtekar and J. Lewandowski, Class. Quant. Grav. 21 (2004) p.R53;
A. Ashtekar, preprint math-ph/0202008;
A. Perez, Class. Quant. Grav. 20 (2003) p.R43.
for a simple introduction see also:
A. Ashtekar, gr-qc/0410054.
C. Rovelli, Phys. World 16 (2003) p.37
L. Smolin, "Three roads to quantum gravity" (Oxford University Press: Oxford) 2000.
some further text books are:
R. Gambini and J. Pullin, "Loops Knots, gauge theories and quantum gravity", (Cambridge University Press : Cambridge) 1996.
C. Rovelli, "Quantum Gravity", (Cambridge University Press : Cambridge) 2004.
T. Thiemann, "Modern Canonical Quantum General Relativity" (Cambridge University Press: Cambridge) *To be published*
15. L. Smolin, hep-th/0408048.
ibid, hep-th/0303185.
16. K. Kiefer, "Quantum Gravity" (Oxford University Press: Oxford) 2004.
17. "Physical Origin of Time asymmetry", eds. J.J. Halliwell, J. Pérez-Mercader W.H. Zurek (Cambridge University Press : Cambridge) 1994.
18. H.D. Zeh, "The Physical basis of the direction of time", (Springer-Verlag: Berlin) 2001.
19. I. Prigogine, "The End of Certainty", (The Free Press:) 1997.
20. "The Future of Theoretical Physics and Cosmology" eds. G.W. Gibbons, E.P.S. Shellard and S.J. Rankin (Cambridge University Press: Cambridge) 2003.

21. “Physics meets Philosophy at the Planck scale”, eds. C. Callender and N. Huggett (Cambridge University Press: Cambridge) 2001.
22. “Science and Ultimate Reality” eds. J.D. Barrow, P.C.W. Davies and C.L. Harper (Cambridge University Press: Cambridge) 2004.
23. R. Penrose, “The Road to Reality”, (Jonathon Cape: London) 2004.
24. R. Penrose, “The Emperor’s New Mind”, (Oxford University Press: Oxford) 1989.
25. E.W. Kolb and M.S. Turner, “The Early Universe” (Addison-Wesley: Reading, MA) 1990.
26. A.D. Linde, “Particle Physics and Inflationary Cosmology”, (Harwood Academic Press: Chur, Switzerland) 1990.
27. S.W. Hawking and G.F.R. Ellis, “The large scale structure of space and time” (Cambridge University Press: Cambridge) 1973.
28. R.M. Wald, “ General Relativity” (Chicago University Press: Chicago) 1984.
29. M. Visser, “Lorentzian Wormholes” (AIP Press: New York) 1996.
30. E. Harrison, “Cosmology 2nd ed.” (Cambridge University Press: Cambridge) 2000
31. J.V. Narlikar, “An Introduction to Cosmology, 3rd. ed.”, (Cambridge University Press: Cambridge) 2002.
32. J.D. Barrow and F.J. Tipler, “The Anthropic Cosmological Principle”, (Oxford University Press: Oxford) 1986.
33. W. Rindler, “Relativity: Special, General and Cosmological” (Oxford university Press: Oxford) 2001.
34. Krasiniński, “Inhomogeneous Cosmological Models” (Cambridge University Press: Cambridge) 1997.
35. S.M. Carroll, astro-ph/0310342.

36. T. Padmanabhan, “Structure Formation in the Universe” (Cambridge University Press: Cambridge) 1993 p.359.
37. L. Liu, F. Zhao and L.X. Li, Phys. Rev. D 52 (1995) p.4752.
38. T. Padmanabhan and M.M. Vasanthi, Phys. Lett. A 89 (1982) p.327.
39. O.M. Moreschi, gr-qc/9911105.
C. Schiller, gr-qc/9610066.
40. J.V. Narlikar and T. Padmanabhan, Phys. Rep. 100 (1983) p.152.
41. S.W. Hawking and D.N. Page, Nucl. Phys. B 298 (1988) p.789.
42. G. Evrard and P. Coles, Class. Quant. Grav. 12 (1995) p. L93.
43. D.H. Coule, Class. Quant. Grav. 13 (1996) p.2029.
44. A. Borde, A.H. Guth and A. Vilenkin, Phys. Rev. Lett. 90 (2003) p.151301.
for earlier proofs see:
A. Borde and A. Vilenkin, in “Proceedings of the Eighth Yukawa Symposium on Relativistic Cosmology” ed M. Sasaki (Universal Academy Press: Japan) 1995.
A. Borde and A. Vilenkin, Int. J. Mod. Phys. D 5 (1996) p.813.
45. E. Calabi and L. Marcus, Ann. Math. 75 (1962) p.63.
see also; H. J. Schmidt, Fortschr. Phys. 41 (1993) p.179.
46. A. Aguirre and S. Gratton, Phys. Rev. D 65 (2002) p.083507.
47. F. Hoyle and J.V. Narlikar, Proc. Roy. Soc. 277 (1964) p.1
48. P.J. Steinhardt and N. Turok, Phys. Rev. D 65 (2002) p.126003.
49. J. Garriga and A. Vilenkin, Phys. Rev. D 57 (1998) p.2230.
50. Y.B. Zeldovich, “My Universe” (Harwood Academic Press) 1992 p.95.
51. V.F. Mukhanov, H.A. Feldman and R.H. Brandenberger, Phys. Rept. 215 no. 5-6 (1992) p.203.
52. R.R. Caldwell, Phys. Lett. B 545 (2002) p.23.

53. F.G. Alvarenga and J.C. Fabris, *Class. Quant. Grav.* 13 (1996) p. L69.
54. D.H. Coule, *Phys. Lett. B* 450 (1999) p.48.
55. S.D.H. Hsu, A. Jenkins and M.B. Wise, *Phys. Lett. B* 597 (2004) p.270.
56. M.C.B. Abdalla, S. Nojiri and S.D. Odintsov, hep-th/0409177.
57. S.M. Carroll, V. Duvvuri, M. Trodden and M.S. Turner, *Phys. Rev. D* 70 (2004) p.043528.
E. Bruning, D.H. Coule and C. Xu, *Gen. Rel. Grav.* 26 (1994) p.1193.
58. P.J. Steinhardt and N. Turok, astro-ph/0404480.
59. C. Molina-Paris and M. Visser, *Phys. Lett. B* 455 (1999) p.90.
B.K. Tippet and K. Lake, gr-qc/0409088.
60. R. Durrer and J. Laukenmann, *Class. Quant. Grav.* 13 (1996) p.1069.
61. A.B. Batista, J.C. Fabris and S.V.B. Goncalves, *Class. Quant. Grav.* 18 (2001) p.1389.
62. N. Pinto Neto, hep-th/0410225.
63. J.D. Barrow and M.P. Dabrowski, *Mon. Not. R. Astron. Soc.* 275 (1995) p.850.
64. F. Hoyle, G. Burbidge and J.V. Narlikar, “A Different Approach to Cosmology”, (Cambridge University Press: Cambridge) 1999.
65. S.K. Banerjee and J.V. Narlikar, *Astrophys. Journ.* 487 (1997) p.69.
66. B.S. De Witt, *Phys. Rev D* 160 (1967) p.1113.
67. J.A. Wheeler, In “Batelle Recontres” eds. C. DeWitt and J.A. Wheeler (Benjamin Press: New York) 1968.
68. P. Ehrenfest, *Proc. Amst. Acad.* 20 (1917) p.200;
ibid, *Ann. Physik* 61 (1920) p. 440.
69. F.R. Tangherlini, *Nuovo. Cim.* 27 (1963) p.636

70. V.P. Frolov, M.A. Markov and V.F. Mukanov, Phys. Lett. B 216 (1989) p.272;
 V. Mukhanov and R. Brandenberger, Phys. Rev. Lett. 68 (1992) p.1969;
 see also: M.A. Markov, Phys. Usp. 37 (1994) p.57.
 V.P. Frolov and I.D. Novikov, "Black Hole Physics" (Kluwer Academic Publishers: Dordrecht) 1998.
71. L. Smolin, Class. Quant. Grav. 9 (1992) p.173.
72. L. Dyson, M. Kleban and L. Susskind, JHEP 0210 (2002) p.011;
 see also, D. Bak, hep-th/0208046.
73. A. Albrecht and L. Sorbo, hep-th/0405270.
74. S.M. Carroll and J. Chen, hep-th/0410270.
75. G. 't Hooft, preprint gr-qc/9310026;
 L. Susskind, J. Math. Phys. 36 (1995) p.6377;
 for reviews of the holography principle see:
 D. Bigatti and L. Susskind, hep-th/0002044.
 L. Smolin, Nucl. Phys. B 601 (2001) p.209.
76. D.H. Coule, Int. J. Mod. Phys. D 12 (2003) p.963.
77. J.D. Barrow, "Cargese Lectures in Physics" eds. W.G. Unruh and J. Hartle (Plenum Press: New York) 1987.
78. C.B. Collins and S.W. Hawking, Astrophys. J. 180 (1973) p.317.
79. D.S. Goldwirth and T. Piran, Phys. Rep. 214 (1992) p.223.
80. W.G. Unruh, in "Critical Dialogues in Cosmology", ed. N. Turok (World Scientific: Singapore) 1997.
81. S.W. Hawking and R. Penrose, "The Nature of Space and Time" (Princeton University Press: Princeton) 1996 p.90.
82. M. Heusler, Phys. Lett. B 253 (1991) p.1991.
83. J. Ibanez and I. Olasagasti, Class. Quant. Grav. 15 (1998) p.1937.

84. O. Iguchi and H. Ishihara, Phys. Rev. D 56 (1997) p.3216.
85. J.K. Erickson, D.H. Wesley, P.J. Steinhardt and N. Turok, Phys. Rev. D 69 (2004) p.063514.
86. P.K.S. Dunsby, N. Goheer, M. Bruni and A. Coley, hep-th/0312174
87. S. Hollands and R.M. Wald, Gen. Rel. Grav. 34 (2000) p.2043.
88. R. Penrose, in “14th Texas symposium”, ed. E.J. Fenyves (New York Academy of Sciences: New York) 1993.
89. R. Maartens, Living Rev. Rel. 7 (2004) p.1.
90. L. Randall and R. Sundrum, Phys. Rev. Lett. 83 (1999) p.4690;
ibid p.3370.
91. M. Bucher, hep-th/0107148;
U.Gen, A. Ishibashi and T. Tanaka, Phys. Rev. D 66 (2002) p.023519.
92. P. Anninos, R.A. Matzner and T. Rothman, Phys. Rev. D 43 (1991) p.3821.
93. C. Barcelo and M. Visser, Phys. Lett. B 482 (2000) p.183.
94. D.H. Coule, Class. Quant. Grav. 18 (2001) p.4265.
95. J.L. Hovdebo and R.C. Myers, JCAP 0311 (2003) p.012.
96. P. Kanti and K. Tamvakis, Phys. Rev. D 68 (2003) p.024014.
97. L.M. Burko and A. Ori, Phys. Rev. Lett. 74 (1995) p.1064.
98. M.G. Brown, K. Freese and W.H. Kinney, astro-ph/0405353.
99. R. Maartens, *private communication*
100. M. Gasperini and G. Veneziano, Astropart. Phys. 1 (1993) p.317.
101. M. Gasperini and G. Veneziano, Phys. Rept. 373 (2003) p.1.
102. D.H. Coule, Class. Quant. Grav. 15 (1998) p.2803.

103. A. Buonanno, T. Damour and G. Veneziano, Nucl. Phys. B 543 (1999) p.275.
104. Y.B. Zeldovich and I.D. Novikov, “The structure and evolution of the universe: relativistic astrophysics vol.2” (Chicago University Press: Chicago)1983 p.666.
105. D. Wands, Phys. Rev D 60 (1999) p.023507.
106. N.D. Birrell and P.C.W. Davies “Quantum fields in curved space” (Cambridge University Press: Cambridge) 1982.
107. P. Martinetti and C. Rovelli, Class. Quant. Grav. 20 (2003) p.4919.
S. Schlicht, Class. Quant. Grav. 21 (2004) p.4647.
108. S. Kawai, M. Sakagami and J. Soda, Phys. Lett. B 437 (1998) p.284.
109. A.D. Linde, Phys. Lett. B 129 (1983) p.177.
110. A.D. Linde, Phys. Lett. B 175 (1986) p.395.
111. V.A. Belinsky, L.P. Grishchuk, Y.B. Zeldovich and I.M. Klatatnikov, Sov. Phys. JEPT 62 (1985) p.195.
V.A. Belinsky and I.M. Khalatnikov, Sov. Phys. JEPT 66 (1987) p.441.
112. M.S. Madsen and P. Coles, Nucl. Phys. B 298 (1988) p. 2757.
113. H.J. Schmidt, Astron. Nachr. 311 (1990) p. 99.
114. L. Kofman, A. Linde and V. Mukhanov, JHEP 0210 (2002) p.057.
115. G.W. Gibbons, S.W. Hawking and J.M. Stewart, Nucl. Phys. B 281 (1987) p.736.
M. Henneaux, Nuovo. Cimento Lett. 38 (1983) p.609.
116. D.N. Page, Phys. Rev. D 36 (1987) p.1607.
117. D.H. Coule, Class. Quant. Grav. 12 (1995) p.455.
118. S. Hollands and R.M. Wald, hep-th/0210001.
119. A. Vilenkin, Phys. Rev. Lett. 74 (1995) p.846.

120. D.H. Coule, Class. Quant. Grav. 10 (1993) p.L25.
121. S.W. Hawking, Nucl. Phys. B 239 (1984) p.257.
122. D.N. Page, Class. Quant. Grav. 1 (1984) p.417.
123. P. Chmielowski and D.N. Page, Phys. Rev. D 38 (1988) p.2392.
124. A. Linde, hep-th/0408164.
125. A.D. Linde, Phys. Lett. B 327 (1994) p.208.
A. Vilenkin, Phys. Rev. Lett. 72 (1994) p.3137.
126. N. Sakai, Class. Quant. Grav. 21 (2004) p.281.
127. A.D. Linde, D.A. Linde and A. Mezhlumian, Phys. Rev. D 49 (1994) p.1783.
128. A.D. Linde and A. Mezhlumian, Phys. Rev. D 52 (1995) p.6789.
129. D.H. Coule, Gen. Rel. Grav. 36 (2004) p.2095.
130. T.M. Davis, P.C.W. Davies and C.H. Lineweaver, Class. Quant. Grav. 20 (2003) p.2753.
P.C.W. Davies and T.M. Davis, Found. of Phys. 32 (2002) p.1877.
131. K. Kunze, Phys. Lett. B 587 (2003) p.1.
132. N. Turok, Class. Quant. Grav. 19 (2002) p.3449.
133. S. Blau, E. Guendelman and A.H. Guth, Phys. Rev. D 35 (1987) p.1747.
134. E.H. Fahri and A.H. Guth, Phys. Lett. B 183 (1987) p.149.
135. K. Sato, M. Sasaki, H. Kodama and K. Maeda, Prog. Theor. Phys. 65 (1981) p.143.
ibid Phys. Lett. B 108 (1982) p.103.
136. T. Vachaspati and M. Trodden, Phys. Rev. D 61 (2000) p. 023502.
ibid, Mod. Phys. Lett. A 14 (1999) p.1661.
137. J.M.M. Senovilla, Gen. Rel. Grav. 30 (1998) p.70.

138. D. Bak and S.J. Rey, Class. Quant. Grav. 17 (2000) p.L83.
139. R. Bousso, JHEP 9907 (1999) p.004.
140. N. Dadhich, J. Astrophysics. Astro. 18 (1997) p.343.
141. E.H. Fahri, A.H. Guth and J. Guven, Nucl. Phys. B 339 (1990) p.417.
142. W. Fischler, D. Morgan and J. Polchinski, Phys. Rev. D 42 (1990) p.4042.
ibid, 42 (199) p.4042.
143. A.D. Linde, Nucl. Phys. B 372 (1992) p.421.
144. J.W. Moffat, Int. J. Mod. Phys. A 2 (1993) p.351.
A. Albrecht and J. Magueijo, Phys. Rev. D 59 (1999) p.043516.
J.D. Barrow, Phys. Rev. D 59 (1999) p.043515.
145. J. Magueijo, Rept. Prog. Phys. 66 (2003) p.2025.
146. D.H. Coule, Mod. Phys. Lett. A 14 (1999) p.2437.
147. U.H. Gerlach, Phys. Rev. 177 (1969) p.1929.
148. I.V. Kanatchikov, Int. J. Theor. Phys. 40 (2001) p.1121.
149. E. Tryon, Nature 246 (1973) p.396.
150. R. Brout, F. Englert and E. Gunzig, Annalen der Physik, 115 (1978) p.78.
151. D. Atkatz and H. Pagels, Phys. Rev. D 25 (1982) p.2065.
152. A. Vilenkin, Phys. Rev. D 27 (1983) p.2848.
153. J.B. Hartle and S.W. Hawking, Phys. Rev. D 28 (1983) p.2960.
154. A.D. Linde, Lett. Nuovo Cim. 39 (1984) p.401.
155. A. Vilenkin, Phys. Rev. D 30 (1984) p.509.
A.D. Linde, Sov. Phys. JEPT 60 (1984) p.211.
V.A. Rubakov, Phys. Lett. B 148 (1984) p.280.

156. Y.B. Zeldovich and A.A. Starobinsky, Sov. Astron. Lett. 10 (1984) p.135.
157. J.B. Hartle and S.W. Hawking, Phys. Rev. D 28 (1983) p.2960.
158. V.A. Rubakov and P.G. Tinyakov, Nucl. Phys. B 342 (1990) p.430.
159. A. Vilenkin, Nucl. Phys. B 252 (1985) p.141.
160. A. Vilenkin, Phys. Rev. D 37 (1988) p.888.
161. A. Vilenkin, Phys. Rev. D 50 (1994) p.2581.
162. D.H. Coule, Mod. Phys. Lett. A 10 (1995) p.1989.
163. G.W. Gibbons and S.W. Hawking, Phys. Rev. D 15 (1977) p.2752.
J. York, Phys. Rev. Lett. 28 (1972) p.1082.
see also: E. Poisson, “A Relativist’s Toolkit” (Cambridge University Press: Cambridge) 2004.
164. S.N. Solodukhin, Phys. Rev. D 62 (2000) p.044016.
165. E. Fahri, Phys. Lett. B 219 (1989) p.403.
166. G.W. Gibbons, S.W. Hawking and M.J. Perry, Nucl. Phys. B 138 (1978) p.141.
K. Schleich, Phys. Rev. D 36 (1987) p.2342.
167. E. Baum, Phys. Lett. B 133 (1989) p.185.
168. S.W. Hawking, Phys. Lett. B 134 (1983) p.403.
169. E. Embacher, Class. Quant. Grav. 13 (1996) p.921;
ibid, “Talk at Alexander Friedmann seminar, St Petersburg” 1995 gr-qc/9507041
170. G.W. Gibbons, Class. Quant. Grav. 15 (1998) p.2605;
G.W. Gibbons, Nucl. Phys. B 472 (1996) p.683.
171. S.W. Hawking, Nucl. Phys. B 144 (1978) p.349.

172. J.P. Luminet, Phys. Reports 254 (1995) p.135;
 Topology of the Universe conference, Cleveland 1997, special issue of
 Class. Quant. Grav. 15 (1998) p.2529.
 for a simpler introduction to compact manifolds in cosmology see:
 W.P. Thurston and J.R. Weeks, Sci. Amer. 251(7) (1984) p.108;
 J.P. Luminet, G.D. Starkman and J.R. Weeks, Sci. Amer. April (1999).
173. S. Coleman, Nucl. Phys. B 310 (1988) p.643.
174. M. Duff, Phys. Lett. B 226 (1989) p.36.
175. A. Linde, Phys. Rev. D 58 (1998) p.083514.
176. H. Firouzjahi, S. Sarangi and S.H. Tye, hep-th/0406107.
177. R.W. Robinett, "Quantum Mechanics", (Oxford University Press: Oxford) 1997.
178. C. Keifer, Class. Quant. Grav. 4 (1987) p.1369.
 J.J. Halliwell, Phys. Rev. D 39 (1989) p.2912.
 F. Fukuyama and M. Morikawa, Phys. Rev D 39 (1989) p.462.
 F. Mellor, Nucl. Phys. B 353 (1991) p.291.
 A.O. Barvinsky, A.Y. Kamenshchik, C. Kiefer and I.V. Mishakov, Nucl.
 Phys. B 551 (1999) p.374.
179. E. Calzetta and E. Verdaguer, Phys. Rev. D 59 (1999) p.083513.
180. J.M. Cline, Phys. Lett. B 224 (1989) p.53;
 M.B. Mijić, Phys. Lett. B 241 (1990) p.242;
 A. Strominger, Nucl. Phys. B 319 (1989) p.722.
181. D. Marolf, Phys. Rev. D 53 (1996) p.6979.
182. O.Y. Shvedov, Surveys High Energy Phys. 10 (1997) p.411.
183. H. Fink and H. Leschke, Found. Phys. Lett. 13 (2000) p.345.
184. D.H. Coule and J. Martin, Phys. Rev. D 61 (2000) p.063501.
185. D.H. Coule, Mod. Phys. Lett. A 13 (1998) p.961.
186. Computer software, MAPLE V release 4.6, Waterloo Maple Inc. 1996.

187. M. Abramowitz and I.A. Stegun, "Handbook of Mathematical Functions" (Dover Press) 1965.
188. S.W. Hawking and D.N. Page, Phys. Rev. D 42 (1990) p.2655.
189. C.M. Bender and S.A. Orszag, "Advanced Mathematical Methods for Scientists and Engineers" (McGraw-Hill:) 1984.
190. S.W. Hawking and D.N. Page, Nucl. Phys. B 264 (1986) p.185.
191. S.P. Kim, Phys. Rev. D 46 (1992) p.3403.
192. N. Kontoleon and D.L. Wiltshire, Phys. Rev. D 59 (1999) p.063513.
193. N.A. Lemos, J. Math. Phys. 37 (1996) p.1449.
J.H. Kung, Gen. Rel. Grav. 27 (1995) p.35.
194. J. Feinberg and Y. Peleg, Phys. Rev. D 52 (1995) p.1988.
195. D.H. Coule, Class. Quant. Grav. 9 (1992) p.2353.
A.K. Sanyal, Int. J. Mod. Phys. A 10 (1995) p.2231.
196. G. Magnano and L.M. Sokolowski, Phys. Rev. D 50 (1994) p.5039.
197. A. Carlini, D.H. Coule and D.M. Solomons, Mod. Phys. Lett. A 18 (1996) p.1453.
198. A. Carlini, D.H. Coule and D.M. Solomons, Int. J. of Mod. Phys. A 12 (1997) p.3517.
199. J.B. Hartle, J. Math. Phys. 30 (1989) p.452.
200. W.M. Suen and K. Young, Phys. Rev. D 39 (1989) p.2201.
201. D.N. Page, Phys. Rev. D 56 (1997) p.2065.
202. V.A. Rubakov, Phys. Lett. B 148 (1984) p.280.
V.A. Rubakov, gr-qc/9910025.
V.T. Gurovich, H.J. Schmidt and I.V. Tokareva, Gen. Rel. Grav. 33 (2001) p.591.
J. Hong, A. Vilenkin and S. Winitzki, Phys. Rev. D 68 (2003) p.023520.
S.P. Kim, gr-qc/0403015

- 203. A. Vilenkin, Phys. Rev. D 58 (1998) p.067301.
A. Vilenkin, gr-qc/9812027.
R. Bousso and S.W. Hawking, Grav. Cosmol. Suppl. 4 (1998) p.28.
- 204. L.P. Grishchuk and Y.V. Sidorov, Sov. Phys. JEPT 67 (1988) p.1533.
- 205. G.W. Gibbons and L.P. Grishchuk, Nucl. Phys. B 313 (1989) p.736.
- 206. L.P. Grischuk, Class. Quant. Grav. 10 (1993) p.2449.
- 207. P.D. D'Eath, "Supersymmetric Quantum Cosmology" (Cambridge University Press: Cambridge) 1995.
- 208. J.R. Gott and Li-Xin Li, Phys. Rev. D 58 (1998) p.023501.
- 209. G.W. Gibbons and J.B. Hartle, Phys. Rev. D 42 (1990) p.2458.
- 210. D. Green and W.G. Unruh, preprint gr-qc/0206068.
- 211. N. Pinto-Neto, gr-qc/0410117.
- 212. D.N. Page, gr-qc/0001001.
- 213. S.B. Giddings and A. Strominger, Nucl. Phys. B 321 (1989) p.481.
M. McGuigan, Phys. Rev. D 38 (1988) p.3031.
- 214. R. Geroch and J.B. Hartle, Found. of Phys. 16 (1986) p.533.
- 215. J.B. Hartle, in "Boundaries and Barriers: On the limits of Scientific Knowledge" eds: J.L. Casti and A. Karlqvist (Addison-Wesley: Reading MA) 1996.
- 216. K. Schleich and D.M. Witt, Phys. Rev. D 60 (1999) p.064013.
J.B. Hartle, Class. Quant. Grav. 2 (1985) p.707.
- 217. S. Carlip, Phys. Rev Lett. 79 (1998) p.4071.
S. Carlip, Class. Quant. Grav. 15 (1998) p.2629.
- 218. M. Anderson, S. Carlip, J.G. Ratcliffe, S. Surya and S.T. Tschantz, Class. Quant. Grav. 21 (2004) p.729.
see also: Y. Fujiwara, S. Higuchi, A. Hosoya, T. Mishima and M. Siino, Phys. Rev. D 44 (1991) p.1756.

219. G.W. Gibbons, in ref.[20].
220. J. Stelmach and I. Jakacka, *Class. Quant. Grav.* 18 (2001) p.2643.
221. J. Yokoyama and K. Maeda, *Phys. Rev. D* 41 (1990) p.1047.
222. V.M. Mostepanenko and N.N. Trunov, “The Casimir effect and its applications” (Oxford University Press: Oxford) 1997.
see also: D. Muller, gr-qc/0403086.
223. C.J. Isham, *Proc. R. Soc. A* 362 (1978) p.383.
224. J.A. Belinchon, *Int. J. Mod. Phys. D* 11 (2002) p.527.
225. R. Mansouri and F. Naseri, *Phys. Rev. D* 60 (1999) p.123512.
226. J. Garriga and M. Sasaki, *Phys. Rev. D* 62 (2000) p.043523.
S. Nojiri, S.D. Odintsov and S. Zerbini, *Class. Quant. Grav.* 17 (2000) p.4855.
A.S. Gorsky and K.G. Selivanov, *Phys. Lett. B* 485 (2000) p.271.
A.S. Gorsky and K.G. Selivanov, *Int. J. of Mod. Phys. A* 16 (2001) p.2243.
K. Koyama and J. Soda, *Phys. Lett. B* 483 (2000) p.043501.
L. Anchordoqui, C. Nunez and K. Olsen, *JHEP* 0010 (2000) p.050.
S.W. Hawking, T. Hertog and H.S. Reall, *Phys. Rev D* 63 (2001) p.083504.
M. Bouhmadi-Lopez, P.F. Gonzalez-Diaz and A. Zhuk, *Class. Quant. Grav.* 19 (2002) p.4863.
S.S. Seahra, H.R. Sepangi and J. Ponce de Leon, *Phys. Rev. D* 68 (2003) p.066009.
R. Cordero and E. Rojas, *Class. Quant. Grav.* 21 (2004) p.4231.
A. Boyarsky, A. Neronov and I. Tkachev, gr-qc/0411144.
227. R. Cordero and A. Vilenkin, *Phys. Rev. D* 65 (2002) p.083519.
228. E.I. Guendelman and A.B. Kaganovich, *Grav. Cosmol.* 1 (1995) p.103.
A. Davidson, *Class. Quant. Grav.* 16 (1999) p.653.
A. Davidson, D. Karasik and Y. Lederer, *Class. Quant. Grav.* 16 (1999) p. 1349.
F. Darabi, W.N. Sajko and P.S. Wesson, *Class. Quant. Grav.* 17

- (2000) p.4357.
P.I. Fomin and Yu.V. Shtanov, Class. Quant. Grav. 19 (2002) p.3139.
229. E. Carugno, M. Litterio, F. Occhionero and G. Pollifrone, Phys. Rev. D 53 (1996) p.6863.
230. K. Aoyanagi and K. Maeda, hep-th/0408008.
231. S. Nojiri and S.D. Odintov, hep-th/0409244.
232. R. Gambini, R.A. Porto and J. Pullin, hep-th/0406260.
ibid, hep-th/0405183.
ibid, gr-qc/0402118.
233. S.W. Hawking, “Talk at GR17 Dublin” 2004
234. Li-Xin Li and J.R. Gott, Phys. Rev. Lett. 80 (1998) p.2980.
235. W.A. Hiscock and D.A. Konkowski, Phys. Rev. D 26 (1982) p.1225.
236. P.F. González-Díaz, Phys. Rev. D 59 (1999) p.123513.
237. J.S. Dowker, Phys. Rev. D 18 (1978) p.1856.
238. S.W. Hawking, Commun. Math. Phys. 43 (1975) p.199.
239. G.W. Gibbons and S.W. Hawking, Phys. Rev. D 15 (1977) p.2738.
240. W.A. Hiscock, preprint gr-qc/0009061.
241. Li-Xin Li, Phys. Rev. D 59 (1999) p.084016.
242. V.L. Ginzburg and V.P. Frolov, Usp. Fiz. Nauk 150 (1986) p.4.
L.P. Grishchuk, Y.B. Zeldovich and L.V. Rozhanski, Sov. Phys. JEPT 65 (1987) p.11.
243. P.R. Anderson, W.A. Hiscock and B.E. Taylor, Phys. Rev. Lett. 85 (2000) p.2438.
244. G.F.R. Ellis and R. Maartens, Class. Quant. Grav. 21 (2004) p.223.
245. J.L. Friedman and A. Higuchi, Phys. Rev. D 52 (1995) p.5687.

- 246. J.D. Barrow, G.F.R. Ellis, R. Maartens and C.G. Tsagas, *Class. Quant. Grav.* 20 (2003) p.L155.
- 247. H.D. Conradi and H.D. Zeh, *Phys. Lett. A* 154 (1991) p.321.
H.D. Conradi, *Phys. Rev. D* 46 (1992) p.612.
- 248. M.B. Mijić, M.S. Morris and W. Suen, *Phys. Rev. D* 39 (1989) p.1496.
- 249. R. Coliste, J.C. Fabris and N. Pinto-Neto, *Phys. Rev. D* 62 (2000) p.83507.
- 250. F.G. Alvarenga, J.C. Fabris, N.A. Lemos and G.A. Monerat, *Gen. Rel. Grav.* 34 (2002) p.651.
- 251. G.F.R. Ellis, A. Sumeruk, D.H. Coule and C. Hellaby, *Class. Quant. Grav.* 9 (1992) p.1535.
G.F.R. Ellis, *Gen. Rel. Grav.* 24 (1992) p.1047.
- 252. F. Embacher, *Phys. Rev. D* 51 (1995) p.6764.
- 253. F. Embacher, *Phys. Rev. D* 52 (1995) p.2150.
- 254. R. Coliste, J.C. Fabris and N. Pinto-Neto, *Phys. Rev. D* 57 (1998) p.4707.
- 255. H. Kodama, *Phys. Rev. D* 42 (1990) p.2548.
- 256. C. Soo, *Class. Quant. Grav.* 19 (2002) p.1051.
E. Witten, gr-qc/0306083.
L. Freidel and L. Smolin, *Class. Quant. Grav.* 21 (2004) p.3831.
- 257. S. Alexander, J. Malecki and L. Smolin, preprint hep-th/0309045.
- 258. V. Husain and O. Winkler, *Phys. Rev. D* 69 (2004) p.084016.
- 259. G.F.R. Ellis and D.R. Matravers, *Gen. Rel. Grav.* 27 (1995) p.777.
R. Zalaletdinov, R. Tavakol and G.F.R. Ellis, *Gen. Rel. Grav.* 28 (1996) p.1251.
- 260. W. Rindler, *Phys. Lett. A* 276 (2000) p.52.
- 261. M. Bojowald, *Class. Quant. Grav.* 19 (2002) p.5113.

- 262. M. Bojowald and F. Hinterleitner, Phys. Rev. D 66 (2002) p.104003.
- 263. M. Bojowald, Class. Quant. Grav. 18 (2001) p.L109.
- 264. M. Bojowald, Phys. Rev. Lett. 89 (2002) p.261301.
- 265. M. Bojowald, Phys. Rev. Lett. 86 (2001) p.5227.
- 266. M. Bojowald and K. Vandersloot, Phys. Rev. D 67 (2003) p.124023.
- 267. M. Bojowald and K. Vandersloot, gr-qc/0303072.
- 268. M. Bojowald and G. Date, Phys. Rev. Lett. 92 (2004) p.071302.
M. Bojowald, G. Date and G.M. Hossain, Class. Quant. Grav. 21 (2004) p.3541.
- 269. J.D. Barrow, Nature 272 (1978) p.211.
- 270. J.D. Barrow, Phys. Rept. 85 (1982) p.1.
- 271. D. Green and W.G. Unruh, gr-qc/0408074.
- 272. J. Kowalski-Glikman and J.C. Vink, Class. Quant. Grav. 7 (1990) p.901.
- 273. S. Mollerach, S. Matarrese and F. Lucchin, Phys. Rev. D 50 (1994) p.4835.
- 274. V.A. Rubakov, M.V. Sazhin and A.V. Veryaskin, Phys. Lett. B 115 (1982) p.189.
L.F. Abbott and M.B. Wise, Nucl. Phys. B 244 (1984) p.541.
- 275. S. Tsujikawa, P. Singh and R. Maartens, preprint astro-ph/0311015
- 276. M. Bojowald, J.E. Lidsey, D.J. Mulryne, P. Singh and R. Tavakol, Phys. Rev. D 70 (2004) p.043530.
- 277. G.V. Vereshchagin, JCAP 0407 (2004) p.013.
- 278. M. Bojowald, R. Maartens and P. Singh, hep-th/0407115.
- 279. G. Date and G.M. Hossain, gr-qc/0407074.
- 280. R. Graham and P. Szepfalusy, Phys. Rev. D 42 (1990) p.2483.

281. A.D. Linde, *private communication*
282. Y.J. Ng and H. van Dam, Found. Phys. 30 (2000) p.795.
283. G. Amelino-Camelia, gr-qc/0204051.
284. C. Simon and D. Jaksch, quant-ph/0406007.
285. P.C.W. Davies, “When Time Began”, New Scientist (special supplement) 9/10/2004.
286. G.F.R. Ellis, in “Modern Cosmology”, eds: S. Bonometto, V. Gorini and U. Moshella, (IOP Press: Bristol) 2001.
G.F.R. Ellis, Class. Quant. Grav. 16 (1999) p.A37.
287. G. ’t Hooft, quant-ph/0212095.
288. J.D. Barrow, New. Astron. 4 (1999) p.333.
see also: P.C.W. Davies in ref.[17].
289. A.A. Grib and Y.V. Pavlov, preprint gr-qc/0206040.
290. B. McInnes, hep-th/0410115.
291. G.L. Kane, M.J. Perry and A.N. Zytlow, New. Astron. 7 (2002) p.45.
292. B.R. Frieden, “Physics from Fischer Information-a unification” (Cambridge University Press: Cambridge) 1998;
see also: H. Christian von Baeyer, “Information the new Language of Science” (Weidenfeld and Nicolson: London) 2003.
293. V. Dzhunushaliev and D. Singleton, Entropy 4 (2002) p.2.
294. S.W. Hawking, Lecture “ Godel and the End of Physics” available at:
www.damtp.cam.ac.uk/strts/-dirac/hawking/
295. J.D. Barrow, “Impossibility” (Oxford University Press: Oxford) 1998.
296. S.W. Hawking, Lecture at KITP, Santa Barbara, USA (2003)