

# On the Energy-Momentum Densities of the Cylindrically Symmetric Gravitational Waves

Ali Havare<sup>1</sup> \*; Mustafa Saltı<sup>1</sup>, Taylan Yetkin<sup>2</sup>

<sup>1</sup>Department of Physics, Mersin University, 33342, Mersin-Turkey

<sup>2</sup>Department of Physics, Çukurova University, 01330, Adana-Turkey

October 23, 2018

## Abstract

In this study, using Møller and Tolman prescriptions we calculate energy and momentum densities for the general cylindrically symmetric spacetime metric. We find that results are finite and well defined in these complexes. We also give the results for some cylindrically symmetric spacetime models.

## 1 Introduction

In general relativity a large number of researches has been devoted considerable attentions to the problem of finding the conserved quantities, such as energy and momentum, associated with various spacetimes. Following Einstein's original pseudotensor for energy-momentum many expressions have introduced in the literature[1, 2, 3, 4, 5, 6, 7, 8, 9, 10].

Einstein's energy momentum complex, used for calculating the energy in general relativistic system, was followed by many prescriptions, e.g. Møller's, Papapetrou's, Landau-Liftshitz's (LL), Tolman's and Weinberg's. Einstein's, Papapetrou's and LL's complexes give meaningful results when the line-element is transformed to the cartesian coordinates. However for the Møller's and Tolman's prescriptions it is not necessary to use of Cartesian coordinates. In the literature Virbhadra and his collaborators have considered many spacetimes and have shown that several energy-momentum complexes give the same and acceptable results[11]. Aguirregabiria *et al.*[12] showed that several energy-momentum complexes give the same result for any Kerr-Shild class metric. Recently, Chang *et al.*[13] showed that every energy-momentum complex can be associated with a particular Hamilton boundary term, therefore the energy-momentum complexes may also be considered as quasi-local.

---

\*Corresponding author. E-mail: alihavare@mersin.edu.tr

The paper organized as follow. In Section 1 We shortly introduce the generic, spatially homogeneous rotating spacetimes and give some kinematical variables for this model. In Section 3 and Section 4 we compute the energy and momentum densities for the line-element which is introduced in Section 2 by using Møller's and Tolman's prescriptions, respectively. Finally in Section 5 we give the results and conclusions. Throughout this paper we choose units such that  $G = 1$ ,  $c = 1$  and follow the convention that Latin indices take values from 0 to 3 and Greek indices take values 1 to 3 otherwise stated.

## 2 Spatially Homogeneous Rotating Spacetimes

After the pioneering works of Gamow[14] and Gödel[15], the idea of global rotation of the universe has become considerably important physical aspect in the calculations of general relativity. A line-element which describes spatially homogenous universes with rotation but with zero shear can be given as follow

$$ds^2 = -dt^2 + dr^2 + A(r)d\phi^2 + dz^2 - 2B(r)dt d\phi. \quad (1)$$

In this study we discuss some special solutions of the Møller and Tolman prescriptions for the energy and momentum densities by using the general model given in (1). For this model the matrices of the  $g_{\mu\nu}$  and  $\Delta^2 g^{\mu\nu}$  are found as follow

$$\begin{pmatrix} -1 & 0 & -B & 0 \\ 0 & 1 & 0 & 0 \\ -B & 0 & A & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} -A & 0 & -B & 0 \\ 0 & \Delta^2 & 0 & 0 \\ -B & 0 & 1 & 0 \\ 0 & 0 & 0 & \Delta^2 \end{pmatrix} \quad (2)$$

where  $\Delta(r) = \sqrt{A + B^2}$ . Using the Christoffel symbols which are defined by

$$\Gamma_{\mu\nu}^\alpha = \frac{1}{2}g^{\alpha\beta}(g_{\mu\beta,\nu} + g_{\nu\beta,\mu} - g_{\mu\nu,\beta}). \quad (3)$$

we obtain the nonvanishing Christoffel symbols as

$$2\Delta^2\Gamma_{\mu\nu}^0 = BB'(\delta_\mu^0\delta_\nu^1 + \delta_\mu^1\delta_\nu^0) + (AB' - BA')(\delta_\mu^1\delta_\nu^2 + \delta_\mu^2\delta_\nu^1) \quad (4)$$

$$2\Gamma_{\mu\nu}^1 = B'(\delta_\mu^0\delta_\nu^2 + \delta_\mu^2\delta_\nu^0) - A'\delta_\mu^2\delta_\nu^2 \quad (5)$$

$$2\Delta^2\Gamma_{\mu\nu}^2 = (A' + BB')(\delta_\mu^1\delta_\nu^2 + \delta_\mu^2\delta_\nu^1) - B'(\delta_\mu^0\delta_\nu^1 + \delta_\mu^1\delta_\nu^0). \quad (6)$$

One can introduces the tetrad basis as follow

$$\theta^0 = dt + Bd\phi, \quad \theta^1 = dr, \quad \theta^2 = \Delta d\phi, \quad \theta^3 = dz. \quad (7)$$

With the comoving tetrad formalism, the kinematical variables of this model can be expressed solely in terms of the structure coefficients of the tetrad basis which are defined as [16]

$$d\theta^\alpha = \frac{1}{2}c_{\beta\gamma}^\alpha\theta^\beta\wedge\theta^\gamma. \quad (8)$$

By taking the exterior derivatives of the tetrad basis which we introduced and the following kinematics formulas [17]

$$\begin{aligned}
\text{four-acceleration vector:} & \quad a_i = c_{i0}^0 \\
\text{vorticity tensor:} & \quad \omega_{ij} = \frac{1}{2}c_{ij}^0 \\
\text{expansion(deformation) tensor:} & \quad \theta_{ij} = \frac{1}{2}(c_{i0j} + c_{j0i}) \\
\text{expansion scalar:} & \quad \theta = c_{01}^1 + c_{02}^2 + c_{03}^3 \\
\text{vorticity vector:} & \quad \omega^1 = \frac{1}{2}c_{23}^0, \quad \omega^2 = \frac{1}{2}c_{31}^0, \quad \omega^3 = \frac{1}{2}c_{12}^0 \\
\text{vorticity scalar:} & \quad \omega = \frac{1}{4}[(c_{23}^0)^2 + (c_{31}^0)^2 + (c_{12}^0)^2]^{1/2} \\
\text{shear tensor:} & \quad \sigma_{ij} = \theta_{ij} - \frac{1}{3}\theta\delta_{ij}.
\end{aligned} \tag{9}$$

we find the following quantities for Eq.(1)

$$\omega^1 = \omega^2 = 0, \quad \omega^3 = \frac{B'}{2\Delta}, \quad a_i = \theta_{ij} = \sigma_{ij} = 0 \tag{10}$$

where prime indicates derivative with respect to  $r$ . The model given in Eq.(1) is shear-free and has non-vanishing vorticity. We also note that this frame has a vanishing four-acceleration. The line-element (1) can be reduced to well-known spacetime models in the literature by imposing some conditions on metric functions:

1. *The Rebouças spacetime*[19]

Defining  $A(r) = -(1 + 3c^2)$ ,  $B(r) = 2c$  with  $c = \cosh 2r$  the line-element 1 reduces to

$$ds^2 = -dt^2 + dr^2 - (1 + 3c^2)d\phi^2 + dz^2 + 4cdtd\phi \tag{11}$$

2. *The Som-Raychaudhuri spacetime*[19]

Defining  $A(r) = r^2(1 - r^2)$ ,  $B(r) = r^2$  the line-element 1 reduces to line-element

$$ds^2 = -dt^2 + dr^2 + r^2(1 - r^2)d\phi^2 + dz^2 - 2r^2dtd\phi \tag{12}$$

3. *The Hoenselaers-Vishveshwara spacetime*[19]

Defining  $A(r) = -\frac{1}{2}(c - 1)(c - 3)$ ,  $B(r) = c - 1$  with  $c = \cosh r$  the line-element 1 reduces to

$$ds^2 = -dt^2 + dr^2 - \frac{1}{2}(c - 1)(c - 3)d\phi^2 + dz^2 - 2(c - 1)dtd\phi \tag{13}$$

4. *The Gödel-Friedman spacetime*[19]

Defining  $A(r) = s^2(1 - s^2)$ ,  $B(r) = \sqrt{2}s^2$  with  $x^\mu = a\tilde{x}^\mu$  the line-element 1 reduces to

$$ds^2 = -dt^2 + dr^2 + dz^2 + s^2(1 - s^2)d\phi^2 - 2\sqrt{2}s^2dtd\phi \tag{14}$$

where  $x^\mu = (t, r, \theta, z)$ ,  $\tilde{x}^\mu = (\eta, \varepsilon, \phi, y)$ ,  $a = (1, 1) \times 10^{25}m$  and  $s = \sinh r$

5. *The Stationary Gödel spacetime*[15]

Defining  $A(r) = -e^{2ar}/2$ ,  $B(r) = e^{ar}$ ,  $a = \text{constant}$ , the line-element 1 reduces to

$$ds^2 = -(dt + e^{ar}d\theta)^2 + dr^2 + \frac{1}{2}(e^{ar}d\theta)^2 + dz^2 \quad (15)$$

### 3 Energy and Momentum Densities in Møller's Prescription

Since the Møller complex is not restricted to the use of Cartesian coordinates we can perform the computations in  $t, r, \theta, z$  coordinates. The computations in these coordinates are easier compared to those in  $t, x, y, z$  coordinates.

The following is the Møller's energy-momentum complex  $\Theta_a^b$  [21]:

$$\Theta_i^k = \frac{1}{8\pi}\chi_{i,l}^{kl} \quad (16)$$

where

$$\chi_i^{kl} = -\chi_i^{lk} = \sqrt{-g}[g_{in,m} - g_{im,n}]g^{km}g^{nl} \quad (17)$$

here  $\Theta_0^0$  is the energy density,  $\Theta_\alpha^0$  are the momentum density components. The Møller energy-momentum obeys the local conservation laws

$$\partial_k \Theta_i^k = 0 \quad (18)$$

The energy-momentum components are expressed by

$$P_i^M = \iiint \Theta_i^k dx^1 dx^2 dx^3 \quad (19)$$

where  $\mu_\beta$  are the components of a normal vector over an infinitesimal surface element  $dS$ .  $P_\alpha^M$  give the momentum components  $P_1^M, P_2^M, P_3^M$  and  $P_0^M$  gives the energy. For the line element (1) the non-vanishing components of  $\chi_i^{kl}$  are found as

$$\chi_0^{01} = -\frac{BB'}{\Delta} \quad (20)$$

$$\chi_2^{01} = \frac{BA' - AB'}{\Delta} \quad (21)$$

$$\chi_1^{01} = \chi_3^{01} = 0 \quad (22)$$

Using these results in equation (3), we obtain energy and momentum densities in Møller's prescription as

$$\Theta_i^0 = \frac{1}{8\pi}\chi_{i,1}^{01} \quad (23)$$

$$\Theta_j^0 = 0. \quad (24)$$

where  $i = 0, 2$  and  $j = 1, 3$ . By using the metric functions which are given in Section 2 it is easy to calculate the energy and momentum densities. The results obtained in Møller's prescription are given in Table 1.

## 4 Energy and Momentum in Tolman's Prescription

Tolman's energy and momentum complex [5] is given by

$$\Phi_i^k = \frac{1}{8\pi} \Lambda_{i,l}^{kl} \quad (25)$$

where  $\Phi_0^0$  and  $\Phi_\alpha^0$  are the energy and momentum components. We have

$$\Lambda_i^{kl} = \sqrt{-g}[-g^{pk}\Omega_{ip}^l + \frac{1}{2}g_k^i g^{pm}\Omega_{pm}^i] \quad (26)$$

with

$$\Omega_{jk}^i = -\Gamma_{jk}^i + \frac{1}{2}g_j^i \Gamma_{mk}^m + \frac{1}{2}g_k^i \Gamma_{mj}^m. \quad (27)$$

Also, the energy and momentum complex  $\Phi_i^k$  satisfies the local conservation laws

$$\partial_k \Phi_i^k = 0. \quad (28)$$

The energy and momentum in Tolman's prescription are given by

$$P_i = \iiint \Phi_i^k dx^1 dx^2 dx^3 \quad (29)$$

where  $n_\alpha$  are the components of a normal vector over an infinitesimal surface element  $dS$ .  $P_\alpha$  give the momentum components  $P_1, P_2, P_3$  and  $P_0$  gives the energy. For the line element (1) the non-vanishing components of  $U_i^{kl}$  are found as

$$\Lambda_0^{01} = \frac{A' + BB'}{2\Delta} \quad (30)$$

$$\Lambda_2^{01} = \frac{BA' - AB'}{2\Delta} \quad (31)$$

$$\Lambda_1^{01} = \Lambda_3^{01} = 0 \quad (32)$$

using these results in Eq.(27), we obtain energy and momentum densities in Tolman's prescription

$$\Phi_i^0 = \frac{1}{8\pi} \Lambda_{i,1}^{01} \quad (33)$$

$$\Phi_j^0 = 0. \quad (34)$$

where  $i = 0, 2$  and  $j = 1, 3$ . Again with the metric functions' definitions which are given in Section 2 it is easy to calculate the energy and momentum densities. The results obtained in Tolman's prescription are given in Table 1.

Space-time/Components	Møller's Prescription		Tolman's Prescription	
	$\Theta_0^0$	$\Theta_2^0$	$\Phi_0^0$	$\Phi_2^0$
Rebouças	$-\frac{s}{\pi}$	$-\frac{6sc^2}{r^3}$	$-\frac{s}{2\pi}$	$-\frac{3sc^2}{r^3}$
Som-Raychaudhri	$-\frac{r}{2\pi}$	$-\frac{r^2s}{2\pi}$	$-\frac{r}{4\pi}$	$-\frac{r^2s}{4\pi}$
Hoenselaers-Vishveshwara	$-\frac{s}{4\sqrt{2}\pi}$	$\frac{s(c-1)}{4\sqrt{2}\pi}$	0	$\frac{s(c-1)}{8\sqrt{2}\pi}$
Gödel-Friedman	$-\frac{cs}{\pi}$	$-\frac{\sqrt{2}cs^3}{\pi}$	0	$-\frac{cs^3}{\sqrt{2}\pi}$
Stationary Gödel	$-\frac{a^2 e^{ar}}{8\pi}$	$-\frac{a^2 e^{2ar}}{4\sqrt{2}\pi}$	0	$-\frac{a^2 e^{2ar}}{8\sqrt{2}\pi}$

Table 1: Energy and momentum densities for the specific space-time models. The  $c$  and  $s$  values in the table are given in Sec. 2.

## 5 Discussions

We used prescriptions of Møller and Tolman to calculate energy and momentum densities for a general form of cylindrically symmetric spacetime. We also gave the energy and momentum densities for the special cases of general cylindrically symmetric spacetime. We found that Møller's prescription give twice the Tolman's prescription,

$$P_i^M = 2P_i^T. \quad (35)$$

Also, for the Hoensalaers-Vishveshwara, Gödel-Friedman and Gödel spacetimes we found that energy components is vanishing in the Tolman's prescription. For this spacetimes one has to perform calculations in cartesian coordinates.

## References

- [1] R.C. Tolman, Phys. Rev. **35**, 875 (1930).
- [2] L.D. Landau and E. M. Lifshitz, *The Classical Theory Of Fields* (Addison-Wesley, Reading, MA, 1951).
- [3] J.N. Goldberg, Phys. Rev. **111**, 315 (1958).
- [4] P.G. Bergmann, Phys. Rev. **112**, 287 (1958).
- [5] C. Møller, Ann. Phys. (N.Y.) **4**, 347 (1958).
- [6] A. Komar, Phys. Rev. **113**, 934 (1959).
- [7] R. Arnowitt. S. Deser, and C.W. Misner, Phys. Rev. **122**, 997 (1961).
- [8] E. Nahmad-Achar and B.F. Schutz, Gen. Relativ. Gravit., **19**, 7 (1987).
- [9] R. Bartnik, Phys. Rev. Lett. **62**, 2346 (1989).
- [10] J.Katz and A. Ori, Class. Quantum Grav. **7**, 5 (1990).

- [11] K.S. Virbhadra, Phys. Rev. **D42**, 1066 (1990); Phys. Rev. **D42**, 2919 (1990); Phys. Lett **A157**, 195 (1991); Mathematics Today **9** 39 (1991); Pramana-J. Phys. **38**, 31 (1992); with N. Rosen, Gen. Rel. Grav. **25**, 429 (1993); Pramana-J. Phys. **45**, 215 (1995); with N. Rosen, Gen. Rel. Grav. **26**, 319 (1994); Phys. Rev. **D60**, 104041 (1999).
- [12] F.I. Cooperstock and S.A. Richardson, in Proc. 4th Canadian Conf. on General Relativity and Relativistic Astrophysics (World Scientific, Singapore, 1991)
- [13] J. M. Aguirregabiria, A. Chamorro, and K.S Virbhadra, Gen. Rel. Grav. **28**, 1393 (1996).
- [14] G. Gamow, Nature (London) **158**, 549 (1946).
- [15] K. Gödel, Rev. Mod. Phys. **21**, 447 (1949).
- [16] C.C. Chang, J. M. Nester and C. Chen, Phys. Rev. Lett. **83**, 1897 (1999).
- [17] O. Gron and H.H. Soleng, Acta Physica Polonica, No:7, Vol. **B20** (1989).
- [18] A. Havare, T. Yetkin, Class. Quantum Grav. **19**, **2783-2791** (2002).
- [19] K.D. Krori, P. Borgohain, P.K. Kar, and Dipali Das (Kar), J. Math. Phys. **29** (7), 1645-1649 (1988).
- [20] S. Carneiro, Phys. Rev. D **61** (2000)
- [21] M. Sharif, Int. J. of Mod. Phys. A, Vol **18**, No.**22**, 4361-4370 (2003).