

Mass and angular momenta of Kerr-anti-de Sitter spacetimes

Nathalie Deruelle¹

Institut des Hautes Etudes Scientifiques, 35 Route de Chartres, 91440, Bures-sur-Yvette, France

Abstract

We briefly review how to compute the mass and angular momenta of rotating, asymptotically anti-de Sitter spacetimes in Einstein-Gauss-Bonnet theory of gravity using superpotentials derived from standard Noether identities. The calculations depend on the asymptotic form of the metrics only and hence take no account of the source of the curvature. When the source of curvature is a black hole, the results can be checked using the first law of black hole thermodynamics.

1 Introduction

Consider Kerr-AdS₄ spacetime (\mathcal{M}) in Boyer-Lindquist coordinates [1]

$$ds^2 = -\frac{\Delta}{\rho^2} \left(dt - \frac{a \sin^2 \theta}{\Xi} d\phi \right)^2 + \frac{\rho^2}{\Delta} dr^2 + \frac{\rho^2}{\Delta_\theta} d\theta^2 + \frac{\Delta_\theta \sin^2 \theta}{\rho^2} \left(a dt - \frac{r^2 + a^2}{\Xi} d\phi \right)^2$$

where $\Delta \equiv (r^2 + a^2)(1 + r^2/l^2) - 2mr$, $\rho^2 \equiv r^2 + a^2 \cos^2 \theta$, $\Delta_\theta \equiv 1 - (a^2/l^2) \cos^2 \theta$ and $\Xi \equiv 1 - a^2/l^2$. It solves the Einstein equations with cosmological constant, $R^\mu_\nu = -(3/l^2)\delta^\mu_\nu$, and depends on two integration constants, the “mass” and “rotation” parameters m and a . \mathcal{M} is asymptotically anti-de Sitter but, as noticed in [2], the Boyer-Lindquist frame is rotating with angular velocity $\Omega_{BL} = a/l^2$ with respect to the standard, static, AdS₄ frame.

\mathcal{M} describes a black hole. Its horizon r_+ , which can be defined as the surface where signals from infalling test particles are measured at infinity with infinite redshift by static observers, is located at

$$\Delta(r_+) = 0 \quad \Longleftrightarrow \quad m = \frac{(r_+^2 + a^2)(1 + r_+^2/l^2)}{2r_+}.$$

Its angular velocity can be defined as the angular velocity of photons orbiting the horizon, as measured at infinity with respect to the static AdS₄ frame, and is

$$\Omega = \frac{a(1 + r_+^2/l^2)}{r_+^2 + a^2}.$$

A Bekenstein entropy can be associated to this black hole, as well as a Hawking black body temperature, respectively related to the area A of the horizon and its surface gravity κ by

$$S = \frac{1}{4}A = \frac{\pi(r_+^2 + a^2)}{\Xi} \quad , \quad T = \frac{\kappa}{2\pi} = \frac{4\pi(r_+^2 + a^2)}{r_+(1 + a^2/l^2 + 3r_+^2/l^2 - a^2/r_+^2)}.$$

Since \mathcal{M} is stationary and axially symmetric, it possesses two Killing vectors. Hence there exists global conserved charges that can be interpreted as the mass M and angular momentum J of the black hole. There are various ways to construct them. As reviewed in [2] there is general agreement in the literature on the definition of J but not on the definition of M . The “right” results are

$$M = \frac{m}{\Xi^2} \quad , \quad J = \frac{ma}{\Xi^2}$$

where the mass is associated with the Killing vector describing time translations in a locally free falling *non rotating* inertial frame in AdS₄. In the rotating Boyer-Lindquist frame its components are $(1, 0, 0, -a/l^2)$.

¹deruelle@ihes.fr

The reason why one can speak of the “right” results for M and J is that, as emphasized in [2], the first law of thermodynamics must hold, that is, one must have

$$T dS = dM - \Omega dJ.$$

Now, T, S, Ω, M, J are known functions of the two parameters m and a (or, computationally more simply, of r_+ and a). Therefore the two equalities

$$T \frac{\partial S}{\partial r_+} = \frac{\partial M}{\partial r_+} - \Omega \frac{\partial J}{\partial r_+} \quad , \quad T \frac{\partial S}{\partial a} = \frac{\partial M}{\partial a} - \Omega \frac{\partial J}{\partial a}$$

must hold and, if they do, provide a consistency check of the interpretation of the Kerr-AdS₄ metric as representing a radiating black hole spacetime, and in particular a check of the pertinence of the definition of its mass.

The object of this contribution is, first, to review in section 2 a method based on standard Noether identities which defines conserved charges associated to a spacetime \mathcal{M} with respect to its asymptotic background $\overline{\mathcal{M}}$ in Einstein-Gauss-Bonnet (EGB) theory [3], irrespective of the source of curvature, that is whether it is a “star” or a black hole.

In Einstein’s theory, when applied to the recently discovered D-Dimensional Kerr-anti-de Sitter Einstein black holes [4], that method yields the right mass, that is, for which the first law holds, as shown in [5] and reviewed in Section 4.

In Einstein-Gauss-Bonnet’s theory the rotating Kerr-like solution is not known. However, on one hand, only the asymptotic form of the metric is required to define the conserved charges and, on the other hand, the Einstein and Einstein-Gauss-Bonnet vacuum equations are the same when linearized on a maximally symmetric background (see section 3) : hence the D-dimensional Kerr-anti-de Sitter solutions to the Einstein field equations found in [4] are also asymptotic solutions to the Einstein-*Gauss-Bonnet* equations. Those two remarks were used in [6] to give an explicit expression for the mass and angular momenta of Kerr-AdS spacetimes in EGB theory, which is reviewed in section 4. When the full geometry of the Kerr-like rotating EGB black holes is eventually obtained, a test of the proposed definitions will be to see if the first law of EGB black hole thermodynamics is satisfied.

2 Noether superpotential and charges in EGB theory

There are various ways to associate conserved charges to a D -dimensional spacetime \mathcal{M} . We shall review here the approach advocated in [7] and [8] and applied to Einstein-Gauss-Bonnet theory of gravity in [3].

Consider the lagrangian

$$\mathcal{L} = L + D_\mu k^\mu \tag{2.1}$$

where L is some function of curvature invariants, functional of the metric $g_{\mu\nu}(x^\rho)$, its first and second derivatives with respect to the coordinates x^ρ and where k^μ is a vector to be appropriately chosen (see below). The variation of the scalar density $\hat{\mathcal{L}}$ with respect to the metric can be written as

$$\delta \hat{\mathcal{L}} = -\hat{\sigma}^{\mu\nu} \delta g_{\mu\nu} + \partial_\mu (\hat{V}^\mu + \delta \hat{k}^\mu). \tag{2.2}$$

(A hat denotes multiplication by $\sqrt{-g}$ where g is the determinant of the metric.) The tensor $\sigma^{\mu\nu}$ is the variational derivative of $\hat{\mathcal{L}}$ and the field equations are $\sigma^{\mu\nu} = 0$, whatever the vector k^μ . As for the vector V^μ it can be written under the form

$$V^\mu = \alpha^{\mu\nu\rho} \delta g_{\nu\rho} + \beta^{\mu\nu\rho}{}_\sigma \delta \Gamma_{\nu\rho}^\sigma \tag{2.3}$$

where $\Gamma_{\nu\rho}^\sigma$ are the Christoffel symbols, where $\delta \Gamma_{\nu\rho}^\sigma = \frac{1}{2} g^{\sigma\lambda} (D_\nu \delta g_{\rho\lambda} + D_\rho \delta g_{\nu\lambda} - D_\lambda \delta g_{\nu\rho})$ and where the tensors $\alpha^{\mu\nu\rho}$ and $\beta^{\mu\nu\rho}{}_\sigma$ depend on the precise dependence of L on the metric and its derivatives.

If the variation $\delta g_{\mu\nu}$ is due to a mere change of coordinates $x^\mu \rightarrow \tilde{x}^\mu = x^\mu - \xi^\mu$ with ξ^μ an infinitesimal vector (so that $\delta \hat{\mathcal{L}} = \partial_\mu (\hat{\mathcal{L}} \xi^\mu)$ and $\delta g_{\mu\nu} = 2D_{(\mu} \xi_{\nu)}$, parentheses denoting symmetrisation), then the variation (2) of the Lagrangian density can be cast in the form :

$$\partial_\mu \hat{j}^\mu = D_\mu \hat{j}^\mu = 0 \quad \text{with} \quad \hat{j}^\mu \equiv (2\hat{\sigma}^{\mu\nu} + g^{\mu\nu} \hat{L}) \xi_\nu - \hat{V}^\mu + 2D_\nu (\xi^{[\mu} \hat{k}^{\nu]}) \tag{2.4}$$

thanks to the (generalized) Bianchi identities. (Brackets stand for anti-symmetrization : $f_{[\mu\nu]} \equiv \frac{1}{2}(f_{\mu\nu} - f_{\nu\mu})$.)

The conservation of j^μ implies that there exists an antisymmetric tensor $j^{[\mu\nu]}$ such that

$$\hat{j}^\mu \equiv D_\nu \hat{j}^{[\mu\nu]} = \partial_\nu \hat{j}^{[\mu\nu]} . \quad (2.5)$$

If we look for an expression of the form

$$j^{[\mu\nu]} = \mathcal{A}^{[\mu\nu]\rho} \xi_\rho + \mathcal{B}^{[\mu\nu]\rho\sigma} D_\rho \xi_\sigma + 2\xi^{[\mu} k^{\nu]} \quad (2.6)$$

then $\mathcal{A}^{[\mu\nu]\rho}$ and $\mathcal{B}^{[\mu\nu]\rho\sigma}$ are obtained in terms of $\alpha^{\mu\nu\rho}$ and $\beta^{\mu\nu\rho\sigma}$ from (4) and (5) by identification.

We obtain similarly $\overline{\hat{j}^\mu}$ and $\overline{\hat{j}^{[\mu\nu]}}$ for the background $\overline{\mathcal{M}}$ (an overline means that the quantity is evaluated on the background spacetime with metric $\overline{g}_{\mu\nu}(x^\rho)$). In practice the only demand on $\overline{\mathcal{M}}$ is that it asymptotes \mathcal{M} . The KBL superpotential density [7] [8] is then defined as

$$\hat{j}^{[\mu\nu]} = -\frac{1}{16\pi} \left(\hat{j}^{[\mu\nu]} - \overline{\hat{j}^{[\mu\nu]}} \right) . \quad (2.7)$$

We now integrate (7) over a region \mathcal{M} of spacetime bounded by $\partial\mathcal{M} = \partial\overline{\mathcal{M}}$, a timelike hypercylinder limited by two spacelike hypersurfaces $x^0 = t_0$ and $x^0 = t_1$. We apply Stokes' theorem and obtain that the “charge”

$$Q = \int_S d^{D-2}x \, n_i \hat{j}^{[0i]} \quad (2.8)$$

is independent of time if $\int_S d^{D-2}x \, n_i (\hat{j}^i - \overline{\hat{j}^i}) = 0$ where S is the $(D-2)$ -dimensional sphere at infinity, with n^i its unit normal vector pointing towards infinity.

In practice we shall restrict our attention to stationary spacetimes and choose ξ^μ to be a Killing vector, so that Q will indeed be time-independent. When that Killing vector is associated with time translations with respect to a non rotating frame at infinity the associated conserved charge is the mass M of \mathcal{M} . If the Killing vector is associated with rotations then the associated charges are the angular momenta J_i .

As for the vector k^μ it is chosen according to circumstances (that is, according to L) in order to cancel as many normal derivatives of the metric variations as possible in the divergence in (2). If they all cancel, the field equations are obtained by keeping only the metric and its tangential derivatives fixed at the boundary, leaving their normal derivatives free, and the variational problem falls within the scope of ordinary Lagrangian field theory.

Let us now specialize to Einstein-Gauss-Bonnet's theory of gravity. The Lagrangian is (see e.g. [9] for a review of its properties)

$$L \equiv -2\Lambda + R + \alpha R^{\mu\nu\rho\sigma} P_{\mu\nu\rho\sigma} \quad \text{with} \quad P_{\mu\nu\rho\sigma} \equiv R_{\mu\nu\rho\sigma} - 2(R_{\mu[\rho} g_{\sigma]\nu} - R_{\nu[\rho} g_{\sigma]\mu}) + R g_{\mu[\rho} g_{\sigma]\nu} \quad (2.9)$$

(so that $R^{\mu\nu\rho\sigma} P_{\mu\nu\rho\sigma} = R^{\mu\nu\rho\sigma} R_{\mu\nu\rho\sigma} - 4R^{\mu\nu} R_{\mu\nu} + R^2$). Λ is a (negative) cosmological constant, α a coupling constant which is zero in Einstein's theory, and $R_{\mu\nu\rho\sigma}$, $R_{\mu\nu}$ and R are the Riemann tensor, Ricci tensor and curvature scalar of the metric $g_{\mu\nu}$.

The equations of motion derived from that Lagrangian are

$$\sigma^{\mu\nu} \equiv \sigma_E^{\mu\nu} + \alpha \sigma_{GB}^{\mu\nu} = 0 \quad (2.10)$$

with

$$\sigma_E^{\mu\nu} \equiv \Lambda g^{\mu\nu} + R^{\mu\nu} - \frac{1}{2} R g^{\mu\nu} \quad (2.11)$$

and

$$\sigma_{GB}^{\mu\nu} \equiv 2R^{\mu\lambda\rho\sigma} P^\nu_{\lambda\rho\sigma} - \frac{1}{2} g^{\mu\nu} R^{\alpha\beta\rho\sigma} P_{\alpha\beta\rho\sigma} . \quad (2.12)$$

The explicit form of the vector V^μ appearing in (2) (3) is

$$V^\mu \equiv V_E^\mu + \alpha V_{GB}^\mu \quad (2.13)$$

with

$$V_E^\mu = -(g^{\mu\nu}\delta\Gamma_{\nu\rho}^\rho - g^{\nu\rho}\delta\Gamma_{\nu\rho}^\mu) \quad (2.14)$$

and

$$V_{GB}^\mu = 4D^{(\mu}\sigma_E^{\nu)\rho}\delta g_{\nu\rho} - 4(P^{\mu\alpha\beta}{}_\gamma - Q^{\mu\alpha\beta}{}_\gamma)\delta\Gamma_{\alpha\beta}^\gamma \quad (2.15)$$

where the tensor $P_{\mu\nu\rho\sigma}$ is defined in (9) and where $Q_{\mu\nu\rho\sigma} \equiv 2(R_{\rho[\mu}g_{\nu]\sigma} + R_{\sigma[\mu}g_{\nu]\rho})$. The tensors $\mathcal{A}^{[\mu\nu]\rho}$ and $\mathcal{B}^{[\mu\nu]\rho\sigma}$ in (6) are then easily calculated and yield the following expression for the EGB superpotential

$$\hat{J}^{[\mu\nu]} \equiv \hat{J}_E^{[\mu\nu]} + \alpha \hat{J}_{GB}^{[\mu\nu]} \quad (2.16)$$

where the Einstein contribution is given by, see [7-8]

$$-8\pi\hat{J}_E^{[\mu\nu]} \equiv D^{[\mu}\hat{\xi}^{\nu]} - \overline{D^{[\mu}\hat{\xi}^{\nu]}} + \hat{\xi}^{[\mu}k_E^{\nu]} \quad (2.17)$$

As for the Gauss-Bonnet contribution $\hat{J}_{GB}^{[\mu\nu]}$ it is, when ξ^μ is a Killing vector [3]

$$-8\pi\hat{J}_{GB}^{[\mu\nu]} \equiv 2\left[P^{\mu\nu\alpha\beta}D_{[\alpha}\hat{\xi}_{\beta]} - \overline{P^{\mu\nu\alpha\beta}D_{[\alpha}\hat{\xi}_{\beta]}}\right] + \hat{\xi}^{[\mu}k_{GB}^{\nu]} \quad (2.18)$$

The last step is to choose the vectors k_E^μ and k_{GB}^μ . The choice made in [7] and [8], that is

$$k_E^\nu = g^{\mu\nu}\Delta_{\nu\rho}^\rho - g^{\nu\rho}\Delta_{\nu\rho}^\mu \quad \text{with} \quad \Delta_{\nu\rho}^\mu \equiv \Gamma_{\nu\rho}^\mu - \overline{\Gamma_{\nu\rho}^\mu} \quad (2.19)$$

has passed all tests with flying colours (see e.g. [10]). The following proposal was made in [3] for the Gauss-Bonnet contribution

$$k_{GB}^\mu = 4(P^{\mu\alpha\beta}{}_\gamma - Q^{\mu\alpha\beta}{}_\gamma)\Delta_{\alpha\beta}^\gamma \quad (2.20)$$

where it was shown that such a definition yielded the correct formula for the mass of the static EGB-Schwarzschild-like spacetimes, that is, which is consistent with the first law of EGB black hole thermodynamics.

More generally, when the vector ξ_t^μ is the Killing vector of time translations with respect to a non rotating frame at infinity the associated conserved charge is the mass M . If the vector ξ_i^μ is a Killing vector for rotations with respect to the i -axis then the associated charges is the angular momentum J_i :

$$M = \int_S d^{D-2}x \hat{J}_t^{[01]} \quad , \quad J_i = \int_S d^{D-2}x \hat{J}_i^{[01]} \quad (2.21)$$

Equipped with those formulae it is then a matter of straightforward calculation to obtain the mass M and angular momenta J_i for a given metric which solves the field equations (10-12). It is important to note that only the asymptotic form of the metric is needed and that the mass and angular momenta thus obtained do not depend on the source of curvature, which can be a “star” or a black hole.

3 The asymptotic Kerr-AdS solutions of the EGB equations

The Einstein-Gauss-Bonnet equations of motion were given in the previous section, equations (2.10-2.12). In order to compute the mass and angular momenta of stationary and axi-symmetric spacetimes which solve those field equations, only the asymptotic form of the metric is required. We therefore linearize the EGB equations, that is, we set

$$g_{\mu\nu} = \bar{g}_{\mu\nu} + h_{\mu\nu} \quad (3.1)$$

We choose the background $\bar{g}_{\mu\nu}$ to be the metric of a maximally symmetric spacetime $\overline{\mathcal{M}}$ solving the EGB equations. Since they are quadratic in the Riemann tensor there are two solutions, defined by

$$\bar{R}_{\mu\nu\rho\sigma} = -\frac{1}{\mathcal{L}^2}(\bar{g}_{\mu\rho}\bar{g}_{\nu\sigma} - \bar{g}_{\mu\sigma}\bar{g}_{\nu\rho}) \quad \text{where} \quad \frac{1}{\mathcal{L}^2} \equiv \frac{1}{2\tilde{\alpha}} \left(1 \pm \sqrt{1 - \frac{4\tilde{\alpha}}{l^2}}\right) \quad (3.2)$$

with $\tilde{\alpha} \equiv (D-3)(D-4)\alpha$ and $l^2 \equiv -\frac{(D-1)(D-2)}{2\Lambda}$, D being the dimension of spacetime. We shall choose the lower sign for \mathcal{L}^2 so that the limit when $\alpha = 0$ solves the Einstein equations. (When Λ and α are such that $\sqrt{1-4\tilde{\alpha}/l^2} = 0$ the two roots coalesce into a single double root.)

Using the fact that

$$R^\mu_{\nu\rho\sigma} = \overline{R^\mu_{\nu\rho\sigma}} + \frac{1}{2}\overline{D}_\rho(\overline{D}_\nu h^\mu_\sigma + \overline{D}_\sigma h^\mu_\nu - \overline{D}^\mu h_{\sigma\nu}) - \frac{1}{2}\overline{D}_\sigma(\overline{D}_\nu h^\mu_\rho + \overline{D}_\rho h^\mu_\nu - \overline{D}^\mu h_{\rho\nu}) + \mathcal{O}(h^2) \quad (3.3)$$

(\overline{D}_μ being the covariant derivative associated to $\overline{g}_{\mu\nu}$) it is an exercise to compute $R^{\mu\lambda\rho\sigma}P_{\nu\lambda\rho\sigma}$ at linear order and see that the EGB equations read (a result first obtained in [11] for $\Lambda = 0$)

$$\sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \left[R^\mu_\nu - \frac{1}{2}\delta^\mu_\nu R + \delta^\mu_\nu \Lambda \frac{l^2}{\mathcal{L}^2} \right] = \mathcal{O}(h^2) \quad (3.4)$$

where

$$R^\mu_\nu = -\frac{(D-1)}{\mathcal{L}^2}(\delta^\mu_\nu - h^\mu_\nu) + \frac{1}{2} \left[\overline{D}_\alpha(\overline{D}_\nu h^{\alpha\mu} + \overline{D}^\mu h^\alpha_\nu - \overline{D}^\alpha h^\mu_\nu) - \overline{D}^\mu \overline{D}_\nu h^\rho_\rho \right] + \mathcal{O}(h^2) \quad (3.5)$$

is the linearized Ricci tensor on $\overline{\mathcal{M}}$.

Hence, when linearized on a maximally symmetric background, the Einstein-Gauss-Bonnet equations are the same as the linearized pure Einstein equations ($\alpha = 0$) with effective cosmological constant $\Lambda_e \equiv \Lambda \frac{l^2}{\mathcal{L}^2}$, and up to the overall factor $\sqrt{1-4\tilde{\alpha}/l^2}$ that we shall assume not to be zero.

Now, in [4] Gibbons, Lu, Page and Pope found D-dimensional Kerr-anti-de Sitter exact solutions of Einstein's equations. Since the linearized Einstein-Gauss-Bonnet equations are the same as the linearized pure Einstein equations, those solutions also solve the Einstein-Gauss-Bonnet equations at linear order.

We shall write them in Kerr-Schild coordinates. In odd dimensions $D = 2n + 1$ for instance, the line element reads

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = d\overline{s}^2 + \frac{2m}{U} (h_\mu dx^\mu)^2 \quad \text{with } \mu = \{0, 1, \mu_i, \phi_i\} \quad (i = 1, \dots, n) \quad (3.6)$$

where $d\overline{s}^2$ is the metric of the anti-de Sitter background and where the coordinates μ_i are subject to the constraint $\sum_{i=1}^{i=n} \mu_i^2 = 1$. The mass parameter m is an integration constant, h_μ is some null vector (denoted k_μ in [2] [4]) depending on n rotation parameters a_i and U is a scalar function. Explicit forms of those functions can be found in Section 2 of ref [4].

As an example, the 5-dimensional AdS line element in Kerr-Schild ellipsoidal coordinates $x^\mu = (t, r, \theta, \phi, \psi)$ reads :

$$d\overline{s}^2 = -\frac{(1+r^2/\mathcal{L}^2)\Delta_\theta}{\Xi_a\Xi_b} dt^2 + \frac{r^2\rho^2}{(1+r^2/\mathcal{L}^2)(r^2+a^2)(r^2+b^2)} dr^2 + \frac{\rho^2}{\Delta_\theta} d\theta^2 + \frac{r^2+a^2}{\Xi_a} \sin^2\theta d\phi^2 + \frac{r^2+b^2}{\Xi_b} \cos^2\theta d\psi^2 \quad (3.7)$$

where $\Delta_\theta \equiv \Xi_a \cos^2\theta + \Xi_b \sin^2\theta$, where $\rho^2 \equiv r^2 + \mathcal{L}^2(1 - \Delta_\theta)$, and where Ξ_a and Ξ_b are related to the rotation parameters a and b by $\Xi_a \equiv 1 - a^2/\mathcal{L}^2$, $\Xi_b \equiv 1 - b^2/\mathcal{L}^2$. As for the function U and the null vector h_μ they are given by $U = \rho^2$ and

$$h_\mu dx^\mu = \frac{\Delta_\theta}{\Xi_a\Xi_b} dt + \frac{r^2\rho^2}{(1+r^2/\mathcal{L}^2)(r^2+a^2)(r^2+b^2)} dr + \frac{a \sin^2\theta}{\Xi_a} d\phi + \frac{b \cos^2\theta}{\Xi_b} d\psi. \quad (3.8)$$

Note that we shall only need their asymptotic behaviours at leading order. Note also that the full Ricci tensor of the Kerr-Schild metric (6-8) is linear in $h_{\mu\nu} = \frac{2m}{U} h_\mu h_\nu$ and hence exactly given by (5), see e.g. [4].

The background admits a timelike Killing vector $\xi = \xi_t$ and n plane rotation Killing vectors $\xi = \xi_i$ associated with one parameter displacements, say τ . ξ_t is normalized in such a way that $\xi_t^\mu \delta\tau$ is an infinitesimal time translation in a locally free falling *non rotating* inertial frame in the background. Thus

for the background metric in its conventional static form when $D = 2n + 1$ for instance (equations (2.11) and (2.12) of Ref [4])

$$d\bar{s}^2 = - \left(1 + \frac{y^2}{l^2}\right) dt^2 + \frac{dy^2}{1 + y^2/l^2} + y^2 d\sigma_{(D-2)}^2 \quad \text{where} \quad d\sigma_{(D-2)}^2 = \sum_{i=1}^{i=n} (d\tilde{\mu}_i^2 + \tilde{\mu}_i^2 d\phi_i^2), \quad (3.9)$$

with $\Sigma_i \tilde{\mu}_i^2 = 1$, the components of the timelike Killing vector ξ_t corresponding to time translations are $(1, 0, 0, 0, \dots)$. In Kerr-Schild ellipsoidal coordinates ξ_t has the same components. If the vector ξ_i^μ is the Killing vector associated with rotations, for example with respect to the ϕ (resp. ψ) axis in five dimensions, that is $\xi_a^\mu = (0, 0, 0, 1, 0)$ (resp. $\xi_b^\mu = (0, 0, 0, 0, 1)$), then the associated charges are the angular momenta.

4 Kerr-AdS masses and angular momenta in EGB theory

To obtain the mass and angular momenta of the 5D-Kerr-like solution of EGB theory we inserted the metric (3.6-3.8) of section 3 in the definitions (2.16-2.21) of section 2 and took the large r limit (using *Maple* and *GR tensor*). The result was found to be [6]

$$M_{(5)} = \sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \left[\frac{m\pi}{4(\Xi_a \Xi_b)^2} (2\Xi_a + 2\Xi_b - \Xi_a \Xi_b) \right], \quad J_{(5)a} = \sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \left[\frac{\pi m a}{2\Xi_a^2 \Xi_b} \right]. \quad (4.1)$$

We calculated similarly the mass and angular momentum in $D = 6, 7, 8$ dimensions in the case when there is only one non zero rotation parameter. The results could be cast under the form ($D = 5, 6, 7, 8$) [6]

$$M_D = \sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \frac{m \mathcal{V}_{D-2}}{4\pi \Xi_a^2} \left[1 + \frac{(D-4)}{2} \Xi_a \right], \quad J_D = \sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \frac{m a \mathcal{V}_{D-2}}{4\pi \Xi_a^2} \quad (4.2)$$

where \mathcal{V}_{D-2} is the volume of the $(D-2)$ -unit sphere. Finally, we extrapolated to all dimensions the asymptotic expressions for the Kerr-AdS Riemann tensor obtained with *Maple* and *GR tensor* in $D = 5, 6, 7, 8$ dimensions, and found (by hand) the mass and angular momenta of the general D-dimensional Kerr-like solutions of Einstein-Gauss-Bonnet theory as [6]

$$M_{2n} = \sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \left[\frac{m \mathcal{V}_{2n-2}}{4\pi \Xi} \sum_{i=1}^{i=n-1} \frac{1}{\Xi_i} \right], \quad M_{2n+1} = \sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \left[\frac{m \mathcal{V}_{2n-1}}{4\pi \Xi} \left(\sum_{i=1}^{i=n} \frac{1}{\Xi_i} - \frac{1}{2} \right) \right] \\ J_{(D)i} = \sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \left[\frac{m \mathcal{V}_{D-2}}{4\pi \Xi} \frac{a_i}{\Xi_i} \right], \quad (4.3)$$

where $\Xi = \prod_{i=1}^{i=(n-1)} \Xi_i$ for $D = 2n$ and $\Xi = \prod_{i=1}^{i=n} \Xi_i$ for $D = 2n + 1$.

Let us fill up some details of the calculation.

We worked in Kerr-Schild coordinates in which the metric is given for example by (3.6-3.8). Some useful properties are that the metric coefficients do not depend on x^0 and ϕ_i ; the $g_{\mu_i \mu_j}$ coefficients do not depend on m ; the other off-diagonal components are linear in m and hence do not enter the calculation. The fact that the vector h^μ is null further simplifies the calculation since $\sqrt{-g} = \sqrt{-\bar{g}}$. Using those informations one thus arrives for example at

$$D^{[0] \hat{\xi}_t^{[1]}] - \overline{D^{[0] \hat{\xi}_t^{[1]}]} = \overline{D^{[0] \hat{\xi}_t^{[1]}]} \left(\frac{\delta g'_{tt}}{\bar{g}'_{tt}} - \frac{\delta g_{tt}}{\bar{g}_{tt}} - \frac{\delta g_{rr}}{\bar{g}_{rr}} \right) + \mathcal{O}(m^2) \quad \text{with} \quad \overline{D^{[0] \hat{\xi}_t^{[1]}]} = - \frac{\sqrt{-\bar{g}} \bar{g}'_{tt}}{2\bar{g}_{tt} \bar{g}_{rr}} \quad (4.4)$$

where $\xi_t^\mu = (1, 0, \dots, 0)$, where $\delta g_{\mu\nu} \equiv g_{\mu\nu} - \bar{g}_{\mu\nu}$ and where a prime denotes differentiation with respect to r . A similar expression is obtained for $\hat{\xi}_t^{[0] k_E^{[1]}}$ with k_E^μ given by equation (2.19) of section 2.

Using then the asymptotic form of the D-dimensional Kerr-AdS metric found in [4] at leading order in r one then arrives for example at

$$\hat{\xi}_t^{[0] k_E^{[1]}} = - \frac{m}{\Xi} \sqrt{g_{(D-2)}} + \mathcal{O}\left(\frac{m^2}{r}\right), \quad (4.5)$$

and

$$\hat{J}_{Et}^{[01]} = \frac{m}{8\pi\Xi} [(D-1)W - 1]\sqrt{g_{(D-2)}} + \mathcal{O}\left(\frac{m^2}{r}\right) \quad \text{with} \quad W \equiv \sum_{i=1}^{i=n} \frac{\mu_i^2}{\Xi_i} \quad (4.6)$$

where $g_{(D-2)}$ is the determinant of the metric $d\sigma_{(D-2)}^2$ in (3.9) with tildes dropped and with $a_n = 0$ in even dimensions $D = 2n$.

The calculation of $\hat{\xi}_t^{[0]} k_{GB}^{[1]}$ is straightforward since

$$k_{GB}^\mu = -\frac{2}{\mathcal{L}^2}(D-2)(D-3)k_E^\mu + \mathcal{O}(m^2). \quad (4.7)$$

The calculation of $-4\pi\hat{J}_{P,t}^{[01]} \equiv (P^{01\alpha\beta}D_{[\alpha}\hat{\xi}_{t\beta]} - \overline{P^{01\alpha\beta}D_{[\alpha}\hat{\xi}_{t\beta]}})$ at leading order yields in a first step

$$-4\pi\hat{J}_{P,t}^{[01]} = \frac{\sqrt{g_{(D-2)}}}{\Xi\mathcal{L}^2} \left[-\frac{2}{\mathcal{L}^2} \lim_{r \rightarrow \infty} r^{D+1} \delta R_{101}^0 + mW(D-1)(D-2)(D-3) - 2m(D^2 - 5D + 8) \right] \quad (4.8)$$

and requires the knowledge of the asymptotic form of the Riemann tensor. The structure of the relevant component for the calculation of the mass is easily guessed and depends on two constants only :

$$\delta R_{101}^0 \equiv \lim_{r \rightarrow \infty} (R_{101}^0 - \overline{R_{101}^0}) = \frac{m\mathcal{L}^2}{r^{D+1}}(D-1)[c_1(D-3)W - c_2] \quad (4.9)$$

where, recall, $W \equiv \sum_{i=1}^{i=[D/2]} \frac{\mu_i^2}{\Xi_i}$, with $\sum_{i=1}^{i=[D/2]} \mu_i^2 = 1$ and $\Xi_{D/2} = 1$ if D is even. The *Maple* and *GRtensor* calculations in $D = 5$ determine the constants c_1 and c_2 to be one, similar calculations in $D = 6, 7, 8$ confirm them, so that the results can confidently be extended to any D .

After gathering all the previous information the remaining task is to integrate the superpotential on the sphere at infinity. While the “multi-cylindrical” pairs of coordinates (μ_i, ϕ_i) were convenient in [4] to find the Kerr-anti-de Sitter metric as well as in our previous calculations, the explicit integrations are more easily performed in spherical coordinates, say, φ_k with $(k = 1, 2, \dots, D-2)$.

The final results are given above, formulae (3).

The calculation of the angular momenta proceeds along similar lines. Note that neither the background nor the vectors k_E^μ and k_{GB}^μ enter the calculations. However, in Einstein-Gauss-Bonnet theory, they are not given simply by Komar’s integrals because the first term in (2.18) (that is $\hat{J}_{P,i}^{[01]}$) is not zero.

5 Conclusion

Formulae (4.3) for the mass and angular momenta of Kerr-AdS spacetimes in Einstein-Gauss-Bonnet theory have the right limits. When $\alpha = 0$ (Einstein’s theory) they reduce to the results obtained in [2] and [5] and, as shown in [2], they are consistent with the first law of Einstein black hole thermodynamics.

When there is no rotation ($a_i = 0, \Xi_i = 1$) the masses (4.3) reduce to the result obtained in [12-13] and [3] (noting that the coefficients of the metric (3.6) then tend to $g_{tt} \simeq 1/g_{rr} \simeq r^2/\mathcal{L}^2 - 2m/r^{D-3}$) and are also consistent with the first law of Einstein-Gauss-Bonnet black hole thermodynamics (where the entropy is no longer proportional to the surface area of the horizon, see e.g. [12-13]).

Since the overall factor in (4.3) is the same as the one which appears in (3.4) one may conjecture that in the general Lovelock theory in D dimensions (whose Lagrangian is the sum of the first $[D/2]$ dimensionally continued Euler forms) the expressions for the masses and angular momenta will also be proportional to their values in Einstein’s theory with an overall coefficient which is zero when the equation for the maximally symmetric space curvature has one single root of maximal multiplicity $[D/2]$.

Finally, the decisive check of formulae (4.3), that is of the pertinence of the proposal (2.20) made in [3] for the vector k_{GB}^μ , will be possible when the full geometry of the Kerr-like rotating *black holes* is known, by seeing if, with such definitions, the first law of thermodynamics is still satisfied.

6 Acknowledgements

Most of the work summarized here is the result of greatly enjoyable collaborations with Joseph Katz, Yoshiyuki Morisawa and Sachiko Ogushi.

References

- [1] B. Carter, *Hamilton-Jacobi and Schrödinger separable solutions of Einstein's equations*, Comm. Math. Phys. **10** (1968) 280
- [2] G.W. Gibbons, M.J. Perry and C.N. Pope, *The First Law of Thermodynamics for Kerr-de Sitter Metrics*, hep-th/0408217
- [3] N. Deruelle, J. Katz and S. Ogushi, *Conserved Charges in Einstein- Gauss-Bonnet theory*, Class. Quantum Grav., **21** (2004) 1971, gr-qc/0310098
- [4] G.W. Gibbons, H. Lu, Don N. Page and C.N. Pope, *The General Kerr-de Sitter Metrics in All Dimensions*, hep-th/0404008
- [5] N. Deruelle and J.Katz, *On the mass of a Kerr-anti-de Sitter spacetime in D dimensions*, Class. Quantum Grav. **22** (2005) 421-424, gr-qc/0411035
- [6] N. Deruelle and Y. Morisawa, *Mass and angular momenta of Kerr-anti-de Sitter spacetimes in Einstein-Gauss-Bonnet theory*, accepted by Class. and Quant. Grav, gr-qc/0411135
- [7] J. Katz, *A note on Komar's anomalous factor*, Class. Quantum Grav., **2** (1985) 423
- [8] J. Katz, J. Bičák, D. Lynden-Bell, *Relativistic conservation laws and integral constraints for large cosmological perturbations*, Phys. Rev. **D55** (1997) 5759
- [9] N. Deruelle and J. Madore, *On the quasi-linearity of the Einstein "Gauss-Bonnet" gravity field equations*, in "Progress in General Relativity and Quantum Cosmology", gr-qc/0305004
- [10] B. Julia and S. Silva, *Currents and superpotentials in classical gauge theories : II. Global aspects and the extension of affine gravity*, Class. Quan. Grav., **17** (2000) 4733, gr-qc/0005127
- [11] D.G. Boulware and S. Deser, *String generated gravity models*, Phys. Rev. Lett. **55** (1985) 2656
- [12] Rong-Gen Cai, *Gauss-Bonnet Black Holes in AdS Spaces*, Phys.Rev. **D65** (2002) 084014, hep-th/0109133
- [13] I. Neupane, *Black hole entropy in string-generated gravity models*, Phys.Rev.**D67** (2003) 061501, hep-th/0212092