Volume elements and torsion

Ricardo A. Mosna* and Alberto Saa[†]

Instituto de Matemática, Estatística e Computação Científica,
Universidade Estadual de Campinas, C.P. 6065, 13083-859, Campinas, SP, Brazil.

We reexamine here the issue of consistency of minimal action formulation with the minimal coupling procedure (MCP) in spaces with torsion. In Riemann-Cartan spaces, it is known that a proper use of the MCP requires that the trace of the torsion tensor be a gradient, $T_{\mu} = \partial_{\mu}\theta$, and that the modified volume element $\tau_{\theta} = e^{\theta} \sqrt{g} \ dx^{1} \wedge \cdots \wedge dx^{n}$ be used in the action formulation of a physical model. We rederive this result here under considerably weaker assumptions, reinforcing some recent results about the inadequacy of propagating torsion theories of gravity to explain the available observational data. The results presented here also open the door to possible applications of the modified volume element in the geometric theory of crystalline defects.

PACS numbers: 02.40.Hw, 04.50.+h, 61.72.Lk

I. INTRODUCTION

The use of modified volume elements in physical models has been receiving considerable interest in recent years. In the context of the Einstein-Cartan theory of gravity[1], for instance, the introduction of a new volume element gave rise to some models possessing several interesting characteristics, such as: propagation of torsion[2, 3], interaction between torsion and gauge fields without violation of the gauge symmetry[4], and a geometrical interpretation of string-theory inspired models of gravity[5]. Non-canonical volume elements have also proved to be useful in other contexts. For recent applications in field and string theory, see, for instance, [6].

Geometries with torsion, on the other hand, have a long tradition in Physics. For instance, in Einstein-Cartan theory of gravity[1], spacetime is assumed to be a Riemann-Cartan manifold, *i.e.*, a manifold endowed with a Lorentzian metric and a non-symmetrical metric-compatible linear connection. The anti-symmetrical part of the connection (the torsion tensor) is also relevant to field theory, mainly with respect to renormalization issues[7]. The very active area of defects in crystal microstructures also takes advantage of these non-Riemannian geometries: disclinations and dislocations in a crystal can be described by means of curvature and torsion (for a revision, see [8, 9]). Dislocations, in particular, are described by the Burgers vector, which has the torsion tensor as its surface density (see Fig. 1).


The motivation for the introduction of the modified volume element proposed in [10] was the observation that, in the presence of torsion, the minimal coupling procedure (MCP) performed at the action level is not equivalent to that performed directly at the corresponding field equations. As shown originally in [10], in Riemann-Cartan spaces the equivalence between the MCP performed at the action and at the field equations is recovered provided that the trace of the torsion tensor be a gradient, $T_{\mu} = \partial_{\mu}\theta$, and that the usual volume element $\tau_0 = \sqrt{g} \, dx^1 \wedge \cdots \wedge dx^n$ be deformed to

$$\tau_{\theta} = e^{\theta} \tau_{0}. \tag{1}$$

In [3], Fiziev reinterpreted such model as a Transposed-Equi-Affine Theory of Gravity, since, as we will see, the covariant derivative of densities used in [10] can be related to a certain "transposed" connection. Such a transposed connection, however, is a highly artificial object, being generically not metric-compatible even when the initial connection is. Besides, at first glance, the very motivation of demanding equivalence between the MCP as applied to the action level and to the field equations may be contested on physical grounds. In fact, even in the case of General Relativity (where the connection is torsion-free), the naive minimal coupling between the electromagnetic field and gravity does not satisfy such equivalence requirement if one does not choose carefully in which set of equations the MCP should be applied. In such a case, the field equations contain partial derivatives of higher order, which renders the MCP, if applied directly to some field equations, ill-defined [11]. The usual solution for this kind of difficulty is to assume that the MCP should be always applied at the Lagrangian level.

The aim of this paper is to show that the modified volume element (1) follows from a more fundamental hypothesis, reinforcing its mathematical and physical relevance. We will show that the existence of the modified volume element

^{*}Electronic address: mosna@ime.unicamp.br †Electronic address: asaa@ime.unicamp.br






FIG. 1: We show, for illustrative purposes only, a lattice with two edge dislocation defects (upper right) and a reference one (upper left). A closed rectangular path in the reference lattice becomes unclosed in the presence of the defects. The vector $\overline{BB'}$, the so-called Burgers vector, measures how the path fails to close. In the associated geometrical description of the path generated by the (infinitesimal) vectors X^{μ} and Y^{μ} at the point A, the vector \overline{CB} is obtained by parallel transport of Y^{μ} along X^{μ} , and $\overline{DB'}$ is obtained by transporting X^{μ} along Y^{μ} . The Burgers vector in this case is given by $\overline{BB'} = T_{\nu\kappa}{}^{\mu}X^{\nu}Y^{\kappa}$.

(1) is a necessary and sufficient condition to render the MCP well-defined even when considered solely at the action level. In particular, we will show that the results presented in [10] can be obtained without recourse to transposed connections. Our conclusions apply to any physical model where torsion is regarded as a dynamical entity that couples to the remaining fields via the MCP. In this way, one can expect that the results presented here can be also useful in the geometric theory of crystalline defects[9, 12], since the expressions for the free and interaction energies of defects and the equations for phonons and other test fields can be obtained by arguments very similar to the MCP.

We observe that the model proposed in [2] was carefully studied in [13, 14]. By considering solutions describing compact (neutron) stars and some solar system experiments, it was shown[13] that the predictions of the model [2] are in contradiction with General Relativity and, more importantly, with the available observational data. The propagating torsion model proposed in [2], therefore, is not a viable alternative to General Relativity. As we will see, the results presented here make the conclusions of [13] much stronger, since we will show that the existence of (1) is required to ensure the very consistency of the action formulation via the MCP. In other words, the results of [13] would allow us to conclude that the very reasonable requirement of consistency of the action formulation with the MCP imply, on account of the observational data, that the trace of the torsion tensor of spacetime should vanish.

This work is organized as follows. The next section is a brief review of the mathematical framework necessary to set up the problem. Section III contains our main results, and the last section is devoted to some final remarks. Mathematical proofs are left to the Appendix, where we also discuss how the results presented here relate to the

earlier approach of [2, 3, 4, 5, 10].

II. NOTATION

In order to introduce the problem, we briefly recall some definitions. For the sake of clarity, we employ here a more mathematically oriented notation, which slightly differs from that used in the original works [2, 3, 4, 5, 10]. Let \mathcal{M} be an n-dimensional manifold. In the following, the components of the covariant derivative of a vector field $A = A^{\nu}e_{\nu}$, with respect to a local basis in $T\mathcal{M}$, are denoted by $D_{\mu}A^{\nu}$, so that $\nabla_{\mu}A = (D_{\mu}A^{\nu})e_{\nu}$, with $D_{\mu}A^{\nu} = \partial_{\mu}A^{\nu} + \Gamma_{\mu\sigma}{}^{\nu}A^{\sigma}$. The components of the torsion tensor associated with the connection ∇ ,

$$T(X,Y) = \nabla_X Y - \nabla_Y X - [X,Y],\tag{2}$$

are given, in a coordinate basis, by

$$T_{\mu\nu}{}^{\kappa} = \Gamma_{\mu\nu}{}^{\kappa} - \Gamma_{\nu\mu}{}^{\kappa}. \tag{3}$$

Whenever the manifold \mathcal{M} is endowed with a metric $g_{\mu\nu}$, the connection coefficients $\Gamma_{\mu\nu}{}^{\kappa}$ can be decomposed as

$$\Gamma_{\mu\nu}{}^{\kappa} = \begin{Bmatrix} \kappa \\ \mu\nu \end{Bmatrix} + K_{\mu\nu}{}^{\kappa} + V_{\mu\nu}{}^{\kappa}, \tag{4}$$

where $\begin{Bmatrix} \kappa \\ \mu\nu \end{Bmatrix}$ are the Christoffel symbols associated with the underlying Levi-Civita connection (uniquely defined metric-compatible connection without torsion on (\mathcal{M}, g)),

$$K_{\mu\nu\kappa} = -\frac{1}{2} \left(T_{\nu\kappa\mu} + T_{\mu\kappa\nu} - T_{\mu\nu\kappa} \right) \tag{5}$$

is the contorsion tensor associated with ∇ , and

$$V_{\mu\nu\kappa} = -\frac{1}{2} \left(D_{\mu} g_{\nu\kappa} + D_{\nu} g_{\kappa\mu} - D_{\kappa} g_{\mu\nu} \right) \tag{6}$$

provides a measure of the nonmetricity associated with ∇ . Relevant quantities here are the trace of the above defined tensors and their associated 1-forms

$$T_{\nu} = T_{\mu\nu}{}^{\mu}, \quad t = T_{\nu} dx^{\nu},$$

 $V_{\nu} = V_{\mu\nu}{}^{\mu}, \quad v = V_{\nu} dx^{\nu}.$ (7)

A volume element on a n-dimensional manifold \mathcal{M} is a nowhere vanishing n-form on \mathcal{M} [15]. If \mathcal{M} is endowed with a metric, an arbitrary volume element τ can be written as

$$\tau = f\tau_0,\tag{8}$$

where f is a nowhere vanishing smooth function on \mathcal{M} and τ_0 is the canonical volume element

$$\tau_0 = \theta^1 \wedge \dots \wedge \theta^n = \sqrt{g} \, dx^1 \wedge \dots \wedge dx^n, \tag{9}$$

where $\{\theta^i\}$ is a local orthonormal basis.

III. MINIMAL COUPLING AND EQUIVALENT LAGRANGIANS

Let \mathcal{L} be a Lagrangian density describing a given physical model defined on flat space or spacetime \mathcal{M} . There is a natural equivalence relation \sim on the set of such Lagrangians, defined by $\mathcal{L}_1 \sim \mathcal{L}_2$ if and only if \mathcal{L}_1 and \mathcal{L}_2 give rise to *identical* field equations on M. Except for scale transformations, we then have

$$\mathcal{L}_1 \sim \mathcal{L}_2 \iff \mathcal{L}_1 - \mathcal{L}_2 = \operatorname{div}(X)$$
 (10)

for some vector field X, where, for flat \mathcal{M}

$$\operatorname{div}(X) = \partial_{\mu} X^{\mu}. \tag{11}$$

This follows trivially from Gauss theorem, since integration of the Lagrangian density immediately yields

$$\int \operatorname{div}(X) \, \tau_0 = [\text{surface term}], \tag{12}$$

with no contribution to the Euler-Lagrange equations. We emphasize that, in the context of flat space (and of Special Relativity), any choice \mathcal{L} in a given class $[\mathcal{L}]$ of Lagrangians leads to completely equivalent physical models.

Suppose now that the metric and connection on \mathcal{M} become dynamical, describing in this way physical fields interacting with test particles or fields on \mathcal{M} . This would be the case, for instance, of General Relativity, in which gravity is described by a dynamical metric $g_{\mu\nu}$ with its associated Levi-Civita connection. Other examples are the Einstein-Cartan theory of gravity[1], where torsion becomes an additional dynamical quantity, and the geometric theory of defects[8, 9], where curvature and torsion are related to the density of dislocations and disclinations of the crystalline lattice structure.

The minimal coupling procedure (MCP) is the standard prescription to obtain the curved space equations from the flat space ones. It states that one needs merely to substitute ordinary derivatives by covariant ones and the Minkowskian metric tensor by its non-flat counterpart. It is worth noting that the MCP is used with success in nearly all physically relevant gauge theories, including General Relativity [16].

Let us denote the Lagrangian density obtained from \mathcal{L} , via MCP, by \mathcal{L}^* . Similarly to the flat-space case discussed above, the condition for \mathcal{L}_1^* and \mathcal{L}_2^* to be equivalent (i.e., to yield the same Euler-Lagrange equations) is given by

$$\mathcal{L}_1^* - \mathcal{L}_2^* = \operatorname{div}_{\tau}(X) \tag{13}$$

for some vector field X, with the definition of divergence being given now by [15]

$$(\operatorname{div}_{\tau} X) \tau = \pounds_X \tau, \tag{14}$$

where τ is a volume form on \mathcal{M} and \mathcal{L}_X denotes the Lie derivative along the vector field X. This is the most natural definition of divergence in this context, since $\mathcal{L}_X \tau = d i_X \tau$ (where d denotes the exterior derivative and i_X the contraction by X [15]) leads directly to Gauss theorem

$$\int_{\mathcal{M}} \operatorname{div}_{\tau}(X)\tau = \int_{\partial \mathcal{M}} i_X \tau = [\text{surface term}]. \tag{15}$$

If \mathcal{L}_1 and \mathcal{L}_2 are equivalent Lagrangians in flat-space, we get, after applying the MCP,

$$\mathcal{L}_1 - \mathcal{L}_2 = \partial_\mu X^\mu \xrightarrow{\text{MCP}} \mathcal{L}_1^* - \mathcal{L}_2^* = D_\mu X^\mu. \tag{16}$$

Therefore, if the equivalence class of Lagrangians $[\mathcal{L}]$ is required to be preserved under the MCP, we need to have

$$D_{\mu}X^{\mu} = \operatorname{div}_{\tau}(X),\tag{17}$$

or, equivalently,

$$\pounds_X \tau = (D_\mu X^\mu) \tau. \tag{18}$$

The necessary and sufficient conditions to the existence of solutions for (17) can be obtained from the following proposition, whose (simple) proof is left to the Appendix.

Proposition 1 Let \mathcal{M} be a differentiable manifold endowed with a metric g and a connection ∇ . Let $\tau = f\tau_0$ be a volume form on \mathcal{M} . Then, for every vector field X on \mathcal{M} ,

$$D_{\mu}X^{\mu} = \operatorname{div}_{\tau} X + t(X) + v(X) - X(\ln f),$$

where t and v are defined in eqs. (7).

A. MCP in Affine manifolds

In order to illustrate the previous results, let us recall their application to the case of the metric-affine theory of gravity[19], where spacetime is represented by a manifold endowed with a connection with possibly nonzero curvature,

torsion and nonmetricity. We note that this contains, as a particular case, the Einstein-Cartan theory of gravity, where the associated connection presents curvature and torsion but is still metric-compatible (i.e., $D_{\kappa}g_{\mu\nu} = 0$).

For this case, Proposition 1 implies that the canonical volume form τ_0 yields

$$D_{\mu}X^{\mu} - \operatorname{div}_{\tau_0} X = (T_{\mu} + V_{\mu})X^{\mu}. \tag{19}$$

It follows from our previous discussion that equivalent Lagrangian densities of Special Relativity will be mapped, via MCP, to nonequivalent Lagrangian densities in the corresponding non-flat spacetime. Therefore, since there is no canonical representative of $[\mathcal{L}]$ in Special Relativity, the MCP turns out to be essentially ill-defined in this context.

This difficulty can be circumvented with the help of Proposition 1 itself. The above incompatibility arises only when $D_{\mu}X^{\mu} \neq \operatorname{div}_{\tau} X$. However, admitting a more general volume form $\tau = f\tau_0$ on \mathcal{M} , one can get $D_{\mu}X^{\mu} = \operatorname{div}_{\tau} X$ by choosing f such that

$$t(\partial_{\mu}) + v(\partial_{\mu}) - \partial_{\mu} \ln f = 0, \tag{20}$$

which is equivalent to

$$T_{\mu} + V_{\mu} = \partial_{\mu}\theta, \quad \text{and } \tau = e^{\theta}\tau_0.$$
 (21)

Therefore, the MCP is well defined (preserves equivalence classes of Lagrangians) precisely when:

- 1. t + v is an exact form, say $t + v = d\theta$, and
- 2. the volume element on \mathcal{M} is given by $\tau = e^{\theta} \tau_0$.

Again, since in the context of flat space (Special Relativity) there is no canonical choice of representative of a given class of Lagrangians, our results imply that the MCP is well defined only in the presence of the modified volume element (1).

This is our main result: in order to preserve a given class $[\mathcal{L}]$ of Lagrangians under the MCP, one must have $T_{\mu} + V_{\mu} = \partial_{\mu}\theta$ and, obligatorily, to use the modified volume element (1). If the sum of traces $T_{\mu} + V_{\mu}$ is not a gradient, the MCP turns out to be ill-defined, since it produces nonequivalent theories out of equivalent flat-space Lagrangians, even at the classical level.

B. The λ -symmetry

In an affine manifold, the Riemann tensor is invariant under the transformation

$$\Gamma_{\mu\nu}^{\ \kappa} \to \Gamma_{\mu\nu}^{\ \kappa} + \delta^{\kappa}_{\nu} \partial_{\mu} \lambda,$$
 (22)

where λ is an arbitrary function. This is the so-called λ -symmetry [17], introduced by Einstein in his pioneering work on unified field theory, which has been proved to be very useful in the analysis of affine theories of gravity[18]. In a λ -invariant theory, the function λ can be properly chosen in order to cancel some parts of the torsion tensor or/and of the non-metricity, simplifying the overall analysis. Under (22), the torsion tensor and $V_{\mu\nu\kappa}$ are changed as

$$T_{\mu\nu\kappa} \to T_{\mu\nu\kappa} + (g_{\kappa\nu}\partial_{\mu}\lambda - g_{\kappa\mu}\partial_{\nu}\lambda),$$

$$V_{\mu\nu\kappa} \to V_{\mu\nu\kappa} + (g_{\kappa\nu}\partial_{\mu}\lambda + g_{\kappa\mu}\partial_{\nu}\lambda - g_{\mu\nu}\partial_{\kappa}\lambda),$$
(23)

implying to their traces

$$T_{\nu} \to T_{\nu} - (n-1)\partial_{\nu}\lambda$$
 and $V_{\nu} \to V_{\nu} + n\partial_{\nu}\lambda$. (24)

One sees from (23) that a λ -transformation does not preserve the spaces of symmetric nor metric-compatible connections. However, the condition ensuring the consistency of MCP, namely that $T_{\mu} + V_{\mu}$ be a gradient, is indeed preserved under (22). The Einstein-Hilbert action is, obviously, also preserved under (22). Hence, provided that the matter content action is also λ -invariant, the λ -transformation will map distinct solutions in a metric-affine theory of gravity governed by the Einstein-Hilbert action, preserving all the isometries if λ is properly chosen.

An interesting consequence arises from the transformation of the modified volume element under a λ -transformation,

$$e^{\theta}\tau_0 \to e^{\theta+\lambda}\tau_0.$$
 (25)

Starting with, for instance, a Riemann-Cartan manifold with a gradient torsion trace and an Einstein-Hilbert action with the modified volume element, it is possible to choose $\lambda = -\theta$ and to get an equivalent theory, where the MCP is also well-defined, formulated in an affine manifold (the connection is not metric-compatible anymore) but with the usual volume element. In other words, in this new manifold, $D_{\mu}X^{\mu} = \operatorname{div}_{\tau_0} X$ for any vector field X.

IV. FINAL REMARKS

Before discussing possible applications of the results presented here to the study of crystalline defects, let us recall how they reinforce the results of [13] about the inadequacy of the propagating torsion model proposed in [2] to explain the available observational data. The modified volume element (1) was introduced in the model of [2] by means of exclusively geometrical arguments, and, hence, there is no free parameter associated with the field θ , which is responsible for the propagating components of the torsion. Moreover, by employing the MCP, no extra parameters, besides the Newtonian constant G, should arise in any gravitational model. Thus, the only way of fitting the available observational data described in [13] is to have θ constant, leading to the already mentioned conclusion that the results of [13] would allow us to deduce that the observational data and the very reasonable requirement of consistency of the action formulation with the MCP imply that the trace of the torsion tensor of spacetime should vanish.

Condensed matter is a vast subject but, to the best of our knowledge, the ideas presented here were not yet applied to the study of crystalline defects, which possess a geometrical description since the seminal works of K. Kondo in the fifties (see, for references, [8]). The physics of crystalline defects is a promising arena to text experimentally the ideas of the last sections, with special emphasis to new effects that might be associated with the modified volume element (1). The first conclusion about the modified volume element in this context is that, in principle, it has no relation to the real volume (area, in this case) of the polygon depicted in Fig. 1. As it is discussed in [8], contributions to the elementary area coming from the Burgers vector are of higher order and should vanish in the linear continuum limit. Incidentally, for X^{μ} and Y^{μ} orthogonal in a Riemann-Cartan manifold ($D_{\kappa}g_{\mu\nu}=0$), the area of the polygon of Fig. 1 is given, up to higher order terms, by

$$A = |X||Y| \left(1 - \frac{1}{2} (X^{\mu} + Y^{\mu}) T_{\mu}\right), \tag{26}$$

from where one sees that, despite of being unclosed, its area coincides with the area of the parallelogram with sides X^{μ} and Y^{μ} if $T_{\mu}=0$. However, the modified volume element (1) cannot represent a physical area since any geometrical property of the crystalline lattice should be invariant under a transformation of the type $\theta \to \theta + \Lambda$, with constant Λ , and the modified volume element, under this transformations, behaves as $\tau \to e^{\Lambda}\tau$. This behavior, on the other hand, is allowed in the minimal action formulation, since, as it was already noted, a scale transformation of a Lagrangian preserves the dynamics. The dynamics of defects, on the other hand, are governed by their free energy, which can be expressed by means of a Lagrangian density related to the Einstein-Hilbert Lagrangian[12, 20]. The use of the modified volume element in the action of the free energy would define a new dynamics to the torsion and metric tensors, analogous to that proposed in [2] in the context of Einstein-Cartan theory of gravity. An immediate consequence would be the propagation of the trace of the torsion tensor and, consequently, of the associated Burgers vector.

The free energy plays the role of the kinetic term in the geometrical action formulation of defects. However, kinetic terms are not, a priori, specified by the MCP. Our analysis seems more suitable to the study of fields/particles propagating on the crystalline lattice, such as phonons, impurities and vacancies [12]. If the perturbation energy associated with these objects is small enough, their backreaction on the lattice may be ignored, and they can be considered test fields. Phonons are nothing more than small elastic (sound) excitations in the crystal; in the geometrical description, they correspond to gauge independent linearized perturbations of the metric [9]. On the other hand, one can also consider impurities or vacancies moving on the background medium defined by the solid with crystalline defects. The effect of the background on such test fields/particles can be gotten from the MCP by demanding that the corresponding equations for phonons, impurities and vacancies be obtained, from the equations for the regular lattice, by changing the ordinary derivative to the covariant one and the Euclidean metric to the metric associated with the background medium. With use of MCP and the modified volume element, testable predictions for the propagation of phonons, impurities and vacancies could arise. For instance, the modified volume element should affect the wavefunctions and energy spectrum of impurities and vacancies [12] in the presence of a crystalline defect. The occurrence of bound states due to dislocations[21] can also be studied in the light of the present results. One probably, however, needs to go beyond the eikonal approximation for the test fields, since in this case the trajectory of the associated particle is assumed to be a geodesic curve [12], a good approximation for high-frequency fields, but with no contributions from the torsion tensor. These points are now under investigation.

Acknowledgments

This work was supported by FAPESP and CNPq. The authors are grateful to Dr. C. Miranda for helpful discussions about crystalline defects, and to the Abdus Salam International Centre for Theoretical Physics, Trieste, Italy, where this work was partly done, for the hospitality.

APPENDIX

Let us first discuss how the results of this paper relate to the earlier approach of [2, 3, 4, 5, 10]. Unlike the approach presented here, the analysis of these works explicitly uses the concept of covariant derivative of densities, which are defined as follows. Let h be a scalar density on \mathcal{M} . One can define the covariant derivative of h in two different ways:

$$D_{\mu}h = \partial_{\mu}h - \Gamma_{\mu\kappa}{}^{\kappa}h \tag{27}$$

or

$$\tilde{D}_{\mu}h = \partial_{\mu}h - \Gamma_{\kappa\mu}{}^{\kappa}h. \tag{28}$$

The definition (27) can be called the *usual* covariant derivative, since, as we show in the following,

$$\nabla_X(h \, dx^1 \wedge \dots \wedge dx^n) = [X^{\mu}D_{\mu}h] \, dx^1 \wedge \dots \wedge dx^n. \tag{29}$$

The covariant derivative (28), used also, for instance, in [22], is the so-called transposed derivative[3], since it can be interpreted as the covariant derivative with respect to the transposed connection $\tilde{\Gamma}_{\mu\nu}{}^{\kappa} = \Gamma_{\nu\mu}{}^{\kappa}$ (see, in this context, the volume-preserving connection introduced in [19]). Note that, according to eq. (3), $\tilde{D}_{\mu}h = D_{\mu}h$ provided that the trace of the torsion tensor vanishes so that, in this case, no ambiguity concerning the definition of the covariant derivative of densities arises. The covariant derivative (28) is closer to the *Lie derivative* on \mathcal{M} , since it satisfies, as we will see,

$$\pounds_X(h \, dx^1 \wedge \dots \wedge dx^n) = [(D_\mu X^\mu) \, h + X^\mu (\tilde{D}_\mu h)] dx^1 \wedge \dots \wedge dx^n. \tag{30}$$

The mathematical property that distinguishes the volume element (1) in [10] is that, whereas the canonical volume element (9) is covariantly constant with respect to the usual derivative,

$$D_{\mu}\sqrt{g} = 0, (31)$$

the volume element (1) is covariantly constant with respect to the transposed derivative,

$$\tilde{D}_{\mu}(e^{\theta}\sqrt{g}) = 0, \tag{32}$$

provided that

$$T_{\mu} + V_{\mu} = \partial_{\mu}\theta. \tag{33}$$

By comparing with eq. (30), we see that (18) holds if and only if one chooses a volume form $\tau = f\tau_0$ such that

$$\tilde{D}_{\mu}(f\sqrt{q}) = 0, \tag{34}$$

which has a solution only if $T_{\mu} + V_{\mu} = \partial_{\mu}\theta$ and $f = e^{\theta}$ [10]. This shows how the earlier approach of [2, 3, 4, 5, 10] relates to the present work, which, as discussed in the previous sections, is based on weaker physical assumptions.

We end by providing proofs of identities (29) and (30), and of Proposition 1. Identity (29) can be proved as follows:

$$\nabla_X (h dx^1 \wedge \dots \wedge dx^n) = (\nabla_X h) dx^1 \wedge \dots \wedge dx^n + h \sum_{\kappa} dx^1 \wedge \dots \wedge \nabla_X dx^{\kappa} \wedge \dots \wedge dx^n$$

$$= X(h) dx^1 \wedge \dots \wedge dx^n + h \sum_{\kappa} dx^1 \wedge \dots \wedge (-X^{\mu} \Gamma_{\mu\nu}{}^{\kappa} dx^{\nu}) \wedge \dots \wedge dx^n$$

$$= X^{\mu} (\partial_{\mu} h - \Gamma_{\mu\kappa}{}^{\kappa}) dx^1 \wedge \dots \wedge dx^n.$$
(35)

As for (30), since \mathcal{L}_X and d commute, we have

$$\pounds_X dx^{\kappa} = d\pounds_X x^{\kappa} = dX^{\kappa} = \partial_{\nu} X^{\kappa} dx^{\nu}, \tag{36}$$

which leads to

$$\pounds_X(hdx^1 \wedge \dots \wedge dx^n) = \pounds_X(h) dx^1 \wedge \dots \wedge dx^n + h \sum_{\kappa} dx^1 \wedge \dots \wedge \pounds_X dx^{\kappa} \wedge \dots \wedge dx^n$$
 (37)

$$= [X(h) + h\partial_{\kappa}X^{\kappa}] dx^{1} \wedge \dots \wedge dx^{n}$$
(38)

$$= [hD_{\mu}X^{\mu} + X^{\mu}(\partial_{\mu}h - h\Gamma_{\kappa\mu}{}^{\kappa})] dx^{1} \wedge \dots \wedge dx^{n}.$$
(39)

Finally, Proposition 1 can be proved as follows. We first notice that

$$D_{\kappa}X^{\kappa} = \partial_{\kappa}X^{\kappa} + \Gamma_{\kappa\mu}{}^{\kappa}X^{\mu} \tag{40}$$

$$= \partial_{\kappa} X^{\kappa} + \left(\begin{Bmatrix} \kappa \\ \kappa \mu \end{Bmatrix} + K_{\kappa \mu}{}^{\kappa} + V_{\kappa \mu}{}^{\kappa} \right) X^{\mu} \tag{41}$$

$$= \partial_{\kappa} X^{\kappa} + (\partial_{\kappa} \ln \sqrt{g} + T_{\kappa} + V_{\kappa}) X^{\kappa}. \tag{42}$$

On the other hand, it follows from eq. (38) that

$$\pounds_X \left(f \sqrt{g} \, dx^1 \wedge \dots \wedge dx^n \right) = \left(X^{\kappa} \partial_{\kappa} \ln(f \sqrt{g}) + \partial_{\kappa} X^{\kappa} \right) \tau. \tag{43}$$

Thus, from definition (14),

$$\operatorname{div}_{\tau} X = X^{\kappa} \partial_{\kappa} \ln(f \sqrt{g}) + \partial_{\kappa} X^{\kappa}, \tag{44}$$

which finally yields

$$D_{\kappa}X^{\kappa} = \operatorname{div}_{\tau}X + (T_{\kappa} + V_{\kappa} - \partial_{\kappa}\ln f)X^{\kappa}. \tag{45}$$

- F.W. Hehl, P. Von Der Heyde, G.D. Kerlick and J.M. Nester, Rev. Mod. Phys. 48, 393 (1976); R.T. Hammond, Rept. Prog. Phys. 65, 599 (2002).
- [2] A. Saa, Gen. Rel. Grav. 29, 205 (1997), e-Print: gr-qc/9609011.
- [3] P.P. Fiziev, Gen. Rel. Grav. 30, 1341 (1998), e-Print: gr-qc/9712004.
- [4] A. Saa, Mod. Phys. Lett. A9, 971 (1994), e-Print: gr-qc/9306026.
- [5] A. Saa, Class. Quantum Grav. 12, L85 (1995), e-Print: hep-th/9307095.
- [6] E. Guendelman, A. Kaganovich, E. Nissimov and S. Pacheva, New Physics from a dynamical volume element, e-Print: hep-th/0411122.
- [7] I.L. Shapiro, Phys. Rept. 357, 113 (2001), e-Print: hep-th/0103093; I.L. Buchbinder, S.D. Odintsov and I.L. Shapiro, Effective action in quantum gravity, IOP (UK) (1992).
- [8] E. Kröner, Continuum theory of defects, in Les Houches 1980, Physics of Defects, Ed. R. Balian, M. Kleman and J.P. Poirier, pp. 315-315, North-Holland, 1980.
- [9] H. Kleinert, Gauge fields in condensed matter, Vol. 2, World Scientific (1989).
- [10] A. Saa, Mod. Phys. Lett. A8, 2565 (1993); J. Geom. Phys. 15, 102 (1995), e-Print: hep-th/9308087; Grav. Cosmol. 2, 89 (1996), e-Print: gr-qc/9405054.
- [11] C.W. Misner, K.S. Thorne and J.A. Wheeler, Gravitation, W. H. Freeman and Company (1973).
- [12] M.O. Katanaev and I.V. Volovich, Ann. Phys. 216, 1 (1992); 271, 203 (1999), e-Print: gr-qc/9801081; M.O. Katanaev, Geometric theory of defects, e-Print: cond-mat/0407469.
- [13] T. Boyadjiev, P.P. Fiziev and S. Yazadjiev, Class. Quant. Grav. 16, 2359 (1999), e-Print: gr-qc/9803084; P. Fiziev and S. Yazadjiev, Mod. Phys. Lett. A14, 511 (1999), e-Print: gr-qc/9807025.
- [14] S. Manoff, Gen. Rel. Grav. 32, 1559 (2000), e-Print: gr-qc/0010048; Class. Quant. Grav. 18, 1111 (2001), e-Print: gr-qc/9908061.
- [15] S. Kobayashi and K. Nomizu, Foundations of Differential Geometry, vol. I, John Wiley & Sons (1963).
- [16] We note that the coupling prescription of Teleparallel Gravity is obtained by a different procedure, see R.A. Mosna and J.G. Pereira, Gen. Rel. Grav. **36**, 2525 (2004), e-Print: gr-qc/0312093.
- [17] A. Einstein, The Meaning of Relativity, Princeton University Press (1970).
- [18] E. I. Guendelman, Mod. Phys. Lett. A 14, 1043 (1999), e-Print:gr-qc/9901017; E. I. Guendelman and A. B. Kaganovich, Phys. Rev. D 60, 065004 (1999), e-Print:gr-qc/9905029.
- [19] F.W. Hehl, J.D. McCrea, E.W. Mielke and Y. Neeman, Phys. Rept. 258, 1 (1995).
- [20] M.L. Ruggiero and A. Tartaglia, Am. J. Phys. 71, 1303 (2003), e-Print: gr-qc/0306029.
- [21] R. Bausch, R. Schmitz, and L.A. Turski, Phys. Rev. Lett. 80, 2257 (1998); Phys. Rev. B59, 13491 (1999).
- [22] D. Lovelock and H. Rund, Tensors, differential forms, and variational principles, John Wiley & Sons (1975).