

Noncommutative geometry inspired Schwarzschild black hole

Piero Nicolini*

*Dipartimento di Matematica e Informatica, Università degli Studi di Trieste,
Dipartimento di Matematica, Politecnico di Torino, Turin,
Istituto Nazionale di Fisica Nucleare, Sezione di Trieste, Institut Jožef Stefan, Ljubljana*

Anais Smailagic†

Istituto Nazionale di Fisica Nucleare, Sezione di Trieste

Euro Spallucci‡

*Dipartimento di Fisica Teorica, Università degli Studi di Trieste,
and Istituto Nazionale di Fisica Nucleare, Sezione di Trieste*

We investigate the behavior of a noncommutative radiating Schwarzschild black hole. It is shown that coordinate noncommutativity cures usual problems encountered in the description of the terminal phase of black hole evaporation. More in detail, we find that:
the evaporation end-point is a zero temperature extremal black hole even in the case of electrically neutral, non-rotating, objects;
there exists a finite *maximum* temperature that the black hole can reach before cooling down to *absolute zero*;
there is no curvature singularity at the origin, rather we obtain a regular *DeSitter* core at short distance.

The theoretical discovery of radiating black holes [1] disclosed the first physically relevant window on the mysteries of quantum gravity. After thirty years of intensive research in this field (see [2] for a recent review with an extensive reference list) various aspects of the problem still remain under debate. For instance, a fully satisfactory description of the late stage of black hole evaporation is still missing. The string/black hole correspondence principle [3] suggests that in this extreme regime stringy effects cannot be neglected. This is just one of many examples of how the development of string theory has affected various aspects of theoretical physics. Among different outcomes of string theory, we focus on the result that target spacetime coordinates become *noncommuting* operators on a *D*-brane [4]. Thus, string-brane coupling has put in evidence the necessity of *spacetime quantization*. This indication gave a new boost to reconsider older ideas of similar kind pioneered in a, largely ignored, paper by Snyder [5]. The noncommutativity of spacetime can be encoded in the commutator

$$[\mathbf{x}^\mu, \mathbf{x}^\nu] = i \theta^{\mu\nu} \quad (1)$$

where $\theta^{\mu\nu}$ is an anti-symmetric matrix which determines the fundamental cell discretization of spacetime much in the same way as the Planck constant $\hbar$ discretizes the phase space. The modifications of quantum field theory implied by (1) has been recently investigated to a large extent. Currently the approach to noncommutative quantum field theory follows two distinct paths: one is based on the Weyl-Wigner-Moyal *-product and the other on coordinate coherent state formalism [6], [7]. In a recent paper, following the second formulation, it has been shown that controversial questions of Lorentz invariance and unitarity [8], raised in the *-product approach, can be solved assuming $\theta^{\mu\nu} = \theta \text{diag}(\epsilon_{ij}, \epsilon_{ij} \dots)$ [9], where θ is a constant with dimension of length squared. Furthermore, the coordinate coherent state approach profoundly modifies the structure of the Feynman propagator rendering the theory ultraviolet finite, i.e. curing the short distance behavior of pointlike structures. It is thus reasonable to believe that noncommutativity could as well cure divergences that appear, under various forms, in General Relativity. In such a framework, it would be of particular interest to study the effects of noncommutativity on the terminal phase of black hole evaporation. In the standard formulation, the temperature

*Electronic address: nicolini@cmfd.univ.trieste.it

†Electronic address: anais@ictp.trieste.it

‡Electronic address: spallucci@trieste.infn.it

diverges and an arbitrarily large curvature state is reached. A preliminary analysis both in $(1+1)$ dimensions [10] and in $(3+1)$ linearized General Relativity [11], [12] suggests that noncommutativity can cure this pathological short distance behavior.

In order to prove the above conjecture, and in view of spherical symmetry, it is sufficient to consider a r - t section of the noncommutative Schwarzschild metric. At first glance, one could think of modifying the $4D$ Einstein action to incorporate noncommutative effects. To analyze black hole evaporation in this new framework one has to solve corresponding field equations. Fortunately, we shall argue that it is not necessary to change the Einstein tensor part of the field equations, and that the noncommutative effects can be implemented acting only on the matter source. The line of reasoning is the following. Metric field is a geometrical structure defined over an underlying manifold. Curvature measures the strength of the metric field, i.e. is the response to the presence of a mass-energy distribution. What we know for sure, is that noncommutativity is an *intrinsic property* of the manifold itself, rather than a super-imposed geometrical structure. Under this respect, it affects gravity in a subtle, indirect way. By studying quantum mechanics and quantum field theory in simple non-commutative manifolds, e.g. a plane, one sees that noncommutativity influences matter energy and momentum distribution and propagation [6], [7], [13]. On the other hand, energy-momentum density determines spacetime curvature. Thus, we conclude that in General Relativity the effects of noncommutativity can be taken into account by keeping the standard form of the Einstein tensor in the l.h.s. of the field equations and introducing a modified energy-momentum tensor as a source in the r.h.s. This is exactly the gravitational analogue of the NC modification of quantum field theory [6], [7].

It has been shown [6], [7] that noncommutativity eliminates point-like structures in favor of smeared objects in flat spacetime. The effect of smearing is mathematically implemented as a “substitution rule”: position Dirac-delta function is replaced everywhere with a Gaussian distribution of minimal width $\sqrt{\theta}$. Inspired by this result, we choose the mass density of a static, spherically symmetric, smeared, particle-like gravitational source as

$$\rho_{\theta}(r) = \frac{M}{(4\pi\theta)^{3/2}} \exp(-r^2/4\theta) \quad (2)$$

The particle mass M , instead of being perfectly localized at the point is, instead, *diffused* throughout a region of linear size $\sqrt{\theta}$. This is due to the intrinsic uncertainty encoded in the coordinate commutator (1).

Phenomenological results, so far, imply that noncommutativity is not visible at presently accessible energies, i.e. $\sqrt{\theta} < 10^{-16} \text{ cm}$. Thus, at “large distances” one expects minimal deviations from standard vacuum Schwarzschild geometry. On the other hand, new physics is expected at distance $r \simeq \sqrt{\theta}$ where a non negligible density of energy and momentum is present.

Guided by the above considerations, we are going to look for a static, spherically symmetric, asymptotically Schwarzschild solution of the Einstein equations, with (2) describing the energy density of the system. In order to completely define the energy-momentum tensor, we rely on the covariant conservation condition $T^{\mu\nu}{}_{;\nu} = 0$. For spherically symmetric metric this equation turns out to be

$$\partial_r T^r_r = -\frac{1}{2} g^{00} \partial_r g_{00} (T^r_r - T^0_0) - g^{\theta\theta} \partial_r g_{\theta\theta} (T^r_r - T^{\theta}_{\theta}) \quad (3)$$

To preserve Schwarzschild-like property: $g_{00} = -g_{rr}^{-1}$, we require $T^r_r = T^0_0 = -\rho_{\theta}(r)$. With this requirement the divergence free equation allows a solution for T^{θ}_{θ} which reads

$$T^{\theta}_{\theta} \equiv \partial_r (r T^r_r) = -\rho_{\theta}(r) - r \partial_r \rho_{\theta}(r) \quad (4)$$

Rather than a massive, structureless point, a source turns out to a *self-gravitating, droplet of anisotropic fluid* of density ρ_{θ} , radial pressure $p_r = -\rho_{\theta}$ and *tangential pressure*

$$p_{\perp} = -\rho_{\theta} - r \partial_r \rho_{\theta}(r) \quad (5)$$

On physical grounds, a non-vanishing radial pressure is needed to balance the inward gravitational pull, preventing droplet to collapse into a matter point. This is the basic physical effect on matter caused by spacetime noncommutativity and the origin of all new physics at distance scale of order $\sqrt{\theta}$. By solving Einstein equations with (2) as a matter source. we find the line element [20]:

$$ds^2 = \left(1 - \frac{4M}{r\sqrt{\pi}} \gamma(3/2, r^2/4\theta) \right) dt^2 - \left(1 - \frac{4M}{r\sqrt{\pi}} \gamma(3/2, r^2/4\theta) \right)^{-1} dr^2 - r^2 (d\vartheta^2 + \sin^2 \vartheta d\phi^2) \quad (6)$$

where $\gamma(3/2, r^2/4\theta)$ is the lower incomplete Gamma function:

$$\gamma(3/2, r^2/4\theta) \equiv \int_0^{r^2/4\theta} dt t^{1/2} e^{-t} \quad (7)$$




FIG. 1: g_{rr}^{-1} vs r , for various values of $M/\sqrt{\theta}$. Intercepts on the horizontal axis give radii of the event horizons. $M = \sqrt{\theta}$, (cyan curve) no horizon; $M = 1.9\sqrt{\theta}$, (yellow curve) one *degenerate* horizon $r_0 \approx 3.0\sqrt{\theta}$, *extremal* black hole; $M = 3\sqrt{\theta}$ (magenta curve) two horizons.

The classical Schwarzschild metric is obtained from (6) in the limit $r/\sqrt{\theta} \rightarrow \infty$. Eq.(2) leads to the mass distribution $m(r) = 2M \gamma(3/2, r^2/4\theta) / \sqrt{\pi}$ where M is the total mass of the source. We recall the following properties of matter density ρ_θ :

1. near the origin, i.e. $r \ll \sqrt{\theta}$, $\frac{d\rho_\theta}{dr} \simeq 0 \rightarrow \rho_\theta \simeq \rho_\theta(0)$;
2. few standard deviations from the origin, say $r \geq 4\sqrt{\theta}$, $\frac{d\rho_\theta}{dr} \simeq 0 \rightarrow \rho_\theta \simeq \text{const.} \ll \rho_\theta(0)$
3. asymptotically far away, i.e. $r \gg 2M$, $\rho_\theta = 0$.

The function $p_\perp(r)$ defined in (5) is plotted in **Fig.(3)**

As far as the sign of the pressure is concerned, it is worth to remark that we are *not* dealing with the inward pressure of outer layers of matter on the core of a “star”, but with a totally different kind of “quantum” pressure. It is the outward push, which is conventionally defined to be negative, induced by noncommuting coordinate quantum fluctuations. In a simplified picture, quantum pressure is akin to the cosmological constant in DeSitter universe. As a consistency check of this interpretation we are going to show that line element (6) is well described near the origin by a DeSitter geometry.

The line element (6) describes the geometry of a noncommutative black hole and should give us useful insights about possible noncommutative effects on Hawking radiation.

Eventual event horizon(s) can be found where $g_{00}(r_H) = 0$, that is

$$r_H = \frac{4M}{\sqrt{\pi}} \gamma(3/2, r_H^2/4\theta) \quad (8)$$

Equation (8) cannot be solved in closed form. However, by plotting g_{00} one can read intersections with the r -axis and determine numerically the existence of horizon(s) and their radius. Figure (1) shows that noncommutativity introduces new behavior with respect to standard Schwarzschild black hole. Instead of a single event horizon, there are different possibilities:




FIG. 2: Plot of $\theta \rho$ vs $r/\sqrt{\theta}$ for different values of $M/\sqrt{\theta}$: $M/\sqrt{\theta} = 1$ cyan curve; $M/\sqrt{\theta} = 1.9$ yellow curve; $M/\sqrt{\theta} = 3$ magenta curve. Matter distribution is *flat* both near the origin and for $r \geq 4\sqrt{\theta}$.




FIG. 3: Plot of the tangential pressure p_θ vs $r/\sqrt{\theta}$ for different values of $M/\sqrt{\theta}$. $M/\sqrt{\theta} = 1$ cyan curve; $M/\sqrt{\theta} = 1.9$ yellow curve; $M/\sqrt{\theta} = 3$ magenta curve. The pressure is always negative, and it is flat both near the origin and for $r \geq 5\sqrt{\theta}$. The scale has been chosen to magnify the behavior in the range $2\sqrt{\theta} \leq r \leq 5\sqrt{\theta}$.

1. two distinct horizons for $M > M_0$ (yellow curve) ;
2. one degenerate horizon in $r_0 = 3.0 \times \sqrt{\theta}$, with $M = M_0 = 1.9 \times (\sqrt{\theta})/G$ corresponding to *extremal black hole* (cyan curve);
3. no horizon for $M < M_0$ (violet curve) .

Notice that the outer horizon has always a radius larger than the extremal limit, i.e. $r_H \geq 3\sqrt{\theta}$, thus our previous approximation $r_H \geq 4\sqrt{\theta}$ is quite acceptable to get a qualitative picture of black hole evaporation. A more refined treatment is only required close to the extremal limit if one need precise quantitative estimates.

In view of these results, there can be no black hole if the original mass is less than the *minimal mass* M_0 . Furthermore,

contrary to the usual case, there can be *two horizons* for large masses. By increasing M , i.e. for $M \gg M_0$, the *inner horizon* shrinks to zero, while the *outer* one approaches the Schwarzschild value $r_H = 2M$.

Equation (8) can be conveniently rewritten in terms of the upper incomplete Gamma function as

$$r_H \approx 2M \left[1 - \frac{2}{\sqrt{\pi}} \Gamma(3/2, M^2/\theta) \right] \quad (9)$$

The first term in (9) is the Schwarzschild radius, while the second term brings in θ -corrections. In the “large radius” regime $r_H^2/4\theta \gg 1$ equation (9) can be solved by iteration. At the first order in $M/\sqrt{\theta}$, we find

$$r_H = q2M \left(1 - \frac{M}{\sqrt{\pi\theta}} e^{-M^2/\theta} \right) \quad (10)$$

The effect of noncommutativity is exponentially small, which is reasonable to expect since at large distances spacetime can be considered as a smooth classical manifold. On the other hand, at short distance one expects significant changes due to the spacetime fuzziness, i.e. the “quantum geometry” becomes important $r_H \simeq \sqrt{\theta}$. Let us now consider the black hole temperature:

$$\begin{aligned} T_H &\equiv \left(\frac{1}{4\pi} \frac{dg_{00}}{dr} \right)_{r=r_H} \\ &= \frac{1}{4\pi r_H} \left[1 - \frac{r_H^3}{4\theta^{3/2}} \frac{e^{-r_H^2/4\theta}}{\gamma(3/2; r_H^2/4\theta)} \right] \end{aligned} \quad (11)$$

where, we expressed M in terms of r_H from the horizon equation (8). For large black holes, i.e. $r_H^2/4\theta \gg 1$, and one recovers the standard result for the Hawking temperature

$$T_H = \frac{1}{4\pi r_H} \quad (12)$$

At the initial state of radiation the black hole temperature increases while the horizon radius is decreasing. It is crucial to investigate what happens as $r_H \rightarrow \sqrt{\theta}$. In the standard (commutative) case T_H diverges and this puts limit on the validity of the conventional description of Hawking radiation. Against this scenario, temperature (11) includes noncommutative effects which are relevant at distances comparable to $\sqrt{\theta}$. Behavior of the temperature T_H as a function of the horizon radius is plotted in **Fig.(4)**.

In the region $r_H \simeq \sqrt{\theta}$, T_H deviates from the standard hyperbola (12). Instead of exploding with shrinking r_H , T_H reaches a maximum[21] in $r_H \simeq 4.7\sqrt{\theta}$ corresponding to a mass $M \approx 2.4\sqrt{\theta}/G_N$, then quickly drops to zero for $r_H = r_0 = 3.0\sqrt{\theta}$ corresponding to the radius of the extremal black hole in figure (1). In the region $r < r_0$ there is no black hole and the corresponding temperature cannot be defined. As a summary of the results, the emerging picture of non commutative black hole is that for $M \gg M_0$ the temperature is given by the Hawking temperature (12) with negligibly small exponential corrections, and increases, as the mass is radiated away. T_H reaches a maximum value at $M = 2.4\sqrt{\theta}$ and then drops as M approaches M_0 . When $M = M_0$, $T_H = 0$, event horizon is degenerate, and we are left with a “frozen” extremal black hole.

The temperature behavior shows that noncommutativity plays the same role in General Relativity as in Quantum Field Theory, i.e. removes short distance divergences.

At this point, important issue of Hawking radiation back-reaction should be discussed. In commutative case one expects relevant back-reaction effects during the terminal stage of evaporation because of huge increase of temperature [15], [16]. As it has been shown, the role of noncommutativity is to cool down the black hole in the final stage. As a consequence, there is a suppression of quantum back-reaction since the black hole emits less and less energy. Eventually, back-reaction may be important during the maximum temperature phase. In order to estimate its importance in this region, let us look at the thermal energy $E = T_H \simeq 0.015/\sqrt{\theta}$ and the total mass $M \simeq 2.4\sqrt{\theta} M_{Pl}^2$. In order to have significant back-reaction effect T_H^{Max} should be of the same order of magnitude as M . This condition leads to the estimate

$$\sqrt{\theta} \approx 0.2 l_{Pl} \sim 10^{-34} \text{ cm} \quad (13)$$




FIG. 4: Plot of T_H vs r_H , in $\sqrt{\theta}$ units. yellow curve is the plot of (11): $T_H = 0$ for $r_H = r_0 = 3.0\sqrt{\theta}$, i.e. for the extremal black hole, while the maximum temperature $T_H \simeq 0.015 \times 1/\sqrt{\theta}$ corresponds to a mass $M \simeq 2.4 \times \sqrt{\theta}$. For comparison, we plotted in yellow the standard Hawking temperature. The two temperatures coincide for $r_H > 6\sqrt{\theta}$.

Expected values of $\sqrt{\theta}$ are well above the Planck length l_{Pl} , and (13) indicates that back-reaction effects are suppressed even if $\sqrt{\theta} \approx 10 l_{Pl}$, and $T_H^{Max} \approx 10^{16} \text{ GeV}$. For this reason we can safely use unmodified form of the metric (6) during all the evaporation process.

Finally, we would like to clarify what happens if the starting object has mass smaller than M_0 . It has been shown that in this case there is no event horizon. Could it be a *naked singularity*? We are going to show that this is not the case. In order to establish the geometrical picture for sub-minimally massive objects we are going to study the curvature scalar near $r = 0$. In case of naked singularity one should obtain divergent curvature, $R \rightarrow \infty$. Short distance behavior of R is obtained by tracing the Einstein equations. One finds

$$R(0) = \frac{4M}{\sqrt{\pi}\theta^{3/2}} \quad (14)$$

For $r \ll \sqrt{\theta}$ the curvature is actually *constant and positive*. An eventual naked singularity is replaced by a DeSitter, *regular* geometry around the origin. Earlier attempts to avoid curvature singularity at the origin of Schwarzschild metric have been made by matching DeSitter and Schwarzschild geometries both along time-like [17], and space-like matter shells [18], or by cleverly guessing regular black hole geometries and deriving the corresponding energy momentum tensor from the Einstein equations [19]. Our approach is “dual” to the one in [19] as we start from an energy-momentum tensor originating from noncommutative geometry and with this source we solve Einstein equation. As a conclusion, the results derived in this letter show that the coordinate coherent state approach to noncommutative effects can cure the singularity problems at the terminal stage of black hole evaporation.

We have shown that noncommutativity, which is an intrinsic property of the manifold itself, can be introduced in General Relativity by modifying the matter source. Energy-momentum tensor needed for this description is of the ideal fluid form and requires a non-trivial pressure. In spite of complicated equation of state it can be studied in the regions of interest and new black hole behavior is discovered in the region $r \simeq \sqrt{\theta}$. Specifically, we have shown that there is a minimal mass $M_0 = 1.9\sqrt{\theta}$ to which a black hole can decay through Hawking radiation.

The reason why it does not end-up into a naked singularity is due to the finiteness of the curvature at the origin. The everywhere regular geometry and the residual mass M_0 are both manifestations of the Gaussian de-localization of the source in the noncommutative spacetime. On the thermodynamic side, the same kind of regularization takes place eliminating the divergent behavior of Hawking temperature. As a consequence there is a maximum temperature that the black hole can reach before cooling down to absolute zero. As already anticipated in the introduction, noncommutativity regularizes divergent quantities in the final stage of black hole evaporation in the same way it cured UV infinities in noncommutative quantum field theory.

We have also estimated that back-reaction does not modify the original metric in a significant manner.

Acknowledgments

We would like to thank Dr. Thomas G. Rizzo for a careful reading of the paper and for pointing out a couple of misprints.

One of us, P.N., thanks the “Dipartimento di Fisica Teorica, Università di Trieste” and PRIN-COFIN project 2004 “Metodi matematici per le teorie cinetiche” for financial support.

-
- [1] S. W. Hawking, *Comm. Math. Phys.* **43** 199 (1975)
 - [2] T. Padmanabhan, *Phys. Rep.* **406** 49 (2005)
 - [3] L. Susskind, *Phys. Rev. Lett.* **71** 2367 (1993)
 - [4] E. Witten, *Nucl. Phys. B* **460** 335 (1996); N. Seiberg, E. Witten, *JHEP* **9909** 032 (1999)
 - [5] H. S. Snyder, *Phys. Rev.* **71** 38 (1947)
 - [6] A. Smailagic, E. Spallucci, *J. Phys.* **A36** L467 (2003)
 - [7] A. Smailagic, E. Spallucci, *J. Phys.* **A36** L517 (2003)
 - [8] M. Chaichian, A. Demichev, P. Presnajder, *Nucl. Phys.* **B567** (2000) 360; S. Cho, R. Hinterding, J. Madore, H. Steinacker, *Int. J. Mod. Phys. D9* (2000) 161
 - [9] A. Smailagic, E. Spallucci, *J. Phys.* **A37** 7169 (2004)
 - [10] P. Nicolini, A. Smailagic and E. Spallucci, “*The fate of radiating black holes in noncommutative geometry* Proceedings of EPS 13 General Conference ”Beyond Einstein, Physics for the 21st Century” July 11-15, 2005, University of Bern, Bern, Switzerland; e-Print Archive: hep-th/0507226
 - [11] A. Gruppuso, *J. Phys.* **A38** 2039 (2005)
 - [12] P. Nicolini *J. Phys.* **A38** L631 (2005)
 - [13] A. Smailagic, E. Spallucci, *Phys. Rev.* **D65** (2002) 107701
A. Smailagic, E. Spallucci, *J. Phys.* **A35** L363 (2002)
 - [14] R. C. Tolmann, *Phys. Rev.* **55**, 364 (1939);
J. R. Oppenheimer, G. M. Volkoff *Phys. Rev.* **55**, 374 (1939)
 - [15] R. Balbinot, A. Barletta, *Class. Quant. Grav.* **6** 195, 1989;
 - [16] R. Balbinot, A. Fabbri, V. Frolov, P. Nicolini, P. J. Sutton, A. Zelnikov, *Phys. Rev.* **D63** 084029 (2001);
R. Balbinot, A. Fabbri, P. Nicolini, P. J. Sutton, *Phys. Rev.* **D66** 024014 (2002)
 - [17] A. Aurilia, G. Denardo, F. Legovini, E. Spallucci *Phys. Lett.* **B147** 258 (1984) A. Aurilia, G. Denardo, F. Legovini, E. Spallucci *Nucl. Phys. B* **252** 523 (1985)
A. Aurilia, R.S. Kissack, R. Mann *Phys. Rev.* **D35** 2961 (1987)
 - [18] V. P. Frolov, M.A. Markov, V.F. Mukhanov *Phys. Rev.* **D41** 383 (1990)
 - [19] D. A. Easson *JHEP* **0302** 037 (2003)
S. A. Hayward “*Formation and evaporation of regular black holes*” e-Print Archive: gr-qc/0506126
 - [20] We use convenient units $G_N = 1$, $c = 1$.
 - [21] We are in the range of distances, $r_H \geq 4\sqrt{\theta}$, where our approximation is physically reliable.