Non-Metric Gravity I: Field Equations

Kirill Krasnov *

v3: Oct 15, 2007

Abstract

We describe and study a certain class of modified gravity theories. Our starting point is Plebański formulation of gravity in terms of a triple B^i of 2-forms, a connection A^i and a "Lagrange multiplier" field Ψ^{ij} . The generalisation we consider stems from presence in the action of an extra term proportional to a scalar function of Ψ^{ij} . As in the usual Plebański general relativity (GR) case, a certain metric can be constructed from B^i . However, unlike in GR, the connection A^i no longer coincides with the self-dual part of the metric-compatible spin-connection. Field equations of the theory are shown to be relations between derivatives of the metric and components of field Ψ^{ij} , as well as its derivatives, the later being in contrast to the GR case. The equations are of second order in derivatives. An analog of the Bianchi identity is still present in the theory, as well as its contracted version tantamount to energy conservation equation.

1 Introduction

Modified gravity theories have become popular recently, motivated mainly by the fact that Einstein's general relativity (GR) interprets the available observational data in terms of most of the content of the universe being in the form of some dark energy and matter that have never been observed directly. Modified gravity theories are essentially of two main types. One type introduces new fields. Some examples of these are: Brans-Dicke [1] (a generalisation of GR in which the gravitational constant is replaced by a scalar field), a scalar-vector-tensor theory of Bekenstein [2] that is designed to give modified Newtonian dynamics (MOND) as its non-relativistic limit, the non-symmetric gravitation theory of Moffat [3] that is a theory with non-metric gravitational degrees of freedom, a scalar-vector-tensor theory of Moffat, see e.g. [4], brane-world theories [5], [6] that introduce an extra spacetime dimension (an thus an infinite number of new fields). Another interesting recent proposal [7] calls for gravity to be described by the so-called area metric instead of the metric in the usual sense. One of the major challenges for all these models is to show how Einstein's GR arises in "usual circumstances". The other type of modifications considered is that of pure GR, i.e. with no new fields added, but with modified field equations. The examples of such modifications considered so far all involved higher derivatives. Indeed, it is well-known that Einstein's GR is the only diffeomorphism-invariant theory in which the main dynamical field is the spacetime metric and which has at most second order derivatives in the action. Therefore, in order to modify GR without introducing new fields there is no other way but to introduce higher derivative terms in the action. These modifications typically also increase the number of degrees of freedom described by the theory, as one now needs to specify more initial data (more than just first time derivatives of the spatial metric). The quantum gravity induces modifications (quantum corrections) are precisely of this type, which is one of the reasons for interest in this models. Some examples of these are: theories used for string cosmology, see e.g. [8], Weyl conformal gravity considered in e.g. [9], the so-called f(R) theories [10]. There are other modified gravity theories considered in the literature, and the above list makes no attempt to be complete. The two types described are not necessarily mutually exclusive, as integrating the new fields out in a theory of the first type one typically gets a theory of the second type (typically with an infinite number of derivatives - a non-local theory). One of the lessons learnt from all the above developments is that it is rather hard to modify Einstein's general relativity in any interesting way and to remain consistent with what is observed.

In this paper we introduce and study a new class of modified gravity theories which in a certain sense falls simultaneously into both of the categories described above. Thus, to formulate the theory certain new fields are added to the gravity action. However, this is done in such a way that the new fields do not propagate. Another, more precise way to describe the class of theories considered in this paper is as follows. General

^{*}School of Mathematical Sciences, University of Nottingham, Nottingham, NG7 2RD, UK and Perimeter Institute for Theoretical Physics, Waterloo, N2L 2Y5, Canada.

relativity can be described [11] as a certain topological field theory called BF theory (with no propagating degrees of freedom) in which some of the large set of symmetries that make this theory topological have been "removed" or "gauge fixed" by adding to the action certain Lagrange multiplier terms. The theories we consider are exactly of the same type, the difference with GR being that the "topological symmetries" are removed in a certain modified fashion. In other words, a more precise description of the theories we consider is not that new non-propagating degrees of freedom are added as compared to gravity, but that the topological symmetries that are to be removed to get a theory with propagating degrees of freedom are "gauge fixed" in a modified fashion. Thus, what this paper describes is the set of generally covariant modifications of four-dimensional GR that propagate exactly two degrees of freedom. Only very few such generalizations were known in the past, see the end of this Introduction for references. This work provides a construction of an infinite parameter family of such theories. The above description makes it clear that the type of modification we consider is unlike what is done in most if not all previously considered modified gravity scenarios.

Yet another description of the effect of the modification is as follows. As in Plebański formulation of GR [11], a certain new field, referred to as the Lagrange multiplier field Ψ is introduced. In the GR case some of the equations of motion of the theory express Ψ in terms of second derivatives of the metric (or, in terms of the curvature of g). In modified theories we consider Ψ is no longer so simply expressible through g. The equation that relates g and Ψ is, schematically

$$D^2g + \tilde{D}^2f(\Psi) = \Psi + \phi(\Psi), \tag{1}$$

where D^2, \tilde{D}^2 are certain second-order differential operators, and ϕ, f are certain functions of Ψ . This equation can be solved, at least in principle, for Ψ , with the solution being, again schematically

$$\Psi = D^2 g + \sum_{n,m=1}^{\infty} a_{nm} (\tilde{D}^2)^n (D^2 g)^m,$$
(2)

where a_{nm} are some coefficients. Substituted back into the action this gives the usual Einstein-Hilbert functional corrected by an infinite series of higher order terms in the curvature of g.

At the very least, the class of models we consider is an interesting compact (and useful for quantum considerations, see more comments on this below) way of re-writing such infinite expansions. As is clear from (1) the coupled equations for g, Ψ are second order, which guarantees that many unpleasant problems with higher derivative theories are automatically avoided. Moreover, as we have already mentioned above, the new fields Ψ are added in such a way that the structure of the phase space of the theory is unchanged. Thus, there are no new degrees of freedom as compared to the usual GR case, even though this is not at all obvious by examining (1), as it might seem that one has to specify the initial data for both g and Ψ . As a detailed analysis shows, see the recent work of Bengtsson [12], and as we have already commented on above, this is not the case: the count of the degrees of freedom is unchanged as compared to GR. Thus, the class of theories we consider is quite interesting in that it does not increase the number of derivatives in the field equations, as well as does not introduce any new degrees of freedom apart from those already present in GR. The price one has to pay for this is that the theory becomes non-metric.

"Renormalizable" gravity. Most optimistically, this formulation of (quantum corrected) gravity could allow one to come to terms with the behaviour of gravity under renormalization, as we shall now review. It was argued in [13] that in the Plebański formulation [11] the behaviour of gravity under renormalization is more transparent and simple than in the usual metric based scheme. In particular, it was argued that the renormalization group flow is a flow in the space of scalar functions of two complex variables. Thus, using certain field redefinition arguments as well as direct computation (based on a certain gauge fixing procedure) it was argued that the class of theories defined by the action:

$$S[B, A, \Psi] = \frac{1}{8\pi G} \int_M B^i \wedge F^i - \frac{1}{2} \left(\Lambda \delta^{ij} + \Psi^{ij} + \phi \delta^{ij} \right) B^i \wedge B^j, \qquad (3)$$
$$\phi = \phi(\operatorname{Tr}(\Psi^2), \operatorname{Tr}(\Psi^3))$$

is closed under the renormalization group flow. It is in this sense that the theory is referred to as "renormalizable". As explained above, for any ϕ the theory (3) is actually equivalent to a metric theory given by an infinite expansion in powers of the curvature of this metric. Thus, the statement that (3) is closed under the renormalization group flow does not contradict what is known from the usual perturbative quantum gravity. In this action B^i is a (complex) $\mathfrak{su}(2)$ Lie-algebra valued 2-form (indices $i, j, \ldots = 1, 2, 3$ are $\mathfrak{su}(2)$ Lie-algebra ones), $F^i = dA^i + (1/2)[A, A]^i$ is the curvature of an $\mathfrak{s}u(2)$ Lie-algebra valued connection A^i, Ψ^{ij} is a traceless symmetric "Lagrange multiplier" field, that on shell gets related to the "curvature", see more on this below. Let us note that instead of requiring Ψ^{ij} to be traceless one could add to the action a Lagrange multiplier enforcing this condition. Continuing with our description of the above action, Λ is a multiple of the cosmological constant, the "usual" cosmological constant $\tilde{\Lambda}$ appearing in the EH action in the combination $(R-2\tilde{\Lambda})$ is related to the one in (3) via $\Lambda = 3\Lambda$, and G is the Newton's constant. Finally, the function $\phi(\text{Tr}(\Psi^2), \text{Tr}(\Psi^3))$ is (as its arguments indicate) a function of two scalars that can be constructed from Ψ^{ij} . The fact that this function can only depend on two scalar invariants, not three as it would be for a general 3×3 matrix, is a consequence of the tracelessness of Ψ . The function ϕ is zero for Plebański formulation of GR [11]. Equivalently, one can view the usual cosmological constant as a function $\phi(\Psi)$ that is independent of Ψ (constant). As is shown in [13], the additional ϕ -term is generated by quantum corrections, and, importantly, it is argued that this is the only term that gets generated. The renormalization group flow is then a flow in the space of functions ϕ , which makes this function scale dependent: $\phi = \phi_{\mu}$, where μ is energy scale. The asymptotic safety scenario of Weinberg [14] can be reformulated in this context as a conjecture that there exists a non-trivial limit $\phi^* = \lim_{\mu \to \infty} \phi_{\mu}$, and that the theory that describes gravity at our energy scales is on a renormalization group flow trajectory that leads to ϕ^* . However, gravity being a diffeomorphism invariant theory, and there being no way to define what "energy scale" means in a diffeomorphism invariant context (at least in the absence of other fields that can provide one with a reference background), it is not unreasonable to suppose that the theory that describes gravity at all scales is the one with $\phi = \phi^*$, i.e. that the theory at all scales coincides with the UV fixed point one. The main aim of this paper is to study the theory (3) assuming that ϕ is a given function (equal to the unknown ϕ^*). The asymptotic safety conjecture and the problem of determining ϕ^* is not addressed here.

Quantum corrections or a new scale? Before we embark on a systematic study of the theory defined by (3) it is worth emphasising one important conceptual difference between (3) and the usual "quantum corrected" Einstein-Hilbert action. Thus, let us remind the reader that Einstein-Hilbert action receives quantum corrections from counterterms necessary to cancel the divergences arising in perturbation theory. This counterterms have the form of various invariants constructed from Riemann curvature tensor integrated over the spacetime. At leading (first) order the quantum corrections are of the form $(curvature)^2$ integrated over the spacetime. As curvature has dimensions 1/L, L being length, such terms are dimensionless. To give them the dimension $M \cdot L$, M being mass, required from the action one has to multiply these terms by a dimensionfull parameter - the Planck constant \hbar . Thus, the terms in the quantum corrections. This power of \hbar agrees with the fact that these terms arise at one loop order of perturbation theory. Containing a prefactor of \hbar , these terms should be ignored when considering the classical gravity theory, as this is obtained via $\hbar \to 0$ limit. This nicely complements the fact that, had we considered such terms in the classical action, the resulting theory would have fourth derivatives of the dynamical fields, and as such would have various unpleasant problems. This provides us with an "explanation" of why Einstein-Hilbert action is the correct action for the classical theory of gravity.¹

Let us now consider the theory defined by (3). In spite of the seeming similarity to the usual metric based quantum gravity in that all powers of the "curvature field' Ψ arise in counterterms (in (3) the function ϕ may be thought of as a powers series expansion in both of its arguments), the theory (3) is quite different. Indeed, in the usual metric based quantum gravity higher powers of the curvature field imply higher derivatives in the equations of motion. In the case of (3) the field Ψ becomes related to the curvature only on-shell, and the action is second-order for any choice of the function ϕ . This suggests that the ϕ -term of the action (3) should be considered not a quantum correction, but instead an additional term in the classical action. Note, also that the two expansions: one in powers of the curvature in the usual metric based gravity, and the other in powers of Ψ in theory (3) are quite different. Indeed, as is clear from (1) even a seemingly innocuous modification with $\phi = q \operatorname{Tr}(\Psi)^2$ leads to an infinite expansion in powers of the curvature when interpreted in metric terms!

Let us supplement this "second derivatives only" argument with dimensional analysis. The function $\phi(\operatorname{Tr}(\Psi^2), \operatorname{Tr}(\Psi^3))$ has the dimension of the other terms in the brackets in the second term of (3), i.e. $1/L^2$. This term becomes equally important as the term Ψ^{ij} when the curvature field Ψ^{ij} is of the same order as ϕ . This can be phrased in a more meaningful way by saying that the effects of the ϕ -term become significant for curvatures Ψ^{ij} such that the eigenvalues of the matrix Ψ^{ij} are of the same order as the function ϕ of these eigenvalues: $\lambda_{1,2} \sim \phi(\lambda_1, \lambda_2)$. This equation defines a new length scale $l_* : \lambda_{1,2} \sim 1/l_*^2$.

For "small" curvatures the term proportional to $Tr(\Psi)^2$ in the expansion of ϕ is dominant. The coefficient in front of this term has dimensions L^2 . It is this coefficient that can be used as a definition of the length scale

¹There is another, more standard, and actually related explanation of this that uses the Wilsonian renormalization group flow arguments, see e.g. [14]. We will not use such Wilsonian way of reasoning in the present "classical" paper.

$$\phi(\mathrm{Tr}(\Psi^2), \mathrm{Tr}(\Psi^3)) = \pm l_*^2 \mathrm{Tr}(\Psi)^2 + O(\Psi^3), \tag{4}$$

where, in principle, both signs are possible in the above formula.

Now, to determine whether the ϕ -term in the action (3) is a classically important term or just a quantum correction one must ask whether l_* goes to zero when $\hbar \to 0$. In the usual metric based scheme $l_* \sim l_p$, the Planck length and does go to zero in the classical limit. In the pure gravity case, i.e. when no matter degrees of freedom is present, there is no other scale but l_p , so l_* can only be a multiple of l_p . In this case there is no other choice but to interpret the theory (3) as a quantum corrected gravity theory, with the length scale l_* being the Planck scale. The theory then describes how the familiar notions of geometry get modified when one approaches the Planck scale. This interpretation has to be taken with caution, however, as there can only be a limited applicability of a classical description in the essentially quantum regime close to the Planckian scale. Now, when matter degrees of freedom are present in the theory they do affect the running of ϕ and will generally introduce other scales. It is then possible that the modification of the type we consider survives the classical limit. Indeed, the fact that (3) is second-order in derivatives suggests that l_* may remain finite even when \hbar is sent to zero. In other papers of the series we will follow up on this idea and see what kind of modifications arise when the scale l_* is taken to be large (astrophysical).

"Non-metric" modified gravity theory. Thus, if one is to take the quantum gravity scenario reviewed above seriously, one is led to consider the modified gravity theory (3) (with some yet unknown function $\phi(\operatorname{Tr}(\Psi^2), \operatorname{Tr}(\Psi^3))$ depending on some length scales defined by the matter) as the classical limit $\hbar \to 0$ of the quantum corrected theory of gravity. The major difference between (3) as a classical theory of gravitation and GR is that the theory (3) is no longer about metric in any obvious way. Indeed, as in Plebański formulation of GR, the gravity theory (3) is formulated in such a way that the metric never appears. However, in Plebański gravity one of the equation implies "metricity", which then in particular implies that A in (3) coincides with the metric-compatible connection for some metric g. In contrast, when $\phi \neq 0$ it is no longer true that A is metric-compatible. It is for this reason that we propose to refer to this theory (or, rather to the class of theories defined by (3)) as *non-metric gravity*.

In order to avoid a possible confusion we would like to stress that the term "non-metric" is used here in the sense that no spacetime metric appears in the formulation of the theory, not in the sense that some additional "non-metric gravitational" degrees of freedom are present as in e.g. [3].

As it should have become clear from arguments above, Einstein's GR is a very good approximation to theories (3) when curvatures are smaller than $1/l_*^2$. However, when curvatures are large so are deviations from general relativity. In particular, deviations from Einstein's GR are expected to be large near spacetime singularities. As we shall see in one of the subsequent papers, the behaviour of all the fields of theory (3) near e.g. a would be singularity inside a black hole is much less dramatic than that in Einstein's GR. This gives yet another motivation to take the theories (3) seriously.

More on the nature of modification. It is worth explaining the main results of this paper in simple terms, so that the reader is not lost in the rather technical discussion of the main body of the paper. In order to explain what is the main feature that makes gravity (3) different from the usual GR let us recall some basic facts about the self-dual formulation of gravity due to Plebański [11]. In this approach, the main field that replaces the usual metric of GR is a 2-form B^i with values in the Lie algebra of (complexified) $\mathfrak{su}(2)$. One of the equations of the theory states:

$$B^i \wedge B^j = \frac{1}{3} \delta^{ij} B^k \wedge B_k, \tag{5}$$

or, when written in the spinor form

$$B^{AB} \wedge B^{CD} = \frac{1}{6} \epsilon^{A(C} \epsilon^{|B|D)} B^{EF} \wedge B_{EF}.$$
(6)

It is not hard to show that this equation implies that there exists a quadruple of one-forms $\theta^{AA'}$ (A, A' are spinor indices, see the main text for more details) such that B^{AB} is given by:

$$B^{AB} = \frac{1}{2} \theta^{AA'} \wedge \theta^B_{A'}.$$
(7)

In turn, the tetrad $\theta^{AA'}$ allows one to construct a metric $ds^2 = \theta^{AA'} \otimes \theta_{AA'}$. It is in this sense that the equation (5) implies "metricity" of B. One can also show that B^{AB} given by (7) is self-dual as a 2-form, where the self-duality is defined with respect to the above metric. Thus, after the "metricity" equation (5) of Plebański formulation of GR is solved, the field B^{AB} becomes a self-dual 2-form, valued in $\mathfrak{su}(2)$. As such it can be thought of as a map identifying the space of self-dual 2-forms (at a point of spacetime) with the Lie algebra $\mathfrak{su}(2)$. The spinor formalism, which is used to convert spacetime indices into spinor ones, then interprets B^{AB} as the identity map.

In gravity theory (3) most of the above is still true. The triple B^i of 2-forms spans a 3-dimensional subspace in the space of 2-forms, and this subspace can be called the space of self-dual 2-forms. Demanding that there is a metric such that this subspace is the space of self-dual 2-forms with respect to this metric defines the metric modulo conformal factor, see below for more on the ambiguities associated to this procedure of determining the metric via B^i . Thus, B^i becomes a map from $\mathfrak{su}(2)$ to the space of self-dual 2-forms of some metric. However, this map is no longer the identity, see formula (57) for the corresponding expression. There are now at least two different, non-coinciding ways of normalizing this map, which is related to the ambiguity in the choice of the conformal factor for the metric. The action of the metric-compatible derivative operator ∇ on 2-forms and the A-compatible derivative operator \mathcal{D} on $\mathfrak{su}(2)$ -valued quantities is different, see formula (79), (78) for the relations between two derivative operators. This difference is quantified by departure of B^{AB} from the identity map, which is in turn related to certain derivatives of the field ϕ , see relations (39).

The content of field equations of the theory is also similar to that in GR. The equations are given by (89), (90) and, similarly to the usual GR case, state that the anti-self-dual part of the curvature 2-form F(A) is proportional to the (traceless part of the) "stress-energy" 2-form T, while the self-dual part of F(A) is related to the Lagrange multiplier field Ψ as well as to the trace part of T in a certain way. What is different is that one cannot anymore simply solve for the components of Ψ in terms of the second derivatives of the tetrad, as the corresponding equations become more involved and contain second derivatives of Ψ . One now has to solve the combined system of equations for both the tetrad and the components of Ψ simultaneously.

A purely metric formulation? Given the fact that a certain metric (defined by the tetrad) does appear in the theory one may question if its legitimate to refer to the theory as "non-metric". Indeed, as we have sketched above, one might imagine solving for the components of the field Ψ in terms of the derivatives of the tetrad (one will generate an infinite expansion in derivatives this way, as a derivative operator will have to be inverted), and then substituting the solution back into the action (3). One would obtain a generally covariant action for the tetrad-defined metric, which would have the form of an infinite expansion in terms of curvature invariants, with the first term being the usual Einstein-Hilbert action. This is indeed possible in principle, even though rather hard in practise. It would be of interest to see which exactly subclass of such infinite expansions gets produced by actions of the form (3) when ϕ is varied. This procedure seems to suggest that the formulation (3) is just a rather compact way of re-writing a certain class of non-local (containing all powers of the curvature) gravity actions. The critic may argue that this may be interesting by itself, but probably does not call for a name "non-metric" gravity, which suggests a much more profound change of conceptual framework. What we think makes this purely metric interpretation misleading are two things: (i) the metric (or the tetrad) that is obtained in the process of solving the equations for B is defined only up to a conformal factor. We shall return to this point in the main text. (ii) It is both possible and natural to couple matter fields not to a metric (which only arises when one of the equations of motion is solved) but directly to the B field. For example, the action describing a coupling of Maxwell field to non-metric B is given by [15]:

$$S_{EM}[B,a,\phi] = \int_M f(a) \wedge \phi^i B^i - \frac{1}{2} \phi^i \phi^j B^i \wedge B^j, \qquad (8)$$

where a is the electromagnetic potential, f(a) = da, and ϕ^i is a new 0-form field that is introduced to make the coupling directly to B possible, see the above cited paper for more details on this action. Note that this action was introduced in [15] to describe the coupling of Maxwell field to the usual metric gravity in the 2-form formalism. However, this action does make sense even in the non-metric case, as we shall demonstrate in another paper from the series. One could, of course, couple the electromagnetic field directly to the metric arising from B in the usual way, but this seems highly unnatural given the possibility of direct coupling (8). Moreover, such a direct coupling to the metric defined using B would make unavailable the possibility of shifts in B that was heavily used in [13] to describe renormalization. On the contrary, the action (8) being quadratic in B allows for such shifts. This argument suggest a strong criterion for possible matter couplings: the matter should only couple to the B field, and at most quadratic in B couplings are allowed. Work is currently in progress to find a description of the fermionic matter interacting with gravity that couples fermions directly to the B field. **Energy conservation.** The theories we consider, like GR, have Bianchi identity that can be used to get energy conservation equations. In the retrospect, this is not surprising, as energy conservation follows from general covariance of the theory. However, given the rather complicated nature of the generalisation, it is somewhat miraculous that the energy conservation takes the form familiar from GR, as we shall see below.

Relation to "neighbours of GR". An interesting class of modified gravity theories motivated by the pure connection formulation of gravity [16] has been reviewed in e.g. [17]. As described in [12], the class of theories we consider in the present paper is closely related to the theories in [17], our theories giving the most general modification of GR along these lines.

Organisation of the paper. We start in the next section by discussing the content of the modified "metricity" equations. Then, in section 3, we develop spinor techniques for dealing with spacetime forms, which will be of immense value for us in the following sections. Section 4 solves the equations for the gauge field A, with the result being expressed in terms of the usual metric-compatible spin-connection as well as the derivatives of the functions controlling non-metricity. We discuss field equations in section 5. Finally, section 6 derives and analyses the "Bianchi" identity. Appendices discuss some technical details on "metricity" and "compatibility" equations.

2 Modified "metricity" equations

Our first goal is to study what the presence of the ϕ -term in (3) implies for the field B^i . We remind the reader that in the absence of this term the equation one obtains by varying (3) with respect to Ψ^{ij} guarantees "metricity", see (7). When ϕ -term is present, this "metricity" equation gets modified, and it is our goal in this section to study what this modification implies.

The "metricity" condition. Thus, let us consider the equation one gets by varying (3) with respect to Ψ^{ij} . Special care should be taken in view of the fact that Ψ^{ij} is traceless. The equation one gets is as follows:

$$B^{i} \wedge B^{j} + \left(2\partial_{1}\phi\Psi^{ij} + 3\partial_{2}\phi((\Psi^{2})^{ij} - \frac{1}{3}\delta^{ij}\mathrm{Tr}(\Psi)^{2})\right)B^{k} \wedge B_{k} = \frac{1}{3}\delta^{ij}B^{k} \wedge B_{k},\tag{9}$$

where $\partial_{1,2}\phi$ are the partial derivatives of ϕ with respect to the first and second arguments correspondingly. When $\phi = 0$ we get the "usual" metricity equations of Plebański formulation of GR. Before we can continue with our quest on what this equation implies we need to review some facts about the field Ψ^{ij} . The simplest way to describe this field, and relations satisfied by it is via the so-called spinor techniques, which we therefore have to review.

Spinors. We will not attempt a comprehensive introduction to spinors, describing only what is relevant for us here. There are many excellent sources, see e.g. [18], on the subject, to which we refer the reader for more details.

The action (3), as well as the metricity equation (9) has been written using $\mathfrak{s}u(2)$ Lie-algebra valued fields. One can pass to the spinorial language by considering the fundamental representation of this Lie-algebra in terms of traceless 2×2 matrices. Thus, each $\mathfrak{s}u(2)$ index i, j, \ldots gets replaced by a pair of spinor indices AB where $A, B, \ldots = 1, 2$. These indices are raised-lowered using the tensor ϵ^{AB} , which is skew, i.e. $\epsilon^{BA} = -\epsilon^{AB}$. We will use conventions of [18]:

$$\psi^A = \epsilon^{AB} \psi_B, \qquad \psi_B = \psi^A \epsilon_{AB}, \tag{10}$$

i.e. the index that is contracted is always located on top in the first spinor and at the bottom in the second spinor. Due to the fact that ϵ^{AB} is skew, the condition that a second rank spinor X^{AB} is traceless is equivalent to the condition that it is symmetric. Thus, each $\mathfrak{su}(2)$ index i, j, \ldots is equivalent to a symmetric pair of spinor indices AB. Note that the fact that ϵ^{AB} is skew also implies that the product of any spinor with itself is zero: $\lambda^A \lambda_A = 0$.

Spinorial representation of the curvature field Ψ **.** The field Ψ^{ij} is represented in the spinor form by a rank 4 spinor Ψ^{ABCD} . This spinor is symmetric in each of the two pairs of indices AB and CD and is symmetric under the exchange of these two pairs (as Ψ^{ij} is a symmetric matrix). It is easy to see that the condition that Ψ^{ij} is traceless is equivalent to the condition that Ψ^{ABCD} is completely symmetric.

It is often convenient to choose a basis in the space of spinors. Let the basic spinors be denoted by \tilde{i}^A, \tilde{o}^A . The reason why we put a tilde over the basic spinors will become clear in the next section, where we contrast the "internal" spinors that we are dealing with in this section with "spacetime" spinors. The untilded notation is reserved for the "spacetime" spinor basis.

It is customary to normalize the basic spinors as:

$$\tilde{i}^A \tilde{o}_A = 1. \tag{11}$$

The curvature field Ψ can then be decomposed as follows.

$$\Psi^{ABCD} = \Psi_0 \tilde{i}^A \tilde{i}^B \tilde{i}^C \tilde{i}^D + \Psi_1 \tilde{i}^{(A} \tilde{i}^B \tilde{i}^C \tilde{o}^D) + \Psi_2 \tilde{i}^{(A} \tilde{i}^B \tilde{o}^C \tilde{o}^D) + \Psi_3 \tilde{i}^{(A} \tilde{o}^B \tilde{o}^C \tilde{o}^D) + \Psi_4 \tilde{o}^A \tilde{o}^B \tilde{o}^C \tilde{o}^D,$$
(12)

where the five quantities Ψ_0, \ldots, Ψ_4 are the (basis dependent) spinor curvature components.

Basis of rank 2 spinors. It is convenient to introduce the following 3 basic rank 2 spinors:

$$\tilde{i}^{A}\tilde{i}^{B} = \tilde{X}_{-}^{AB}, \qquad \tilde{o}^{A}\tilde{o}^{B} = \tilde{X}_{+}^{AB}, \qquad \tilde{i}^{(A}\tilde{o}^{B)} = \tilde{X}^{AB}.$$
(13)

The following commutational relations are easy to obtain:

$$[\tilde{X}_{-}, \tilde{X}_{+}] = 2\tilde{X}, \qquad [\tilde{X}, \tilde{X}_{+}] = \tilde{X}_{+}, \qquad [\tilde{X}, \tilde{X}_{-}] = -\tilde{X}_{-},$$
(14)

where our convention for the commutator is $[X, Y]_A^B = X_A^C Y_C^B - Y_A^C X_C^B$. This allows us to identify:

$$\tilde{X}_{-} = \begin{pmatrix} 0 & 0 \\ -1 & 0 \end{pmatrix}, \qquad \tilde{X}_{+} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \qquad \tilde{X} = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$
(15)

The rank 2 spinors $\tilde{X}_{\pm}, \tilde{X}$ thus span the $\mathfrak{s}u(2)$ Lie algebra. The Killing-Cartan form evaluated on the basic rank 2 spinors is:

$$(\tilde{X}_+, \tilde{X}_-) = 1, \qquad (\tilde{X}, \tilde{X}) = -\frac{1}{2},$$
(16)

where $(X, Y) := -\operatorname{Tr}(XY) = -X_A^B Y_B^A = X^{AB} Y_{AB}.$

Field Ψ in terms of the rank 2 spinors. The following representation of the field Ψ^{ABCD} in terms of the rank 2 spinors $\tilde{X}_{\pm}, \tilde{X}$ turns out to be convenient:

$$\Psi = \Psi_0 \tilde{X}_- \otimes \tilde{X}_- + \frac{\Psi_1}{2} (\tilde{X}_- \otimes \tilde{X} + \tilde{X} \otimes \tilde{X}_-) + \frac{\Psi_2}{6} (\tilde{X}_+ \otimes \tilde{X}_- + \tilde{X}_- \otimes \tilde{X}_+ + 4\tilde{X} \otimes \tilde{X})$$

$$+ \frac{\Psi_3}{2} (\tilde{X}_+ \otimes \tilde{X} + \tilde{X} \otimes \tilde{X}_+) + \Psi_4 \tilde{X}_+ \otimes \tilde{X}_+.$$
(17)

A proof is by an elementary computation.

Principal spinors and Petrov classification. Any completely symmetric rank 4 spinor Ψ^{ABCD} can be represented as:

$$\Psi^{ABCD} = k_1^{(A} k_2^B k_3^C k_4^{D)}, \tag{18}$$

where k_1^A, \ldots, k_4^A are referred to as *principal spinors*. The *Petrov type* of the curvature field Ψ^{ABCD} is determined according to coincidence relations between the principal spinors. This goes from type I - general, all 4 principal spinors are distinct to type N - all 4 principal spinors coincide.

A convenient gauge. An elementary computation shows that by an SU(2) rotation of the basis \tilde{i}^A, \tilde{o}^A the curvature field Ψ^{ABCD} can be brought into the form:

$$\Psi = \alpha (\tilde{X}_{-} \otimes \tilde{X}_{-} + \tilde{X}_{+} \otimes \tilde{X}_{+}) + \beta (\tilde{X}_{+} \otimes \tilde{X}_{-} + \tilde{X}_{-} \otimes \tilde{X}_{+} + 4\tilde{X} \otimes \tilde{X}),$$
(19)

i.e. the spinor curvature components Ψ_1, Ψ_3 can be eliminated and the curvature components Ψ_0, Ψ_4 can be made equal. This is possible always except in the case when Ψ is of type *III*, i.e. when 3 of the principal spinors coincide and do not coincide with the fourth principal spinor. A local SU(2) transformation allows to bring the field Ψ into the form (19) in any region of spacetime where the Petrov type of Ψ does not change. Let us start by discussing the general case. The other (algebraically special) cases (except type III) can be obtained from the general case sending one of the parameters to zero, or making them equal. The type III case will be treated below separately.

Matrix products. Using the gauge (19) it is easy to evaluate the quantities $Tr(\Psi)^2$, $Tr(\Psi)^3$ that we need as arguments of the function ϕ in terms of the quantities α, β . We have:

$$\operatorname{Tr}(\Psi)^2 = 6\beta^2 + 2\alpha^2, \qquad \operatorname{Tr}(\Psi)^3 = -6\beta^3 + 6\beta\alpha^2.$$
 (20)

Note that these formulae imply that ϕ is a function of α^2 only. Therefore, $\phi_{\alpha}|_{\alpha=0} = 0$, where ϕ_{α} is a partial derivative with respect to α . This fact will be important when we consider (in another paper) the spherically symmetric solution of the theory.

We will also need the traceless part of the matrix Ψ^2 , as it appears in the metricity equation (9). We have:

$$\Psi^2 - \frac{1}{3} \operatorname{Id} \operatorname{Tr}(\Psi)^2 = 2\alpha\beta(\tilde{X}_- \otimes \tilde{X}_- + \tilde{X}_+ \otimes \tilde{X}_+) + \frac{1}{3}(\alpha^2 - 3\beta^2)(\tilde{X}_+ \otimes \tilde{X}_- + \tilde{X}_- \otimes \tilde{X}_+ + 4\tilde{X} \otimes \tilde{X}).$$
(21)

Here Id is the identity tensor, which has the expression

$$\mathrm{Id} = \tilde{X}_{+} \otimes \tilde{X}_{-} + \tilde{X}_{-} \otimes \tilde{X}_{+} - 2\tilde{X} \otimes \tilde{X}.$$
(22)

The tensor that appears in brackets on the left-hand-side of (9) is then given by:

$$(2\alpha\partial_1\phi + 6\alpha\beta\partial_2\phi)(\tilde{X}_- \otimes \tilde{X}_- + \tilde{X}_+ \otimes \tilde{X}_+) + (2\beta\partial_1\phi + (\alpha^2 - 3\beta^2)\partial_2\phi)(\tilde{X}_+ \otimes \tilde{X}_- + \tilde{X}_- \otimes \tilde{X}_+ + 4\tilde{X} \otimes \tilde{X}).$$
(23)

Change of coordinates. It is convenient to use α, β instead of the quantities $\text{Tr}(\Psi)^2, \text{Tr}(\Psi)^3$ as arguments of the function ϕ . The change of coordinates is elementary. Thus, we replace

$$\partial_1 \phi = \partial_\alpha \phi \partial_1 \alpha + \partial_\beta \phi \partial_1 \beta, \qquad \partial_2 \phi = \partial_\alpha \phi \partial_2 \alpha + \partial_\beta \phi \partial_2 \beta, \tag{24}$$

The following identities are then checked by an elementary computation:

$$2\alpha\partial_1\alpha + 6\alpha\beta\partial_2\alpha = 1/2, \qquad 2\beta\partial_1\beta + (\alpha^2 - 3\beta^2)\partial_2\beta = 1/6$$

$$2\alpha\partial_1\beta + 6\alpha\beta\partial_2\beta = 0, \qquad 2\beta\partial_1\alpha(\alpha^2 - 3\beta^2)\partial_2\alpha = 0.$$
(25)

Thus, we get:

$$\frac{\partial\phi}{\partial\Psi} = \frac{\phi_{\alpha}}{2} (\tilde{X}_{-} \otimes \tilde{X}_{-} + \tilde{X}_{+} \otimes \tilde{X}_{+}) + \frac{\phi_{\beta}}{6} (\tilde{X}_{+} \otimes \tilde{X}_{-} + \tilde{X}_{-} \otimes \tilde{X}_{+} + 4\tilde{X} \otimes \tilde{X}).$$
(26)

where $\phi_{\beta}, \phi_{\alpha}$ are the partial derivatives of the function $\phi = \phi(\alpha, \beta)$ viewed as a function of α, β .

Metricity equation. Having obtained a convenient representation (26) for the derivative of the function ϕ with respect to the curvature field Ψ we are ready to write the metricity equations (9) in a simple form. Let us decompose the 2-form $B^i = B^{AB}$ into a basis of rank 2 spinors:

$$B = B_{+}\tilde{X}_{+} + B_{-}\tilde{X}_{-} + B\tilde{X}, \tag{27}$$

where B_{\pm} , B are some 2-forms. The trace of the product of two of these 2-forms is easy to compute:

$$(B \wedge B) = 2B_+ \wedge B_- - \frac{1}{2}B \wedge B.$$
⁽²⁸⁾

Taking into account (26), after some algebra, one obtains for the non-trivial part of the metricity equations (9):

$$B_{+} \wedge B_{+} = B_{-} \wedge B_{-} = -\phi_{\alpha} \left(B_{+} \wedge B_{-} - \frac{1}{4} B \wedge B \right), \qquad (29)$$
$$B_{+} \wedge B = B_{-} \wedge B = 0,$$
$$2B_{+} \wedge B_{-} + B \wedge B = -2\phi_{\beta} \left(B_{+} \wedge B_{-} - \frac{1}{4} B \wedge B \right).$$

In the metric case (usual GR) the right hand side of all these equations is zero.

Interpretation. Let us first discuss how the equations (29) should be interpreted. These equations were obtained by varying the action with respect to field Ψ . As such, they should be interpreted as equations on Ψ , not on B. In the case of usual Plebański gravity these equations are independent of Ψ , and this interpretation is impossible. However, in the case considered the right hand side of these equations is a (non-trivial) function of Ψ , and so they can be used to solve for components of Ψ in terms of components of B, at least in principle.

In order to obtain such relations between B and Ψ in a more explicit form it is desirable to be able to characterise the 2-forms B^i by some scalar functions. These functions could then be related algebraically to components of Ψ via the "metricity" equations (29). To this end it is desirable to choose some basis of oneforms, which can then be used to decompose the 2-forms B_{\pm}, B . Functions appearing as coefficients in this decomposition could then be used as "components" of B_{\pm}, B , and related to the components of Ψ .

It is in the process of constructing such a decomposition of B into some basic two-forms that the notion of a metric appears. The idea is as follows. The 2-forms B_{\pm} , B span (generically) a 3-dimensional subspace in the 6-dimensional space of (complex) 2-forms. Let us denote this subspace by S. The idea is to declare S to be the subspace of self-dual 2-forms. The notion of self-duality depends on a metric. As we shall soon see, declaring S to be the subspace of self-dual 2-forms determines the metric with respect to which these forms are self-dual modulo conformal factor. The above idea is, of course, motivated by the fact that in the usual Plebański GR case, the 2-forms B_{\pm} , B are self-dual 2-forms of some metric will be instrumental in selecting the scalar function that can later be related to the components of Ψ .

To specify a triple of 2-forms B_{\pm} , B one needs $3 \times 6 = 18$ numbers. As one is free to perform internal SL(2) rotations, 3 of these numbers are pure gauge, which leaves one with 15 "degrees of freedom" contained in B_{\pm} , B. As we have described, demanding that this triple of 2-forms is self-dual with respect to some metric determines this metric, up to the ambiguity in the choice of the conformal factor, see more on this below. To specify a metric in 4 spacetime dimensions one needs 10 numbers. Thus, the triple of 2-forms B_{\pm} , B carries 5 more (complex) "degrees of freedom" as compared to a metric. These 5 degrees of freedom can be translated into 5 degrees of freedom of Ψ via the metricity equations. The above considerations quantify the difference between the usual metric theory of gravity, with the main dynamical field being a metric, and the non-metric theory (3), where the main dynamical field is a triple of 2-forms.

Metric. Now, to see how a (complex) metric arises let us choose an arbitrary volume form vol, and using this form define a 3×3 symmetric matrix h^{ij} via:

$$B^i \wedge B^j = h^{ij} vol. \tag{30}$$

We note that in the usual metric case the metricity equations state that the matrix h^{ij} is a multiple of the identity matrix. When eigenvectors of h^{ij} are all linearly independent the matrix h^{ij} can be diagonalised by an SO(3, \mathbb{C}) transformation. The matrix of this transformation is (a_1, a_2, a_3) , where $a_I, I = 1, 2, 3$ are the eigenvectors of h^{ij} normalized such that $a_I^i a_J^i = \delta_{IJ}$. When all the eigenvalues are distinct the eigenvectors a_I^i are uniquely defined (modulo sign). Consider the following two-forms:

$$B_I := \frac{a_I^i}{\sqrt{\lambda_I}} B^i,\tag{31}$$

where both signs of the square root can be taken. The three 2-forms so-defined satisfy:

$$B_I \wedge B_J = \delta_{IJ} \, vol. \tag{32}$$

Modulo the coefficient on the right-hand-side, this is the relation satisfied by the two-forms B^i in the metric case. Thus, the two-forms B_I can be used to construct a metric. This metric is subject to the same discrete ambiguities as in the usual metric case. An additional ambiguity not present in the metric case is that of the conformal factor. Indeed, it is clear that the above construction determines the metric only modulo conformal factor. To fix this ambiguity one has to fix the volume form to be used. A natural choice seems to be:

$$vol = \frac{1}{3}B^i \wedge B^i,\tag{33}$$

which would make the two-forms B_I satisfy exactly the same metricity equation (5) as in the case of Plebański theory. However, other choices are possible, see more on this below.

Reality conditions. In order for the metric defined above to be real Lorentzian the two-forms B^i have to satisfy certain reality conditions. To see what these conditions are we recall the reality conditions that must be satisfied by B^i in the usual metric case. These are discussed in some length in [15], so we simply state them here. They read:

$$B^i \wedge (B^i)^* = 0, \tag{34}$$

where the star denotes the complex conjugation, plus the condition that the volume form (33) is real. Since in the general case the metric is constructed from two-forms B_I in exactly the same way as one proceeds in Plebański theory, to guarantee that a real Lorentzian metric arises one must impose exactly the same reality conditions on B_I . However, since we have assumed that the eigenvectors of h^{ij} are all linearly independent, the two-forms B_I and B^i span the same three-dimensional subspace in the space of two-forms. The conditions on B_I then simply state that this space must be orthogonal to the complex conjugate space. Thus, the reality conditions can be imposed directly on the forms B^i and take exactly the same form as in the case of Plebański theory.

The reader should note, however, that previous to a discussion of how matter couples to this theory of gravity it is unclear what the correct reality conditions are. Indeed, the matter is anticipated to couple directly to the B field, and it is not clear why the metric constructed from B – rather secondary object – should be real. This can only be understood after matter propagation in the background of a general B is considered, something we hope to return to in another paper of this series.

Metricity equations explicitly. Our idea is thus to declare the subspace S spanned by B_{\pm} , B to be the space of self-dual 2-forms of a certain metric. To this end, let us choose a basis of one-forms, which we will denote by l, n, m, \overline{m} , so that the space S coincides with the one spanned by the triple

$$m \wedge l, \qquad n \wedge \bar{m}, \qquad l \wedge n - m \wedge \bar{m}.$$
 (35)

These are exactly the two-forms self-dual with respect to the metric

$$ds^2 = 2l \otimes n - 2m \otimes \bar{m},\tag{36}$$

so finding the metric according to the above construction one also finds the basis l, n, m, \bar{m} . This basis is, of course, only defined modulo SL(2) rotations. Reality conditions imposed on B^i (or, equivalently, B_I , see above) then translate into the condition that l, n are real and $m^* = \bar{m}$.

Once the basis l, n, m, \bar{m} is found, one can use the available freedom in choosing l, n, m, \bar{m} to make sure that the 2-forms B_{\pm} have decompositions of the form:

$$B_{+} = m \wedge l + an \wedge \bar{m}, \qquad B_{-} = n \wedge \bar{m} + a'm \wedge l, \tag{37}$$

which is demonstrated in the Appendix. One can now substitute these expressions for the forms B_{\pm} , as well as a general decomposition for the form B into the metricity equations (29) to find that, in the special spinor basis chosen (in which $\Psi_1 = \Psi_3 = 0$ and $\Psi_0 = \Psi_4$), the field B has the following form:

$$B = \tilde{X}_{-}(m \wedge l + a n \wedge \bar{m}) + \tilde{X}_{+}(n \wedge \bar{m} + a m \wedge l) + \tilde{X}c(l \wedge n - m \wedge \bar{m}),$$
(38)

with a, c related to $\phi_{\alpha}, \phi_{\beta}$ as follows:

$$2a + \phi_{\alpha}(1 + a^2 + c^2/2) = 0, \qquad (1 + a^2 - c^2) + \phi_{\beta}(1 + a^2 + c^2/2) = 0.$$
(39)

These equations should be interpreted as equations for the functions α, β in terms of the functions a, c the later appearing as components of the field B. Note that the fact that only two functions both in B and in Ψ survived is related to the fact that we have chosen a very special basis in the spinor space. The choice of this basis encodes the remaining 3 degrees of freedom of both B and Ψ . One could have repeated the analysis similar to that above working in the general spinor basis. In that case the 2-forms B_{\pm} would be still given by expressions (37), and the 2-form B would have all 3 components, with 3 functions, say c_{\pm}, c appearing in front of the 2-forms (35). The 5 functions a, a', c_{\pm}, c would then be related to 5 components of the field Ψ via the metricity equations. The corresponding formulae are rather cumbersome, and we will refrain from giving them here. The approach that does not rely on a special choice of the spinor basis may be convenient for some developments, but we will not follow it in the present paper.

Let us note that when $\phi = 0$ the solution to (39) is a = 0, c = 1, which, when substituted into (38), gives the usual "metric" 2-form:

$$B = \tilde{X}_{+}m \wedge l + \tilde{X}_{-}n \wedge \bar{m} + \tilde{X}(l \wedge n - m \wedge \bar{m}).$$

$$\tag{40}$$

Different possible metric structures. As we have seen above, the fundamental field of the theory, the B field, defines a metric structure, only modulo ambiguity in the choice of the conformal factor. There are at least a few natural metrics in the same conformal class that can be constructed from B. One of such metrics is the one known as Urbantke metric [19]. The corresponding expression reads:

$$\sqrt{|g^{\mathrm{U}}|} g^{\mathrm{U}}_{\mu\nu} = \frac{1}{3} \epsilon^{\alpha\beta\gamma\delta} \mathrm{Tr} \left(B_{\mu\alpha} B_{\beta\gamma} B_{\delta\nu} \right) \,. \tag{41}$$

As it is not hard to check, this metric constructed directly from B, is in the same conformal class as the one introduced earlier via (36) based on the choice of the tetrad l, n, m, \bar{m} that leads to say (38). However, the conformal factor relating the two metrics is quite non-trivial.

Another natural possibility, already advocated above, is to require that the volume form as defined by the metric coincides with the volume form

$$\frac{1}{3}\mathrm{Tr}\left(B\wedge B\right).\tag{42}$$

This leads to yet a different possible mechanism for fixing the metric. Note that in the usual metric case when B is of the form (40) all the above choices of the conformal factor agree and there is no ambiguity in what the metric is.

B as a map. Above we have seen that the difference between the usual metric case (40) and the general one (38) is due to appearance of 5 more dynamical fields as compared to those described by a metric. The components of Ψ can be expressed in terms of these degrees of freedom via the metricity equations. Note however that in spite of hiving introduced more degrees of freedom into the theory the number of physical degrees of freedom has not changed — the theory (3) still propagates 2 polarisations of the graviton. This e.g. follows from the analysis of the structure of the phase space of the theory performed in [12]. It is important to understand the geometrical interpretation of the new scalar fields introduced. One viewpoint that will prove quite instrumental below is to regard B as a map from the $\mathfrak{su}(2)$ Lie algebra spanned by $\tilde{X}_{\pm}, \tilde{X}$ to the space of self-dual 2-forms. This map thus defines the notion of self-dual 2-forms, and thus defines a metric with respect to which these two-forms are self-dual. However, this map contains more information than just that about a metric, which, for example, follows from the counting of degrees of freedom contained in B that we performed above. In the metric case the field B carries information just about a metric, and can be identified with the identity map after certain additional techniques are developed. The general non-metric B corresponds to a map that is not an identity.

Type III case. When Ψ is Petrov type III the above analysis is not applicable. In this case the spinor basis can be chosen so that the field Ψ becomes

$$\Psi = \alpha (\tilde{X}_+ \otimes \tilde{X} + \tilde{X} \otimes \tilde{X}_+). \tag{43}$$

We will also need the expression for its square:

$$\Psi^2 = -\frac{\alpha^2}{2}\tilde{X}_+ \otimes \tilde{X}_+. \tag{44}$$

Note that this is traceless, so no need to subtract the trace part.

The metricity equations become:

$$B_{-} \wedge B_{-} = B \wedge B_{-} = 2B_{+} \wedge B_{-} + B \wedge B = 0, \tag{45}$$
$$B_{+} \wedge B \sim \alpha (B_{+} \wedge B_{-} - (1/4)B \wedge B), \qquad B_{+} \wedge B_{+} \sim \alpha^{2} (B_{+} \wedge B_{-} - (1/4)B \wedge B).$$

A solution of these equations is:

$$B = \tilde{X}_{-}(m \wedge l) + \tilde{X}_{+}(n \wedge \bar{m} + a(l \wedge n - m \wedge \bar{m})) + \tilde{X}(l \wedge n - m \wedge \bar{m}),$$
(46)

where a is proportional to α . Thus, a "canonical" quadruple of one-forms appears even in the type III case, and the field B is still purely self-dual. We will not consider the Petrov type III case further in the present paper.

3 Spinor techniques

So far we only dealt with algebraic equations for B involving $\mathfrak{su}(2)$ matrices, and we saw that these are much easier to solve if one employs spinor notations. The spinors we have introduced above are "internal" ones, in that they simply replace the internal $\mathfrak{su}(2)$ indices by a pair of spinor ones. In this section we will introduce different spinors, the ones that will allow us to simply computations with forms and vector fields considerably. In usual gravity, the "internal" and "spacetime" spinors coincide, and this statement is equivalent to the statement that B has the form (40). In other words, after spinor methods are employed, B of the metric case is the identity map that identifies the "internal" and "spacetime" spinors. In our more general case we will need to introduce two different types of spinors, one for dealing with "internal" indices, one for spacetime indices. In this section we will remind the reader how the spacetime indices can be converted into spinor ones, and how this simplifies certain computations. The material reviewed here is standard, see e.g. [18]. The only non-standard point is that the spinors we are going to deal with in this section are different from the ones considered above. For this reason, we have denoted the "internal" spinors by tilded letters and reserved the usual notation for the "usual" spacetime spinors. The corresponding "spacetime" spinor basis will be denoted by i^A , o^A .

There are several different ways to introduce spinor techniques. We will first present what we think is a more geometric way, and then give the usual one of e.g. [18].

Two-forms as maps. It will be extremely useful to think of the 2-forms from space S as maps acting on one-forms, and sending them to vector fields. This way of thinking about 2-forms will allow us to develop a spinor approach to forms. What we are about to describe is standard in the usual GR.

Let us consider a given 2-form B. Any one-form k can be exterior multiplied by B with the result being a 3-form $k \wedge B$. Now, given a basis l, n, m, \overline{m} in the space of one-forms we can use this basis to produce 4 numbers out of the 3-form $k \wedge B$. Indeed, multiplying $k \wedge B$ (from say the right) by each of the basis one-forms one obtains a 4-form that must be proportional to the "volume" 4-form $l \wedge n \wedge \overline{m} \wedge \overline{m}$. Let us introduce a special notation for this proportionality coefficient. Thus, we write:

$$k \wedge B \wedge e^{I} := (k \wedge B \wedge e^{I})l \wedge n \wedge m \wedge \bar{m}, \tag{47}$$

which defines the quantity $(k \wedge B \wedge e^I)$. Here e^I is any of the basis one-forms. One can use the 4 quantities $(k \wedge B \wedge e^I)$ to form a vector field $v^a := \sum_I (k \wedge B \wedge e^I) e^{Ia}$. Here e^{Ia} denotes the basis of vector fields dual to $e^I = e^I_a$ one forms. By definition: $l^a n_b + n^a l_b - m^a \bar{m}_b - \bar{m}^a m_b = \delta^a_b$, where δ^a_b is the Kronecker delta - a mixed tensor of rank 2, and $l^a n_a = 1, m^a \bar{m}_a = -1$, while all other scalar products are zero.

The above discussion allows us to interpret any 2-form B as a map from the space of one-forms to the space of vector fields. It is useful to work out this map for the self-dual 2-forms forming a basis in S. An elementary computation gives:

$$m \wedge l: \begin{array}{c} n_a \to -m^a \\ \bar{m}_a \to -l^a \end{array}, \qquad n \wedge \bar{m}: \begin{array}{c} l_a \to \bar{m}^a \\ m_a \to n^a \end{array}, \qquad l \wedge n - m \wedge \bar{m}: \begin{array}{c} l_a \to -l^a \\ n_a \to n^a \\ m_a \to -m^a \\ \bar{m}_a \to \bar{m}^a \end{array}$$
(48)

A similar map for the anti-self-dual forms is given by:

$$l \wedge \bar{m}: \begin{array}{cc} n_a \to -\bar{m}^a \\ m_a \to -l^a \end{array}, \qquad n \wedge m: \begin{array}{cc} l_a \to -m^a \\ \bar{m}_a \to -n^a \end{array}, \qquad l \wedge n + m \wedge \bar{m}: \begin{array}{cc} l_a \to l^a \\ n_a \to -n^a \\ m_a \to -m^a \\ \bar{m}_a \to \bar{m}^a \end{array}$$
(49)

Spinor interpretation. The above described way of viewing two-forms as maps from one-forms to vector fields will allow us to develop a completely form-free approach, in which forms will get replaced by spinorial objects. Let us first deal with the basic one-forms (tetrad-forming) l, n, m, \bar{m} . To replace them by spinor quantities we will have to introduce the so-called primed spinors, in addition to the unprimed spinors we have considered so far. Primed spinors form a different representation of the Lorentz group. The vector representation is obtained by tensoring together the primed and unprimed representations. This is the reason why the primed spinors are necessary to describe one-forms.

The primed spinors are denoted by a symbol with a primed spinorial index next to it, e.g. $\lambda^{A'}$ is a rank one primed spinor. Primed indices are similarly raised and lowered with their corresponding $\epsilon^{A'B'}$ anti-symmetric

rank 2-spinor. The raising-lowering conventions are the same (10) as for the unprimed spinors. It is convenient to introduce a basis in the space of primed spinors. We get the primed spinor basis $i^{A'}, o^{A'}$.

Let us now introduce an important set of relations which are to provide the identification between rank 2 mixed primed-unprimed spinors and one-forms. We write:

$$l_a = o_A o_{A'}, \qquad n_a = i_A i_{A'}, \qquad m_a = o_A i_{A'}, \qquad \bar{m}_a = i_A o_{A'}. \tag{50}$$

As it is clear, these relations depend both on the spinor basis chosen as well as on the basis in the space of one-forms. These relations can be combined into a single quantity

$$\theta_a^{AA'} := l_a i^A i^{A'} + n_a o^A o^{A'} - m_a i^A o^{A'} - \bar{m}_a o^A i^{A'}, \tag{51}$$

which can be viewed as a map from rank 2 mixed spinors into one-forms, the map precisely given by (50).

Using the above relations we can give a description of self-dual 2-forms in terms of symmetric rank 2 unprimed spinors. Indeed, let us identify:

$$m \wedge l = X_+, \qquad n \wedge \bar{m} = X_-, \qquad l \wedge n - m \wedge \bar{m} = -2X$$

$$(52)$$

for the self-dual 2-forms and

$$l \wedge \bar{m} = X'_{+}, \qquad n \wedge m = -X'_{-}, \qquad l \wedge n + m \wedge \bar{m} = 2X'$$
(53)

for the anti-self-dual ones. Here the unprimed rank 2 spinors are essentially the same to the ones we already encountered in (13), and the primed spinors are similar:

$$i^{A}i^{B} := X_{-}^{AB}, \qquad o^{A}o^{B} := X_{+}^{AB}, \qquad \frac{1}{2}(i^{A}o^{B} + o^{A}i^{B}) = X^{AB},$$

$$i^{A'}i^{B'} := X_{-}^{\prime AB}, \qquad o^{A'}o^{B'} := X_{+}^{\prime AB}, \qquad \frac{1}{2}(i^{A'}o^{B'} + o^{A'}i^{B'}) = X^{\prime AB}.$$
(54)

It is then easy to check that the action of the rank 2 spinors on the mixed spinors (50) is exactly as in (48). Indeed, let us consider e.g. $(m \wedge l)(n)$. The corresponding spinor quantity is $X_{+}^{AB}i_{B}i^{A'} = -o^{A}i^{A'} = -m^{a}$, as in (48). Similarly, $(l \wedge \bar{m})(n)$ is given by $X'_{+}^{A'B'}i_{B'}i^{A} = -i^{A}o^{A'} = -\bar{m}^{a}$ as in (49).

Another way to describe two-forms as rank 2-spinors. The following is a more standard approach to spinors, see e.g. [18]. One starts by postulating the relations (50) between the 4 basic one-forms and spinors of the spinor basis. This relation allows one to replace any spacetime index by a mixed pair of spinor indices. It is then easy to see that e.g. self-dual two forms becomes rank 2 unprimed spinors. Indeed, let us consider a 2 form $B = v \wedge w$ obtained by an exterior product of two one-forms v, w. Introducing indices this reads: $B_{ab} = 2v_{[a}w_{b]}$. The last quantity here is given by:

$$v_{[a}w_{b]} = v_{[AA'}w_{BB']} = -\frac{1}{2}(v \wedge w)_{AB}\epsilon_{A'B'} + \frac{1}{2}(v \wedge w)_{A'B'}\epsilon_{AB},$$
(55)

where the quantities $(v \wedge w)_{AB}, (v \wedge w)_{A'B'}$ are defined as:

$$(v \wedge w)_{AB} = v_{(A}^{A'} w_{B)A'}, \qquad (v \wedge w)_{A'B'} = -v_{(A'}^A w_{A|B')}.$$
 (56)

The choice of minus signs in these formulae is so that they agree with (52), (53).

B as a rank 4 spinor. The above description of self-dual two-forms as rank 2 unprimed symmetric spinors allows us to interpret the B field (38) as a rank 4 unprimed spinor. Indeed, using (52) we write:

$$B = a(\tilde{X}_{-} \otimes X_{-} + \tilde{X}_{+} \otimes X_{+}) + \tilde{X}_{-} \otimes X_{+} + \tilde{X}_{+} \otimes X_{-} - 2c\tilde{X} \otimes X.$$

$$(57)$$

We see, therefore, that the *B* field interpreted as a map is the identity map (22) in the metric case a = 0, c = 1 (in this case the "internal" and "spacetime" spinors coincide, as we shall see below), and deviates from identity in the non-metric case. This gives one way to understand what the non-metric deformation does to *B*. For completeness, let us also give the expression for *B* as a rank 4 spinor in the type III case:

$$B = -2a\tilde{X}_{+} \otimes X + Id. \tag{58}$$

Metric compatible derivative operator. Having a basis of one-forms l, n, m, \bar{m} one has the metric (36), and can consider the derivative operator compatible with this metric. We will denote this operator by ∇ . Spinor representation $\nabla_{AA'}$ of this derivative operator will be very convenient for us in what follows. In order to describe the action of this operator on forms, we just have to understand its action on basic spinors, as we have already translated one- and two-forms that are of interest for us into spinor form. Thus, we introduce what is known as *spin-coefficients* according to the following formulae:

$$\gamma_{AA'}^{+} := -i^{B} \nabla_{AA'} i_{B}, \qquad \gamma_{AA'}^{-} := -o^{B} \nabla_{AA'} o_{B}, \qquad \frac{1}{2} \gamma_{AA'} := i^{B} \nabla_{AA'} o_{B} = o^{B} \nabla_{AA'} i_{B}. \tag{59}$$

The derivatives of the basic spinors are then given by the following formulae:

$$\nabla_{AA'}i_B = -\frac{1}{2}\gamma_{AA'}i_B - \gamma^+_{AA'}o_B, \qquad \nabla_{AA'}o_B = \gamma^-_{AA'}i_B + \frac{1}{2}\gamma_{AA'}o_B.$$
(60)

Let us note that the anti-symmetric tensor $\epsilon_{AB} = o_A i_B - i_A o_B$ that serves the role of the metric in the spinor space is preserved by the derivative operator $\nabla : \nabla \epsilon_{AB} = 0$, which is quite easy to check using (60). This fact is related to the fact that ∇ preserves the metric constructed from the basis of one-forms.

Using (60) it is easy to work out the action of the derivative operator ∇ on the basic two-forms. We will only need these results for the self-dual 2-forms which appear in the decomposition of B. Thus, using (52) and (60) we get:

$$\nabla_{AA'} X^{BC}_{-} = -\gamma_{AA'} X^{BC}_{-} - 2\gamma^{+}_{AA'} X^{BC}_{+}, \qquad \nabla_{AA'} X^{BC}_{+} = \gamma_{AA'} X^{BC}_{+} + 2\gamma^{-}_{AA'} X^{BC}_{+}, \qquad (61)$$
$$\nabla_{AA'} X^{BC}_{-} = \gamma^{-}_{AA'} X^{BC}_{-} - \gamma^{+}_{AA'} X^{BC}_{+}.$$

We now note that these derivatives of self-dual 2-forms can be converted into vector fields. This is done simply by contracting the unprimed index of $\nabla_{AA'}$ with an unprimed index of the tensors X_{\pm}, X . Thus, we introduce the vector fields $(\nabla X_{\pm}), (\nabla X)$ corresponding to $\nabla X_{\pm}, \nabla X$, and get for the corresponding mixed spinors:

$$(\nabla X_{-})^{AA'} = -X_{-}^{AB} \gamma_{B}^{A'} - 2X^{AB} \gamma_{B}^{+A'}, \qquad (\nabla X_{+})^{AA'} = X_{+}^{AB} \gamma_{B}^{A'} + 2X^{AB} \gamma_{B}^{-A'}, \qquad (62)$$
$$(\nabla X)^{AA'} = X_{-}^{AB} \gamma_{B}^{-A'} - X_{+}^{AB} \gamma_{B}^{+A'}.$$

These formulae can be rewritten much more compactly by suppressing the spinor indices. Thus, implying matrix multiplication, we can re-write:

$$(\nabla X_{-}) = -X_{-}\gamma - 2X\gamma^{+}, \quad (\nabla X_{+}) = X_{+}\gamma + 2X\gamma^{-}, \quad (\nabla X) = X_{-}\gamma^{-} - X_{+}\gamma^{+}.$$
 (63)

4 Compatibility equations

Having understood what the modified "metricity" equations imply, and developed spinor techniques for dealing with forms and their derivatives, we are ready to analyse another set of equations that follow from the action (3) - the compatibility equations between the 2-form B and the connection A. As we shall see in this section, these equations allow one to find the connection A in terms of the derivatives of the quantities that appear in the expression (38) for B. Spinor techniques developed in the previous section will be of great help here.

The compatibility condition. The condition relating B and A is obtained by varying the action (3) with respect to the connection A. It reads $\mathcal{D}B^i = 0$, where \mathcal{D} is the covariant derivative with respect to the connection A. In spinorial notations this equation reads dB + [A, B] = 0, or, with indices:

$$dB^{AB} + A^{AC} \wedge B^B_C - B^{AC} \wedge A^B_C = 0.$$
⁽⁶⁴⁾

To rewrite this in a more manageable form let us decompose the connection into the basis of "internal" rank 2 spinors:

$$A = \tilde{X}_{-}A_{-} + \tilde{X}_{+}A_{+} + \tilde{X}A.$$
 (65)

The commutator present in (64) is then easy to compute:

$$[A,B] = \tilde{X}_{-}(A_{-} \wedge B - A \wedge B_{-}) + \tilde{X}_{+}(A \wedge B_{+} - A_{+} \wedge B) + 2\tilde{X}(A_{-} \wedge B_{+} - A_{+} \wedge B_{-}).$$
(66)

Derivatives of the basis spinors. To compute the derivative dB we need to act on the spinors \tilde{i}^A, \tilde{o}^A . We could have chosen a constant basis such that the basis spinors do not depend on a point in spacetime. However, to write the field Ψ is its most convenient for computations form we have chosen to adapt the spinor basis at every point to the field Ψ and its proper spinors. Thus, the basis in which the field Ψ has the simple form (19) is generally not constant. Thus, we have to allow for no-vanishing derivatives of \tilde{i}^A, \tilde{o}^A . Let us proceed similarly to what was done in the previous section and decompose these derivatives into the basic spinors:

$$d\tilde{i}^A := h\tilde{i}^A + g\tilde{o}^A, \qquad d\tilde{o}^A := f\tilde{i}^A - h\tilde{o}^A.$$
(67)

Here, h, g, f are one-forms, and to write the second relation we have used the normalization condition $\tilde{i}^A \tilde{o}_A = 1$, which implies that the one-form coefficient in front of \tilde{o}^A in the second relation is minus the one-form coefficient in front of \tilde{i}^A in the first.

Gauge transformations. It is useful to discuss what gauge transformations that act on the basis \tilde{i}^A, \tilde{o}^A translate to when they act on the one-forms h, g, f. Choosing directions of the spinors \tilde{i}^A, \tilde{o}^A and requiring them to be normalized does not fix them completely. Indeed, there is still a freedom of rescaling: $\tilde{i}^a \to \alpha \tilde{i}^A, \tilde{o}^A \to (1/\alpha)\tilde{o}^A$, where α is a complex number different from zero. It is easy to see that under this gauge transformation:

$$h \to h + \alpha^{-1} d\alpha, \qquad g \to \alpha^2 g, \qquad f \to \alpha^{-2} f,$$
 (68)

and so h transforms as a U(1) connection, while the other one-forms g, f transform as Higgs fields.

Derivatives of the basis rank 2 spinors. Using (67) it is easy to compute the derivatives of the basic rank 2 spinors $\tilde{X}_{\pm}, \tilde{X}$. We get:

$$d\tilde{X}_{-} = 2h\tilde{X}_{-} + 2g\tilde{X}, \qquad d\tilde{X}_{+} = 2f\tilde{X} - 2h\tilde{X}_{+}, \qquad d\tilde{X} = f\tilde{X}_{-} + g\tilde{X}_{+}.$$
 (69)

These formulae are quite similar to (61) except that "internal" spinors are considered.

The compatibility equations. Combining (66) and (69) we can write down the compatibility equations that follow from (64):

$$2h \wedge B_{-} + dB_{-} + f \wedge B + A_{-} \wedge B - A \wedge B_{-} = 0,$$

$$g \wedge B + dB_{+} - 2h \wedge B_{+} + A \wedge B_{+} - A_{+} \wedge B = 0,$$

$$2g \wedge B_{-} + dB + 2f \wedge B_{+} + 2(A_{-} \wedge B_{+} - A_{+} \wedge B_{-}) = 0.$$
(70)

Each of these equations is a condition that a certain 3-form vanishes. Therefore, each gives rise to 4 equations when projected onto a basis of 3-forms. Thus, the compatibility equations are $3 \times 4 = 12$ equations for 3 one-forms A_{\pm} , A. We see that the number of equations matches the number of unknowns. Below we will write down an explicit solution for A in terms of the B field of the form (38).

Solving the compatibility equations. To solve the equations (70) we decompose the one-forms A_{\pm} , A into the same basis l, n, m, \bar{m} that was used to write down the expression (38) for B. We then multiply each of the equations (70) by one-forms l, n, m, \bar{m} in turn, and thus extract components of these equations. This procedure allows us to find A in terms of B.

A very convenient way of doing this is to use the spinor method developed in the previous section. In view of (50) decomposing A_{\pm} , A into basic one-forms is equivalent to introducing the rank 2 mixed spinors $A_{\pm AA'}$, $A_{AA'}$. The field B can in turn be represented by a rank 4 unprimed spinor (57), or, equivalently, each of the 2-forms B_{\pm} , B can be represented by a rank 2 unprimed spinor. We have:

$$B_{-} = X_{+} + aX_{-}, \qquad B_{+} = X_{-} + aX_{+}, \qquad B = -2cX, \tag{71}$$

where we have omitted spinor indices for compactness. Note that it is the "spacetime" rank 2 spinors X_{\pm}, X that are used here. These should not be confused with the "internal" ones. To solve compatibility conditions we will need the inverses of these matrices, where the inverse B^{-1} of B is defined so that $(B^{-1})^{AC}B_C^B = \epsilon^{AB}$. These are easy to find, we get:

$$B_{-}^{-1} = -X_{-} - (1/a)X_{+}, \qquad B_{+}^{-1} = -X_{+} - (1/a)X_{-}, \qquad B^{-1} = -(2/c)X.$$
(72)

In preparation for solving the equations (70), let us introduce special notations for the commutators that appear in (70):

$$C_{-} = A_{-} \wedge B - A \wedge B_{-}, \qquad C_{+} = A \wedge B_{+} - A_{+} \wedge B, \qquad C = A_{-} \wedge B_{+} - A_{+} \wedge B_{-}.$$
(73)

Using the spinorial notations for all the quantities, and converting the 3-forms that appear here into vector fields, these expressions can be rewritten as:

$$C_{-}^{FF'} = B^{FE}A_{-E}^{F'} - B_{-}^{FE}A_{E}^{F'}, \qquad C_{+}^{FF'} = B_{+}^{FE}A_{E}^{F'} - B^{FE}A_{+E}^{F'}, \qquad C^{FF'} = B_{+}^{FE}A_{-E}^{F'} - B_{-}^{FE}A_{+E}^{F'}, \tag{74}$$

or, in a more compact form, implying matrix multiplication:

$$C_{-} = BA_{-} - B_{-}A, \qquad C_{+} = B_{+}A - BA_{+}, \qquad C = B_{+}A_{-} - B_{-}A_{+}.$$
 (75)

Now, assuming C_{\pm} , C are known, it is easy to solve for A_{\pm} , A. Thus, for example, multiplying the first equation by $B_{+}B^{-1}$ and the second by $B_{-}B^{-1}$, adding the results and then subtracting the third equation we get:

$$(B_{+}B^{-1}B_{-} - B_{-}B^{-1}B_{+})A = C - B_{+}B^{-1}C_{-} - B_{-}B^{-1}C_{+}.$$
(76)

One then checks that

$$B_{+}B^{-1}B_{-} - B_{-}B^{-1}B_{+} = \frac{1-a^{2}}{c}\mathrm{Id},$$
(77)

and therefore

$$A = \frac{c}{1 - a^2} \left(C - B_+ B^{-1} C_- - B_- B^{-1} C_+ \right).$$
(78)

By similar manipulations

$$A_{-} = \frac{a}{c(1-a^{2})} \left(-BB_{-}^{-1}C + B_{+}B_{-}^{-1}C_{-} + C_{+} \right), \quad A_{+} = \frac{a}{c(1-a^{2})} \left(-BB_{+}^{-1}C + C_{-} + B_{-}B_{+}^{-1}C_{+} \right).$$
(79)

Dependence on one-forms h, g, f. One should now simply take the expressions for C_{\pm}, C from the compatibility conditions (70) and substitute these into the above expressions to find A. It can now be checked that the one-forms h, g, f appear in the connection A in the following simple way:

$$A = -\tilde{X}_{-}f + \tilde{X}_{+}g + 2\tilde{X}h + \tilde{A}, \tag{80}$$

where \hat{A} is independent of h, g, f. This is as expected, for

$$-\tilde{X}_{-}f + \tilde{X}_{+}g + 2\tilde{X}h = \begin{pmatrix} h & g \\ f & -h \end{pmatrix} = dG^{-1} \cdot G,$$
(81)

where G is an SL(2) transformation that sends the basis \tilde{i}^A, \tilde{o}^A into some constant basis.

We note that the components of \tilde{A} have a very simple meaning:

$$\tilde{i}^A \mathcal{D}\tilde{i}_A = -\tilde{A}_+, \qquad \tilde{o}^A \mathcal{D}\tilde{o}_A = -\tilde{A}_-, \qquad \tilde{o}^A \mathcal{D}\tilde{i}_A = \tilde{i}^A \mathcal{D}\tilde{o}_A = \frac{1}{2}\tilde{A},$$
(82)

where \mathcal{D} is the covariant derivative with respect to the connection A. These relations are easy to check using (80). Thus, the components of \tilde{A} are the spin-coefficients, but for the "internal" spinor basis instead of the "spacetime" one. As we shall see very soon, in the metric case the two sets of spin-coefficients coincide. However, in a more general non-metric situation the two sets are different. Correspondingly, the covariant derivatives that act on "internal" and "spacetime" spinor indices are different.

Spin-coefficients. To compute \tilde{A} we have to substitute

$$\tilde{C}_{-} = -(\nabla B_{-}), \qquad \tilde{C}_{+} = -(\nabla B_{+}), \qquad \tilde{C} = -(1/2)(\nabla B),$$
(83)

where B_{\pm} , B are given by (71) into the formulae (79), (78). Note that we are free to replace the usual derivatives here by the covariant ones, as their (exterior product) action on forms is the same. Thus, using (63) we have:

$$\tilde{C}_{-} = X_{-}(a\gamma - \nabla a) - X_{+}\gamma + 2X(a\gamma^{+} - \gamma^{-}), \qquad \tilde{C}_{+} = X_{-}\gamma - X_{+}(a\gamma + \nabla a) + 2X(\gamma^{+} - a\gamma^{-}), \qquad (84)$$
$$\tilde{C} = cX_{-}\gamma^{-} - cX_{+}\gamma^{+} + X\nabla c.$$

This gives:

$$\tilde{A}_{-} = \frac{1}{2c(1-a^{2})} \Big(c(X_{+} + aX_{-})\nabla c - 2(X_{-} + aX_{+})\nabla a + (1+c^{2}-a^{2})\gamma^{-} - 2(1-c^{2}+3a^{2})X\gamma^{-} \\ -(1-c^{2}-a^{2})a\gamma^{+} + 2a(3-c^{2}+a^{2})X\gamma^{+} + 4a(X_{-} - aX_{+})\gamma \Big), \\ \tilde{A}_{+} = \frac{1}{2c(1-a^{2})} \Big(c(X_{-} + aX_{+})\nabla c - 2(X_{+} + aX_{-})\nabla a + (1+c^{2}-a^{2})\gamma^{+} + 2(1-c^{2}+3a^{2})X\gamma^{+} \\ -(1-c^{2}-a^{2})a\gamma^{-} - 2a(3-c^{2}+a^{2})X\gamma^{-} + 4a(aX_{-} - X_{+})\gamma \Big), \\ \tilde{A} = \frac{1}{1-a^{2}} \Big(cX\nabla c - 2aX\nabla a + (1+a^{2})\gamma + (1-c^{2}+a^{2})(X_{+}\gamma^{+} - X_{-}\gamma^{-}) + 2a(X_{-}\gamma_{+} - X_{+}\gamma^{-}) \Big).$$

$$(85)$$

As a check of these results, let us note that in the metric case a = 0, c = 1 and the computed "internal" spin-coefficients coincide with the "spacetime" ones.

For practical computations, e.g. in the case of spherical symmetry, many of the quantities we have been discussing simplify, and the spinor method is an overkill. A different method of computing the spin-coefficients that avoids spinors and uses only exterior product of forms is given in the Appendix.

5 Field equations

In this section, using the results we have obtained above, we write down the field equations for theory (3). As we shall see, these become equations for the derivatives of the one-forms l, n, m, \bar{m} as well as for the curvature functions α, γ and their derivatives.

The field equations. The equations we are to discuss in this section are obtained by varying the action with respect to B^i . We get:

$$F^{i} + (\Lambda \delta^{ij} + \Psi^{ij} + \phi \delta^{ij})B^{j} = 8\pi G T^{i}, \tag{86}$$

where the right hand side is obtained by varying the matter part of the action with respect to B^i . Note that in the discussion above we have tacitly assumed that matter only couples to the B field and not to A, Ψ . This is true for the gauge fields, Maxwell field in particular, which will serve as a principal example of test matter in the subsequent papers of the series.

Spinor form. As before, it is useful to write these equations using spinors. We will make use of the gauge in which Ψ has the form (19) and *B* has the form (38). Let us first compute the matrix that appears in front of B^i in the second term of (86):

$$(\Lambda + \phi) \operatorname{Id} + \Psi = \alpha (\tilde{X}_{-} \otimes \tilde{X}_{-} + \tilde{X}_{+} \otimes \tilde{X}_{+}) +$$

$$(\beta + \Lambda + \phi) (\tilde{X}_{+} \otimes \tilde{X}_{-} + \tilde{X}_{-} \otimes \tilde{X}_{+}) + 2(2\beta - (\Lambda + \phi)) \tilde{X} \otimes \tilde{X}.$$
(87)

The second term on the left hand-side of (86) can now be easily computed:

$$((\Lambda + \phi)Id + \Psi)B = X_{-}(\alpha B_{+} + (\beta + \Lambda + \phi)B_{-}) + X_{+}(\alpha B_{-} + (\beta + \Lambda + \phi)B_{+}) + (\Lambda + \phi - 2\beta)XB.$$
(88)

To get the field equations it remains to split each of the 2-forms F^{AB} , T^{AB} into their self-dual and anti-self-dual parts. Thus, the field equations can be split as:

$$F' = 8\pi GT',\tag{89}$$

$$F + \ddot{X}_{-}(\alpha B_{+} + (\beta + \Lambda + \phi)B_{-}) + \ddot{X}_{+}(\alpha B_{-} + (\beta + \Lambda + \phi)B_{+}) + (\Lambda + \phi - 2\beta)\ddot{X}B = 8\pi GT.$$
(90)

We note that for conformally invariant matter (e.g. Yang-Mills fields) the self-dual part of the stress-energy 2-form vanishes (this is equivalent to the statement that the trace of the stress-energy tensor vanishes). We thus see that field equations for theory (3) have interpretation quite similar to that in usual GR. The curvature 2-form F^{AB} splits into two parts: "Weyl" part F and "Ricci" part F'. The first "Weyl" part gets related to the "curvature field" Ψ components α, γ , as well as to the trace part of the stress-energy tensor, via equation (90), while the second "Ricci" part is proportional to the traceless part T' of the stress-energy.

Thus, modifications due to non-metricity are not so dramatic after all. The main difference with the usual GR case is that the connection components (and thus curvature F components) depend not only on the frame l, n, m, \bar{m} and its derivatives but also on the other, non-metric components of B and their derivatives. The later can be expressed in terms of Ψ components via the metricity equations, and it is in this form that the field equations are most convenient for analysis. We thus get the schematic structure described by equation (1).

Curvature. The curvature of A is easy to compute. We have:

$$F_{-} = dA_{-} + A_{-} \wedge A, \qquad F_{+} = dA_{+} + A \wedge A_{+}, \qquad F = dA + 2A_{-} \wedge A_{+}. \tag{91}$$

One should now substitute (80) into these expressions. The result does not seem to be very illuminating, and we shall refrain from giving it here.

6 Bianchi identity

In this section we derive Bianchi identities for theory (3) and discuss the energy conservation. Spinor method is most effective for this purpose, and will be used heavily in this section.

"Bianchi" identity. A very important identity, analogous to the Bianchi identity in usual GR, is obtained by applying the operator of covariant derivative with respect to A to the equation (86). In view of $\mathcal{D}F^i = 0$ identically, we get

$$\mathcal{D}(\Lambda\delta^{ij} + \Psi^{ij} + \phi\delta^{ij})B^j = 8\pi G \mathcal{D}T^i,\tag{92}$$

In usual GR, contracting this equation in a certain way gets rid of the left-hand-side, and gives the equation of energy conservation. We would like to obtain an analog of this equation for theory (3).

Computation. To compute the covariant derivative of the quantity on the left-hand-side of (92) we need to know how \mathcal{D} acts on the basic rank 2 internal spinors $\tilde{X}_{\pm}, \tilde{X}$ as well as on the 2-forms B_{\pm}, B . The action of \mathcal{D} on tilded "internal" rank 2 spinors is given by:

$$\mathcal{D}\tilde{X}_{-} = -\tilde{A}\tilde{X}_{-} - 2\tilde{A}_{+}\tilde{X}, \qquad \mathcal{D}\tilde{X}_{+} = \tilde{A}\tilde{X}_{+} + 2\tilde{A}_{-}\tilde{X}, \qquad \mathcal{D}\tilde{X} = \tilde{A}_{-}\tilde{X}_{-} - \tilde{A}_{+}\tilde{X}_{+}, \tag{93}$$

which is easy to show using the formulae (82) for covariant derivatives of the basic spinors.

The action of \mathcal{D} on 2-forms B_{\pm} , B is that of the derivative operator ∇ compatible with the metric defined by l, n, m, \bar{m} . The most economic way to compute this action is using spinor representation. In this representation the self-dual 2-forms B_{\pm} , B become the matrices (71). The action of the operator ∇ on them can be obtained from relations (75) and (83). We have the following formulae:

$$(\nabla B_{-}) = B_{-}\tilde{A} - B\tilde{A}_{-}, \qquad (\nabla B_{+}) = B\tilde{A}_{+} - B_{+}\tilde{A}, \qquad (\nabla B) = 2(B_{-}\tilde{A}_{+} - B_{+}\tilde{A}_{-}),$$
(94)

where matrix multiplication on the right-hand-side is implied. Using these formulae, and passing into the spinor representation by replacing 3-forms of the type $A \wedge B$ by the spinor quantity BA (matrix multiplication implied) we get the following 3 components of the "Bianchi" identities:

$$B_{+}\nabla\alpha + B_{-}\nabla(\beta + \phi) + \alpha(BA_{+} - 2B_{+}A) - 3\beta BA_{-} = 8\pi G(\mathcal{D}T)_{-}, \qquad (95)$$

$$B_{-}\nabla\alpha + B_{+}\nabla(\beta + \phi) + \alpha(2B_{-}\tilde{A} - B\tilde{A}_{-}) + 3\beta B\tilde{A}_{+} = 8\pi G(\mathcal{D}T)_{+}, \qquad (95)$$

$$B\nabla(\phi - 2\beta) + 2\alpha(B_{-}\tilde{A}_{-} - B_{+}\tilde{A}_{+}) - 6\beta(B_{-}\tilde{A}_{+} - B_{+}\tilde{A}_{-}) = 8\pi G(\mathcal{D}T), \qquad (95)$$

where on the right hand side of these equations one finds the projections of the quantity $(\mathcal{D}T)^{AB}$ onto the 3 rank 2 spinors $\tilde{X}_{\pm}, \tilde{X}$.

Contracted "Bianchi" identity. Let us now multiply the first of the equations in (95) by B_+ , the second of them by B_- and add the results. We get:

$$(B_{+}B_{+} + B_{-}B_{-})\nabla\alpha + (B_{+}B_{-} + B_{-}B_{+})\nabla(\beta + \phi) + \alpha(B_{+}B\tilde{A}_{+} - B_{-}B\tilde{A}_{-}) - 3\beta(B_{+}B\tilde{A}_{-} - B_{-}B\tilde{A}_{+}) = (96)$$

$$8\pi G(B_{+}(\mathcal{D}T)_{-} + B_{-}(\mathcal{D}T)_{+}).$$

Let us multiply the third equation in (95) by B/2. We get:

$$(1/2)BB\nabla(\phi - 2\beta) + \alpha(BB_{-}\tilde{A}_{-} - BB_{+}\tilde{A}_{+}) - 3\beta(BB_{-}\tilde{A}_{+} - BB_{+}\tilde{A}_{-}) = 8\pi G((1/2)B(\mathcal{D}T)).$$
(97)

Now, subtracting this equation from (96), and using the fact that $BB_+ + B_+B = 0$, $BB_- + B_-B = 0$ we get:

$$(B_{+}B_{+} + B_{-}B_{-})\nabla\alpha + (B_{+}B_{-} + B_{-}B_{+})\nabla(\beta + \phi) - (1/2)BB\nabla(\phi - 2\beta) = 8\pi G(B_{+}(\mathcal{D}T)_{-} + B_{-}(\mathcal{D}T)_{+} - (1/2)B(\mathcal{D}T)).$$
(98)

It is now elementary to check that the quantity on the left-hand-side of this equation is zero. Indeed, using $B_+B_+ + B_-B_- = -2a$, $B_+B_- + B_-B_+ = -(1 + a^2)$, $BB = c^2$ we get for the left-hand-side of (98):

$$-2a\nabla\alpha - (1+a^2)\nabla(\beta+\phi) - (c^2/2)\nabla(\phi-2\beta) = -2a\nabla\alpha - (1+a^2-c^2)\nabla\beta - (1+a^2-c^2/2)\nabla\phi.$$
 (99)

However, this obviously vanishes in view of (39). Thus, we have established the following of "energy-momentum" conservation law for theory (3):

$$B_{+}(\mathcal{D}T)_{-} + B_{-}(\mathcal{D}T)_{+} - (1/2)B(\mathcal{D}T) = 0.$$
(100)

Using the fact that $\mathcal{D}B = 0$, we can rewrite this identity as:

$$\nabla BT = 0,\tag{101}$$

or, re-introducing all indices:

$$\nabla^{BB'} B^{CD}_{AB} T_{CD\,A'B'} = 0. \tag{102}$$

Thus, we see that the energy conservation law in this theory takes the usual form, with the stress-energy tensor that is conserved being $T_{ab} = (BT)_{ab}$. The 2-form *B* appears in the expression for the quantity T_{ab} as it is necessary to convert the pair *AB* of "internal" indices of $T_{A'B'}^{AB}$ into a pair of spacetime spinor indices. After this is done one gets a quantity with only spacetime indices, and it is this quantity that is conserved in the usual way.

Acknowledgements

It is a pleasure to thank Laurent Freidel, Andrei Mikhailov, Yuri Shtanov and Sergey Winitzki for stimulating discussions. The author was partially supported by an EPSRC advanced fellowship and by Perimeter Institute for Theoretical Physics.

Appendix: The "metricity" equations

The main novelty of the theory under consideration as compared to Plebański one is in the content of "metricity" equations that follow by varying the action with respect to the Lagrange multiplier field Ψ . As we have already discussed in the main text, these equations should be interpreted as determining the components of Ψ in terms of the components of the *B* field. For practical considerations, however, it is more convenient to interpret these equations as determining the 5 components of the *B* field (that *B* describes in addition to a metric) in terms of the Ψ field. For this interpretation one needs to introduce a metric. The most natural way to do this, also suggested by what happens in the case of Plebański gravity theory, is to declare the triple of two-forms B_{\pm}, B to be the self-dual two-forms with respect to some metric, which then determines the metric up to a conformal factor, in the way discussed in the main text.

In this appendix we prove that using the SL(2) freedom in choosing the basis l, n, m, \bar{m} , any solution of the equations (29) can be brought into the form (38).

As a preliminary, let us consider the question of how a general self-dual 2-form

$$C = c_1 m \wedge l + c_2 n \wedge \bar{m} + c_3 (l \wedge n - m \wedge \bar{m}) \tag{103}$$

can be simplified by rotating the basis l, n, m, \bar{m} . Thus, let us consider transformations that act on l, m and involve the other one-forms n, \bar{m} :

$$l \to l + \alpha n + \beta \bar{m}, \qquad m \to m + \gamma n + \delta \bar{m}.$$
 (104)

Simple analysis shows that there is a way to perform this transformation that no anti-self-dual part is introduced. This requires: $\alpha = \delta$, $\beta = \gamma = 0$. Under such a transformation the 2-form C changes to

$$C \to c_1 m \wedge l + (c_1 \alpha^2 + c_2 - 2\alpha c_3)n \wedge \bar{m} + (c_3 - c_1 \alpha)(l \wedge n - m \wedge \bar{m}).$$

$$(105)$$

Thus, choosing $\alpha = c_3/c_1$ we can eliminate the last term and (using an additional rescaling of the one-forms l, m) bring a general self-dual 2-form into the form:

$$C = m \wedge l + cn \wedge \bar{m},\tag{106}$$

We emphasise that in order to achieve this form only the transformations of the one-forms l, m were used. One can show by a similar argument that any 2-form C' can be brought into the form: $C' = n \wedge \overline{m} + c'm \wedge l$ by acting on the one-forms n, \overline{m} only.

Let us now consider three self-dual 2-forms B_{\pm} , B satisfying (29). We can use the available freedom in choosing the basis l, n, m, \bar{m} to bring these 3-forms into the desired form (38). To achieve this we first bring the self-dual 2-forms B_{\pm} into their "canonical" forms (as discussed above) by acting on the pairs l, m via $l \to l + \alpha n, m \to m + \alpha \bar{m}$ and on n, \bar{m} via $n \to n + \alpha' l, \bar{m} \to \bar{m} + \alpha' m$ correspondingly, as well as rescaling l, n, m, \bar{m} . This way we achieve

$$B_{+} = m \wedge l + a n \wedge \bar{m}, \qquad B_{-} = n \wedge \bar{m} + a' m \wedge l.$$
(107)

Importantly, under these transformations the self-dual 2-form B remains self-dual, as is clear from (105). Given the 2-forms (107) together with a general self-dual 2-form B it is easy to see that the metricity equations of the second line of (29) imply that B is of the form

$$B = c(l \wedge n - m \wedge \bar{m}). \tag{108}$$

The two equations of the first line of (29) imply, in particular, that a = a'. This finishes our proof of the validity of the representation (38) for B.

Appendix: Connection A without spinors

Sometimes it is more convenient to avoid spinors, and do computations directly with forms. Here we give the expressions for the spin-coefficients that are obtained this way. Thus, using the expression (38) for B and decomposing A into the basis of one-forms, we can easily compute the projections of C_{\pm}, C onto the basis one-forms. We have:

$$(C_{-} \wedge l) = cA_{-n} - aA_{m}, \qquad (C_{-} \wedge n) = -cA_{-l} + A_{\bar{m}}, \tag{109}$$
$$(C_{-} \wedge m) = -cA_{-\bar{m}} + aA_{l}, \qquad (C_{-} \wedge \bar{m}) = cA_{-m} - A_{n}.$$

$$(C_{+} \wedge l) = A_{m} - cA_{+n}, \qquad (C_{+} \wedge n) = -aA_{\bar{m}} + cA_{+l},$$
(110)
$$(C_{+} \wedge m) = -A_{l} + cA_{+\bar{m}}, \qquad (C_{+} \wedge \bar{m}) = aA_{n} - cA_{+m}.$$

$$(C \wedge l) = A_{-m} - aA_{+m}, \qquad (C \wedge n) = A_{+\bar{m}} - aA_{-\bar{m}}, \tag{111}$$
$$(C \wedge m) = -A_{-l} + aA_{+l}, \qquad (C \wedge \bar{m}) = -A_{+n} + aA_{-n}.$$

Assuming C_{\pm} , C are given, it is easy to write down the solution for A. We get:

$$A_{-l} = -\frac{1}{1-a^2} \left((C \wedge m) - \frac{a}{c} ((C_+ + aC_-) \wedge n) \right), \quad A_{-n} = -\frac{1}{1-a^2} \left(a(C \wedge \bar{m}) - \frac{1}{c} ((C_- + aC_+) \wedge l) \right), (112)$$
$$A_{-m} = \frac{1}{1-a^2} \left((C \wedge l) - \frac{a}{c} ((C_+ + aC_-) \wedge \bar{m}) \right), \quad A_{-\bar{m}} = \frac{1}{1-a^2} \left(a(C \wedge n) - \frac{1}{c} ((C_- + aC_+) \wedge m) \right).$$

$$A_{+l} = -\frac{1}{1-a^2} \left(a(C \wedge m) - \frac{1}{c} ((C_+ + aC_-) \wedge n) \right), \quad A_{+n} = -\frac{1}{1-a^2} \left((C \wedge \bar{m}) - \frac{a}{c} ((C_- + aC_+) \wedge l) \right), (113)$$
$$A_{+m} = \frac{1}{1-a^2} \left(a(C \wedge l) - \frac{1}{c} ((C_+ + aC_-) \wedge \bar{m}) \right), \quad A_{+\bar{m}} = \frac{1}{1-a^2} \left((C \wedge n) - \frac{a}{c} ((C_- + aC_+) \wedge m) \right).$$

$$A_{l} = \frac{1}{1 - a^{2}} \left(c(C \wedge n) - \left((C_{+} + aC_{-}) \wedge m \right) \right), \quad A_{n} = \frac{1}{1 - a^{2}} \left(c(C \wedge l) - \left((C_{-} + aC_{+}) \wedge \bar{m} \right) \right), \quad (114)$$
$$A_{m} = -\frac{1}{1 - a^{2}} \left(c(C \wedge \bar{m}) - \left((C_{+} + aC_{-}) \wedge l \right) \right), \quad A_{\bar{m}} = -\frac{1}{1 - a^{2}} \left(c(C \wedge m) - \left((C_{-} + aC_{+}) \wedge n \right) \right).$$

These relations should be compared to (79), (78).

To obtain the spin coefficients \tilde{A} we can use the following identities:

$$-(\tilde{C}_{+} + a\tilde{C}_{-}) = (1 + a^{2})d(n \wedge \bar{m}) + 2ad(m \wedge l) + da \wedge B_{-},$$

$$-(\tilde{C}_{-} + a\tilde{C}_{+}) = (1 + a^{2})d(m \wedge l) + 2ad(n \wedge \bar{m}) + da \wedge B_{+}.$$
(115)

This gives:

$$-((\tilde{C}_{+}+a\tilde{C}_{-})\wedge l) = (1+a^{2})((d\bar{m})_{m\bar{m}} - (dn)_{nm}) - 2a(dl)_{n\bar{m}} + a(da)_{m},$$
(116)

$$-((\tilde{C}_{+}+a\tilde{C}_{-})\wedge n) = -(1+a^{2})(dn)_{ml} + 2a((dm)_{m\bar{m}} + (dl)_{l\bar{m}}) - (da)_{\bar{m}},$$

$$-((\tilde{C}_{+}+a\tilde{C}_{-})\wedge m) = (1+a^{2})(-(d\bar{m})_{l\bar{m}} - (dn)_{ln}) - 2a(dm)_{n\bar{m}} - a(da)_{l},$$

$$-((\tilde{C}_{+}+a\tilde{C}_{-})\wedge \bar{m}) = -(1+a^{2})(d\bar{m})_{ml} + 2a((dm)_{nm} - (dl)_{ln}) + (da)_{n}.$$

$$-((\tilde{C}_{-}+a\tilde{C}_{+})\wedge l) = -(1+a^{2})(dl)_{n\bar{m}} + 2a(-(dn)_{nm} + (d\bar{m})_{m\bar{m}}) + (da)_{m},$$
(117)

$$-((\tilde{C}_{-}+a\tilde{C}_{+})\wedge n) = (1+a^{2})((dm)_{m\bar{m}} - (dl)_{n\bar{m}}) - 2a(dn)_{ml} - a(da)_{\bar{m}},$$

$$-((\tilde{C}_{-}+a\tilde{C}_{+})\wedge m) = -(1+a^{2})(dm)_{n\bar{m}} + 2a(-(dn)_{ln} - (d\bar{m})_{l\bar{m}}) - (da)_{l},$$

$$-((\tilde{C}_{-}+a\tilde{C}_{+})\wedge \bar{m}) = (1+a^{2})((dm)_{nm} - (dl)_{ln}) - 2a(d\bar{m})_{ml} + a(da)_{n}.$$

The notation that we used here is almost self-explanatory. Thus, e.g. $dm = (dm)_{nm}n \wedge m + \ldots$, where the dots stand for the other terms. Finally, we get for the projections of the 3-form \tilde{C} :

$$-2(\tilde{C} \wedge l) = -c(dl)_{m\bar{m}} + c(dm)_{nm} + c(d\bar{m})_{n\bar{m}} + (dc)_n,$$
(118)

$$-2(\tilde{C} \wedge n) = -c(dn)_{m\bar{m}} + c(dm)_{ml} - c(d\bar{m})_{l\bar{m}} - (dc)_l,$$

$$-2(\tilde{C} \wedge m) = c(dl)_{l\bar{m}} + c(dn)_{n\bar{m}} + c(dm)_{ln} - (dc)_{\bar{m}},$$

$$-2(\tilde{C} \wedge \bar{m}) = c(dl)_{ml} - c(dn)_{nm} + c(d\bar{m})_{ln} + (dc)_m.$$

References

- [1] R. H. Dicke, "The theoretical significance of experimental relativity," Gordon and Breach, New York, 1964.
- [2] J. D. Bekenstein, "Relativistic gravitation theory for the MOND paradigm," Phys. Rev. D 70, 083509 (2004) [Erratum-ibid. D 71, 069901 (2005)] [arXiv:astro-ph/0403694].
- [3] J. W. Moffat, "Nonsymmetric gravitational theory," J. Math. Phys. 36, 3722 (1995) [Erratum-ibid. 36, 7128 (1995)].
- M. A. Clayton and J. W. Moffat, "Scalar-Tensor Gravity Theory For Dynamical Light Velocity," Phys. Lett. B 477, 269 (2000) [arXiv:gr-qc/9910112].
- [5] G. R. Dvali, G. Gabadadze and M. Porrati, "4D gravity on a brane in 5D Minkowski space," Phys. Lett. B 485, 208 (2000) [arXiv:hep-th/0005016].
- [6] Yu. V. Shtanov, "On brane-world cosmology," arXiv:hep-th/0005193.
- [7] R. Punzi, F. P. Schuller and M. N. R. Wohlfarth, "Area metric gravity and accelerating cosmology," JHEP 0702, 030 (2007) [arXiv:hep-th/0612141].

- [8] T. Biswas, R. Brandenberger, A. Mazumdar and W. Siegel, "Non-perturbative gravity, Hagedorn bounce and CMB," arXiv:hep-th/0610274.
- [9] P. D. Mannheim, "Linear Potentials And Galactic Rotation Curves," Astrophys. J. 419, 150 (1993) [arXiv:hep-ph/9212304].
- [10] S. M. Carroll, V. Duvvuri, M. Trodden and M. S. Turner, Phys. Rev. D 70, 043528 (2004) [arXiv:astro-ph/0306438].
- [11] J. Plebański, "On the separation of Einsteinian substructures," Journ. Math. Phys. 18, 2511 (1977).
- [12] I. Bengtsson, arXiv:gr-qc/0703114.
- [13] K. Krasnov, "Renormalizable non-metric quantum gravity?," arXiv:hep-th/0611182.
- [14] S. Weinberg, "Ultraviolet divergence in quantum theories of gravitation," in Hawking, S.W., Israel, W.: General Relativity, 790-831.
- [15] R. Capovilla, J. Dell, T. Jacobson, L. Mason, "Self-dual 2-forms and gravity", Class. Quant. Grav. 8, 41 (1991).
- [16] R. Capovilla, T. Jacobson and J. Dell, "A Pure spin connection formulation of gravity," Class. Quant. Grav. 8, 59 (1991).
- [17] P. Peldan, "Actions for gravity, with generalizations: A Review," Class. Quant. Grav. 11, 1087 (1994) [arXiv:gr-qc/9305011].
- [18] R. Penrose and W. Rindler, "Spinors and space-time," Cambridge University Press, 1984.
- [19] H. Urbantke, J. Math. Phys. 25, 2321 (1983).