

Conserving Integrals in Metric Theories of Gravitation

Dimitry Palatnik *

January 2009

Abstract

In this work I present location to first mentioning to result of Robert Geroch(?) Preprint deals with conserving quantities of metric gravitational theories They are constructed from Killing vector fields (if any exists) and symmetric tensors of arbitrary rank with vanishing divergence I also suggest alternative approach by introducing spinorial fields allowing to construct conserved integrals of energy-momentum etc even if timelike Killing vector doesn't necessarily exists

1 Conservation of energy-momentum and angular momenta

First off result presented in this section (ie formula (5) below also see [6]) was reported on seminar in EFI at University of Chicago and belongs probably to Robert Geroch

Problem of conserving integrals in general relativity is as old as theory itself and one should recognise effort put by people in dealing with it Most famous attempt is that of Landau and Lifshits [1] who constructed symmetric pseudo-tensor describing energy-momentum of gravitational field One should acknowledge works of Tolman [2] Papapetrou [3] Bergmann [4] Møller [5] and Weinberg [7] Details of that story one may find in [9]

Consider n -dimensional metric manifold (M, g_{ab}) I work with class of theories specified by action

$$S = -k^{-1} \int d\Omega \sqrt{|g|} R + \int d\Omega \sqrt{|g|} \Lambda [g_{ab}; \psi] \quad (1)$$

Here $d\Omega$ and R are volume element and scalar curvature respectively g is metric determinant and Λ is lagrangian density of matter depending on metrics and other fields specified for short by ψ If T_{ab} is stress-energy corresponding to Λ then

$$T^{ab} = T^{(ab)}; \quad (2)$$

$$\nabla_a T^{ab} = 0 \quad (3)$$

Let ξ_c be Killing vector field satisfying equations

$$\nabla_c \xi_b + \nabla_b \xi_c = 0 \quad (4)$$

If solution to (4) exists then from (2)-(4) follows

$$\nabla_a \{T^{ab} \xi_b\} = \xi_b \nabla_a T^{ab} + T^{ab} \nabla_{(a} \xi_{b)} = 0 \quad (5)$$

Then integrating (5) over n -dimensional volume and using respective Gauss theorem with appropriate boundary of integration one obtains conserving integral

$$Q = \int d\Sigma_a \sqrt{|g|} T^{ab} \xi_b \quad (6)$$

Here $d\Sigma_a$ is element of integration over $(n-1)$ -dimensional hypersurface Consider limit of weak gravitational field $k \rightarrow 0$ In flat spacetime and Cartesian coordinates (x^a) solution of (4) is

$$\xi_c = p_c + \omega_{cb} x^b \quad (7)$$

*The Waterford 07445 N Sheridan Chicago IL 60626-1818 USA

Here n components of p_c and $n(n-1)/2$ components of $\omega_{cb} = \omega_{[cb]}$ are arbitrary Substituting (7) into (6) one obtains

$$Q = p_c P^c + \omega_{cb} M^{cb} \quad (8)$$

Here

$$P^c = \int d\Sigma_b T^{cb}; \quad (9)$$

$$M^{cb} = \int d\Sigma_e \{T^{ec} x^b - T^{eb} x^c\} \quad (10)$$

Arbitrariness of coefficients p_c and ω_{cb} in (8) causes conservation of integrals (9)–(10) individually In 4-dimensional spacetime one may select $d\Sigma_a = (dV, 0, 0, 0)$ where dV is 3-dimensional element of integration over space and recognise (9) as energy-momentum and (10) as angular momenta (including boosts) What if solution of (4) for say timelike Killing vector doesn't exist? Well then respective integral corresponding to energy doesn't exist too Situation is similar to one taking place in quantum mechanics where existence of energy (conservation of hamiltonian) depends on existence of translational symmetry of system in time

2 Generalization on tensors of arbitrary rank

Suppose Killing equations have q independent vectors as solutions $\xi_c^{(n)}$ where $n = 1, 2, \dots, q$ I may consider two separate cases 1 Let $T^{cc_1 c_2 \dots c_p}$ be symmetric rank $(p+1)$ tensor with vanishing divergence

$$T^{cc_1 c_2 \dots c_p} = T^{(cc_1 c_2 \dots c_p)}; \quad (11)$$

$$\nabla_c T^{cc_1 c_2 \dots c_p} = 0 \quad (12)$$

Construct vector

$$T^{c\{m_1 m_2 \dots m_p\}} = T^{cc_1 \dots c_p} \xi_{c_1}^{(m_1)} \xi_{c_2}^{(m_2)} \dots \xi_{c_p}^{(m_p)} \quad (13)$$

then it follows from Killing equations (4) and from (11) (12) that divergence of $T^{c\{m_1 m_2 \dots m_p\}}$ vanishes

$$\nabla_c T^{c\{m_1 m_2 \dots m_p\}} = 0 \quad (14)$$

and in analogy with (6) conserving integrals are

$$Q^{\{m_1 m_2 \dots m_p\}} = \int d\Sigma_c \sqrt{|g|} T^{cc_1 \dots c_p} \xi_{c_1}^{(m_1)} \xi_{c_2}^{(m_2)} \dots \xi_{c_p}^{(m_p)} \quad (15)$$

Here $Q^{\{m_1 m_2 \dots m_p\}} = Q^{\{(m_1 m_2 \dots m_p)\}}$ where $m_1, m_2, \dots, m_p = 1, 2, \dots, q$ so one obtains $N = \frac{(p+q-1)!}{p!(q-1)!}$ conserving integrals for that case Here N is number of components of symmetric object of rank p with each index running q values¹ 2 Take symmetric tensor of rank $(p+1)$ with vanishing divergence $T^{cc_1 c_2 \dots c_p}$ so that formulae (11) (12) still hold but assume now that it traceless

$$g_{c_1 c_2} T^{cc_1 c_2 \dots c_p} = 0 \quad (16)$$

Consider set of solutions $\{\chi_c^{(l)}\}$ $l = 1, 2, \dots, r$ to second type of Killing equations

$$\nabla_b \chi_c^{(l)} + \nabla_c \chi_b^{(l)} = 2g_{cb} \quad (17)$$

I should use notation for combined set of vectors $\{Y_c^B\} = \{\xi_c^{(n)}, \chi_c^{(l)}\}$ $B = 1, 2, \dots, q+r$ Then analogously to previous pattern I construct conserving (according to (4) (11) (12) (16) and (17)) currents $J^{c\{B_1 B_2 \dots B_p\}}$

$$\nabla_c J^{c\{B_1 B_2 \dots B_p\}} = 0 \quad (18)$$

Here

$$J^{c\{B_1 B_2 \dots B_p\}} = T^{cc_1 c_2 \dots c_p} Y_{c_1}^{B_1} Y_{c_2}^{B_2} \dots Y_{c_p}^{B_p} \quad (19)$$

Respective conserving integrals are

$$Q^{\{B_1 B_2 \dots B_p\}} = \int d\Sigma_c \sqrt{|g|} T^{cc_1 c_2 \dots c_p} Y_{c_1}^{B_1} Y_{c_2}^{B_2} \dots Y_{c_p}^{B_p} \quad (20)$$

¹ I wish to thank Boris Tsirelson for updating me on the formula for N

3 Examples

1 Introduce first example in order to test vector χ_c Consider 4-dimensional flat spacetime and take Cartesian coordinates $x^a = (ct, x^\alpha)$ Metric is $g_{cb} = \eta_{cb} = \text{diag}(1, -1, -1, -1)$ Take stress-energy of electromagnetic field with no charges $T^{cb} = \frac{1}{4\pi} \{-F_e^c F^{eb} + \frac{1}{4} g^{cb} F_{ef} F^{ef}\}$ which is traceless symmetric and divergence free I found from (17) solution for χ_c

$$\chi_c = \eta_{cb} x^b \quad (21)$$

Then taking $d\Sigma_c = (dV, 0, 0, 0)$ one obtains conserving integral

$$ct \int dV T_0^0 + \int dV T_\alpha^0 x^\alpha = \text{constant} \quad (22)$$

Equation (22) describes propagation of electromagnetic radiation (Static cases which refer to configurations with non-zero charges are not valid since charges contribute their own stress-energy with nonvanishing trace)

2 Second example is Bel-Robinson tensor [6] [8] which in vacuum ($R_{cb} = 0$) in 4-dimensional spacetime is

$$T^{cbag} = R_e^c{}^a R^{begf} + \tilde{R}_e^c{}^a \tilde{R}^{begf} \quad (23)$$

where

$$\tilde{R}_{aecf} = \frac{1}{2} e_{aebd} R^{bd}{}_{cf} \quad (24)$$

Here e_{abce} is absolute skew unit tensor Bel-Robinson (23) is symmetric traceless and divergence free (non verified by me)

$$T^{cbag} = T^{(cbag)}; \quad (25)$$

$$g_{cb} T^{cbag} = 0; \quad (26)$$

$$\nabla_c T^{cbag} = 0 \quad (27)$$

Using formula (20) one obtains conserving integrals

$$Q^{\{C_1 C_2 C_3\}} = \int d\Sigma_c \sqrt{-g} T^{cbag} Y_b^{C_1} Y_a^{C_2} Y_g^{C_3} \quad (28)$$

3 Third example is Lanczos-Lovelock tensor $S^{ab} = 2\tau^{ab}$ which exists if dimension of spacetime $n \neq 4$

$$\tau^{ab} = R_{cde}^a R^{bcde} - 2R_{ce} R^{caeb} - 2R_c^a R^{bc} + R R^{ab} - \frac{1}{4} g^{ab} (R_{cdef} R^{cdef} - 4R_{ce} R^{ce} + R^2) \quad (29)$$

Tensor (29) is symmetric and divergence free so conserving integrals are

$$Q^{\{m\}} = \int d\Sigma_c \sqrt{|g|} \tau^{ce} \xi_e^{(m)} \quad (30)$$

4 As fourth example I suggest another conserving integral for 4-dimensional empty spacetime ($R_{cb} = 0$) Let ξ_c be Killing vector field First introduce twist

$$\omega_a = e_{abcd} \xi^b \nabla^c \xi^d \quad (31)$$

As it's known [6] twist satisfies equations

$$\nabla_{[c} \omega_{b]} = 0; \quad (32)$$

$$\omega_c \xi^c = 0 \quad (33)$$

Construct vector

$$R^a = R^{abcd} \omega_b \nabla_c \xi_d \quad (34)$$

I intend to prove that

$$\nabla_c R^c = 0 \quad (35)$$

Really

$$\nabla_a R^a = (\nabla_a R^{abcd}) \omega_b (\nabla_c \xi_d) + R^{abcd} (\nabla_{[a} \omega_{b]}) (\nabla_c \xi_d) + R^{abcd} \omega_b (\nabla_a \nabla_c \xi_d) \quad (36)$$

First term on rhs of (36) vanishes due to Bianchi identity

$$\nabla_a R^{abcd} = \nabla^c R^{bd} - \nabla^d R^{bc} \quad (37)$$

second term vanishes due to (32) and for third term I use known [6] equation

$$\nabla_a \nabla_b \xi_c = -R_{bcad} \xi^d \quad (38)$$

so it follows

$$\begin{aligned} R^{abcd} \omega_b (\nabla_a \nabla_c \xi_d) &= -R^{abcd} R_{aecd} \xi^e \omega_b \\ &= -\frac{1}{4} R^{efgh} R_{efgh} \xi^c \omega_c \\ &= 0 \end{aligned} \quad (39)$$

In second line of (39) I used (non verified by me) fact that Lanczos-Lovelock tensor (29) vanishes identically in 4-dimensional spacetime and third line follows from equation (33) This completes proof of (35) Thus conserving integral is

$$Q = \int d\Sigma_a \sqrt{-g} R^{abcd} \omega_b \nabla_c \xi_d \quad (40)$$

4 Alternative approach: spinorial fields

Problem with approach suggested above is that Killing vector exists only for very restricted class of solutions of Einstein equations Aim of this section is to demonstrate how this problem can be solved by introducing (physical and classical) spinorial fields for case of 4-dimensional spacetime Denote Einstein tensor

$$G^{cb} = \frac{c^4}{8\pi k} \{R^{cb} - \frac{1}{2} g^{cb} R\} \quad (41)$$

Here c is speed of light and k is Newton constant of gravitational interaction Dimensional coefficient in (41) is introduced for simplicity For reference I write

$$\nabla_c G^{cb} = 0 \quad (42)$$

Einstein equations are

$$G^{cb} = T^{cb} \quad (43)$$

here T^{cb} is stress-energy in standard units I introduce (complex) spinorial fields $\bar{\xi}$ and ξ By standard definition

$$\nabla_c \xi = \partial_c \xi - \Gamma_c \xi ; \quad (44)$$

$$(\bar{\xi} \nabla_c) = \partial_c \bar{\xi} + \bar{\xi} \Gamma_c \quad (45)$$

For covariantly constant Dirac matrices γ_c defined in standard way

$$\gamma_c \gamma_e + \gamma_e \gamma_c = 2g_{ec}$$

one obtains

$$\nabla_c \gamma_e = \partial_c \gamma_e - \Gamma_{ec}^b \gamma_b - \Gamma_c \gamma_e + \gamma_e \Gamma_c = 0 \quad (46)$$

I take action of spinorial fields as

$$S_\xi = \frac{\ell}{2c} \Re \int d\Omega \sqrt{-g} G^{ce} \{ \bar{\xi} \gamma_c \nabla_e \xi - (\bar{\xi} \nabla_e) \gamma_c \xi \} \quad (47)$$

Here ℓ is characteristic length considered to be small parameter Respective stress-energy

$$\begin{aligned} T_\xi^{(ab)} &= K \{ \nabla_c \nabla^c H^{ab} + g^{ab} \nabla_c \nabla_e H^{ce} - 2 \nabla_c \nabla^{(a} H^{b)c} \} \\ &\quad + 2\ell \{ 2G_c^{(a} H^{b)c} - \frac{1}{2} G^{ab} H - \frac{1}{2} G H^{ab} - \frac{1}{2} g^{ab} (G^{ce} H_{ce} - \frac{1}{2} G H) \} \\ &\quad + \frac{\ell}{2} \Re \nabla_c \{ G^{ab} \bar{\xi} \gamma^c \xi - G^{c(a} \bar{\xi} \gamma^{b)} \xi + \frac{1}{2} G^{ea} \bar{\xi} \gamma^{[b} \gamma^{c]} \gamma_e \xi + \frac{1}{2} G^{eb} \bar{\xi} \gamma^{[a} \gamma^{c]} \gamma_e \xi \} \\ &\quad - \frac{\ell}{2} G^{c(a} \Re \{ \bar{\xi} \gamma^{b)} \nabla_c \xi - (\bar{\xi} \nabla_c) \gamma^{b)} \xi \} ; \end{aligned} \quad (48)$$

$$T_\xi^{[ab]} = -\frac{\ell}{2} G^{c[a} \Re \{ \bar{\xi} \gamma^{b]} \nabla_c \xi - (\bar{\xi} \nabla_c) \gamma^{b]} \xi \} - \frac{\ell}{2} \Re \nabla_e \{ G^{e[a} \bar{\xi} \gamma^{b]} \xi + \frac{1}{2} G^{ce} \bar{\xi} \gamma^{[a} \gamma^{b]} \gamma_c \xi \} \quad (49)$$

Here $K = \frac{\ell c^4}{8\pi k}$ and

$$H_{ab} = \frac{1}{2} \Re \{ \bar{\xi} \gamma_{(a} \nabla_{b)} \xi - (\bar{\xi} \nabla_{(b} \gamma_{a)} \xi - \frac{1}{2} g_{ab} [\bar{\xi} \gamma^c \nabla_c \xi - (\bar{\xi} \nabla_c) \gamma^c \xi] \} \quad (50)$$

Here $G = G_c^c$ and $H = H_c^c$ Stress-energy is obtained by varying action in components of tetrad vector specified below Since spinor fields are introduced one should use tetrad formalism for solving Einstein equations I use following formulae

$$\gamma_c = (e_c)^C \Delta_C ; \quad (51)$$

$$\Gamma_c = \frac{1}{4} (e_b)^B \nabla_c (e^b)^A \Delta_{[A} \Delta_{B]} \quad (52)$$

Here $(e_c)^E$ $E = 0...3$ is tetrad and Δ_E are standard constant Dirac matrices

Suppose lagrangian density of other fields (with respective action S_f) depends on tensor

$$c_{cb...f} = \Re \bar{\xi} C_{cb...f} \xi \quad (53)$$

where $C_{cb...f}$ is algebraic operator ie it doesn't contain derivative operators Eg $c_{cb...f} = 1; \gamma_c$ etc Denote $M^{cb...f}$ covariant analog of functional derivative of S_f in $c_{cb...f}$

$$\delta S_f = \frac{\ell}{c} \int d\Omega \sqrt{-g} M^{cb...f} \delta c_{cb...f} \quad (54)$$

Varying action of all fields in rhs of Einstein equation $S_m = S_\xi + S_f$ one obtains equations for spinorial fields

$$G^{cb} \gamma_c \nabla_b \xi + M^{cb...f} C_{cb...f} \xi = 0 ; \quad (55)$$

$$G^{cb} (\bar{\xi} \nabla_b) \gamma_c - M^{cb...f} \bar{\xi} C_{cb...f} = 0 \quad (56)$$

Introduce vector

$$J^c = G^{ce} \bar{\xi} \gamma_e \xi \quad (57)$$

One may write

$$\nabla_c J^c = (\nabla_c G^{ce}) \bar{\xi} \gamma_e \xi + G^{ce} (\bar{\xi} \nabla_c) \gamma_e \xi + G^{ce} \bar{\xi} (\nabla_c \gamma_e) \xi + G^{ce} \bar{\xi} \gamma_e \nabla_c \xi = 0 \quad (58)$$

Conservation of current J^c follows from (42) (46) (55) and (56) Integrating (58) over 4-dimensional volume with appropriate boundary and using Gauss theorem one obtains (generally complex) conserving integral

$$E = \int d\Sigma_c \sqrt{-g} T^{ce} \bar{\xi} \gamma_e \xi \quad (59)$$

In (59) Einstein tensor (41) is substituted by stress-energy T^{ce} I believe that theory works only if E is real which should be done by proper selection of spinors in limit $k \rightarrow 0$ Eg if $\bar{\xi} \gamma_c \xi$ corresponds to timelike Killing vector responsible for energy conservation then say in asymptotically flat spacetime it should have limiting value $(1, 0, 0, 0)$

Introduce additional 'epsilon-term' as action S_f

$$S_f = -\frac{\ell^2 \epsilon^2}{c} \int d\Omega \sqrt{-g} \Re \{ \bar{\xi} \xi \} \quad (60)$$

Respective stress-energy is

$$T_\epsilon^{cb} = \ell^2 \epsilon^2 g^{cb} \Re \{ \bar{\xi} \xi \} \quad (61)$$

Here $[\epsilon^2] = g \cdot cm^{-3} \cdot sec^{-2}$ Complete set of equations is

$$G^{cb} \gamma_c \nabla_b \xi - \ell \epsilon^2 \xi = 0 ; \quad (62)$$

$$G^{cb} (\bar{\xi} \nabla_b) \gamma_c + \ell \epsilon^2 \bar{\xi} = 0 ; \quad (63)$$

$$G^{cb} - T_\epsilon^{(cb)} - T_\epsilon^{cb} = 0 ; \quad (64)$$

$$T_\epsilon^{[cb]} = 0 \quad (65)$$

One should treat extra 6 equations (65) because in tetrad formalism applied here one has 16 components of $(e_c)^C$ in stead of 10 components of g_{cb} Note that Palatini method shouldn't be used because equations (42) (46) don't hold for that case

5 Flat spacetime case

I intend to solve equations (62)–(65) Expand all fields in powers of ϵ and ℓ and in 0-order approximation consider flat spacetime and use Cartesian coordinates As it is shown below $\tau^{cb} \propto \epsilon$ where τ^{cb} is defined in (66) so that

$$G^{cb} = T_{\xi}^{(cb)} + T_{\epsilon}^{cb} = K\tau^{cb} + O(\epsilon\ell^2)$$

where

$$\tau^{ab} = \nabla_c \nabla^c H^{ab} + g^{ab} \nabla_c \nabla_e H^{ce} - 2\nabla_c \nabla^{(a} H^{b)c} \quad (66)$$

In 0-order approximation take standard Lorentzian metric $g_{cb} = \text{diag}(1, -1, -1, -1)$ By inspecting stress-energy (48) (49) one may rewrite equations for spinors as

$$\tau^{cb} \gamma_c \nabla_b \xi - \frac{8\pi k \epsilon^2}{c^4} \xi = O(\ell\epsilon); \quad (67)$$

$$\tau^{cb} (\bar{\xi} \nabla_b) \gamma_c + \frac{8\pi k \epsilon^2}{c^4} \bar{\xi} = O(\ell\epsilon) \quad (68)$$

Search for solutions for spinors as expansions

$$\xi = \sum_{n=0}^{\infty} \epsilon^n \cdot \xi_n; \quad \bar{\xi} = \sum_{n=0}^{\infty} \epsilon^n \cdot \bar{\xi}_n$$

where

$$\xi_0 = a^0 + b_c^0 x^c; \quad (69)$$

$$\xi_1 = a^1 + b_c^1 x^c + \frac{1}{2} c_{cb}^1 x^c x^b; \quad (70)$$

$$\xi_2 = a^2 + b_c^2 x^c + \frac{1}{2} c_{cb}^2 x^c x^b + \frac{1}{3!} f_{cbe}^2 x^c x^b x^e; (etc) \quad (71)$$

similarly

$$\bar{\xi}_0 = \bar{a}^0 + \bar{b}_c^0 x^c; \quad (72)$$

$$\bar{\xi}_1 = \bar{a}^1 + \bar{b}_c^1 x^c + \frac{1}{2} \bar{c}_{cb}^1 x^c x^b; \quad (73)$$

$$\bar{\xi}_2 = \bar{a}^2 + \bar{b}_c^2 x^c + \frac{1}{2} \bar{c}_{cb}^2 x^c x^b + \frac{1}{3!} \bar{f}_{cbe}^2 x^c x^b x^e; (etc) \quad (74)$$

Here $a^0 \dots \bar{f}_{cbe}^2 \dots$ are constant spinors with complex components Doing calculation I obtained

$$\tau^{cb} = \epsilon \cdot \tau_1^{cb} + \epsilon^2 \cdot \tau_2^{cb} + \dots$$

where $\tau_1^{cb}(\bar{\xi}_0; \bar{\xi}_1; \xi_0; \xi_1)$ (constant in x^c) $\tau_2^{cb}(\bar{\xi}_0 \dots \bar{\xi}_2; \xi_0 \dots \xi_2)$ (first order polynomial in x^c) $\tau_3^{cb}(\bar{\xi}_0 \dots \bar{\xi}_3; \xi_0 \dots \xi_3)$ (second order polynomial in x^c) etc One may rewrite equations (67) and (68) as

$$\tau_1^{cb} \gamma_c \nabla_b \xi_0 = 0; \quad (75)$$

(0-order polynomial in x^c)

$$\tau_1^{cb} \gamma_c \nabla_b \xi_1 + \tau_2^{cb} \gamma_c \nabla_b \xi_0 - \frac{8\pi k}{c^4} \xi_0 = 0; \quad (76)$$

(first order polynomial in x^c)

$$\tau_1^{cb} \gamma_c \nabla_b \xi_2 + \tau_2^{cb} \gamma_c \nabla_b \xi_1 + \tau_3^{cb} \gamma_c \nabla_b \xi_0 - \frac{8\pi k}{c^4} \xi_1 = 0; \quad (77)$$

(second order polynomial in x^c etc)

$$\tau_1^{cb} (\bar{\xi}_0 \nabla_b) \gamma_c = 0; \quad (78)$$

(0-order polynomial in x^c)

$$\tau_1^{cb} (\bar{\xi}_1 \nabla_b) \gamma_c + \tau_2^{cb} (\bar{\xi}_0 \nabla_b) \gamma_c + \frac{8\pi k}{c^4} \bar{\xi}_0 = 0; \quad (79)$$

(first order polynomial in x^c)

$$\tau_1^{cb}(\bar{\xi}_2 \nabla_b) \gamma_c + \tau_2^{cb}(\bar{\xi}_1 \nabla_b) \gamma_c + \tau_3^{cb}(\bar{\xi}_0 \nabla_b) \gamma_c + \frac{8\pi k}{c^4} \bar{\xi}_1 = 0 ; \quad (80)$$

(second order polynomial in x^c etc)

Equations (75) (78) may be solved if

$$g_{cb} \tau_1^{cb} = 2\Re\{\bar{b}_c^0 \gamma^c c_b^{1b} - \bar{c}_b^{1b} \gamma^c b_c^0 + \bar{c}_{cb}^1 \gamma^c b^{0b} - \bar{b}^{0b} \gamma^c c_{cb}^1\} ; \quad (81)$$

select

$$b_c^0 = \gamma_c c ; \quad \bar{b}_c^0 = \gamma_c \bar{c} \quad (82)$$

where c , $\bar{c} = c^* \gamma_0$ are constant complex spinors and c^* has complex conjugate coefficients relatively to c
Substituting (82) into (81) one obtains

$$g_{cb} \tau_1^{cb} = 6\Re\{\bar{c} c_e^{1e} - \bar{c}_e^{1e} c\} \quad (83)$$

Equation (83) satisfies identically if

$$\bar{c}_e^{1e} = (c_e^{1e})^* \gamma_0$$

where $(c_e^{1e})^*$ has complex conjugate coefficients relatively to c_e^{1e} In limit $\epsilon \rightarrow 0$

$$\bar{\xi} \gamma_a \xi = \bar{\xi}_0 \gamma_a \xi_0 = p_a + \Omega_{ab} x^b + \omega g_{ab} x^b + 2\lambda_c g_{ab} x^c x^b - \lambda_a g_{cb} x^c x^b \quad (84)$$

where

$$p_c = \bar{a}^0 \gamma_c a^0 ; \quad \Omega_{ab} = \bar{a}^0 \gamma_{[a} \gamma_{b]} c - \bar{c} \gamma_{[a} \gamma_{b]} a^0 ; \quad \omega = \bar{c} a^0 + \bar{a}^0 c ; \quad \lambda_b = \bar{c} \gamma_b c$$

One obtains here $4 + 6 + 1 + 4 = 15$ independent real coefficients p_a , $\Omega_{cb} = \Omega_{[cb]}$, ω , λ_c From (59) (84) etc it follows (if to neglect terms of order ϵ^2) conservation of integrals ($T^{cb} = K \epsilon \tau_1^{cb}$)

$$P^a = \int dV T^{0a} ; \quad (85)$$

$$M^{ab} = \int dV \{T^{0a} x^b - T^{0b} x^a\} ; \quad (86)$$

$$Q = \int dV T_c^0 x^c ; \quad (87)$$

$$S^a = \int dV \{T_b^0 x^b x^a - \frac{1}{2} T^{0a} x_b x^b\} \quad (88)$$

Integrals (85) and (86) correspond to energy-momentum and angular momenta Integrals (87) and (88) conserve only if $T_c^c = 0$ Really introduce currents

$$\tilde{J}^c = T_b^c x^b ; \quad (89)$$

$$J^{ec} = T^{eb} x_b x^c - \frac{1}{2} T^{ec} x_b x^b \quad (90)$$

One obtains (due to $\nabla_c T^{cb} = 0$)

$$\nabla_c \tilde{J}^c = T_c^c = 0 ; \quad (91)$$

$$\nabla_c J^{cb} = T_c^c x^b = 0 \quad (92)$$

Integrating last two equations over 4-dimensial spacetime (with adequate boundary) and using Gauss theorem one obtains conservation of integrals (87) and (88) Approach suggested in that section makes sense in limit ℓ and $\epsilon \rightarrow 0$

6 Static spherically symmetric case

I intend to solve equations (62)–(65) anew Expand all fields in powers of ϵ and ℓ

$$\xi = \sum_{n=0}^{\infty} \epsilon^n \cdot \xi_n ; \quad \bar{\xi} = \sum_{n=0}^{\infty} \epsilon^n \cdot \bar{\xi}_n$$

and in 0-order approximation consider Ricci-flat spacetime and use spherical coordinates One may show (as in case of flat spacetime)

$$\tau^{cb} = \epsilon \cdot \tau_1^{cb} + \epsilon^2 \cdot \tau_2^{cb} + \dots$$

where τ^{cb} is defined in (66) As above

$$G^{cb} = T_{\xi}^{(cb)} + T_{\epsilon}^{cb} = K\tau^{cb} + O(\epsilon\ell^2)$$

One might take standard Schwarzschild metric I prefer though to use metric in isotropic coordinates [6] Equations for spinorial fields are

$$\tau^{cb}\gamma_c\nabla_b\xi - \frac{8\pi k\epsilon^2}{c^4}\xi = O(\ell\epsilon) ; \quad (93)$$

$$\tau^{cb}(\bar{\xi}\nabla_b)\gamma_c + \frac{8\pi k\epsilon^2}{c^4}\bar{\xi} = O(\ell\epsilon) \quad (94)$$

Interval is

$$ds^2 = \left(\frac{r-r_o}{r+r_o}\right)^2 dt^2 - \left(\frac{r+r_o}{r}\right)^4 \{dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2)\} \quad (95)$$

here $r_o = \frac{r_g}{4}$ where r_g is Schwarzschild parameter [1] Then relevant Dirac matrices are

$$\gamma_0 = \left(\frac{r-r_o}{r+r_o}\right) \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} ; \quad (96)$$

$$\gamma_1 = \left(\frac{r+r_o}{r}\right)^2 \begin{bmatrix} 0 & \sigma_1 \\ -\sigma_1 & 0 \end{bmatrix} ; \quad (97)$$

$$\gamma_2 = \frac{(r+r_o)^2}{r} \begin{bmatrix} 0 & \sigma_2 \\ -\sigma_2 & 0 \end{bmatrix} ; \quad (98)$$

$$\gamma_3 = \frac{(r+r_o)^2}{r} \sin\theta \begin{bmatrix} 0 & \sigma_3 \\ -\sigma_3 & 0 \end{bmatrix} \quad (99)$$

Following notations are used for linear combinations of standard Pauli matrices

$$\sigma_1 = \begin{bmatrix} \cos\theta & \sin\theta \exp(-i\phi) \\ \sin\theta \exp(i\phi) & -\cos\theta \end{bmatrix} ; \quad (100)$$

$$\sigma_2 = \begin{bmatrix} -\sin\theta & \cos\theta \exp(-i\phi) \\ \cos\theta \exp(i\phi) & \sin\theta \end{bmatrix} ; \quad (101)$$

$$\sigma_3 = \begin{bmatrix} 0 & -i \exp(-i\phi) \\ i \exp(i\phi) & 0 \end{bmatrix} \quad (102)$$

Non-zero components of Christoffel symbol are

$$\Gamma_{01}^0 = \frac{2r_o}{r^2 - r_o^2} ; \Gamma_{11}^1 = -\frac{2r_o}{r(r+r_o)} ; \Gamma_{33}^2 = -\sin\theta \cos\theta ; \Gamma_{22}^1 = -r \left(\frac{r-r_o}{r+r_o}\right) ;$$

$$\Gamma_{00}^1 = \frac{2r_or^4(r-r_o)}{(r+r_o)^7} ; \Gamma_{12}^2 = \Gamma_{13}^3 = \frac{1}{r} \left(\frac{r-r_o}{r+r_o}\right) ; \Gamma_{23}^3 = \cot\theta ; \Gamma_{33}^1 = -r \left(\frac{r-r_o}{r+r_o}\right) \sin^2\theta$$

Solving equations (46) for Γ_c one may find spinorial connections

$$\Gamma_0 = a_0\gamma_0\gamma_1 ; \Gamma_1 = 0 ; \Gamma_2 = a_2\gamma_1\gamma_2 ; \Gamma_3 = a_2\gamma_1\gamma_3$$

Here

$$a_0 = \frac{rr_o^4}{(r-r_o)(r+r_o)^5} ; a_2 = \frac{r_or^3}{(r+r_o)^5}$$

Suppose spinors $\bar{\xi}_0$ and ξ_0 have components $(\bar{f}, 0, 0, 0)$ and $[f, 0, 0, 0]^T$ respectively Calculating (50) I obtained only one non-zero component

$$H_{01} = \frac{1}{4} \{ \bar{\xi}_0 \gamma_0 \xi'_0 - \bar{\xi}'_0 \gamma_0 \xi_0 \} \quad (103)$$

Here prime denotes derivative over r Focusing attention on eqs (93) and (94) with τ^{cb} having only one non-zero component $\tau^{01}(r)$ I rewrite these eqs in following way

$$\tau^{01} \{ \xi'_0 + a_0 g_{11} \xi_0 \} - \frac{8\pi k \epsilon^2}{c^4} \xi_0 = 0 ; \quad (104)$$

$$\tau^{01} \{ \bar{\xi}'_0 + a_0 g_{11} \bar{\xi}_0 \} + \frac{8\pi k \epsilon^2}{c^4} \bar{\xi}_0 = 0 \quad (105)$$

or

$$f' + a_0 g_{11} f = \epsilon Y f + O(\epsilon \ell) ; \quad (106)$$

$$\bar{f}' + a_0 g_{11} \bar{f} = -\epsilon Y \bar{f} + O(\epsilon \ell) \quad (107)$$

Here $Y = Y(r)$ is solution of second order nonlinear differential equation

$$\tau^{01}(r; Y) \cdot Y = \frac{8\pi k \epsilon}{c^4} \quad (108)$$

From (103)–(105) follows

$$H_{01} = \frac{\epsilon}{2} Y(r) \left(\frac{r-r_o}{r+r_o} \right) \bar{f} f + O(\epsilon \ell) \quad (109)$$

which manifests (due to (66)) that τ^{01} contains second order derivatives of Y and doesn't have term of order ϵ^0 in its decomposition Thus one should solve eqs (66) (106)–(109) If solution of (108) for Y exists then one may write solution for spinors as

$$f = \left(\frac{r-r_o}{r+r_o} \right)^{\frac{1}{2}} \exp \left(-\epsilon \int_r^\infty Y dr \right) + O(\epsilon \ell) ; \quad (110)$$

$$\bar{f} = \left(\frac{r-r_o}{r+r_o} \right)^{\frac{1}{2}} \exp \left(+\epsilon \int_r^\infty Y dr \right) + O(\epsilon \ell) \quad (111)$$

It follows

$$\bar{\xi} \neq \xi^* \gamma^0$$

where superscript $*$ stands for complex conjugate Transferring to Schwarzschild coordinates one obtains

$$\bar{\xi} \gamma_c \xi = \left(1 - \frac{r_g}{r} + O(\epsilon \ell), 0, 0, 0 \right) \quad (112)$$

In limit $\epsilon \cdot \ell \rightarrow 0$ vector (112) coincides with appropriate Killing vector For black hole stress-energy only non-zero component one obtains

$$T^{00} = \frac{\mu c^2}{g_{00}}$$

where $\mu(x^\alpha) = M \delta^3(x^\alpha)$ is density of mass defined in Cartesian coordinates Using formulae (59) and (112) one obtains for energy

$$E = c^2 \int \mu(x^\alpha) r^2 dr \sin \theta d\theta d\phi = M c^2 + O(\epsilon \ell)$$

(I transferred back to Cartesian coordinates to compute integral)

7 Conclusion

In this work I introduced pair of 4-dim spinorial fields $\bar{\xi}$ and ξ allowing (if theory is consistent) to construct conserving (in time) integral(s) at least for flat spacetime and possibly for static spherically symmetric Ricci-flat case (if eq (108) possesses solution) One observes that $\bar{\xi}\gamma_c\xi = \xi_c + O(\epsilon\ell)$ It should be noted that suggested approach admits generalization for higher (even-)dimensional spacetime

References

- [1] L D Landau E M Lifshits *The Classical Theory of Fields* Pergamon Press 1975
- [2] R C Tolman *Relativity Thermodynamics and Cosmology* Oxford U Press London 1934
- [3] A Papapetrou Proc R Ir Acad A52 11 (1948)
- [4] P G Bergmann R Thompson Phys Rev 89 400 (1953)
- [5] C Møller Ann Phys (NY) 4 347 (1958)
- [6] Robert M Wald *General Relativity* U of Chicago Press 1984
- [7] S Weinberg *Gravitation and Cosmology* John Wiley and Sons Inc NY 1972
- [8] S Deser arXiv:gr-qc/9901007
- [9] Ragab M Gad arXiv:gr-qc/0403039