## A no-go theorem for polytropic spheres in Palatini f(R) gravity

Enrico Barausse<sup>1</sup>, Thomas P. Sotiriou<sup>1</sup> and John C.  $Miller^{1,2}$ 

<sup>1</sup>SISSA, International School for Advanced Studies, Via Beirut 2-4, 34014 Trieste, Italy and INFN, Sezione di Trieste <sup>2</sup>Department of Physics (Astrophysics), University of Oxford, Keble Road, Oxford OX1 3RH, England

E-mail: barausse@sissa.it,sotiriou@sissa.it,miller@sissa.it

**Abstract.** Non-vacuum static spherically-symmetric solutions in Palatini f(R) gravity are examined. It is shown that for generic choices of f(R), there are commonly-used equations of state for which no satisfactory physical solution of the field equations can be found within this framework, apart from in the special case of General Relativity, casting doubt on whether Palatini f(R) gravity can be considered as giving viable alternatives to General Relativity.

The search for theories of gravity which can serve as an alternative to General Relativity (GR) has received a powerful stimulus from current developments in observational cosmology. If one accepts GR as the correct theory of gravity, then it would seem that the energy density of the universe must be dominated at the present time by a cosmological constant, or by an unknown form of energy (dark energy) that mimics the behaviour of a cosmological constant [1]. The various problems associated with this [2] have prompted many authors to question whether GR is indeed a completely correct theory of gravity on the classical level and to investigate possible alternatives which would not require the inclusion of dark energy.

One of these alternatives is f(R) gravity in the Palatini formalism, or simply Palatini f(R) gravity [3], and we will be examining some of its properties in this paper. We first recall how this class of theories comes about as a generalization of GR. The Einstein equations can be derived from the Einstein-Hilbert action not only by the standard metric variation but also by making independent variations with respect to the metric and the connections (see, for example, [4]). This is called Palatini variation and, if one proceeds in this way, the form of the connections is a derived property rather than being specified separately (*e.g.* by specifying the Levi-Civita connection, as in standard GR). In Palatini f(R) gravity, this extended form of variation is applied to an action which is a general function of the scalar curvature f(R):

$$S = \frac{1}{16\pi} \int d^4x \sqrt{-g} f(R) + S_M(g_{\mu\nu}, \psi),$$
(1)

where  $R = g^{\mu\nu}R_{\mu\nu}$ , g is the determinant of the metric  $g_{\mu\nu}$ ,  $S_M$  is the matter action and  $\psi$  collectively denotes the matter fields. (We are using, throughout, units in which c = G = 1.) Note that here  $R_{\mu\nu}$  is not the Ricci tensor of the metric  $g_{\mu\nu}$ , but is constructed from the independent connections  $\Gamma^{\lambda}{}_{\mu\nu}$ . Independent variation with respect to the metric and the connections gives

$$F(R)R_{\mu\nu} - \frac{1}{2}f(R)g_{\mu\nu} = 8\pi T_{\mu\nu},$$
(2)

$$\nabla_{\sigma}(\sqrt{-g}F(R)g^{\mu\nu}) = 0, \qquad (3)$$

where  $F(R) = \partial f/\partial R$ ,  $T_{\mu\nu} \equiv -2(-g)^{-1/2} \delta S_M / \delta g^{\mu\nu}$  is the usual stress-energy tensor of the matter and  $\nabla_{\mu}$  is the covariant derivative with respect to the connections  $\Gamma^{\lambda}_{\mu\nu}$ . Setting f(R) = R leads to the standard GR Einstein equations while (3) in this case incorporates the definition of the Levi-Civita connection. Note that in order to derive (2) and (3), one assumes that the matter action does not depend on the independent connections [see (1)]. Using Palatini variation while allowing the matter to couple to the independent connections leads to metric-affine f(R) gravity [5] which is a different theory with enriched phenomenology [6, 7].

Since it was shown that Palatini f(R) gravity might be able to address darkenergy problems [8], a number of studies have been made of its cosmological aspects and of its consistency with cosmological constraints [9] as well as of the Newtonian and post-Newtonian limits and consistency with Solar System constraints [10]. However, the question of obtaining solutions describing stars and compact objects has received only a small amount of attention so far. In this paper, we focus on the problem of finding consistent solutions for static spherically-symmetric stars when  $f(R) \neq R$ .

We first note that doing this is helped by the fact that Palatini f(R) gravity retains a useful characteristic of GR: the exterior spherically symmetric solution is unique (*Birkhoff's theorem*). To see this one must take the trace of (2),

$$F(R)R - 2f(R) = 8\pi T,$$
 (4)

where  $T \equiv g^{\mu\nu}T_{\mu\nu}$ . For a given f(R), this is an algebraic equation in R and therefore it can be solved to give R as a function of T. We will not consider cases where this equation has no roots, since those do not give viable classical gravity theories [11]. Equation (4) also implies that if T = 0, R must be constant. If we denote the value of R when T = 0 by  $R_0$  and insert it into (3), this equation reduces to the covariant conservation of  $g^{\mu\nu}$ , fixing the connection to be the Levi-Civita one. This will be the situation in vacuum and in this case (2) reduces to

$$R_{\mu\nu} - \Lambda(R_0)g_{\mu\nu} = 0, \tag{5}$$

where  $\hat{R}_{\mu\nu}$  is now indeed the Ricci tensor of the metric and  $\Lambda(R_0) = R_0/4$ . According to whether  $R_0$  is zero or not, which of course depends on the choice of f(R), the theory reduces in vacuum to GR without or with a cosmological constant. The vacuum spherically symmetric solution is unique in either case, being either Schwarzschild or Schwarzschild-(anti-)de Sitter.

Having determined an exterior solution, we then need to find an interior solution and perform a matching between the two. Recently, the generalisation of the Tolman–Oppenheimer–Volkoff (TOV) hydrostatic equilibrium equation for Palatini f(R) gravity has been derived [12], opening the way for finding equilibrium interior solutions. We will consider here the matching of such interior solutions with exterior ones.

We begin by reviewing the formulae that we will need. Denoting the Ricci scalar of  $g_{\mu\nu}$  by  $\tilde{R} \equiv g^{\mu\nu}\tilde{R}_{\mu\nu}$  and setting  $\tilde{G}_{\mu\nu} = \tilde{R}_{\mu\nu} - g_{\mu\nu}\tilde{R}/2$ , the field equations (2) and (3) can be rewritten as a single one

$$\widetilde{G}_{\mu\nu} = \frac{8\pi}{F} T_{\mu\nu} - \frac{1}{2} g_{\mu\nu} \left( R - \frac{f}{F} \right) + \frac{1}{F} \left( \widetilde{\nabla}_{\mu} \widetilde{\nabla}_{\nu} - g_{\mu\nu} \widetilde{\Box} \right) F -$$

A no-go theorem for polytropic spheres in Palatini f(R) gravity

$$-\frac{3}{2}\frac{1}{F^2}\left((\widetilde{\nabla}_{\mu}F)(\widetilde{\nabla}_{\nu}F) - \frac{1}{2}g_{\mu\nu}(\widetilde{\nabla}F)^2\right),\tag{6}$$

3

where  $\widetilde{\nabla}_{\mu}$  is the covariant derivative with respect to the Levi-Civita connection of  $g_{\mu\nu}$ and  $\widetilde{\Box} \equiv g^{\mu\nu}\widetilde{\nabla}_{\mu}\widetilde{\nabla}_{\nu}$ . To arrive at this equation, one has to solve (3) for  $\Gamma^{\lambda}_{\mu\nu}$ , insert this into (2) and express the resulting equation in terms only of metric quantities (for an alternative method, see [6]).

Using the static spherically symmetric ansatz

$$ds^{2} \equiv -e^{A(r)} dt^{2} + e^{B(r)} dr^{2} + r^{2} d\Omega^{2},$$
(7)

in (6), considering perfect-fluid matter with  $T_{\mu\nu} = (\rho + p)u_{\mu}u_{\nu} + pg_{\mu\nu}$  (where  $\rho$  is the energy density, p is the pressure and  $u^{\mu}$  is the fluid 4-velocity) and representing d/dr with a prime, one arrives at the equations

$$A' = \frac{-1}{1+\gamma} \left( \frac{1-e^B}{r} - \frac{e^B}{F} 8\pi G r p + \frac{\alpha}{r} \right),\tag{8}$$

$$B' = \frac{1}{1+\gamma} \left( \frac{1-e^B}{r} + \frac{e^B}{F} 8\pi G r \rho + \frac{\alpha+\beta}{r} \right),\tag{9}$$

$$\alpha \equiv r^2 \left( \frac{3}{4} \left( \frac{F'}{F} \right)^2 + \frac{2F'}{rF} + \frac{e^B}{2} \left( R - \frac{f}{F} \right) \right), \tag{10}$$

$$\beta \equiv r^2 \left( \frac{F''}{F} - \frac{3}{2} \left( \frac{F'}{F} \right)^2 \right), \qquad \gamma \equiv \frac{rF'}{2F}.$$
 (11)

Defining  $m_{\text{tot}}(r) \equiv r(1 - e^{-B})/2$  and using Euler's equation, one gets the generalized TOV equations [12]:

$$p' = -\frac{1}{1+\gamma} \frac{(\rho+p)}{r(r-2m_{\rm tot})} \left( m_{\rm tot} + \frac{4\pi r^3 p}{F} - \frac{\alpha}{2}(r-2m_{\rm tot}) \right), (12)$$

$$m'_{\text{tot}} = \frac{1}{1+\gamma} \left( \frac{4\pi r^2 \rho}{F} + \frac{\alpha+\beta}{2} - \frac{m_{\text{tot}}}{r} (\alpha+\beta-\gamma) \right).$$
(13)

We consider here matter which can be described by a one-parameter equation of state (EOS)  $p = p(\rho)$ . When this is specified, one can in principle solve the above equations and derive an interior solution. However, this is hard to do in practice because the equations are implicit, their right-hand sides effectively including through F' and F'' both first and second derivatives of the pressure, e.g., F' = d/dr [F(R(T))] = (dF/dR) (dR/dT) (dT/dp) p'. We therefore first put them in an explicit form, which allows us not only to solve them numerically, but also to study their behaviour at the stellar surface.

Multiplying (12) by dF/dp and using the definitions of  $\alpha$  and  $\gamma$ , we get a quadratic equation in F', whose solution is

$$F' = \frac{-4rF(\mathcal{C} - F)(r - 2m_{\text{tot}}) + D\sqrt{2\Delta}}{r^2(3\mathcal{C} - 4F)(r - 2m_{\text{tot}})}$$
(14)

where  $D = \pm 1$  and where we have defined

$$\mathcal{C} = \frac{dF}{dp}(p+\rho) = \frac{dF}{d\rho}\frac{d\rho}{dp}(p+\rho), \tag{15}$$

$$\Delta = Fr^{2}(r - 2m_{\text{tot}}) \left[ 8F(\mathcal{C} - F)^{2}(r - 2m_{\text{tot}}) - \mathcal{C}(4F - 3\mathcal{C}) \left( (16\pi p - FR + f)r^{3} + 4Fm_{\text{tot}}) \right].$$
(16)

We will now focus on polytropic EOSs given by  $p = \kappa \rho_0^{\Gamma}$ , where  $\rho_0$  is the restmass density and  $\kappa$  and  $\Gamma$  are constants, noting that this can be rewritten as  $\rho = (p/\kappa)^{1/\Gamma} + p/(\Gamma - 1)$ , giving a direct link between p and  $\rho$ . In (15), we have written C in terms of  $dF/d\rho$  because this is finite at the stellar surface  $(r = r_{out}$  where  $p = \rho = 0$ ). In fact,  $dF/d\rho = (dF/dR) (dR/dT) (3dp/d\rho - 1)$ , where dF/dR and dR/dT are in general finite even when  $T = 3p - \rho$  goes to zero [check for instance the  $R^2$  or 1/R models] and  $dp/d\rho \to 0$  for  $p \to 0$ . Note also that while  $d\rho/dp$  diverges when  $p \to 0$ , the product  $(p + \rho) d\rho/dp$  goes to zero for  $p \to 0$  if  $\Gamma < 2$ . Therefore, for a polytrope with  $\Gamma < 2$ , C = 0 at the surface.

We now match the interior solution to the exterior one. For the latter, the general solution to (5) is  $\exp(-B(r)) = \ell \exp(A(r)) = 1 - 2m/r - R_0 r^2/12$ , where  $\ell$  and m are integration constants to be fixed by requiring continuity of the metric coefficients across the surface and  $R_0$  is again the vacuum value of R. Using the definition of  $m_{\text{tot}}(r)$  this gives, in the exterior,  $m_{\text{tot}}(r) = m + r^3 R_0/24$ . Besides continuity of the metric, one has to impose some junction conditions for A'. The exterior solution evaluated at the surface gives

$$A'(r_{\rm out}) = \frac{2\left(r_{\rm out}^3 R_0 - 12m\right)}{r_{\rm out}\left(R_0 r_{\rm out}^3 - 12r_{\rm out} + 24m\right)},\tag{17}$$

whereas the value of  $A'(r_{out})$  for the interior solution can be calculated with (8). For this we need  $F'(r_{out})$ . Evaluating (14) at the surface, where  $\mathcal{C} = p = 0$  and R, F and f take their constant vacuum values  $R_0$ ,  $F_0$  and  $f_0 = F_0 R_0/2$ , we get  $F'(r_{out}) = -(1 + \tilde{D})F_0/r_{out}$ , where  $\tilde{D} = D \operatorname{sign}(r_{out} - 2m_{tot}) \ddagger$ . Choosing  $\tilde{D} = 1$  implies  $\gamma = -1$  at the surface [see (11)] giving  $A' \to \infty$  for  $r \to r_{out}^-$  [see (8)], whereas A' keeps finite for  $r \to r_{out}^+$  [see (17)]. Because  $\tilde{G}_{\mu\nu}$  involves A'', this infinite discontinuity leads to the presence of Dirac deltas in the field equations. These Dirac deltas cannot be cancelled by the derivatives of F' on the right-hand side, because the discontinuity of F' is only a finite one, and one should therefore invoke an infinite surface density at  $r = r_{out}$ . Since this is unreasonable, we focus only on  $\tilde{D} = -1$ , for which  $F'(r_{out}) = 0$  when  $r \to r_{out}^-$ , making both F' and A' continuous across the surface.

In order to study the behaviour of  $m_{\text{tot}}$  at the surface, we need first to derive an explicit expression for F''. If we take the derivative of (14), F'' appears on the left-hand side and also on the right-hand side [through  $m'_{\text{tot}}$ , calculated from (13) and the definition of  $\beta$ , (11)], giving a linear equation in F''. The solution to this, evaluated at the surface, is

$$F''(r_{\rm out}) = \frac{\left(R_0 r_{\rm out}^3 - 8m_{\rm tot}\right)\mathcal{C}'}{8r_{\rm out}(r_{\rm out} - 2m_{\rm tot})} \tag{18}$$

Evaluating  $\alpha$ ,  $\beta$  and  $\gamma$  at the surface using F' = 0 and F'' given by (18), and inserting into (13) gives

$$m'_{\rm tot}(r_{\rm out}) = \frac{2F_0 R_0 r_{\rm out}^2 + (r_{\rm out}^3 R_0 - 8m_{\rm tot}) \mathcal{C}'}{16F_0} \,. \tag{19}$$

For  $1 < \Gamma < 3/2$ ,  $\mathcal{C}' = d\mathcal{C}/dp \, p' \propto d\mathcal{C}/dp \, (p+\rho) \to 0$  at the surface so that expression (19) is finite and it even gives continuity of  $m'_{\text{tot}}$  across the surface. However, for  $3/2 < \Gamma < 2$ ,  $\mathcal{C}' \to \infty$  as the surface is approached, provided that  $dF/dR(R_0) \neq 0$ 

<sup>&</sup>lt;sup>‡</sup> Unlike in GR, one cannot prove that  $r_{\text{out}} > 2m_{\text{tot}}$  from (12) because p' is not necessarily positive, although one may expect  $r_{\text{out}} > 2m_{\text{tot}}$  in sensible solutions.

and  $dR/dT(T_0) \neq 0$  (note that these conditions are satisfied by generic forms of f(R), *i.e.* whenever an  $R^2$  term or a term inversely proportional to R is present). While  $m_{\text{tot}}$  keeps finite [as can be shown using  $p \sim (r_{\text{out}} - r)^{\Gamma/(\Gamma-1)}$ , which can be derived by integrating (12) near the surface], the divergence of  $m'_{\text{tot}}$  drives to infinity the Riemann tensor of the metric,  $\tilde{R}_{\mu\nu\sigma\lambda}$ , and curvature invariants, such as  $\tilde{R}$  or  $\tilde{R}^{\mu\nu\sigma\lambda}\tilde{R}_{\mu\nu\sigma\lambda}$ , as can easily be checked §. This singular behaviour would cause unphysical phenomena, such as infinite tidal forces which would destroy anything present at the surface [cf. the geodesic deviation equation]. We can then conclude that no physically relevant solution exists for any polytropic EOS with  $3/2 < \Gamma < 2$ . Of course, polytropes give only simplified models for stars and the EOS in the outer layers is critical for the behaviour of  $m'_{\text{tot}}$  at the surface in the non-GR case. One would like to use a more accurate EOS, but while this can give regular solutions in many cases (*e.g.* if  $p \propto \rho_0$ near the surface), the existence of counter-examples is worrying for the viability of the theory.

Setting aside the surface singularity, we next turn to the behaviour in the interior, focusing on models of neutron stars constructed using an analytical approximation to the FPS EOS [14]. Adding positive powers of R to the Einstein-Hilbert action produces significant effects for compact stars while adding negative ones predominantly affects more diffuse stars. Generically, though, one would expect terms of both types to be added if there is a deviation away from GR. Since the 1/R term commonly used in cosmology would have a negligible effect in the interior of a neutron star, we used here  $f(R) = R + \epsilon R^2$ . As can be seen from (6), the metric will be sensitive to derivatives of the matter fields, since R is a function of  $T \parallel$ .

This can be seen in Figure 1:  $m_{\rm tot}$ , which in GR has a smooth profile, now develops peculiar features when  $d\rho/dp$  and  $d^2\rho/dp^2$  change rapidly in going from the core to the inner crust and from the inner crust to the outer crust. If  $m_{\rm tot}$  were plotted against the radius, these features would look much more abrupt, because they occur in a small range of radii close to the surface. While  $m_{\rm tot}$  does not represent a real mass in the interior, such a strong dependence of the metric on the derivatives of the matter fields is not very plausible and could have dramatic consequences.

We have therefore found two unappealing characteristics of Palatini f(R) gravity as applied to stellar models, each of which arises because of the dependence of the metric on higher order derivatives of the matter field. First: whether or not a regular solution can be found depends crucially on the microphysics, through the EOS, with polytropic EOSs having  $3/2 < \Gamma < 2$  being ruled out for generic f(R). Second: even if an EOS does allow a regular solution at the surface, the interior metric depends on the first and second derivatives of the density with respect to the pressure, giving a problematic behaviour. While polytropic EOSs are highly idealized, we note that  $\Gamma = 5/3$ , corresponding to an isentropic monatomic gas or a degenerate non-relativistic particle gas, falls within the range not giving a regular solution. The fact that the gravity theory cannot provide a consistent description for these cases, casts doubt on whether it should be considered as a viable alternative to GR.

Since the problems discussed here arise due to the dependence of the metric on higher order derivatives of the matter fields, one can expect that they will also

<sup>§</sup> This seems to have been missed in [15].

<sup>||</sup> The unusual behaviour of this class of theories has been mentioned in a different context in [16]. However, we disagree with the claims made there about the violation of the equivalence principle, because they seem to be based on an ill-posed identification of the metric whose geodesics should coincide with free-fall trajectories.



Figure 1. Profiles of  $m_{\rm tot}$  (in  $M_{\odot}$ ) and other associated quantities plotted against density in the interior of a neutron-star with central density  $10^{15}$ g/cm<sup>3</sup> and p' = 0 in the centre as required by local flatness. We have used the FPS EOS [14] and  $f(R) = R + \epsilon R^2$ . The dot-dashed line shows  $m_{\rm tot}$  as calculated with  $\epsilon = 0.1$  and the dotted line shows the equivalent curve in GR ( $\epsilon = 0$ ); the solid line shows  $dm_{\rm tot}/dr$  (in  $M_{\odot}/\rm km$ ) for  $\epsilon = 0.1$  (this value is orders of magnitude lower than Solar System constraints [13]). Note the bumps in the  $dm_{\rm tot}/dr$ curve resulting from rapid composition changes in the EOS (the corresponding features in the  $m_{\rm tot}$  curve for  $\epsilon = 0.1$  are less apparent but a noticeable dip is seen at  $\rho \sim 10^{14} \rm g/cm^3$ ). To make evident the influence of composition changes, we also show comparisons between the FPS EOS and a polytrope (with  $\Gamma = 4/3$  and  $\kappa = 10^{15}$  cgs): the long-dashed and short-dashed curves show  $0.1 \times (d\rho/dp)_{\rm FPS}/(d\rho/dp)_{\rm polytrope}$  and  $0.01 \times (d^2\rho/dp^2)_{\rm FPS}/(d^2\rho/dp^2)_{\rm polytrope}$ , respectively.

appear in other gravity theories having these characteristics. Any theory having a representation in which the field equations include second derivatives of the metric and higher than first derivatives of the matter fields will face similar problems because having a higher differential order in the metric than in the matter field is what guarantees that the metric depends in a cumulative way on the matter. If this is not the case then the metric loses its immunity to rapid changes in matter gradients since it is directly related to them instead of being an integral over them.

The same problem should be expected for any theory which includes fields other than the metric for describing the gravitational interaction (e.g. scalar fields) which are algebraically related to matter rather than dynamically coupled. In this case one can always solve the field equations for the extra field and insert the solution into the field equation for the metric, inducing a dependence of the metric on higher derivatives of the matter fields. An example of such a theory is a scalar-tensor theory with Brans-Dicke parameter  $\omega = -3/2$ , which is anyway an equivalent representation of Palatini f(R) gravity [6]. One should mention that this problem could probably be addressed in Palatini f(R) gravity by adding higher order curvature invariants in the action [e.g.  $f(R, R^{\mu\nu}R_{\mu\nu})$ ], since this would introduce more dynamics and break the non-dynamical coupling between matter and the extra gravitational degrees of freedom.

In conclusion, we suggest that our results cast doubt on the viability of theories including higher order derivatives of the matter fields in one of their representations, such as Palatini f(R) gravity or  $\omega = -3/2$  scalar-tensor theory.

*Note added:* Prior to publication, a paper by Kainulainen *et al.* appeared [17], which agreed with the validity of our results but criticized our interpretation of them. A response to this criticism, as well as an additional analysis of the issues presented here, can be found in [18].

## Acknowledgements

We wish to thank G. Olmo and S. Liberati for helpful discussions.

## References

- [1] Astier P et al. 2006 Astron. Astrophys. 447 31;
- Spergel D N et al. [WMAP Collaboration] 2007 Astrophys. J. Suppl. 170 377
- [2] Carroll S M 2001 Living Rev. Rel. 4 1
- [3] Buchdahl H A 1970 Mon. Not. Roy. Ast. Soc. 150 1;
- Ferraris M, Francaviglia M and Reina C 1982 Gen. Rel. Grav. 14 243
- [4] Wald R M 1984 General Relativity, (University of Chicago Press, Chicago)
- [5] Sotiriou T P and Liberati S 2007 Ann. Phys. 322 935;
   Sotiriou T P and Liberati S 2007 J. Phys. Conf. Ser. 68 012022
- [6] Sotiriou T P 2006 Class. Quant. Grav. 23 5117
- [7] Sotiriou T P 2006 arXiv:gr-qc/0611158
- [8] Vollick D N 2003 Phys. Rev. D 68 063510
- [9] Barraco D, Hamity V H and Vucetich H 2002 Gen. Rel. Grav. 34 533; Allemandi G, Borowiec A and Francaviglia M 2004 Phys. Rev. D 70 043524; Meng X H and Wang P 2004 Class. Quant. Grav. 21 951; Meng X H and Wang P 2004 Phys. Lett. B 584 1; Meng X H and Wang P 2005 Class. Quant. Grav. 22 23; Allemandi G et al. 2005 Phys. Rev. D 72 063505; Sotiriou T P 2006 Phys. Rev. D 73 063515; Amarzguioui M et al. 2006 Astron. Astrophys. 454 707; Koivisto T and Kurki-Suonio H 2006 Class. Quant. Grav. 23 2355; Koivisto T 2006 Phys. Rev. D 73 083517; Li B, Chan K C and Chu M C 2007 Phys. Rev. D 76 024002
  [10] Olmo G J 2005 Phys. Rev. Lett. 95 261102;
- Allemandi G et al. 2005 Gen. Rel. Grav. 37 1891;
  Cembranos J A R 2006 Phys. Rev. D 73 064029;
  Sotiriou T P 2006 Gen. Rel. Grav. 38 1407;
  Sotiriou T P 2007 Phys. Lett. B 645 389;
  Allemandi G and Ruggiero M L 2007 Gen. Rel. Grav. 39 1381
  [11] Ferraris M, Francaviglia M and Volovich I 1993 arXiv: gr-qc/9303007.
- [11] Ferraris M, Francavigna M and Volovich T 1995 arXiv. gr-qc/9505007.
- [12] Kainulainen K, Reijonen V and Sunhede D 2007 Phys. Rev. D 76 043503
- [13] Sotiriou T P 2006 Class. Quant. Grav. 23 1253
- [14] Haensel P and Potekhin A Y 2004 Astron. Astroph. 428 191
- [15] Barraco D E and Hamity V H 2000 Phys. Rev. D 62 044027
- [16] Olmo G J 2007 Phys. Rev. Lett. 98 061101
- [17] Kainulainen K, Piilonen J, Reijonen V and Sunhede D 2007 Phys. Rev. D 76 024020
- [18] Barausse E, Sotiriou T P and Miller J C 2007 arXiv:0712.1141 [gr-qc].