

# Stability of Closed Timelike Geodesics.

V. M. Rosa\*

Departamento de Matemática, Universidade Federal de Viçosa  
36570-000 Viçosa, M.G., Brazil

P. S. Letelier†

Departamento de Matemática Aplicada-IMECC,  
Universidade Estadual de Campinas,  
3083-970, Campinas, S.P., Brazil

February 27, 2019

## Abstract

The existence and stability under linear perturbations of closed timelike geodesics (CTG) in Bonnor-Ward spacetime is studied in some detail. Regions where the CTG exist and are linearly stable are exhibited.

In 1949 Gödel found a solution to the Einstein field equation with nonzero cosmological constant that admits closed timelike curves (CTC) [1]. It could be argued that the Gödel solution is without physical significance, since it corresponds to a rotating, stationary cosmology, whereas the actual universe is expanding and apparently non rotating. The van Stockum solution [2], that also contain CTC, is physically inadmissible since refers to an infinitely long cylinder. But there exist examples of solutions of vacuum Einstein's equations which contains CTC that can represent the exterior of physically admissible sources [3] [4].

In [3] it is described the case of a massless spinning rod of finite length. In [4] it is analyzed the CTC in Kerr-Newman spacetime and in a solution of the Einstein equations for a source named Perjeon, due to Perjés [5], which represents a single charged, rotating, magnetic object. This solution was also studied independently by Israel and Wilson [6], and it is referred as a PIW spacetime. In these three cases one expect that the CTC region be covered by the source. The same does not happen when we work with two Perjeons [4].

The PIW metric is given by

$$ds^2 = -f^{-1} h_{mn} dx^m dx^n + f(\omega_m dx^m + dt)^2, \quad (1)$$

where the three dimensional positive definite tensor  $h_{mn}$  has zero Ricci tensor and it will be taken as the usual three dimensional Euclidean metric in cylindrical coordinates, the electromagnetic field is given in terms of two scalar potentials:

$$F_{4n} = \Phi_{,n}, \quad F^{ab} = \eta^{abm} f \psi_{,m}, \quad (2)$$

---

\*vmrosa@ufv.br

†letelier@ime.unicamp.br

$\eta^{abm}$  being the Levi-Civita tensor related to  $h_{mn}$ , and  $(\cdot)_{,n} = \partial/\partial x^n$ . The entire solution is generated by two functions  $L$  and  $M$  that are harmonic with respect to  $h_{mn}$  by means of the equations,

$$\begin{aligned} f &= \frac{1}{4L^2+4M^2}, \\ \psi &= \frac{1}{2}\epsilon\frac{M}{L^2+M^2}, \\ \Phi &= -\frac{1}{2}\epsilon\frac{L}{L^2+M^2}, \\ \omega_{a,b} - \omega_{b,a} &= 8(ML_{,c} - LM_{,c})\eta_{abm}h^{mc}. \end{aligned} \quad (3)$$

The Bonnor-Ward (BW) solution [7] refers to two Perjeons, with masses  $m_1$  and  $m_2$ , placed on the  $z$ -axis at  $z = \pm a$ , ( $a > 0$ ), with magnetic moments ( $\mu_1$  and  $\mu_2$ ) also parallel to the  $z$ -axis, and

$$\begin{aligned} L &= (1 + m_1/r_1 + m_2/r_2)/2 \\ M &= (\mu_1(z-a)/r_1^3 + \mu_2(z+a)/r_2^3)/2 \\ r_1 &= \sqrt{\rho^2 + (z-a)^2} \\ r_2 &= \sqrt{\rho^2 + (z+a)^2} \\ \omega &= (-\rho^2\phi)\Omega. \end{aligned} \quad (4)$$

We shall consider the particular case of BW solution [4],

$$\begin{aligned} \Omega &= \frac{\mu_1}{r_1^3}\left(2 + \frac{m_1}{r_1}\right) + \frac{m_1\mu_2}{\rho^2}\left(\frac{r_1}{2a^2r_2} + \frac{(z+a)(\rho^2 + z^2 - a^2)}{ar_1r_2^3}\right) \\ &+ \frac{\mu_2}{r_2^3}\left(2 + \frac{m_2}{r_2}\right) + \frac{m_2\mu_1}{\rho^2}\left(\frac{r_2}{2a^2r_1} - \frac{(z-a)(\rho^2 + z^2 - a^2)}{ar_2r_1^3}\right) \\ &+ \frac{\alpha}{\rho^2} \end{aligned} \quad (5)$$

with  $\alpha$  restricted to the values  $\alpha = \pm(m_1\mu_2 + m_2\mu_1)/(2a^2)(\neq 0)$  only. When  $\alpha = (m_1\mu_2 + m_2\mu_1)/(2a^2)$  the spacetime has a torsion line singularity (TLS) on  $\rho = 0$  for  $z^2 > a^2$  and in the other case the singularity appears between the sources. In both cases there is no strut singularity [4]. The presence of the TLSs is unavoidable in this case, they keep the two stationary spinning, charged, magnetic sources rotating in equilibrium (for torsions lines see [8][9]).

Let us denote by  $\gamma$  a closed curve in BW spacetime given in its parametric form by,

$$t = t_0, \quad \rho = r_0, \quad \phi \in [0, 2\pi], \quad z = z_0, \quad (6)$$

where  $t_0$ ,  $\rho_0$  and  $z_0$  are constants. The closed curve is timelike when  $g_{\phi\phi} > 0$ , i.e.,

$$\Omega^2 f^2 \rho^2 - 1 > 0. \quad (7)$$

The four-acceleration of  $\gamma$  is

$$a^t = 0, \quad (8)$$

$$a^\rho = \frac{\partial_\rho f \Omega^2 \rho^3 f^2 + 2f^3 \Omega \rho^3 \partial_\rho \Omega + 4f^3 \Omega^2 \rho^2 - 2f + \rho \partial_\rho f}{2\rho(f^2 \Omega^2 \rho^2 - 1)} \dot{\phi}^2, \quad (9)$$

$$a^\phi = 0, \quad (10)$$

$$a^z = \frac{\partial_z f \Omega^2 \rho^2 f^2 + 2f^3 \Omega \rho^2 \partial_z \Omega + \partial_z f}{2(f^2 \Omega^2 \rho^2 - 1)} \dot{\phi}^2. \quad (11)$$

We shall restrict us to the special case,

$$m_1 = m_2 = m, \mu_1 = \mu_2 = \mu. \quad (12)$$

Furthermore, we take  $z_0 = 0$  to have  $a^z = 0$ . The condition  $a^\rho = 0$  gives us to different cases, depending on the position of the TLS.

First we analyse the case  $\alpha = -(m_1\mu_2 + m_2\mu_1)/(2a^2)$  (TSL between the particles). By assuming the restriction (12) and doing  $a^\rho = 0$  we obtain the following relation for  $\mu$ ,  $a$ ,  $\rho$  and  $m$ ,

$$\begin{aligned} \mu = -1/4 \big( & -2(4\rho^4 m^3 + 8\sqrt{\rho^2 + a^2 m^2} \rho^4 + 2\sqrt{\rho^2 + a^2} \rho^6 - 16a^2 \rho^2 m^3 \\ & - 19a^4 m \rho^2 + 6\rho^6 m - 2a^2 \sqrt{\rho^2 + a^2} \rho^4 + 7a^4 m^3 - 15a^2 m \rho^4 + 2a^6 m \\ & - 32a^2 \sqrt{\rho^2 + a^2 m^2} \rho^2 + 8a^4 \sqrt{\rho^2 + a^2 m^2} - 4a^4 \sqrt{\rho^2 + a^2} \rho^2)(a^6 \sqrt{\rho^2 + a^2} \\ & + 10a^6 m + 3a^4 \sqrt{\rho^2 + a^2} \rho^2 + 28a^4 m \rho^2 + 80a^4 m^3 + 40a^4 \sqrt{\rho^2 + a^2 m^2} \\ & + 80a^2 \sqrt{\rho^2 + a^2 m^4} + 64a^2 \sqrt{\rho^2 + a^2 m^2} \rho^2 + 3a^2 \sqrt{\rho^2 + a^2} \rho^4 + 32a^2 m^5 \\ & + 112a^2 \rho^2 m^3 + \sqrt{\rho^2 + a^2} \rho^6 + 24\sqrt{\rho^2 + a^2 m^2} \rho^4 + 8\rho^6 m + 32\rho^4 m^3 \\ & + 26a^2 m \rho^4 + 16\sqrt{\rho^2 + a^2 m^4} \rho^2) \big)^{1/2} (\rho^2 + a^2) / \big( 4\rho^4 m^3 + 8\sqrt{\rho^2 + a^2 m^2} \rho^4 \\ & + 2\sqrt{\rho^2 + a^2} \rho^6 - 16a^2 \rho^2 m^3 - 19a^4 m \rho^2 + 6\rho^6 m - 2a^2 \sqrt{\rho^2 + a^2} \rho^4 \\ & + 7a^4 m^3 - 15a^2 m \rho^4 - 32a^2 \sqrt{\rho^2 + a^2 m^2} \rho^2 + 2a^6 m + 8a^4 \sqrt{\rho^2 + a^2 m^2} \\ & - 4a^4 \sqrt{\rho^2 + a^2} \rho^2 \big). \end{aligned} \quad (13)$$

Using this form for  $\mu$  and doing  $\rho = na$  we obtain for  $g_{\phi\phi}$ ,

$$\begin{aligned} g_{\phi\phi} = & - \left[ 2n^2(n^2 + 1)(2n^2 - 1)a^2 + m\sqrt{n^2 + 1}(6n^4 - 11n^2 - 1)a \right. \\ & + 2m^2(2n^4 - 5n^2 - 1) \big] / \left[ 2n^2 \left( (n^2 + 1)(n^2 - 2)a^2 + 2m\sqrt{n^2 + 1}(n^2 - 4)a \right. \right. \\ & \left. \left. + m^2(2n^2 - 7) \right) \right]. \end{aligned} \quad (14)$$

We can find  $n$ , the distance  $a$ , and the mass  $m$  such that  $g_{\phi\phi} > 0$ . We have closed timelike geodesics for  $n \in [0.708, \sqrt{2}]$  and  $a > Am$ , for  $n \in (\sqrt{2}, 1.6385]$  and  $a \in [Am, Bm]$ , and for  $n \in (1.6385, 1.8872]$  and  $a \in [0, Bm]$ , where

$$\begin{aligned} A &= \frac{\sqrt{1 + 6n^2 + 61n^4 + 60n^6 - 28n^8} - (6n^4 - 11n^2 - 1)}{4\sqrt{n^2 + 1}(2n^2 - 1)}, \\ B &= \frac{4 - n^2 + \sqrt{2 + 3n^2 - n^4}}{\sqrt{n^2 + 1}(n^2 - 2)}. \end{aligned}$$

In the particular case  $a = \rho$ , the condition (7) is written as  $(2\alpha - 2\sqrt{2}m)/(2\alpha + 5\sqrt{2}m) > 0$  and if  $a > \sqrt{2}m$  the closed geodesic is timelike. In [4] Bonnor and Steadman describe a CTG for  $a = \rho = 1/\sqrt{2}$ , in this case the closed curve  $\gamma$  is a geodesic when (13) reduces to  $\mu = (2m + 1)^2/(\sqrt{10m^2 + 22m + 4})$ . The condition (7) is written as  $(1 - 4m)/(1 + 5m) > 0$ , so  $\gamma$  is timelike when  $m < 1/4$ .

Now we analyse the case  $\alpha = (m_1\mu_2 + m_2\mu_1)/(2a^2)$  (two semi-infinity TLS). Assuming that (12) is true and making  $a^\rho = 0$  we obtain the following relation for  $\mu$ ,  $a$ ,  $\rho$  and  $m$ ,

$$\begin{aligned} \mu = & \left( (4a^8 \sqrt{\rho^2 + a^2} + 2a^6 \sqrt{\rho^2 + a^2} \rho^2 - 2a^4 \sqrt{\rho^2 + a^2} \rho^4 + \rho^6 m^3 \right. \\ & + 48a^6 m^2 \sqrt{\rho^2 + a^2} + 24a^8 m - a^2 \rho^6 m - 7a^4 \rho^4 m + 18a^6 \rho^2 m + 32a^6 m^3 \\ & + 4a^2 \rho^4 m^3) (10a^6 m + a^6 \sqrt{\rho^2 + a^2} + 3a^4 \sqrt{\rho^2 + a^2} \rho^2 + 80a^4 m^3 + 28a^4 m \rho^2 \\ & + 40a^4 \sqrt{\rho^2 + a^2} m^2 + 3a^2 \sqrt{\rho^2 + a^2} \rho^4 + 32a^2 m^5 + 26a^2 m \rho^4 + 112a^2 \rho^2 m^3 \\ & + 64a^2 \sqrt{\rho^2 + a^2} m^2 \rho^2 + 80a^2 \sqrt{\rho^2 + a^2} m^4 + 24 \sqrt{\rho^2 + a^2} m^2 \rho^4 + 8\rho^6 m \\ & + 8a^4 \rho^2 m^3 + \sqrt{\rho^2 + a^2} \rho^6 + 16 \sqrt{\rho^2 + a^2} m^4 \rho^2 + 32\rho^4 m^3) \Big)^{1/2} (\rho^2 + a^2) a^2 / \\ & \left( 4 \sqrt{2} (4a^8 \sqrt{\rho^2 + a^2} + 2a^6 \sqrt{\rho^2 + a^2} \rho^2 - 2a^4 \sqrt{\rho^2 + a^2} \rho^4 + \rho^6 m^3 \right. \\ & + 48a^6 m^2 \sqrt{\rho^2 + a^2} + 24a^8 m - a^2 \rho^6 m - 7a^4 \rho^4 m + 18a^6 \rho^2 m \\ & + 32a^6 m^3 \rho^2 m^3 + 8a^4 + 4a^2 \rho^4 m^3) \rho \Big) \end{aligned} \quad (15)$$

Using this form for  $\mu$  and  $\rho = n a$ , as before, we get

$$\begin{aligned} g_{\phi\phi} = & \left[ -2m^2(n^4 - n^2 + 4) + m^2 \sqrt{n^2 + 1}(n^4 + 9n^2 - 8)a + 2(2n^4 + n^2 - 1)a^2 \right], \\ & / \left[ 2(m^2(n^4 + 8) - 2m \sqrt{n^2 + 1}(n^2 - 4)a - (n^2 + 1)(n^2 - 1)a^2) \right]. \end{aligned} \quad (16)$$

Also in this case, we can find  $n$ , a distance  $a$ , and a mass  $m$  such that  $g_{\phi\phi} > 0$ . In particular, we find  $a$  and  $m$  such that  $g_{\phi\phi} > 0$  for  $n > \sqrt{2}$  and  $C m < a < D m$ , where

$$\begin{aligned} C = & \frac{n^2 \sqrt{n^4 + 50n^2 + 17} - n^2 - 9n + 8}{2 \sqrt{n^2 + 1}(2n^2 - 1)}, \\ D = & \frac{n^2 - 4 - n^2 \sqrt{n^2 + 1}}{\sqrt{n^2 + 1}(2 - n^2)}. \end{aligned}$$

For example, we have CTGs for  $n = 1.5$  and  $0.3482 m < a < 9.4644 m$ , for  $n = 2$  and  $0.2725 m < a < 1.5491 m$ , for  $n = 5$  and  $0.2456 m < a < 0.8652 m$ , and for  $n = 10$  and  $0.1704 m < a < 0.9127 m$ .

A generic CTC  $\gamma$  satisfies the system of equations given by

$$\frac{D}{ds} \dot{X}^\mu = F^\mu(X), \quad (17)$$

where  $\frac{D}{ds} W^\alpha$  is the covariant derivative of the vector field  $W^\alpha$  along  $\gamma(s)$  and  $F^\mu$  is an external force per unit of mass.

Let  $\tilde{\gamma}$  be the curve obtained from  $\gamma$  after a perturbation  $\xi$ , i.e.,  $\tilde{X}^\mu = X^\mu + \xi^\mu$ . The system of differential equation satisfied by the perturbation  $\xi$  is [10] [11],

$$\frac{d^2 \xi^\alpha}{ds^2} + 2\Gamma_{\beta\mu}^\alpha \delta u^\beta u^\mu + \Gamma_{\beta\mu,\lambda}^\alpha \xi^\lambda u^\beta u^\mu = F_{,\lambda}^\alpha \xi^\lambda. \quad (18)$$

For the closed curve (6) the system (18), for case where (12) holds,  $z = z_0 = 0$ , and  $\mu$ ,  $a$ ,  $\rho$ , and  $m$  are related in such a way that  $a^\rho = 0$  ( $F^\alpha = 0$ ), reduces to

$$\begin{aligned}\ddot{\xi}^0 + k_1 \dot{\xi}^1 &= 0, \\ \ddot{\xi}^1 + k_2 \dot{\xi}^0 + k_3 \dot{\xi}^1 &= 0, \\ \ddot{\xi}^2 + k_4 \dot{\xi}^1 &= 0, \\ \ddot{\xi}^3 + k_5 \dot{\xi}^3 &= 0,\end{aligned}\tag{19}$$

where  $k_1 = 2\Gamma_{21}^0 \dot{\phi}$ ,  $k_2 = 2\Gamma_{20}^1 \dot{\phi}$ ,  $k_3 = \Gamma_{22,1}^1 \dot{\phi}^2$ ,  $k_4 = 2\Gamma_{21}^2 \dot{\phi}$ ,  $k_5 = \Gamma_{22,3}^3 \dot{\phi}^2$ . The condition for  $\gamma$  to be timelike, when it is parametrized by the proper time  $s$ , is  $\dot{X}^\mu \dot{X}_\mu = 1$ , where the overdot indicates derivation with respect to  $s$ . For the curve  $\gamma$  this last condition gives us

$$\dot{\phi}^2 = \frac{f}{\rho^2(f^2 \rho^2 w^2 - 1)}.\tag{20}$$

The solution of (19) is

$$\begin{aligned}\xi^0 &= -k_1(c_3 \sin(\varpi s + c_4)/\varpi + \lambda s) + c_1 s + c_5, \\ \xi^1 &= c_3 \cos(\varpi s + c_4) + \lambda, \\ \xi^2 &= -k_4(c_3 \sin(\varpi s + c_4)/\varpi + \lambda s) + c_2 s + c_6, \\ \xi^3 &= c_7 \cos(\sqrt{k_5} s + c_8),\end{aligned}\tag{21}$$

where  $c_i$ ,  $i = 1, \dots, 8$  are integration constants,  $\varpi = \sqrt{k_3 - k_1 k_2}$ , and  $\lambda = k_2 c_1$ . The explicit form of  $\varpi$  and  $k_5$  depending on the distance  $a$ , the mass  $m$  and the parameter  $n$  are cumbersome; they will be presented elsewhere.

To describe the regions where the CTG are linearly stable, first, we fix  $n$  ( $\rho = n a$ ) and then we find  $a$  and  $m$  such that  $\varpi^2 > 0$  and  $k_5 > 0$ . When the TLS is between the sources, the closed curve  $\gamma$ , satisfying (12),  $z = z_0 = 0$ , and (13) is a CTG linearly stable when  $n \in [1.5, 1.8]$ . For example, we have  $\varpi^2 > 0$  and  $k_5 > 0$  for  $n = 1.5$  and  $7.4788 m < a < 7.988 m$ , and for  $n = 1.8$  and  $0.1046 m < a < 0.5279 m$ .

For the case  $a = \rho = 1/\sqrt{2}$  given by Bonnor and Steadman [4], we have

$$\frac{\varpi^2}{k_5} = -2 \frac{(84m^4 + 235/2m^2 + 7/2 + 35m + 183m^3)(m+2)}{(1+2m)^2(116m + 33/2 + 53m^3 + 170m^2)}\tag{22}$$

that is always negative. In this case the CTG is not linearly stable. This is a particular case of  $a = \rho$ , where

$$\frac{\varpi^2}{k_5} = \frac{(14\alpha^4 + 70\sqrt{2}m\alpha^3 + 235m^2\alpha^2 + 183\sqrt{2}m^3\alpha + 84m^4)(4\alpha + m\sqrt{2})}{-(\alpha + m\sqrt{2})^2(66\alpha^3 + 232\sqrt{2}m\alpha^2 + 340m^2\alpha + 53m^3\sqrt{2})}\tag{23}$$

When we have two semi-infinite torsion TLS the closed curve  $\gamma$ , satisfying (12),  $z = z_0 = 0$ , and (15), is a CTG linearly stable when  $n \geq 1.5$ . For example, we have  $\varpi^2 > 0$  and  $k_5 > 0$  for  $n = 1.5$  and  $8.7913 m < a < 9.3079 m$ , for  $n = 2$  and  $0.4353 m < a < 1.2147 m$ , for  $n = 5$  and  $0.2904 m < a < 0.4576 m$ , and for  $n = 10$  and  $0.2743 m < a < 0.5014 m$ .

In summary, we found that there are linearly stable closed timelike geodesics in BW spacetime. There are two cases depending whether the torsion line singularity is

between the sources or not. We note that the existence of CTGs or their linear stability does not depend on the value of the mass parameter  $m$ .

The existence and stability of closed timelike geodesics present an evident challenge to the General Relativity Theory.

V.M.R. thanks Departamento de Matemática-UFV for giving the conditions to finish this work which was partially supported by PICDT-UFV/CAPES. PSL thanks the partial financial support of FAPESP and CNPq.

## References

- [1] K. Gödel, *Rev. Mod. Phys.*, **21**, 447 (1949)
- [2] W.J. van Stockum, *Proc. R. Soc. Edin.*, **A21**, 135(1937)
- [3] W.B. Bonnor, *Class. Quantum. Grav.*, **19**, 5951 (2002)
- [4] W.B. Bonnor and B.R. Steadman, *Gen. Rel. Grav.*, **37**, 1833 (2005)
- [5] Z. Perjés, *Phys. Rev. Letters*, **27**, 1668 (1971)
- [6] W. Israel and G.A. Wilson, *J. Math. Phys.*, **13**, 323 (1972)
- [7] W.B. Bonnor and J.P. Ward, *Commun. Math. Phys.*, **28**, 323 (1972)
- [8] P.S. Letelier, *Class. Quantum. Grav.*, **12**, 471 (1995)
- [9] P.S. Letelier and S.R. Oliveira, *Phys. Lett. A*, **238**, 101 (1998)
- [10] M.F. Shirokov, *Gen. Rel. Grav.*, **2**, 131 (1971)
- [11] V.M. Rosa and P.S. Letelier, Stability of Closed Timelike Curves in Gödel Universe, arXiv.org e-print archive: gr-qc/0703100.