Search for Excited Neutrinos at HERA

H1 Collaboration

Abstract

We present a search for excited neutrinos using e^-p data taken by the H1 experiment at HERA at a center-of-mass energy of 318 GeV with an integrated luminosity of 15 pb⁻¹. No evidence for excited neutrino production is found. Mass dependent exclusion limits are determined for the ratio of the coupling to the compositeness scale, f/Λ , independently of the relative couplings to the SU(2) and U(1) gauge bosons. These limits extend the excluded region to higher masses than has been possible in previous searches at other colliders.

Submitted to Physics Letters B

```
C. Adloff<sup>33</sup>, V. Andreev<sup>24</sup>, B. Andrieu<sup>27</sup>, T. Anthonis<sup>4</sup>, V. Arkadov<sup>35</sup>, A. Astvatsatourov<sup>35</sup>,
A. Babaev<sup>23</sup>, J. Bähr<sup>35</sup>, P. Baranov<sup>24</sup>, E. Barrelet<sup>28</sup>, W. Bartel<sup>10</sup>, P. Bate<sup>21</sup>, J. Becker<sup>37</sup>,
A. Beglarian<sup>34</sup>, O. Behnke<sup>13</sup>, C. Beier<sup>14</sup>, A. Belousov<sup>24</sup>, T. Benisch<sup>10</sup>, Ch. Berger<sup>1</sup>,
T. Berndt<sup>14</sup>, J.C. Bizot<sup>26</sup>, J. Boehme, V. Boudry<sup>27</sup>, W. Braunschweig<sup>1</sup>, V. Brisson<sup>26</sup>,
H.-B. Bröker<sup>2</sup>, D.P. Brown<sup>10</sup>, W. Brückner<sup>12</sup>, D. Bruncko<sup>16</sup>, J. Bürger<sup>10</sup>, F.W. Büsser<sup>11</sup>,
A. Bunyatyan<sup>12,34</sup>, A. Burrage<sup>18</sup>, G. Buschhorn<sup>25</sup>, L. Bystritskaya<sup>23</sup>, A.J. Campbell<sup>10</sup>,
J. Cao<sup>26</sup>, S. Caron<sup>1</sup>, F. Cassol-Brunner<sup>22</sup>, D. Clarke<sup>5</sup>, B. Clerbaux<sup>4</sup>, C. Collard<sup>4</sup>,
J.G. Contreras<sup>7,41</sup>, Y.R. Coppens<sup>3</sup>, J.A. Coughlan<sup>5</sup>, M.-C. Cousinou<sup>22</sup>, B.E. Cox<sup>21</sup>,
G. Cozzika<sup>9</sup>, J. Cvach<sup>29</sup>, J.B. Dainton<sup>18</sup>, W.D. Dau<sup>15</sup>, K. Daum<sup>33,39</sup>, M. Davidsson<sup>20</sup>,
B. Delcourt<sup>26</sup>, N. Delerue<sup>22</sup>, R. Demirchyan<sup>34</sup>, A. De Roeck<sup>10,43</sup>, E.A. De Wolf<sup>4</sup>,
C. Diaconu<sup>22</sup>, J. Dingfelder<sup>13</sup>, P. Dixon<sup>19</sup>, V. Dodonov<sup>12</sup>, J.D. Dowell<sup>3</sup>, A. Droutskoi<sup>23</sup>,
A. Dubak<sup>25</sup>, C. Duprel<sup>2</sup>, G. Eckerlin<sup>10</sup>, D. Eckstein<sup>35</sup>, V. Efremenko<sup>23</sup>, S. Egli<sup>32</sup>, R. Eichler<sup>36</sup>,
F. Eisele<sup>13</sup>, E. Eisenhandler<sup>19</sup>, M. Ellerbrock<sup>13</sup>, E. Elsen<sup>10</sup>, M. Erdmann<sup>10,40,e</sup>, W. Erdmann<sup>36</sup>,
P.J.W. Faulkner<sup>3</sup>, L. Favart<sup>4</sup>, A. Fedotov<sup>23</sup>, R. Felst<sup>10</sup>, J. Ferencei<sup>10</sup>, S. Ferron<sup>27</sup>,
M. Fleischer<sup>10</sup>, Y.H. Fleming<sup>3</sup>, G. Flügge<sup>2</sup>, A. Fomenko<sup>24</sup>, I. Foresti<sup>37</sup>, J. Formánek<sup>30</sup>,
G. Franke<sup>10</sup>, E. Gabathuler<sup>18</sup>, K. Gabathuler<sup>32</sup>, J. Garvey<sup>3</sup>, J. Gassner<sup>32</sup>, J. Gayler<sup>10</sup>,
R. Gerhards<sup>10</sup>, C. Gerlich<sup>13</sup>, S. Ghazaryan<sup>4,34</sup>, L. Goerlich<sup>6</sup>, N. Gogitidze<sup>24</sup>, M. Goldberg<sup>28</sup>,
C. Grab<sup>36</sup>, H. Grässler<sup>2</sup>, T. Greenshaw<sup>18</sup>, G. Grindhammer<sup>25</sup>, T. Hadig<sup>13</sup>, D. Haidt<sup>10</sup>,
L. Hajduk<sup>6</sup>, J. Haller<sup>13</sup>, W.J. Haynes<sup>5</sup>, B. Heinemann<sup>18</sup>, G. Heinzelmann<sup>11</sup>,
R.C.W. Henderson<sup>17</sup>, S. Hengstmann<sup>37</sup>, H. Henschel<sup>35</sup>, R. Heremans<sup>4</sup>, G. Herrera<sup>7,44</sup>,
I. Herynek<sup>29</sup>, M. Hildebrandt<sup>37</sup>, M. Hilgers<sup>36</sup>, K.H. Hiller<sup>35</sup>, J. Hladký<sup>29</sup>, P. Höting<sup>2</sup>,
D. Hoffmann<sup>22</sup>, R. Horisberger<sup>32</sup>, S. Hurling<sup>10</sup>, M. Ibbotson<sup>21</sup>, C. İşsever<sup>7</sup>, M. Jacquet<sup>26</sup>,
M. Jaffre<sup>26</sup>, L. Janauschek<sup>25</sup>, X. Janssen<sup>4</sup>, V. Jemanov<sup>11</sup>, L. Jönsson<sup>20</sup>, C. Johnson<sup>3</sup>,
D.P. Johnson<sup>4</sup>, M.A.S. Jones<sup>18</sup>, H. Jung<sup>20,10</sup>, D. Kant<sup>19</sup>, M. Kapichine<sup>8</sup>, M. Karlsson<sup>20</sup>,
O. Karschnick<sup>11</sup>, F. Keil<sup>14</sup>, N. Keller<sup>37</sup>, J. Kennedy<sup>18</sup>, I.R. Kenyon<sup>3</sup>, S. Kermiche<sup>22</sup>,
C. Kiesling<sup>25</sup>, P. Kjellberg<sup>20</sup>, M. Klein<sup>35</sup>, C. Kleinwort<sup>10</sup>, T. Kluge<sup>1</sup>, G. Knies<sup>10</sup>, B. Koblitz<sup>25</sup>,
S.D. Kolya<sup>21</sup>, V. Korbel<sup>10</sup>, P. Kostka<sup>35</sup>, S.K. Kotelnikov<sup>24</sup>, R. Koutouev<sup>12</sup>, A. Koutov<sup>8</sup>,
H. Krehbiel<sup>10</sup>, J. Kroseberg<sup>37</sup>, K. Krüger<sup>10</sup>, A. Küpper<sup>33</sup>, T. Kuhr<sup>11</sup>, T. Kurča<sup>16</sup>, R. Lahmann<sup>10</sup>,
D. Lamb<sup>3</sup>, M.P.J. Landon<sup>19</sup>, W. Lange<sup>35</sup>, T. Laštovička<sup>30,35</sup>, P. Laycock<sup>18</sup>, E. Lebailly<sup>26</sup>,
A. Lebedev<sup>24</sup>, B. Leißner<sup>1</sup>, R. Lemrani<sup>10</sup>, V. Lendermann<sup>7</sup>, S. Levonian<sup>10</sup>, M. Lindstroem<sup>20</sup>,
B. List<sup>36</sup>, E. Lobodzinska<sup>10,6</sup>, B. Lobodzinski<sup>6,10</sup>, A. Loginov<sup>23</sup>, N. Loktionova<sup>24</sup>,
V. Lubimov<sup>23</sup>, S. Lüders<sup>36</sup>, D. Lüke<sup>7,10</sup>, L. Lytkin<sup>12</sup>, H. Mahlke-Krüger<sup>10</sup>, N. Malden<sup>21</sup>,
E. Malinovski<sup>24</sup>, I. Malinovski<sup>24</sup>, R. Maraček<sup>25</sup>, P. Marage<sup>4</sup>, J. Marks<sup>13</sup>, R. Marshall<sup>21</sup>,
H.-U. Martyn<sup>1</sup>, J. Martyniak<sup>6</sup>, S.J. Maxfield<sup>18</sup>, D. Meer<sup>36</sup>, A. Mehta<sup>18</sup>, K. Meier<sup>14</sup>,
A.B. Meyer<sup>11</sup>, H. Meyer<sup>33</sup>, J. Meyer<sup>10</sup>, P.-O. Meyer<sup>2</sup>, S. Mikocki<sup>6</sup>, D. Milstead<sup>18</sup>,
T. Mkrtchyan<sup>34</sup>, R. Mohr<sup>25</sup>, S. Mohrdieck<sup>11</sup>, M.N. Mondragon<sup>7</sup>, F. Moreau<sup>27</sup>, A. Morozov<sup>8</sup>,
J.V. Morris<sup>5</sup>, K. Müller<sup>37</sup>, P. Murín<sup>16,42</sup>, V. Nagovizin<sup>23</sup>, B. Naroska<sup>11</sup>, J. Naumann<sup>7</sup>,
Th. Naumann<sup>35</sup>, G. Nellen<sup>25</sup>, P.R. Newman<sup>3</sup>, T.C. Nicholls<sup>5</sup>, F. Niebergall<sup>11</sup>, C. Niebuhr<sup>10</sup>,
O. Nix<sup>14</sup>, G. Nowak<sup>6</sup>, J.E. Olsson<sup>10</sup>, D. Ozerov<sup>23</sup>, V. Panassik<sup>8</sup>, C. Pascaud<sup>26</sup>, G.D. Patel<sup>18</sup>,
M. Peez<sup>22</sup>, E. Perez<sup>9</sup>, J.P. Phillips<sup>18</sup>, D. Pitzl<sup>10</sup>, R. Pöschl<sup>26</sup>, I. Potachnikova<sup>12</sup>, B. Povh<sup>12</sup>,
K. Rabbertz<sup>1</sup>, G. Rädel<sup>1</sup>, J. Rauschenberger<sup>11</sup>, P. Reimer<sup>29</sup>, B. Reisert<sup>25</sup>, D. Reyna<sup>10</sup>,
C. Risler<sup>25</sup>, E. Rizvi<sup>3</sup>, P. Robmann<sup>37</sup>, R. Roosen<sup>4</sup>, A. Rostovtsev<sup>23</sup>, S. Rusakov<sup>24</sup>, K. Rybicki<sup>6</sup>,
D.P.C. Sankey<sup>5</sup>, J. Scheins<sup>1</sup>, F.-P. Schilling<sup>10</sup>, P. Schleper<sup>10</sup>, D. Schmidt<sup>33</sup>, D. Schmidt<sup>10</sup>,
S. Schmidt<sup>25</sup>, S. Schmitt<sup>10</sup>, M. Schneider<sup>22</sup>, L. Schoeffel<sup>9</sup>, A. Schöning<sup>36</sup>, T. Schörner<sup>25</sup>,
V. Schröder<sup>10</sup>, H.-C. Schultz-Coulon<sup>7</sup>, C. Schwanenberger<sup>10</sup>, K. Sedlák<sup>29</sup>, F. Sefkow<sup>37</sup>,
V. Shekelyan<sup>25</sup>, I. Sheviakov<sup>24</sup>, L.N. Shtarkov<sup>24</sup>, Y. Sirois<sup>27</sup>, T. Sloan<sup>17</sup>, P. Smirnov<sup>24</sup>,
```

- Y. Soloviev²⁴, D. South²¹, V. Spaskov⁸, A. Specka²⁷, H. Spitzer¹¹, R. Stamen⁷, B. Stella³¹,
- J. Stiewe¹⁴, U. Straumann³⁷, M. Swart¹⁴, M. Taševský²⁹, V. Tchernyshov²³,
- S. Tchetchelnitski²³, G. Thompson¹⁹, P.D. Thompson³, N. Tobien¹⁰, D. Traynor¹⁹, P. Truöl³⁷,
- G. Tsipolitis^{10,38}, I. Tsurin³⁵, J. Turnau⁶, J.E. Turney¹⁹, E. Tzamariudaki²⁵, S. Udluft²⁵,
- M. Urban³⁷, A. Usik²⁴, S. Valkár³⁰, A. Valkárová³⁰, C. Vallée²², P. Van Mechelen⁴,
- S. Vassiliev⁸, Y. Vazdik²⁴, A. Vichnevski⁸, K. Wacker⁷, R. Wallny³⁷, B. Waugh²¹, G. Weber¹¹,
- M. Weber¹⁴, D. Wegener⁷, C. Werner¹³, M. Werner¹³, N. Werner³⁷, G. White¹⁷, S. Wiesand³³,
- T. Wilksen¹⁰, M. Winde³⁵, G.-G. Winter¹⁰, Ch. Wissing⁷, M. Wobisch¹⁰, E.-E. Woehrling³,
- E. Wünsch¹⁰, A.C. Wyatt²¹, J. Žáček³⁰, J. Zálešák³⁰, Z. Zhang²⁶, A. Zhokin²³, F. Zomer²⁶,
- J. Zsembery⁹, and M. zur Nedden¹⁰

- ⁴ Inter-University Institute for High Energies ULB-VUB, Brussels; Universitaire Instelling Antwerpen, Wilrijk; Belgium^c
- ⁵ Rutherford Appleton Laboratory, Chilton, Didcot, UK^b
- ⁶ Institute for Nuclear Physics, Cracow, Poland^d
- ⁷ Institut für Physik, Universität Dortmund, Dortmund, Germany^a
- ⁸ Joint Institute for Nuclear Research, Dubna, Russia
- ⁹ CEA, DSM/DAPNIA, CE-Saclay, Gif-sur-Yvette, France
- ¹⁰ DESY, Hamburg, Germany
- ¹¹ II. Institut für Experimentalphysik, Universität Hamburg, Hamburg, Germany^a
- 12 Max-Planck-Institut für Kernphysik, Heidelberg, Germany
- ¹³ Physikalisches Institut, Universität Heidelberg, Heidelberg, Germany^a
- ¹⁴ Kirchhoff-Institut für Physik, Universität Heidelberg, Heidelberg, Germany^a
- ¹⁵ Institut für experimentelle und Angewandte Physik, Universität Kiel, Kiel, Germany
- ¹⁶ Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovak Republic^{e,f}
- ¹⁷ School of Physics and Chemistry, University of Lancaster, Lancaster, UK^b
- 18 Department of Physics, University of Liverpool, Liverpool, UK^b
- ¹⁹ Queen Mary and Westfield College, London, UK^b
- ²⁰ Physics Department, University of Lund, Lund, Sweden^g
- 21 Physics Department, University of Manchester, Manchester, UK^b
- ²² CPPM, CNRS/IN2P3 Univ Mediterranee, Marseille France
- 23 Institute for Theoretical and Experimental Physics, Moscow, Russia l
- ²⁴ Lebedev Physical Institute, Moscow, Russia^{e,h}
- ²⁵ Max-Planck-Institut für Physik, München, Germany
- ²⁶ LAL, Université de Paris-Sud, IN2P3-CNRS, Orsay, France
- ²⁷ LPNHE, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, France
- ²⁸ LPNHE, Universités Paris VI and VII, IN2P3-CNRS, Paris, France
- 29 Institute of Physics, Academy of Sciences of the Czech Republic, Praha, Czech Republic e,i
- ³⁰ Faculty of Mathematics and Physics, Charles University, Praha, Czech Republic^{e,i}
- ³¹ Dipartimento di Fisica Università di Roma Tre and INFN Roma 3, Roma, Italy
- ³² Paul Scherrer Institut, Villigen, Switzerland
- ³³ Fachbereich Physik, Bergische Universität Gesamthochschule Wuppertal, Wuppertal, Germany

¹ I. Physikalisches Institut der RWTH, Aachen, Germany^a

² III. Physikalisches Institut der RWTH, Aachen, Germany^a

³ School of Physics and Space Research, University of Birmingham, Birmingham, UK^b

- ³⁴ Yerevan Physics Institute, Yerevan, Armenia
- ³⁵ DESY, Zeuthen, Germany
- ³⁶ Institut für Teilchenphysik, ETH, Zürich, Switzerland⁹
- ³⁷ Physik-Institut der Universität Zürich, Zürich, Switzerland^j
- ³⁸ Also at Physics Department, National Technical University, Zografou Campus, GR-15773 Athens, Greece
- ³⁹ Also at Rechenzentrum, Bergische Universität Gesamthochschule Wuppertal, Germany
- ⁴⁰ Also at Institut für Experimentelle Kernphysik, Universität Karlsruhe, Karlsruhe, Germany
- ⁴¹ Also at Dept. Fis. Ap. CINVESTAV, Mérida, Yucatán, México^k
- ⁴² Also at University of P.J. Šafárik, Košice, Slovak Republic
- ⁴³ Also at CERN, Geneva, Switzerland
- ⁴⁴ Also at Dept. Fis. CINVESTAV, México City, México^k
- ^a Supported by the Bundesministerium für Bildung und Forschung, FRG, under contract numbers 05 H1 1GUA /1, 05 H1 1PAA /1, 05 H1 1PAB /9, 05 H1 1PEA /6, 05 H1 1VHA /7 and 05 H1 1VHB /5
- ^b Supported by the UK Particle Physics and Astronomy Research Council, and formerly by the UK Science and Engineering Research Council
- ^c Supported by FNRS-NFWO, IISN-IIKW
- ^d Partially Supported by the Polish State Committee for Scientific Research, grant no. 2P0310318 and SPUB/DESY/P03/DZ-1/99, and by the German Federal Ministry of Education and Research (BMBF)
- ^e Supported by the Deutsche Forschungsgemeinschaft
- ^f Supported by VEGA SR grant no. 2/1169/2001
- ^g Supported by the Swedish Natural Science Research Council
- ^h Supported by Russian Foundation for Basic Research grant no. 96-02-00019
- ⁱ Supported by the Ministry of Education of the Czech Republic under the projects INGO-LA116/2000 and LN00A006, by GA AVČR grant no B1010005 and by GAUK grant no 173/2000
- $^{\it j}$ Supported by the Swiss National Science Foundation
- ^k Supported by CONACyT
- ¹ Partially Supported by Russian Foundation for Basic Research, grant no. 00-15-96584

The discovery of excited states of quarks or leptons, as predicted by compositeness models [1, 2], would supply convincing evidence for a new substructure of matter. Electron-proton interactions at very high energies provide ideal conditions to look for excited states of first generation fermions. In particular a magnetic type coupling of the electron would allow for the production of single excited neutrinos (ν^*) through t-channel W boson exchange. The phenomenology of this process is described in [3–5]. In this paper we present a search for ν^* production followed by the electroweak decays $\nu^* \to \nu \gamma$, $\nu^* \to eW$ or $\nu^* \to \nu Z$. The analysis makes use of 15 pb⁻¹ of e^-p data with an electron beam energy of 27.6 GeV and a proton beam energy of 920 GeV collected in 1998 and 1999 with the H1 experiment at HERA. Compared to previous H1 results from e^-p collisions [6] the analysis benefits from an increase in luminosity by a factor of 30 and by an increase of the center-of-mass energy from 300 GeV to 318 GeV. Furthermore, it also improves significantly on results derived from larger luminosity of e^+p data at a center-of-mass energy of 300 GeV [5], due to a much larger cross-section for ν^* production in e^-p scattering as compared to the e^+p case. At a ν^* mass of 200 GeV the ratio of those cross-sections is of the order of 100. Other searches for excited neutrinos have recently been presented by ZEUS [7] and by LEP experiments [8–10].

The production cross section and the decays of excited neutrinos can be calculated using an effective Lagrangian [3,4] which depends on a compositeness mass scale Λ and on form factors (reduced here to parameters) f_s , f and f' allowing for the composite lepton to have arbitrary coupling strengths associated to the gauge groups SU(3), SU(2) and U(1). The excited neutrino can decay into the electroweak gauge bosons via $\nu^* \to \nu \gamma$, $\nu^* \to eW$ and $\nu^* \to \nu Z$. As shown in [4], the decay width of the ν^* is a function of f, f' and Λ and can reach, for part of the accessible mass range, a few hundred GeV, much larger than the detector resolution (10 GeV). For smaller decay widths (corresponding to masses below 200 GeV) the narrow width approximation (NWA) is applicable, in which the assumption is made that the production and decay of a particle factorize. In this range the COMPOS [11] generator is used for cross-section calculations. For masses beyond 200 GeV the full cross-section for ν^* production and decay is evaluated with COMPHEP [12] using the Lagrangian given in [4]. In the overlap region the compatibility of COMPOS and COMPHEP has been verified.

The detector components of the H1 experiment [13] most relevant for this analysis are shortly described in the following. The interaction region is surrounded by a system of drift and proportional chambers covering the polar angular range 1 0 $< \theta < 176^o$. The tracking system is placed inside a finely segmented liquid argon (LAr) calorimeter covering the polar angular range 0 $< \theta < 154^o$ [14]. Energy resolutions of $\sigma_E/E \simeq 12\%/\sqrt{E(GeV)} \oplus 1\%$ for electrons and $\sigma_E/E \simeq 50\%/\sqrt{E(GeV)} \oplus 2\%$ for hadrons have been obtained in test beam measurements [15,16]. The tracking system and calorimeters are surrounded by a superconducting solenoid and an iron yoke instrumented with streamer tubes. Leakage of hadronic showers outside the calorimeter is measured by analogue charge sampling of the streamer tubes with a resolution [17] of $\sigma_E/E \simeq 100\%/\sqrt{E(GeV)}$.

For the decays of the heavy gauge bosons only the dominating hadronic modes are considered. The selection of ν^* events is based on photon or electron identification, missing transverse energy $(E_t^{\rm miss})$ measurement and the requirement for jets, depending on the channel investigated. Electromagnetic clusters are requested to have more than 95% of their energy in the electromagnetic part of the calorimeter and to be isolated from other particles [18]. They are further

¹The polar angle θ is measured with respect to the proton beam direction (+z).

differentiated into electron and photon candidates using the data of associated charged tracks. Jets with a minimum transverse momentum of 5 GeV are reconstructed from the hadronic final state in the LAr calorimeter using a cone algorithm, adapted from the LUCELL scheme in the JETSET package [19].

Background not related to e^-p collisions is rejected by requiring a primary interaction vertex reconstructed within ± 35 cm around the nominal vertex value, by using topological filters and by requiring the event time to coincide with the time of the bunch crossing. Standard Model (SM) backgrounds which could mimic the ν^* signatures are Neutral Current Deep Inelastic Scattering (NC DIS), Charged Current Deep Inelastic Scattering (CC DIS) and photoproduction processes (γp). The background expectation from NC DIS and CC DIS is calculated using the event generator DJANGO [20] which includes first order QED corrections based on HERACLES [21] and QCD radiation based on the Colour Dipole Model [22]. Parton densities are taken from the MRST parameterization [23] which includes constraints from DIS measurements at HERA up to a squared momentum transfer $Q^2 = 5000 \, \mathrm{GeV}^2$ [24–27]. The hadronisation process is simulated in the Lund string fragmentation scheme using JETSET [19]. Direct and resolved γp processes, including prompt photon production, are simulated with PYTHIA [28]. All Monte Carlo samples are subject to a full simulation of the H1 detector.

The $\nu^* \to \nu \gamma$ channel is characterized by missing transverse energy and by an electromagnetic cluster in the calorimeter. The main SM background is expected from CC DIS. Events are selected with an identified photon of transverse momentum (P_t) greater than 16 GeV and total missing transverse energy E_t^{miss} greater than 16 GeV. To reject NC DIS background where the scattered electron (sometimes misinterpreted as a photon) is preferably scattered through small angles, photon candidates are accepted in the forward region of the detector only ($\theta < 1.8 \text{ rad}$). For $E_t^{\text{miss}} > 30$ GeV electromagnetic clusters in the very forward region ($\theta < 1$ rad) are accepted even if they are linked to a track. In this particular region the conversion rate $\gamma \to ee$ and also the number of randomly assigned tracks is expected to be higher due to the high multiplicity of hadronic charged particles from jets. In order to be able to reconstruct the event vertex position from charged particles, the event is required to contain a jet. To further suppress background from events in which hadronic energy fluctuations of jets result in a measured missing transverse momentum, the missing transverse momentum vector of the event is required to have a component of more than 8 GeV perpendicular to the required jet. To reduce the influence of photons coming from QED radiation along the quark line, the jet must be isolated from the photon in azimuth ($\Delta \varphi(\text{jet}, \gamma) > 0.35 \text{ rad}$). In total 2 events are found in this channel for an expected background of 3.0 ± 0.2 (stat.) ± 1.2 (syst.) events. The different sources of systematic errors are discussed below. The background is composed of 2.7 events from CC DIS and 0.3 events from NC DIS with negligible contributions from γp . The resulting selection efficiency ranges between 40% and 65%.

The $\nu^* \to eW_{\to q\bar q}$ channel is characterized by an electromagnetic cluster with an associated track and two jets. The main SM background is NC DIS as photoproduction events do not yield a significant rate of electrons with high transverse momentum (P_t^{ele}) . A cut $P_t^{ele} > 12.5~{\rm GeV}$ is chosen. At very high transverse momentum $P_t^{ele} > 85~{\rm GeV}$ the background from NC DIS is low and no further cuts are applied. In the range 65 GeV $< P_t^{ele} < 85~{\rm GeV}$ two jets with an invariant mass $M_{\rm jj} > 50~{\rm GeV}$ are required as expected for a hadronic W decay. In the range 12.5 GeV $< P_t^{ele} < 65~{\rm GeV}$ three jets are required, where the third jet is supposed to

originate from the quark struck in the W proton interaction. A cut on the electron polar angle is applied which depends on P_t^{ele} and ranges from $\theta_e < 1.20$ to $\theta_e < 2.25$ rad. To reconstruct a W candidate, the dijet-pair with invariant mass closest to the nominal W boson mass is accepted in the range 65 GeV $< M_{\rm jj} < 87$ GeV. The two jets chosen as the W candidate are ordered by their transverse momentum such that $P_t^{\rm jet1} > P_t^{\rm jet2}$. As for many background events jet 2 points in the very forward direction, an additional cut on its polar angle, $\theta^{\rm jet2} > 0.2$ rad, is applied if the transverse momentum of this jet is lower than 30 GeV. After these cuts, 6 events remain in the data. The expected background is $7.0 \pm 0.6 \pm 1.4$ events mainly from NC DIS with negligible contributions from CC DIS and γp . The resulting signal efficiency ranges between 30% and 50%.

The $\nu^* \to \nu Z_{\to q\bar q}$ channel is characterized by two jets and missing transverse energy $E_t^{\rm miss}$. The main background is expected from CC DIS with a moderate contribution from γp , whereas the NC DIS contribution is sufficiently suppressed for large $E_t^{\rm miss}$. A cut $E_t^{\rm miss} > 10$ GeV is chosen. At $E_t^{\rm miss} > 40$ GeV only two jets are required, while at lower $E_t^{\rm miss}$ a third jet is required and events with an electron or photon candidate are rejected. A Z candidate is reconstructed from the combination of 2 jets with invariant mass closest to the nominal Z boson mass provided this mass is greater than 76 GeV. Again these two jets are ordered in P_t . To suppress further the background from CC DIS a cut on the polar angle $\theta^{jet2} > 0.15$ rad is applied. In the region of relatively low missing transverse momentum 10 GeV $< E_t^{\rm miss} < 20$ GeV an additional cut is applied on the transverse momentum of jet 1 ($P_t^{jet1} > 50$ GeV). With these criteria, one candidate event is found in the data, with an expected background of $3.7 \pm 0.2 \pm 0.9$ events. The background consists mainly of CC DIS (2.3 events) and γp (1.3 events). The resulting signal efficiency is above 60% for masses greater than 150 GeV.

Contributions to the systematic uncertainties come from the limited knowledge of the absolute energy scale of the calorimeter and missing higher order corrections in the event generators which are used for the background estimation. The uncertainties of the electromagnetic energy scale amount to 0.7% in the central part of the detector and up to 3% in the forward region. For the hadronic part an uncertainty of 4% is assigned. For the $\nu^* \to \nu \gamma$ channel the lack of QED radiation from the quark line in the DJANGO generator leads to an uncertainty of the CC DIS background expectation which, after applying the $\Delta \varphi(\text{jet},\gamma) > 0.35$ rad cut, is limited to 40% as estimated using [29]. For the $\nu^* \to eW_{\to q\bar{q}}$ and $\nu^* \to \nu Z_{\to q\bar{q}}$ channels the background normalization is varied by 15% to account for differences observed in particular for the 3-jets production between perturbative calculations of the order $O(\alpha_s^2)$ [30–32] and the parton shower approach. The statistical error of the Monte Carlo event samples is taken into account. Finally, the luminosity measurement leads to a normalization uncertainty of 2.25%.

In all three search channels the number of observed and expected events are in good agreement. Upper limits at 95% confidence level on the coupling f/Λ are thus derived as described in [5] following the Bayesian approach [33, 34]. The number of observed and expected events is counted within a sliding mass window which is adopted to the width of the expected excited neutrino signal. Systematic uncertainties are taken into account as in [5].

The resulting limits after combination of all decay channels are given as a function of the ν^* mass in Fig. 1, for the conventional assumptions f=-f' and f=+f'. Note that the decay $\nu^* \to \nu \gamma$ is forbidden for f=+f'. These results improve significantly our limits published earlier in e^-p [6] and e^+p [5] collisions and reach masses up to 240 GeV and couplings f/Λ of




Figure 1: Exclusion limits on the coupling f/Λ at 95% confidence level as a function of the mass of excited neutrinos with the assumptions (a) f=-f' and (b) f=+f'. Exclusion limits are given for H1 e^-p data (full line) with an integrated luminosity of 15 pb $^{-1}$, for H1 e^+p data [5] (dashed line) with an integrated luminosity of 37 pb $^{-1}$ and for L3 [10] (dotted line).

order O(1/100GeV). Using the assumption $f/\Lambda=\frac{1}{M_{\nu^*}}$ excited neutrinos with masses between 50 GeV and 150 GeV (100 GeV and 140 GeV) are excluded by the H1 analysis for f=-f' (f=+f').

Fig. 1 also shows for comparison results obtained by the L3 collaboration in e^+e^- collisions at centre of mass energies up to 202 GeV at LEP II [10]. The H1 limits are more stringent at high masses beyond the kinematic reach of LEP II.

Less model-dependent limits can be derived if arbitrary ratios f'/f are considered. Fig. 2 illustrates how the limits depend on this ratio for various ν^* mass hypothesis. By choosing the point with the worst limit for each mass hypothesis, limits have been derived which are no longer dependent on f'/f in the range -5< f'/f <5. The result is shown in Fig. 3. It deviates from the limits obtained assuming f = +f' only for high ν^* masses. Limits on single ν^* production independent of f'/f also have been shown previously by the OPAL collaboration [9].

In summary, using e^-p data a search for the production of excited neutrinos has been performed and no indication of a signal was found. New limits have been established as function of couplings and excited neutrino masses both for specific relations between the couplings (f = f' and f = -f') and independent of the ratio of f and f'. In comparison to previous analyses the data presented here restrict the existence of excited neutrinos for masses up to 240 GeV and to much smaller couplings.

We are grateful to the HERA machine group whose outstanding efforts have made and continue to make this experiment possible. We thank the engineers and technicians for their

Figure 2: Exclusion limits for excited neutrinos on the coupling f/Λ at 95% confidence level as a function of the value of f'/f. Each curve corresponds to a different ν^* mass. The circles indicate the maximum (worst limit) of each curve. The areas above the lines are excluded.





Figure 3: Exclusion limits on the coupling f/Λ at 95% confidence level as a function of the mass of excited neutrinos. All f'/f values in the interval [-5;+5] have been considered (see figure 2), so this limit is independent of the relation between f and f' in that interval. The area above the line is excluded.



work in constructing and now maintaining the H1 detector, our funding agencies for financial support, the DESY technical staff for continual assistance and the DESY directorate for the hospitality which they extend to the non DESY members of the collaboration.

References

- [1] H. Harari, Phys. Rept. **104** (1984) 159.
- [2] F. Boudjema, A. Djouadi and J. L. Kneur, Z. Phys. C 57 (1993) 425, and references therein.
- [3] K. Hagiwara, D. Zeppenfeld and S. Komamiya, Z. Phys. C **29** (1985) 115.
- [4] U. Baur, M. Spira and P. M. Zerwas, Phys. Rev. D 42 (1990) 815.
- [5] C. Adloff *et al.* [H1 Collaboration], Eur. Phys. J. C **17** (2000) 567, [hep-ex/0007035].
- [6] T. Ahmed et al. [H1 Collaboration], Phys. Lett. B **340** (1994) 205.
- [7] S. Chekanov *et al.* [ZEUS Collaboration], DESY-01-132, Sept 2001, [hep-ex/0109018].
- [8] P. Abreu et al. [DELPHI Collaboration], Eur. Phys. J. C 8 (1999) 41, [hep-ex/9811005].
- [9] G. Abbiendi et al. [OPAL Collaboration], Eur. Phys. J. C 14 (2000) 73, [hep-ex/0001056].
- [10] M. Acciarri et al. [L3 Collaboration], Phys. Lett. B **502** (2001) 37, [hep-ex/0011068].
- [11] T. Köhler, "Hamburg 1991, Proceedings, Physics at HERA, vol. 3" 1526-1541.
- [12] A. Pukhov *et al.*, hep-ph/9908288.
- [13] I. Abt et al. [H1 Collaboration], Nucl. Instrum. Meth. A **386** (1997) 310 and 348.
- [14] B. Andrieu *et al.* [H1 Calorimeter Group Collaboration], Nucl. Instrum. Meth. A **336** (1993) 460.
- [15] B. Andrieu *et al.* [H1 Calorimeter Group Collaboration], Nucl. Instrum. Meth. A **350** (1994) 57.
- [16] B. Andrieu *et al.* [H1 Calorimeter Group Collaboration], Nucl. Instrum. Meth. A **336** (1993) 499.
- [17] W. Braunschweig et al. [H1 Collaboration], Nucl. Instrum. Meth. A 270 (1988) 334.
- [18] A. Schöning, "Untersuchung von Prozessen mit virtuellen und reellen W^{\pm} -Bosonen am H1-Detektor bei HERA", Thesis University of Hamburg, 1996.
- [19] T. Sjöstrand, hep-ph/9508391, LU-TP-95-20, CERN-TH-7112-93-REV, Aug 1995.
- [20] G. A. Schuler and H. Spiesberger, "Hamburg 1991, Proceedings, Physics at HERA, vol. 3" 1419-1432.
- [21] A. Kwiatkowski, H. Spiesberger and H. J. Möhring, Comput. Phys. Commun. **69** (1992) 155.
- [22] B. Andersson, G. Gustafson, L. Lönnblad and U. Pettersson, Z. Phys. C 43 (1989) 625.

- [23] A. D. Martin, R. G. Roberts, W. J. Stirling and R. S. Thorne, Eur. Phys. J. C 4 (1998) 463, [hep-ph/9803445].
- [24] S. Aid et al. [H1 Collaboration], Nucl. Phys. B 470, 3 (1996), [hep-ex/9603004].
- [25] C. Adloff et al. [H1 Collaboration], Nucl. Phys. B 497, 3 (1997), [hep-ex/9703012].
- [26] M. Derrick et al. [ZEUS Collaboration], Z. Phys. C 69, 607 (1996), [hep-ex/9510009].
- [27] M. Derrick et al. [ZEUS Collaboration], Z. Phys. C 72, 399 (1996), [hep-ex/9607002].
- [28] T. Sjöstrand, Comput. Phys. Commun. **82** (1994) 74.
- [29] T. Helbig, H. Spiesberger, Nucl. Phys. B **373**, 73 (1992).
- [30] T. Carli, hep-ph/9906541, Monte Carlo generators for HERA Physics, Proceedings of the Workshop 1998-1999, 185-194.
- [31] P. Bate, "High Transverse Momentum 2-jet and 3-jet Cross-section Measurements in Photoproduction", Thesis University of Manchester, 1999, http://www-h1.desy.de/publications/theses_list.html
- [32] C. Adloff *et al.* [H1 Collaboration], Phys. Lett. B **515** (2001) 17, [hep-ex/0106078].
- [33] R. M. Barnett *et al.* [Particle Data Group Collaboration], Phys. Rev. D **54** (1996) 1.
- [34] O. Helene, Nucl. Instrum. Meth. A **228** (1984) 120.