

GAUGE BOSON SELF COUPLINGS AND FOUR FERMION FINAL STATES AT LEP

PAOLO AZZURRI

Scuola Normale Superiore, Piazza dei Cavalieri 7, 56100 Pisa, Italy

Abstract

Four-fermion productions measured in the LEP2 data are reviewed. The total and differential cross-section yields represent the first clear evidence for the existence of gauge boson self couplings, in support of the non-abelian $SU(2) \otimes U(1)$ structure of the electroweak model, at the percent level.

1 Introduction




Figure 1: *Examples of triple and quartic gauge bosons self-couplings predicted by the $SU(2)$ part of the electroweak model.*

A striking feature of the Standard Model (SM) of electroweak interactions [1], is the presence of gauge bosons self-couplings (GC) due to

the non-abelian nature of the $SU(2)$ group leading to three-line (TGC) and four-line (QGC) boson vertices shown in Fig. 1.

Gauge self couplings influence directly the pair-production of weak bosons, and in particular of W-bosons, so that the LEP2 represents an ideal data sample to verify their structure, through the detection of the resulting four-fermion final states.

2 W-pair and Z-pair productions

The LEP2 data sample of 700pb^{-1} per experiment at $\sqrt{s}=161\text{--}207$ GeV allowed to collect about 10^4 W-pairs and 500 Z-pairs per experiment, identifying these events in all their visible final states. The combined total cross sections, measured as a function of the centre-of-mass energy are shown in Fig. 2. The overall precision of the measured productions rates is at the level of 1% for W-pairs and 5% for Z-pairs, and in agreement with the SM predictions. As it can be seen in Fig. 2, the W-pair cross section measurement represents alone a stunning proof of the presence of both the WWZ and $WW\gamma$ couplings dictated by the electroweak $SU(2)\otimes U(1)$ gauge structure.




Figure 2: Total W-pair and Z-pair cross-sections measured at LEP at $\sqrt{s}=161\text{--}207$ GeV, compared to SM predictions. The two dashed curves show the predicted W-pair cross section in the absence of the WWZ and $WW\gamma$ couplings.

The LEP2 W-pair sample has also proven valuable to provide

the first direct measurement of all leptonic and hadronic W decay branching ratios, resulting in a test of the lepton family universality of charged currents at the level of 1% ($g_\mu/g_e = 1.000 \pm 0.010$, $g_\tau/g_e = 1.026 \pm 0.014$, $g_\tau/g_\mu = 1.026 \pm 0.014$), and of the lepton-quark universality of charged currents at the level of 0.7% ($g_q/g_\ell = 1.010 \pm 0.007$).

3 Single W and single Z productions

Four-fermion final states arising from single W and Z productions have also been measured in the LEP2 data [2], and are shown in Fig. 3. Results are in agreement with SM expectations within the combined experimental precision of 7-8%.




Figure 3: Single W and single Z cross-sections measured at LEP at $\sqrt{s}=161\text{--}207$ GeV, compared to SM predictions.

4 Constraints on Gauge Self Couplings

The structure of charged TGC vertices WWZ and $WW\gamma$ have been studied using W-pair, single W and single γ events, and possible deviations from SM expectations have been searched for. The most common set of parameters to describe the two vertices is g_1^Z , λ_γ and κ_γ , which

conserve both C and P, and obey the SM gauge structure. The combined LEP2 data yields $g_1^Z = 0.991 \pm 0.022$, $\lambda_\gamma = -0.016 \pm 0.022$ and $\kappa_\gamma = 0.984 \pm 0.045$, therefore in agreement with the SM expectations $\lambda_\gamma = 0$ and $\kappa_\gamma = g_1^Z = 1$ at the 2-5% level [2].

Charged QGC as WWZZ, WWZ γ and WW $\gamma\gamma$ are predicted in the SM but are beyond the sensitivity of the LEP2 data. Larger anomalous contributions have been searched for in WW γ and $\nu\nu\gamma\gamma$ final states, leading to limits on the new physics scale of the anomalous contributions $\Lambda > 5-10$ GeV at 95% confidence level [2].

Neutral TGC as ZZZ, ZZ γ and Z $\gamma\gamma$ do not exist in the SM but possible contributions have been searched for in Z pair, Zee and Z γ events. No evidence for neutral TGC has been found and parameterizing the vertices with the h_i^V and f_i^V ($i = 1, 4$ and $V = \gamma, Z$) couplings, limits have been set at the level $h_i^V, f_i^V < 0.05-0.20$ at 95% confidence level [2].

Neutral QGC also do not exist in the SM, but possible contributions of a ZZ $\gamma\gamma$ vertex have been searched for in the LEP2 data. Here the study of ZZ γ and $\nu\nu\gamma\gamma$ events allows to set limits on the new physics scale of the anomalous contributions again with $\Lambda > 5-10$ GeV at 95% confidence level [2].

References

- [1] S.L. Glashow, *Nucl. Phys.* **22**, 579 (1961); A. Salam and J.C. Ward, *Phys. Lett.* **13**, 168 (1964); S. Weinberg, *Phys. Rev. Lett.* **19**, 1264 (1967); A. Salam, in *Elementary Particle Theory: Relativistic groups and Analyticity* (Nobel Symposium 8) p.367 (Stockholm, 1968).
- [2] LEP Collaborations ALEPH, DELPHI, L3, OPAL and Electroweak Working Group, *A Combination of Preliminary Electroweak Measurements and Constraints on the Standard Model*, hep-ex 0312023, and references therein.