

The combination of Wilson Loops into J^{PC} Operators in Lattice Pure Gauge Theory

Da Qing Liu², Ji Min Wu^{1, 2} and Ying Chen²

¹CCAST(World Lab.), P. O. Pox 8730, Beijing 100080, China

²Institute of High Energy Physics, Chinese Academy of Sciences,
P. O. Box 918-4, Beijing, 100039, P. R. China

August 23, 2019

Abstract

To calculate the mass of glueballs with quantum number J^{PC} in lattice pure gauge theory by using Wilson loop operators, we discuss the combination of Wilson loops into irreducible representation of O^{PC} group for arbitrary-link Wilson loops. We present a general computational procedure for this combination which is suitable for any finite group.

1 Introduction

To calculate the glueball mass spectra, one has to construct different J^{PC} glueball operators which act on vacuum to generate states with certain J^{PC} quantum number. These operators should be the correct linear combinations of Wilson loops and transform as certain irreducible representations R^{PC} of group O^{PC} if we also concern about space parity P and charge conjugate C with cubic group O on lattice. Berg and Billoire^[1] has discussed this problem and merged the problem of constructing J^{PC} glueball operators into the combination of Wilson loops with different lengths and types into irreducible representation R^{PC} of group O^{PC} . They also reveal the result of 4-link and 6-link operators^[1].

Therefore, the decomposition of the cubic group is the preliminary step to calculate glueball masses^[5]. On the contrary, since some representation matrices of the group O^{PC} is as large as 96×96 , the calculation is lengthy and some time impossible by hand. It is necessary to find a general and practically easy procedure for the combination of arbitrary-link operators into R^{PC} . We would like to introduce such a procedure here, which can be realized by computer and is also suitable for any finite group.

Section 2 gives some appointments in our calculation and section 3 presents our computational procedure. We present our results in section 4 and a short summary is in section 5. Appendix A shows some property of the cubic group. Appendix B is some lengthy results from our calculation.

2 Some Appointments

We use L-component vector $(f_1, f_2, \dots, f_L)$ ($f_i = \pm 1, \pm 2, \pm 3$), to represent L-link closed Wilson loop in which each component stands for the space direction of the link along the Wilson loop. For instance, we denote a 4-link Wilson operator

$$Tr(U_1(n)U_2(n + a\hat{e}_1)U_1^{-1}(n + a\hat{e}_2)U_2^{-1}(n))$$

by $(1, 2, -1, -2)$. Each vector forms a base of one representation. Therefore, it is obvious that under C transformation:

$$(f_1, f_2, \dots, f_L) \xrightarrow{C} (-f_L, \dots, -f_2, -f_1),$$

and under $M_g \in O^P$: $M_g(f_1, f_2, \dots, f_L) = (M_g f_1, M_g f_2, \dots, M_g f_L)$.

The cubic group O has 24 elements which may be divided into five classes of conjugate elements: E, C_2, C_3, C_4, C_2^2 . We choose the proper ones from each class of the conjugate elements which make the directions transformation as follows under their actions:

$$1 \xrightarrow{C_2} -1, \quad 2 \xrightarrow{C_2} -3, \quad 3 \xrightarrow{C_2} -2; \tag{1a}$$

$$1 \xrightarrow{C_3} -3, \quad 2 \xrightarrow{C_3} -1, \quad 3 \xrightarrow{C_3} 2; \tag{1b}$$

$$1 \xrightarrow{C_4} -3, \quad 2 \xrightarrow{C_4} 2, \quad 3 \xrightarrow{C_4} 1; \quad (1c)$$

$$1 \xrightarrow{C_4^2} -1, \quad 2 \xrightarrow{C_4^2} 2, \quad 3 \xrightarrow{C_4^2} -3. \quad (1d)$$

For generators^[2] C_3 and C_4 of the cubic group O , we appoint their representation matrices as:

$$D^{A_1}(C_4) = 1, \quad D^{A_1}(C_3) = 1; \quad (2a)$$

$$D^{A_2}(C_4) = -1, \quad D^{A_2}(C_3) = 1; \quad (2b)$$

$$D^E(C_4) = \frac{1}{2} \begin{pmatrix} 1 & 3 \\ 1 & -1 \end{pmatrix}, \quad D^E(C_3) = \frac{1}{2} \begin{pmatrix} -1 & -3 \\ 1 & -1 \end{pmatrix}; \quad (2c)$$

$$D^{T_1}(C_4) = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \quad D^{T_1}(C_3) = \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{pmatrix}; \quad (2d)$$

$$D^{T_2}(C_4) = \begin{pmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad D^{T_2}(C_3) = \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{pmatrix}; \quad (2e)$$

Including P and C, group O^{PC} has 4 generators. It is trivial to generalize the result of group O to group O^{PC} .

3 Computational Procedure

Let us consider L-component operators which are classified into N types. Type means assembly of operators with the same shape here(Any group element does not change the shape of the operator). We'll restore all the bases of each type into array A and adopt the computational procedure as follows:

1. To produce a new L-component vector standing for a closed Wilson loop which does not belong to A (by comparing it with the restored operators).
2. To apply 96 elements of group O^{PC} on this Wilson operator(named prototype) to generate all the bases which belong to the same type and append these bases to A. We also can get the representation matrix of these elements in these bases.
3. To construct new bases from the bases obtained from step 2 according to PC=++, - +, + - and - -. Meanwhile, actually, we get a transform matrix between old bases and new bases. Therefore, we get the new representation matrices which are diagonally respect to P and C by making a similar transformation for the old ones obtained from step 2.
4. For certain P and C, to construct new bases which are the combinations of the bases obtained from step 3 in order to decompose the representation matrices into irreducible representation R^{PC} . We get the multiplicity $m_{R^{PC}}$ of irreducible representation R^{PC} by our decomposing for this representation. It is the same as given by group theory:

$$m_{R^{PC}} = \frac{1}{96} \sum_K n_K \chi_K \chi_K^{R^{PC}}, \quad (3)$$

where K stands for the class, n_K is the element number of K^{th} class, χ_K is the character of K^{th} class in this representation, and $\chi_K^{R^{PC}}$ is the character of K^{th} class in the irreducible representation R^{PC} (see table 3);

5. Return to step 1. We can sweep over all the L-component Wilson loops, repeat 4 steps for new type, find out all the types of operators and finish our procedure.

We introduce steps 2, 3 and 4 in some detail here.

(a) Step 2. To find the dimensionality and the bases for certain prototype.

We illuminate it by the cubic group. The group has two generators, which means that all elements $H \in O$ can be written as a product of some powers of two generators,

$$H = C_4^{n_1} C_3^{n'_1} C_4^{n_2} C_3^{n'_2} \dots C_4^{n_m} C_3^{n'_m} \dots,$$

where $n_i = 1, 2, 3, 4$ and $n'_i = 1, 2, 3$.

Applying all possible products of C_4 and C_3 on one prototype, we can find out complete operators in this type. Actually, 24 elements of cubic group produce 24 operators. But, some of them are exactly the same. All these operators have the same shape. They construct the bases of the representation in this type. Therefore, we can extract its dimensionality and the representation matrix of these elements in these bases. The frame of the computer program is described in Fig.1. (b) Step

3. Assuming the operation of space parity P and charge conjugate C on the vector are: $P \vec{a} = \vec{b}, C \vec{d} = \vec{e}$, therefore, we can construct the parity eigenvector: $\vec{a} + \vec{b}$ for $P = +$ and $\vec{a} - \vec{b}$ for $P = -$, and C conjugate eigenvector: $\vec{d} + \vec{e}$ for $C = +$ and $\vec{d} - \vec{e}$ for $C = -$.

Since P and C are commutative, one can find common eigenvector for P and C . In this way, we may classify the bases according to P and C which are the combinations of the bases obtained from step 2.

(c) Step 4. With definite P and C , we reduce the representation into irreducible representation R^{PC} of the O^{PC} group by solving zero eigenvalue of matrix equation.

We exemplify it by decomposing reducible representation into T_1^{PC} representation (3 dimensions).

Denoting the reducible representation matrices of elements C_4, C_3 just by C_4, C_3 themselves, we want to seek three vectors $\vec{a}, \vec{b}, \vec{c}$ to construct the bases in which the representation matrices of C_3, C_4 are given by Eq. (2d). Due to Eq. (2d), we have (we assume that the vector is a row when it is left to the matrix and it is a column when right to the matrix)

$$C_4 \vec{a} = -\vec{c}, \quad C_4 \vec{b} = \vec{b}, \quad C_4 \vec{c} = \vec{a}.$$

That is

$$\vec{a} C_4^T = -\vec{c}, \quad \vec{b} C_4^T = \vec{b}, \quad \vec{c} C_4^T = \vec{a}, \quad (4)$$

here ' T ' stands for the transpose of the matrix.

We can deduce out

$$(\vec{a} \quad \vec{b} \quad \vec{c}) \begin{pmatrix} C_4^T & 0 & 0 \\ 0 & C_4^T & 0 \\ 0 & 0 & C_4^T \end{pmatrix} = (\vec{a} \quad \vec{b} \quad \vec{c}) \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \quad (5a)$$

or

$$(\vec{a} \quad \vec{b} \quad \vec{c}) \begin{pmatrix} C_4^T & 0 & -1 \\ 0 & C_4^T - 1 & 0 \\ 1 & 0 & C_4^T \end{pmatrix} = 0. \quad (5b)$$

For C_3 , we have analogously

$$(\vec{a} \quad \vec{b} \quad \vec{c}) \begin{pmatrix} C_3^T & 1 & 0 \\ 0 & C_3^T & -1 \\ 1 & 0 & C_3^T \end{pmatrix} = 0.$$

So

$$(\vec{a} \quad \vec{b} \quad \vec{c}) \begin{pmatrix} C_4^T & 0 & -1 & C_3^T & 1 & 0 \\ 0 & C_4^T - 1 & 0 & 0 & C_3^T & -1 \\ 1 & 0 & C_4^T & 1 & 0 & C_3^T \end{pmatrix} = 0. \quad (6)$$

Thus we can solve it with mathematical software. After getting the solution $\vec{a}$, $\vec{b}$, $\vec{c}$ from eq. (6), we find out the bases of representation T_1^{PC} which are our expected operators.

We emphasize that above idea is suitable for all the finite groups.

4 Results

We find the numbers of types of 4-,6-,8-,10-link operators are 1, 3, 18, 132, respectively. Table 1 gives the multiplicities of the reducible representation corresponding to definite type for 4-, 6-, 8- and 10-link operators (these reducible representations with the same dimension are not equivalent each other except for the 96-dimensional representations).

dimension of the reducible rep.	6	8	12	24	48	96
4-link	1	0	0	0	0	0
6-link	0	1	1	1	0	0
8-link	2	0	5	5	4	2
10-link	0	0	2	22	43	65

Table 1 The multiplicities of the reducible representations for 4-, 6-, 8- and 10-link operators.

As an example, we only show the results of one type of 10-link operators here, the prototype of which is $(1, 2, -1, 3, 3, 1, -2, -1, -3, -3)$. Its dimensionality is 24.

From step 2, we get bases $\{\vec{o}_i\}(i = 1, \dots, 24)$ which belongs to the same type as follows:

- 1) 1 2 -1 3 3 1 -2 -1 -3 -3 2) -2 -1 2 -3 -3 -2 1 2 3 3 3) 2 1 -2 3 3 2 -1 -2 -3 -3
- 4) -3 -3 -2 -1 2 3 3 -2 1 2 5) 3 3 2 1 -2 -3 -3 2 -1 -2 6) 1 3 -1 -2 -2 1 -3 -1 2 2
- 7) -1 -3 1 2 2 -1 3 1 -2 -2 8) -2 -2 1 3 -1 2 2 1 -3 -1 9) 2 2 -1 -3 1 -2 -2 -1 3 1
- 10) -3 2 3 1 1 -3 -2 3 -1 -1 11) 3 -2 -3 -1 -1 3 2 -3 1 1 12) 1 1 -3 2 3 -1 -1 -3 -2 3
- 13) -1 -1 3 -2 -3 1 1 3 2 -3 14) 3 -1 -3 -2 -2 3 1 -3 2 2 15) -3 1 3 2 2 -3 -1 3 -2 -2
- 16) -2 -2 3 -1 -3 2 2 3 1 -3 17) 2 2 -3 1 3 -2 -2 -3 -1 3 18) 2 3 -2 1 1 2 -3 -2 -1 -1
- 19) -2 -3 2 -1 -1 -2 3 2 1 1 20) 1 1 2 3 -2 -1 -1 2 -3 -2 21) -1 -1 -2 -3 2 1 1 -2 3 2
- 22) -1 2 1 -3 -3 -1 -2 1 3 3 23) 1 -2 -1 3 3 1 2 -1 -3 -3 24) -3 -3 -1 2 1 3 3 -1 -2 1.

From step 3, new bases $\{\vec{a}_i\}(i = 1, 2, \dots, 24)$ decomposed by P, C are listed as follows. They are linear combinations of the old bases $\vec{o}_i$. For instance, we notify $\vec{a}_{13} = \vec{o}_1 - \vec{o}_{23} - \vec{o}_{22} + \vec{o}_{24}$ by $(1, -23, -22, 24)$.

- 1) 1 23 22 24 2) 2 4 3 5 3) 6 8 7 9 4) 10 12 11 13
- 5) 14 16 15 17 6) 18 20 19 21 7) 18 20 -19 -21 8) 14 16 -15 -17
- 9) 10 12 -11 -13 10) 6 8 -7 -9 11) 2 4 -3 -5 12) 1 23 -22 -24
- 13) 1 -23 -22 24 14) 2 -4 3 -5 15) 6 -8 7 -9 16) 10 -12 11 -13
- 17) 14 -16 15 -17 18) 18 -20 19 -21 19) 18 -20 -19 21 20) 14 -16 -15 17
- 21) 10 -12 -11 13 22) 6 -8 -7 9 23) 2 -4 -3 5 24) 1 -23 22 -24.

From step 4, we get the irreducible operators which is the combination of $\vec{a}_i$. the combination coefficients are shown in Table 2.

	$PC = ++$ $\{\vec{a}_i\}, i = 1, \dots, 6$	$PC = -+$ $i = 7, \dots, 12$	$PC = +-$ $i = 13, \dots, 18$	$PC = --$ $i = 19, \dots, 24$
A_1	(1, 1, 1, 1, 1, 1)			(1, 1, 1, 1, 1, 1)
A_2	(1, -1, -1, -1, 1, 1)			(1, 1, -1, -1, -1, 1)
E	$\begin{cases} (-1, 1, 1, -2, 2, -1) \\ (-3, -3, 3, 0, 0, 3) \\ (1, -1, 2, -1, 1, -2) \\ (-3, -3, 0, 3, 3, 0) \end{cases}$			$\begin{cases} (2, -1, 1, -2, 1, -1) \\ (0, -3, -3, 0, 3, 3) \\ (1, 1, -1, -1, 2, -2) \\ (3, -3, -3, 3, 0, 0) \end{cases}$
T_1		$\begin{cases} (0, 0, 0, -1, 0, -1) \\ (-1, 0, 0, 0, 1, 0) \\ (0, -1, 1, 0, 0, 0) \end{cases}$	$\begin{cases} (1, 0, 1, 0, 0, 0) \\ (0, -1, 0, 0, 0, 1) \\ (0, 0, 0, -1, 1, 0) \end{cases}$	
T_2		$\begin{cases} (0, 0, 0, -1, 0, 1) \\ (1, 0, 0, 0, 1, 0) \\ (0, 1, 1, 0, 0, 0) \end{cases}$	$\begin{cases} (1, 0, -1, 0, 0, 0) \\ (0, 1, 0, 0, 0, 1) \\ (0, 0, 0, 1, 1, 0) \end{cases}$	

Table 2. Combination coefficients for the combination of $\{\vec{a}_i\}$

We also present some results of 8-link operators in Appendix B. The calculations for other operators are similar and interested reader can obtain the complete results of 8-link, 10-link operators and computation program for arbitrary-link operators from author D. Q. Liu.

5 Conclusions

Using our method, we can combine arbitrary-link operators into irreducible representation of cubic group easily. If we accept the opinion in Ref.[1], we have known the mapping between arbitrary-link operators and the states with certain $J^{PC}(J \leq 3)$. In this point, we have completed the combination of operators to states with certain $J^{PC}(J \leq 3)$.

But, due to Wigner-Eckart theorem, the mapping is not one-to-one, even is not definite because of the breaking of $SO(3)$ symmetry. The only we can say is, for example, that there maybe exist some projection onto 2^{++} in irreducible representation E^{++} or T_2^{++} . From our calculation, we do not know the projection of which type of the operators onto a definite J^{PC} state are larger^[1]. One way to deal with the problem is the application of variational principle^[3], or continue limits^[4]. We will discuss it more detail in the next paper in which we want to develop a approach named improved-operator one.

References

- [1] B. Berg and A. Billoire, Nucl. Phys. **B221**(1983) 109.
- [2] Jin-quan Chen, Group representation theory for physicists, World Scientific, Singapore, 1989.
- [3] C. J. Morningstar and M. peardon, Phys. Rev.**D56**(1997) 4043.
- [4] M. Luscher and P. Weisz, Comm. Math. Phys.**97**(1985) 59.
- [5] C. Michael, Acta Physica Polonica**B21**(1990) 119.

Appendix A

The cubic group O has 24 elements which may be divided into five classes of conjugate elements: E , C_2 , C_3 , C_4 , C_4^2 , and there are two generators among the elements: C_3 and C_4 . The group has five irreducible representations, named $R = A_1$, A_2 , E , T_1 and T_2 , dimensions of which are 1, 1, 2, 3 and 3, respectively. A_1 is the trivial and T_1 the vector representation in these representations. After generalizing O to O^{PC} , O^{PC} has 96 elements and 20 irreducible representation R^{PC} (P and C have two probabilities: + and -). For the simplification of the discussion, we show character table for irreducible representation of the cubic group O here.

	E	$6C_2$	$8C_3$	$6C_4$	$3C_4^2$
A_1	1	1	1	1	1
A_2	1	-1	1	-1	1
E	2	0	-1	0	2
T_1	3	-1	0	1	-1
T_2	3	1	0	-1	-1

Table 3 The character table for the irreducible representation of the cubic group O .

Appendix B

We only present results for types of 8-link operators with 6-, 12- and 24-dimension representations here.

1. operators with prototype (1 2 3 3 -2 -1 -3 -3).

Bases $\{\vec{o}_i\}$ are

- 1) 1 2 3 3 -2 -1 -3 -3 2) 2 1 3 3 -1 -2 -3 -3 3) -3 -3 -2 -1 3 3 1 2
- 4) 3 3 2 1 -3 -3 -1 -2 5) 1 3 -2 -2 -3 -1 2 2 6) -1 -3 2 2 3 1 -2 -2
- 7) -2 -2 1 3 2 2 -3 -1 8) 2 2 -1 -3 -2 -2 3 1 9) -3 2 1 1 -2 3 -1 -1
- 10) 3 -2 -1 -1 2 -3 1 1 11) 1 1 -3 2 -1 -1 -2 3 12) -1 -1 3 -2 1 1 2 -3
- 13) 3 -1 -2 -2 1 -3 2 2 14) -3 1 2 2 -1 3 -2 -2 15) -2 -2 3 -1 2 2 1 -3
- 16) 2 2 -3 1 -2 -2 -1 3 17) 2 3 1 1 -3 -2 -1 -1 18) -2 -3 -1 -1 3 2 1 1
- 19) 1 1 2 3 -1 -1 -3 -2 20) -1 -1 -2 -3 1 1 3 2 21) -1 2 -3 -3 -2 1 3 3
- 22) 1 -2 3 3 2 -1 -3 -3 23) -3 -3 -1 2 3 3 -2 1 24) 3 3 1 -2 -3 -3 2 -1.

Bases $\{\vec{a}_i\}(i = 1, \dots, 24)$ are

- 1) 1 3 2 4 2) 5 7 6 8 3) 9 11 10 12 4) 13 15 14 16
- 5) 17 19 18 20 6) 21 23 22 24 7) 21 23 -22 -24 8) 17 19 -18 -20
- 9) 13 15 -14 -16 10) 9 11 -10 -12 11) 5 7 -6 -8 12) 1 3 -2 -4
- 13) 1 -3 2 -4 14) 5 -7 6 -8 15) 9 -11 10 -12 16) 13 -15 14 -16
- 17) 17 -19 18 -20 18) 21 -23 22 -24 19) 21 -23 -22 24 20) 17 -19 -18 20
- 21) 13 -15 -14 16 22) 9 -11 -10 12 23) 5 -7 -6 8 24) 1 -3 -2 4

Combination coefficients are presented in table 4.

2. operators with prototype (1 2 -1 3 -1 -2 1 -3).

Bases $\{\vec{o}_i\}$ are

- 1) 1 2 -1 3 -1 -2 1 -3 2) -2 -1 2 -3 2 1 -2 3 3) -3 2 -1 -2 3 -2 1 2
- 4) 1 3 -1 -2 -1 -3 1 2 5) -2 -1 3 1 2 1 -3 -1 6) -3 2 3 1 3 -2 -3 -1
- 7) 1 3 2 -3 -1 -3 -2 3 8) 3 -1 -3 -2 -3 1 3 2 9) -2 -3 -1 3 2 3 1 -3
- 10) 2 3 -2 1 -2 -3 2 -1 11) 1 -2 3 2 -1 2 -3 -2 12) -1 2 1 -3 1 -2 -1 3
- 13) 1 -3 -1 2 -1 3 1 -2 14) 2 -1 -3 1 -2 1 3 -1 15) -3 -2 3 -1 3 2 -3 1
- 16) -1 3 -2 -3 1 -3 2 3 17) -2 1 2 3 2 -1 -2 -3 18) 3 2 1 -2 -3 -2 -1 2
- 19) 1 -2 -1 -3 -1 2 1 3 20) -3 -1 -2 1 3 1 2 -1 21) -3 1 3 -2 3 -1 -3 2
- 22) -2 3 1 -3 2 -3 -1 3 23) -2 -3 2 1 2 3 -2 -1 24) 1 2 -3 -2 -1 -2 3 2.

Bases $\{\vec{a}_i\}(i = 1, \dots, 24)$ are

- 1) 1 12 2) 2 3 3) 4 5 4) 6 7 5) 8 9 6) 10 11
- 7) 13 14 8) 15 16 9) 17 18 10) 19 20 11) 21 22 12) 23 24
- 13) 1 -12 14) 2 -3 15) 4 -5 16) 6 -7 17) 8 -9 18) 10 -11
- 19) 13 -14 20) 15 -16 21) 17 -18 22) 19 -20 23) 21 -22 24) 23 -24

Combination coefficients are shown in table 5.

3. Operators with prototype (1 2 3 3 -1 -2 -3 -3).

Bases $\{\vec{o}_i\}$ are

- 1) 1 2 3 3 -1 -2 -3 -3 2) -2 -1 -3 -3 2 1 3 3 3) -3 -3 -1 -2 3 3 1 2

4) 1 3 -2 -2 -1 -3 2 2 5) -2 -2 3 1 2 2 -3 -1 6) -3 2 1 1 3 -2 -1 -1
7) 1 1 2 -3 -1 -1 -2 3 8) 3 -1 -2 -2 -3 1 2 2 9) -2 -2 -1 3 2 2 1 -3
10) 2 3 1 1 -2 -3 -1 -1 11) 1 1 3 2 -1 -1 -3 -2 12) -1 2 -3 -3 1 -2 3 3
13) -3 -3 2 -1 3 3 -2 1 14) -3 -1 2 2 3 1 -2 -2 15) 2 2 -1 -3 -2 -2 1 3
16) -2 3 -1 -1 2 -3 1 1 17) -1 -1 3 -2 1 1 -3 2 18) 3 3 2 1 -3 -3 -2 -1
19) 1 -3 2 2 -1 3 -2 -2 20) 2 2 -3 1 -2 -2 3 -1 21) -3 -2 -1 -1 3 2 1 1
22) -1 -1 -2 -3 1 1 2 3 23) -2 1 3 3 2 -1 -3 -3 24) 3 3 1 -2 -3 -3 -1 2.

Bases $\{\vec{a}_i\}(i = 1, \dots, 24)$ are

1) 1 18 2) 2 3 3) 4 5 4) 6 7 5) 8 9 6) 10 11
7) 12 13 8) 14 15 9) 16 17 10) 19 20 11) 21 22 12) 23 24
13) 1 -18 14) 2 -3 15) 4 -5 16) 6 -7 17) 8 -9 18) 10 -11
19) 12 -13 20) 14 -15 21) 16 -17 22) 19 -20 23) 21 -22 24) 23 -24
Combination coefficients are given in table 6.

4. Operators with prototype (1 2 -1 -2 1 3 -1 -3).

Bases $\{\vec{o}_i\}$ are

1) 1 2 -1 -2 1 3 -1 -3 2) -2 -1 2 1 -2 -3 2 3 3) 2 1 -2 -1 2 3 -2 -3
4) -3 -2 3 2 -1 -2 1 2 5) 3 2 -3 -2 1 2 -1 -2 6) 1 3 -1 -3 1 -2 -1 2
7) -1 -3 1 3 -1 2 1 -2 8) -2 1 2 -1 3 1 -3 -1 9) 2 -1 -2 1 -3 -1 3 1
10) -3 2 3 -2 -3 1 3 -1 11) 3 -2 -3 2 3 -1 -3 1 12) 1 -3 -1 3 2 -3 -2 3
13) -1 3 1 -3 -2 3 2 -3 14) 2 -1 -2 1 2 3 -2 -3 15) -2 1 2 -1 -2 -3 2 3
16) 3 2 -3 -2 -1 2 1 -2 17) -3 -2 3 2 1 -2 -1 2 18) -1 3 1 -3 -1 2 1 -2
19) 1 -3 -1 3 1 -2 -1 2 20) -2 1 2 -1 -3 1 3 -1 21) 3 2 -3 -2 3 -1 -3 1
22) -3 -2 3 2 -3 1 3 -1 23) -1 3 1 -3 2 3 -2 -3 24) 1 -3 -1 3 -2 -3 2 3.

Bases $\{\vec{a}_i\}(i = 1, \dots, 12)$ are

1) 1 18 19 20 2) 2 4 3 5 3) 6 8 7 9 4) 10 12 11 13
5) 14 16 15 17 6) 21 23 22 24 7) 21 23 -22 -24 8) 14 16 -15 -17
9) 10 12 -11 -13 10) 6 8 -7 -9 11) 2 4 -3 -5 12) 1 18 -19 -20
13) 1 -18 -19 20 14) 2 -4 3 -5 15) 6 -8 7 -9 16) 10 -12 11 -13
17) 14 -16 15 -17 18) 21 -23 22 -24 19) 21 -23 -22 24 20) 14 -16 -15 17
21) 10 -12 -11 13 22) 6 -8 -7 9 23) 2 -4 -3 5 24) 1 -18 19 -20.

Combination coefficients are shown in table 7.

5. Operators with prototype (1 2 -1 2 -1 -2 -2 1).

Bases $\{\vec{o}_i\}$ are

1) 1 2 -1 2 -1 -2 -2 1 2) -2 -1 2 -1 2 1 1 -2 3) 2 1 -2 1 -2 -1 -1 2
4) 2 -1 -1 -2 1 -2 1 2 5) 1 3 -1 3 -1 -3 -3 1 6) -1 -3 1 -3 1 3 3 -1
7) -1 3 3 1 -3 1 -3 -1 8) 1 -3 -3 -1 3 -1 3 1 9) -3 2 3 2 3 -2 -2 -3
10) 3 -2 -3 -2 -3 2 2 3 11) 3 2 2 -3 -2 -3 -2 3 12) -3 -2 -2 3 2 3 2 -3
13) 3 -1 -3 -1 -3 1 1 3 14) -3 1 3 1 3 -1 -1 -3 15) -3 -1 -1 3 1 3 1 -3
16) 3 1 1 -3 -1 -3 -1 3 17) 2 3 -2 3 -2 -3 -3 2 18) -2 -3 2 -3 2 3 3 -2
19) -2 3 3 2 -3 2 -3 -2 20) 2 -3 -3 -2 3 -2 3 2 21) -1 2 1 2 1 -2 -2 -1
22) 1 -2 -1 -2 -1 2 2 1 23) 1 2 2 -1 -2 -1 -2 1 24) -1 -2 -2 1 2 1 2 -1.

Bases $\{\vec{a}_i\}(i = 1, \dots, 12)$ are

1) 1 3 2 4 2) 5 7 6 8 3) 9 11 10 12 4) 13 15 14 16
 5) 17 19 18 20 6) 21 23 22 24 7) 21 23 -22 -24 8) 17 19 -18 -20
 9) 13 15 -14 -16 10) 9 11 -10 -12 11) 5 7 -6 -8 12) 1 3 -2 -4
 13) 1 -3 -2 4 14) 5 -7 6 -8 15) 9 -11 10 -12 16) 13 -15 14 -16
 17) 17 -19 18 -20 18) 21 -23 22 -24 19) 21 -23 -22 24 20) 17 -19 -18 20
 21) 13 -15 -14 16 22) 9 -11 -10 12 23) 5 -7 -6 8 24) 1 -3 2 -4
 Combination coefficients are given in table 8.

6. operators with prototype (1 2 -1 -2 -2 -1 2 1).

Bases $\{\vec{o}_i\}$ are

1) 1 2 -1 -2 -2 -1 2 1 2) 2 1 -2 -1 -1 -2 1 2 3) 1 3 -1 -3 -3 -1 3 1
 4) -1 -3 1 3 3 1 -3 -1 5) -3 2 3 -2 -2 3 2 -3 6) 3 -2 -3 2 2 -3 -2 3
 7) 3 -1 -3 1 1 -3 -1 3 8) -3 1 3 -1 -1 3 1 -3 9) 2 3 -2 -3 -3 -2 3 2
 10) -2 -3 2 3 3 2 -3 -2 11) -1 2 1 -2 -2 1 2 -1 12) 1 -2 -1 2 2 -1 -2 1.

Bases $\{\vec{a}_i\}(i = 1, \dots, 12)$ are

1) 1 2 2) 3 4 3) 5 6 4) 7 8 5) 9 10 6) 11 12
 7) 11 -12 8) 9 -10 9) 7 -8 10) 5 -6 11) 3 -4 12) 1 -2

Combination coefficients are shown in table 9.

7. operators with prototype (1 2 -1 3 1 -2 -1 -3).

Bases $\{\vec{o}_i\}$ are

1) 1 2 -1 3 1 -2 -1 -3 2) -2 -1 2 -3 -2 1 2 3 3) 2 1 -2 3 2 -1 -2 -3
 4) -3 -2 -1 2 3 -2 1 2 5) 3 2 1 -2 -3 2 -1 -2 6) 1 3 -1 -2 1 -3 -1 2
 7) -2 1 3 -1 2 1 -3 -1 8) 2 -1 -3 1 -2 -1 3 1 9) -3 2 3 1 -3 -2 3 -1
 10) 3 -2 -3 -1 3 2 -3 1 11) 1 -3 2 3 -1 -3 -2 3 12) -1 3 -2 -3 1 3 2 -3.

Bases $\{\vec{a}_i\}(i = 1, \dots, 12)$ are

1) 1 8 6 7 2) 2 4 3 5 3) 9 11 10 12 4) 9 11 -10 -12
 5) 2 4 -3 -5 6) 1 8 -6 -7 7) 1 -8 6 -7 8) 2 -4 3 -5
 9) 9 -11 10 -12 10) 9 -11 -10 12 11) 2 -4 -3 5 12) 1 -8 -6 7.

Combination coefficients are given in table 10.

8. operators with prototype (1 2 -1 -1 -1 -2 1 1).

Bases $\{\vec{o}_i\}$ are

1) 1 2 -1 -1 -1 -2 1 1 2) -2 -1 2 2 2 1 -2 -2 3) 2 2 -1 -2 -2 -2 1 2
 4) 1 3 -1 -1 -1 -3 1 1 5) -1 -1 3 1 1 1 -3 -1 6) -3 2 3 3 3 -2 -3 -3
 7) 3 3 2 -3 -3 -3 -2 3 8) 3 -1 -3 -3 -3 1 3 3 9) -3 -3 -1 3 3 3 1 -3
 10) 2 3 -2 -2 -2 -3 2 2 11) -2 -2 3 2 2 2 -3 -2 12) -1 2 1 1 1 -2 -1 -1.

Bases $\{\vec{a}_i\}(i = 1, \dots, 12)$ are

1) 1 12 2) 2 3 3) 4 5 4) 6 7 5) 8 9 6) 10 11
 7) 1 -12 8) 2 -3 9) 4 -5 10) 6 -7 11) 8 -9 12) 10 -11.

Combination coefficients are given in table 11.

9. operators with prototype (1 2 -1 -2 -1 -2 1 2).

Bases $\{\vec{o}_i\}$ are

1) 1 2 -1 -2 -1 -2 1 2 2) -2 -1 2 1 2 1 -2 -1 3) 1 3 -1 -3 -1 -3 1 3
4) -3 -1 3 1 3 1 -3 -1 5) -3 2 3 -2 3 -2 -3 2 6) -2 3 2 -3 2 -3 -2 3
7) 3 -1 -3 1 -3 1 3 -1 8) 1 -3 -1 3 -1 3 1 -3 9) 2 3 -2 -3 -2 -3 2 3
10) -3 -2 3 2 3 2 -3 -2 11) -1 2 1 -2 1 -2 -1 2 12) -2 1 2 -1 2 -1 -2 1.

Bases $\{\vec{a}_i\}(i = 1, \dots, 12)$ are

1) 1 2 2) 3 4 3) 5 6 4) 7 8 5) 9 10 6) 11 12 11 12
7) 1 -2 8) 3 -4 9) 5 -6 10) 7 -8 11) 9 -10 12) 11 -12.

Combination coefficients are given in table 12.

10. operators with prototype (1 2 -1 -2 1 -2 -1 2).

Bases $\{\vec{o}_i\}$ are

1) 1 2 -1 -2 1 -2 -1 2 2) -2 -1 2 1 -2 1 2 -1 3) 2 1 -2 -1 2 -1 -2 1
4) 1 3 -1 -3 1 -3 -1 3 5) -1 -3 1 3 -1 3 1 -3 6) -3 2 3 -2 -3 -2 3 2
7) 3 -2 -3 2 3 2 -3 -2 8) 3 -1 -3 1 3 1 -3 -1 9) -3 1 3 -1 -3 -1 3 1
10) 2 3 -2 -3 2 -3 -2 3 11) -2 -3 2 3 -2 3 2 -3 12) -1 2 1 -2 -1 -2 1 2.

Bases $\{\vec{a}_i\}(i = 1, \dots, 12)$ are

1) 1 12 1 12 2) 2 3 3) 4 5 4 5 4) 6 7 5) 8 9 8 9 6) 10 11
7) 10 -11 10 -11 8) 8 -9 9) 6 -7 6 -7 10) 4 -5 11) 2 -3 2 -3 12) 1 -12.

Combination coefficients are given in table 13.

11. operators with prototype (1 2 2 -1 -1 -2 -2 1).

Bases $\{\vec{o}_i\}$ are

1) 1 2 2 -1 -1 -2 -2 1 2) -2 -1 -1 2 2 1 1 -2 3) 1 3 3 -1 -1 -3 -3 1
4) -1 3 3 1 1 -3 -3 -1 5) -3 2 2 3 3 -2 -2 -3 6) 3 2 2 -3 -3 -2 -2 3.

Bases $\{\vec{a}_i\}(i = 1, \dots, 6)$ are

1) 1 2 2) 3 4 3) 5 6 4) 1 -2 5) 3 -4 6) 5 -6.

Combination coefficients are shown in table 14.

12. operators with prototype (1 2 -1 -2 1 2 -1 -2).

Bases $\{\vec{o}_i\}$ are

1) 1 2 -1 -2 1 2 -1 -2 2) -2 -1 2 1 -2 -1 2 1 3) 1 3 -1 -3 1 3 -1 -3
4) 3 1 -3 -1 3 1 -3 -1 5) -3 2 3 -2 -3 2 3 -2 6) 2 -3 -2 3 2 -3 -2 3.

Bases $\{\vec{a}_i\}(i = 1, \dots, 6)$ are

1) 1 2 2) 3 4 3) 5 6 4) 1 -2 5) 3 -4 6) 5 -6.

Combination coefficients are shown in table 15.

	$PC = ++$ $\{\vec{a}_i\}, i = 1, \dots, 6$	$PC = -+$ $i = 7, \dots, 12$	$PC = +-$ $i = 13, \dots, 18$	$PC = --$ $i = 19, \dots, 24$
A_1	$(1, 1, 1, 1, 1, 1)$			$(1, 1, 1, 1, 1, 1)$
E	$\begin{cases} (0, -1, 1, -1, 1, 0) \\ (2, -1, -1, -1, -1, 2) \end{cases}$			$\begin{cases} (0, 1, -1, 1, -1, 0) \\ (2, -1, -1, -1, -1, 2) \end{cases}$
T_1		$\begin{cases} (1, 0, -1, 0, -1, -1) \\ (1, -1, 0, 1, 0, 1) \\ (0, 1, -1, 1, 1, 0) \end{cases}$	$\begin{cases} (-1, -1, 0, -1, 0, 1) \\ (1, 0, 1, 0, -1, 1) \\ (0, 1, 1, -1, 1, 0) \end{cases}$	
T_2	$\begin{cases} (0, 0, 1, 0, -1, 0) \\ (0, -1, 0, 1, 0, 0) \\ (-1, 0, 0, 0, 0, 1) \end{cases}$	$\begin{cases} (1, 0, 1, 0, 1, -1) \\ (-1, -1, 0, 1, 0, -1) \\ (0, -1, -1, -1, 1, 0) \end{cases}$	$\begin{cases} (1, -1, 0, -1, 0, -1) \\ (1, 0, -1, 0, 1, 1) \\ (0, -1, 1, 1, 1, 0) \end{cases}$	$\begin{cases} (0, 1, 0, -1, 0, 0) \\ (0, 0, -1, 0, 1, 0) \\ (-1, 0, 0, 0, 0, 1) \end{cases}$

Table 4 Combination coefficients for the combination of $\{\vec{a}_i\}$ from type 1.

	$PC = ++$ $\{\vec{a}_i\}, i = 1, \dots, 12$	$PC = +-$ $i = 13, \dots, 24$
A_1	$(1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$	
A_2	$(1, -1, -1, -1, 1, 1, -1, -1, -1, 1, 1, 1)$	
E	$\begin{cases} (-1, 1, 1, -2, 2, -1, 1, -2, 1, -1, 2, -1) \\ (-3, -3, 3, 0, 0, 3, 3, 0, -3, -3, 0, 3) \\ (1, -1, 2, -1, 1, -2, 2, -1, -1, 1, 1, -2) \\ (-3, -3, 0, 3, 3, 0, 0, 3, -3, -3, 3, 0) \end{cases}$	
T_1	$\begin{cases} (0, 1, 0, 0, 1, 0, 0, 0, -1, 0, -1, 0) \\ (-1, 0, 0, -1, 0, 0, 0, 1, 0, 1, 0, 0) \\ (0, 0, -1, 0, 0, -1, 1, 0, 0, 0, 0, 1) \end{cases}$	$\begin{cases} (-1, 0, -1, 0, 0, 0, -1, 0, 0, -1, 0, 0) \\ (0, 1, 0, 0, 0, -1, 0, 0, 1, 0, 0, 1) \\ (0, 0, 0, 1, -1, 0, 0, 1, 0, 0, 1, 0) \\ (0, 0, 0, -1, 0, -1, 0, 1, 0, 0, 0, -1) \\ (0, 0, 1, 0, 1, 0, -1, 0, 0, 0, 1, 0) \\ (-1, 1, 0, 0, 0, 0, 0, 0, -1, 1, 0, 0) \end{cases}$
T_2	$\begin{cases} (0, -1, 0, 0, 1, 0, 0, 0, 1, 0, -1, 0) \\ (-1, 0, 0, 1, 0, 0, 0, -1, 0, 1, 0, 0) \\ (0, 0, 1, 0, 0, -1, -1, 0, 0, 0, 0, 1) \end{cases}$	$\begin{cases} (-1, 0, 1, 0, 0, 0, 1, 0, 0, -1, 0, 0) \\ (0, -1, 0, 0, 0, -1, 0, 0, -1, 0, 0, 1) \\ (0, 0, 0, -1, -1, 0, 0, -1, 0, 0, 1, 0) \\ (0, 0, 0, 1, 0, -1, 0, -1, 0, 0, 0, -1) \\ (0, 0, -1, 0, 1, 0, 1, 0, 0, 0, 1, 0) \\ (-1, -1, 0, 0, 0, 0, 0, 0, 1, 1, 0, 0) \end{cases}$

Table 5 Combination coefficients for the combination of $\{\vec{a}_i\}$ of type 2.

	$PC = ++$ $\{\vec{a}_i\}, i = 1, \dots, 12$	$PC = +-$ $i = 13, \dots, 24$
A_1	$(1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$	
A_2		$(-1, 1, 1, 1, -1, -1, -1, -1, -1, 1, 1, 1)$
E	$\begin{cases} (0, 0, -1, 1, -1, 1, 0, -1, 1, -1, 1, 0) \\ (2, 2, -1, -1, -1, -1, 2, -1, -1, -1, -1, 2) \end{cases}$	$\begin{cases} (2, -2, 1, 1, -1, -1, 2, -1, -1, 1, 1, -2) \\ (0, 0, -3, 3, 3, -3, 0, 3, -3, -3, 3, 0) \end{cases}$
T_1	$\begin{cases} (-1, 1, -1, 0, 1, 0, 1, 1, 0, -1, 0, -1) \\ (-1, 1, 0, -1, 0, -1, -1, 0, 1, 0, 1, 1) \\ (0, 0, -1, 1, -1, -1, 0, 1, -1, 1, 1, 0) \end{cases}$	$\begin{cases} (0, 0, 0, 1, 0, 1, 0, 0, -1, 0, -1, 0) \\ (0, 0, -1, 0, -1, 0, 0, 1, 0, 1, 0, 0) \\ (1, -1, 0, 0, 0, 0, -1, 0, 0, 0, 0, 1) \\ (-1, -1, -1, 0, -1, 0, 1, -1, 0, -1, 0, 1) \\ (1, 1, 0, 1, 0, -1, 1, 0, 1, 0, -1, 1) \\ (0, 0, 1, 1, -1, 1, 0, 1, 1, -1, 1, 0) \end{cases}$
T_2	$\begin{cases} (0, 0, 0, 1, 0, -1, 0, 0, 1, 0, -1, 0) \\ (0, 0, -1, 0, 1, 0, 0, -1, 0, 1, 0, 0) \\ (-1, -1, 0, 0, 0, 0, 1, 0, 0, 0, 0, 1) \\ (1, -1, -1, 0, 1, 0, -1, 1, 0, -1, 0, 1) \\ (-1, 1, 0, 1, 0, 1, -1, 0, -1, 0, -1, 1) \\ (0, 0, 1, 1, 1, -1, 0, -1, -1, -1, 1, 0) \end{cases}$	$\begin{cases} (1, 1, -1, 0, -1, 0, -1, -1, 0, -1, 0, -1) \\ (1, 1, 0, -1, 0, 1, 1, 0, -1, 0, 1, 1) \\ (0, 0, -1, 1, 1, 1, 0, -1, 1, 1, 1, 0) \end{cases}$

Table 6 Combination coefficients for the combination of $\{\vec{a}_i\}$ from type 3.

	$PC = ++$ $\{\vec{a}_i\}, i = 1, \dots, 6$	$PC = -+$ $i = 7, \dots, 12$	$PC = +-$ $i = 13, \dots, 18$	$PC = --$ $i = 19, \dots, 24$
A_1	$(1, 1, 1, 1, 1, 1)$			
A_2				$(-1, -1, -1, -1, -1, 1)$
E	$\begin{cases} (1, -1, 1, 0, -1, 0) \\ (-1, -1, -1, 2, -1, 2) \end{cases}$			$\begin{cases} (2, -1, 2, -1, -1, 1) \\ (0, 3, 0, -3, 3, 3) \end{cases}$
T_1		$\begin{cases} (-1, -1, 1, 0, -1, 0) \\ (1, 0, 1, -1, 0, 1) \\ (0, 1, 0, 1, -1, 1) \end{cases}$	$\begin{cases} (0, 0, 0, -1, 0, -1) \\ (0, -1, 0, 0, 1, 0) \\ (-1, 0, 1, 0, 0, 0) \end{cases}$	$\begin{cases} (0, 0, 0, -1, 0, -1) \\ (0, -1, 0, 0, 1, 0) \\ (-1, 0, 1, 0, 0, 0) \end{cases}$
T_2	$\begin{cases} (-1, 0, 1, 0, 0, 0) \\ (0, -1, 0, 0, 1, 0) \\ (0, 0, 0, -1, 0, 1) \end{cases}$	$\begin{cases} (1, -1, -1, 0, -1, 0) \\ (1, 0, 1, 1, 0, -1) \\ (0, -1, 0, 1, 1, 1) \end{cases}$	$\begin{cases} (0, -1, 0, 1, -1, -1) \\ (-1, 0, -1, 1, 0, 1) \\ (-1, -1, 1, 0, 1, 0) \end{cases}$	

Table 7 Combination coefficients for the combination of $\{\vec{a}_i\}$ from type 4.

	$PC = ++$ $\{\vec{a}_i\}, i = 1, \dots, 6$	$PC = -+$ $i = 7, \dots, 12$	$PC = +-$ $i = 13, \dots, 18$	$PC = --$ $i = 19, \dots, 24$
A_1	$(1, 1, 1, 1, 1, 1)$			
A_2			$(1, -1, -1, 1, 1, 1)$	
E	$\begin{cases} (0, -1, 1, -1, 1, 0) \\ (2, -1, -1, -1, -1, 2) \end{cases}$		$\begin{cases} (-2, -1, -1, 1, 1, -2) \\ (0, 3, -3, -3, 3, 0) \end{cases}$	
T_1		$\begin{cases} (-1, 0, -1, 0, 1, 1) \\ (1, 1, 0, 1, 0, 1) \\ (0, 1, 1, -1, 1, 0) \end{cases}$	$\begin{cases} (0, 0, -1, 0, -1, 0) \\ (0, 1, 0, 1, 0, 0) \\ (-1, 0, 0, 0, 0, 1) \end{cases}$	$\begin{cases} (1, 0, -1, 0, -1, -1) \\ (1, -1, 0, 1, 0, 1) \\ (0, 1, -1, 1, 1, 0) \end{cases}$
T_2	$\begin{cases} (0, 0, 1, 0, -1, 0) \\ (0, -1, 0, 1, 0, 0) \\ (-1, 0, 0, 0, 0, 1) \end{cases}$	$\begin{cases} (-1, 0, 1, 0, -1, 1) \\ (-1, 1, 0, 1, 0, -1) \\ (0, -1, 1, 1, 0, 0) \end{cases}$		$\begin{cases} (1, 0, 1, 0, 1, -1) \\ (-1, -1, 0, 1, 0, -1) \\ (0, -1, -1, -1, 1, 0) \end{cases}$

Table 8 Combination coefficients for the combination of $\{\vec{a}_i\}$ from type 5.

	$PC = ++$ $\{\vec{a}_i\}, i = 1, \dots, 6$	$PC = +-$ $i = 7, \dots, 12$
A_1	$(1, 1, 1, 1, 1, 1)$	
E	$\begin{cases} (0, -1, 1, -1, 1, 0) \\ (2, -1, -1, -1, -1, 2) \end{cases}$	
T_1		$\begin{cases} (1, 0, -1, 0, -1, -1) \\ (1, -1, 0, 1, 0, 1) \\ (0, 1, -1, 1, 1, 0) \end{cases}$
T_2	$\begin{cases} (0, 0, 1, 0, -1, 0) \\ (0, -1, 0, 1, 0, 0) \\ (-1, 0, 0, 0, 0, 1) \end{cases}$	$\begin{cases} (1, 0, 1, 0, 1, -1) \\ (-1, -1, 0, 1, 0, -1) \\ (0, -1, -1, -1, 1, 0) \end{cases}$

Table 9 Combination coefficients for the combination of $\{\vec{a}_i\}$ from type 6.

	$PC = ++$ $\{\vec{a}_i\}, i = 1, \dots, 3$	$PC = -+$ $i = 4, \dots, 6$	$PC = +-$ $i = 7, \dots, 9$	$PC = --$ $i = 10, \dots, 12$
A_1	(1, 1, 1)			
A_2				(-1, -1, 1)
E	$\begin{Bmatrix} (1, -1, 0) \\ (-1, -1, 2) \end{Bmatrix}$			$\begin{Bmatrix} (2, -1, 1) \\ (0, 3, 3) \end{Bmatrix}$
T_1			$\begin{Bmatrix} (-1, 0, 0) \\ (0, 1, 0) \\ (0, 0, 1) \end{Bmatrix}$	
T_2		$\begin{Bmatrix} (0, 0, 1) \\ (0, 1, 0) \\ (1, 0, 0) \end{Bmatrix}$		

Table 10 Combination coefficients for the combination of $\{\vec{a}_i\}$ from type 7.

	$PC = ++$ $\{\vec{a}_i\}, i = 1, \dots, 6$	$PC = +-$ $i = 7, \dots, 12$
A_1	(1, 1, 1, 1, 1, 1)	
A_2	(1, -1, -1, -1, 1, 1)	
E	$\begin{Bmatrix} (-1, 1, 1, -2, 2, -1) \\ (-3, -3, 3, 0, 0, 3) \\ (1, -1, 2, -1, 1, -2) \\ (-3, -3, 0, 3, 3, 0) \end{Bmatrix}$	
T_1		$\begin{Bmatrix} (0, 0, 0, -1, 0, -1) \\ (0, 0, 1, 0, 1, 0) \\ (-1, 1, 0, 0, 0, 0) \end{Bmatrix}$
T_2		$\begin{Bmatrix} (0, 0, 0, -1, 0, 1) \\ (0, 0, 1, 0, -1, 0) \\ (1, 1, 0, 0, 0, 0) \end{Bmatrix}$

Table 11 Combination coefficients for the combination of $\{\vec{a}_i\}$ from type 8.

	$PC = ++$ $\{\vec{a}_i\}, i = 1, \dots, 6$	$PC = +-$ $i = 7, \dots, 12$
A_1	(1, 1, 1, 1, 1, 1)	
A_2		(1, -1, -1, 1, 1, 1)
E	$\begin{Bmatrix} (0, -1, 1, -1, 1, 0) \\ (2, -1, -1, -1, -1, 2) \end{Bmatrix}$	$\begin{Bmatrix} (-2, -1, -1, 1, 1, -2) \\ (0, 3, -3, -3, 3, 0) \end{Bmatrix}$
T_1		$\begin{Bmatrix} (0, 0, -1, 0, -1, 0) \\ (0, 1, 0, 1, 0, 0) \\ (-1, 0, 0, 0, 0, 1) \end{Bmatrix}$
T_2	$\begin{Bmatrix} (0, 0, 1, 0, -1, 0) \\ (0, -1, 0, 1, 0, 0) \\ (-1, 0, 0, 0, 0, 1) \end{Bmatrix}$	

Table 12 Combination coefficients for the combination of $\{\vec{a}_i\}$ from type 9.

	$PC = ++$ $\{\vec{a}_i\}, i = 1, \dots, 6$	$PC = --$ $i = 7, \dots, 12$
A_1	$(1, 1, 1, 1, 1, 1)$	
A_2	$(1, -1, -1, -1, 1, 1)$	
E	$\begin{cases} (-1, 1, 1, -2, 2, -1) \\ (-3, -3, 3, 0, 0, 3) \\ (1, -1, 2, -1, 1, -2) \\ (-3, -3, 0, 3, 3, 0) \end{cases}$	
T_1		$\begin{cases} (0, 0, 0, -1, 0, -1) \\ (-1, 0, 0, 0, 1, 0) \\ (0, -1, 1, 0, 0, 0) \end{cases}$
T_2		$\begin{cases} (0, 0, 0, -1, 0, 1) \\ (1, 0, 0, 0, 1, 0) \\ (0, 1, 1, 0, 0, 0) \end{cases}$

Table 13 Combination coefficients for the combination of $\{\vec{a}_i\}$ from type 10.

	$PC = ++$ $\{\vec{a}_i\}, i = 1, \dots, 3$	$PC = +-$ $i = 4, \dots, 6$
A_1	$(1, 1, 1)$	
E	$\begin{cases} (0, 1, -1) \\ (-2, 1, 1) \end{cases}$	
T_1		$\begin{cases} (0, 0, 1) \\ (0, -1, 0) \\ (1, 0, 0) \end{cases}$

Table 14 Combination coefficients for the combination of $\{\vec{a}_i\}$ from type 11.

	$PC = ++$ $\{\vec{a}_i\}, i = 1, \dots, 3$	$PC = +-$ $i = 4, \dots, 6$
A_1	$(1, 1, 1)$	
E	$\begin{cases} (0, 1, -1) \\ (-2, 1, 1) \end{cases}$	
T_1		$\begin{cases} (0, 0, 1) \\ (0, -1, 0) \\ (1, 0, 0) \end{cases}$

Table 15 Combination coefficients for the combination of $\{\vec{a}_i\}$ from type 12.




Fig 1. The program frame.