MPI-PhT/37-2001 TPJU/7-2001 September, 2001

LATTICE STUDY OF THE SIMPLIFIED MODEL OF M-THEORY FOR LARGER GAUGE GROUPS

P. Bialas^{*a*} and J. Wosiek^{*b*1}

 ^a Inst. of Computer Science Jagellonian University, Nawojki 11, 30-072 Krakow, Poland
 ^b M. Smoluchowski Institute of Physics, Jagellonian University, Reymonta 4, 30-059 Cracow, Poland E-mail: wosiek@th.if.uj.edu.pl

Abstract

Lattice discretization of the supersymmetric Yang-Mills quantum mechanics is discussed. First results of the quenched Monte Carlo simulations, for D=4 and with higher gauge groups ($3 \le N \le 8$), are presented. We confirm an earlier (N=2) evidence that the system reveals different behaviours at low and high temperatures separated by a narrow transition region. These two regimes may correspond to a black hole and elementary excitations phases conjectured in the M-theory. Dependence of the "transition temperature" on N is consistent with 't Hooft scaling and shows a smooth saturation of lattice results towards the large N limit. Is not yet resolved if the observed change between the two regimes corresponds to a genuine phase transition of the lattice model is also proposed and its advantages are briefly discussed.

¹Invited talk presented at the Sixth Workshop on Non-Perturbative QCD, American University of Paris, Paris, June, 2001.

1 Yang-Mills quantum mechanics and its lattice formulation

The hypothesis of M-theory was a culmination of the recent developments in nonperturbative string theories. In the unified scheme, which emerged, all known five string theories are dual images of a single theory, which also contains eleven dimensional supergravity (for a recent review see e.g. [1]). Even though it is not yet known with the full precision, the M-theory has a remarkable potential to unify all interactions and particles. In particular it may offer a topological explanation of such fundamental features as three families and fractional charges. It may lead to a standard model gauge group with $\mathcal{N}=1$ supersymmetry. It provides understanding of the Bekenstein-Hawking entropy puzzle and much more. Due to the proposition of Banks, Fishler, Susskind and Shenker [2], M-theory became amenable to yet more quantitative studies. According to BFSS the spectrum of M-theory is equivalent to that of a supersymmetric Yang-Mills quantum mechanics (SYMQM). The latter is obtained from the dimensional reduction of the 10 dimensional supersymmetric Yang-Mills theory to the one (time) dimension. Such a quantum mechanical system can be quantitatively studied by a host of nonperturbative methods. One of them, which provides steadily increasing understanding of nonperturbative phenomena, is the lattice approach. Therefore we have constructed the Wilson discretization of the above quantum mechanics and proposed to investigate it with the standard lattice methods [3]. The ultimate goal is to study D=10 SYMQM for the large size of the SU(N) matrices. However, even for this relatively simple, one dimensional quantum mechanical system this is still a rather complex task. One of the main difficulty is posed by the complex fermionic determinant (and pfaffian) for addjoint fermions in D=10. Another, although less severe one, is the time consuming Monte Carlo calculation with large nonabelian matrices. On the other hand the system can be simulated at present in other regions of the parameter space D, N, N_f , $(N_f$ being the number of fermions) and such a study may provide a relevant information about its general structure. Therefore we have decided to set up a systematic lattice survey of SYMQM beginning with the simplest case of $D = 4, N = 2, N_f = 0$ and gradually extending it as far as possible towards the BFSS limit. In this talk I will report on the second step along this program, namely the first results for higher N will be presented.

Supersymmetric Yang-Mills quantum mechanics [4, 5, 6] and its zero dimensional counterpart [7, 8] have been intensively studied. Although the

exact solution is still not available, many results are known and a) can be tested and extended, and b) provide us a guidance from a "simple corner" of parameter space to the ultimate "BFSS corner".

The action of the SYMQM reads

$$S = \int dt \left(\frac{1}{2} \operatorname{Tr} F_{\mu\nu}(t)^2 + \bar{\Psi}^a(t) \mathcal{D} \Psi^a(t) \right).$$
 (1)

where $\mu, \nu = 1...D$, and all fields are independent of the space coordinates \vec{x} . The supersymmetric fermionic partners belong to the addjoint representation of SU(N). The discretized system is put on a D dimensional hypercubic lattice $N_1 \times ... \times N_D$ which is reduced in all space directions to $N_i = 1, i = 1...D - 1$. Gauge and fermionic variables are assigned to links and sites of the new elongated lattice in the standard manner. This geometry has two straightforward consequences. First, all space derivatives in bosonic and in fermionc terms vanish, $\partial_i \to 0$. Second, there is a new class of gauge invariant observables, namely $Tr(A_i(t)^2)$ or $Tr(U_i(m))$, since a local gauge transformation of the spatial components $A_i(t)$ reduces to a simple similarity transformation. The gauge part of the action has the usual form

$$S_G = -\beta \sum_{m=1}^{N_t} \sum_{\mu > \nu} \frac{1}{N} Re(\operatorname{Tr} U_{\mu\nu}(m)), \qquad (2)$$

with

$$\beta = 2N/a^3 g^2,\tag{3}$$

and $U_{\mu\nu}(m) = U^{\dagger}_{\nu}(m)U^{\dagger}_{\mu}(m+\nu)U_{\nu}(m+\mu)U_{\mu}(m)$, $U_{\mu}(m) = \exp(iagA_{\mu}(am))$, where *a* denotes the lattice constant and *g* is the gauge coupling in one dimension. The integer time coordinate along the lattice is *m*. Periodic boundary conditions $U_{\mu}(m+\nu) = U_{\mu}(m)$, $\nu = 1 \dots D - 1$, guarantee that Wilson plaquettes $U_{\mu\nu}$ tend, in the classical continuum limit, to the appropriate components $F_{\mu\nu}$ with the space derivatives absent.

Another simple property of the one dimensional gauge system shows up in Eq.(3). The gauge coupling g has a dimension, and as a consequence its numerical value provides directly a scale to all phenomena occuring in this system. In a language familiar from the four dimensional QCD: the dimensionless lattice coupling $\beta = \beta(a)$ is running according to Eq.(3), and g plays a role of Λ_{QCD} .

In the above formulation the projection on the gauge invariant states is naturally implemented. Eq. (2) was the basis of the MC simulations for a simplified SU(2) model (see Sect.(2.1)). Now some results for higher gauge groups N < 9 are also available and will be discussed in Sect.(2.2).

2 Results

$2.1 \quad SU(2)$

One of the remarkable features of the M-theory is the explanation of the problem of the Bekenstein-Hawking entropy[9] in terms of the elementary excitations. Moreover, the theory predicts existence of at least two phases: a low temperature, "black hole", phase and a high temperature phase described by the elementary D0 branes $[10]^2$. Therefore a natural question emerges if the simplified model (i.e. SYMQM), considered at finite temperature, possesses any nontrivial phase structure [11]. It is well established that QCD (or pure Yang-Mills theory) has two different phases (e.g. confinement and deconfinement). Since the action (2) is basically QCD-like, one might naively expect that the dimensionally reduced model may indeed exhibit similar phenomenon. On the other hand the one dimensional model with local interactions cannot have a phase transition for finite N. However at infinite N the sharp singularity may occur [12]. It follows that, contrary to the first expectations, the mechanism which may eventually lead to such a transition will have to be rather different that for, e.g., N = 3 QCD. Hence the problem is even more interesting, since a) it is relevant for the M-theory builders and b) the nature of the transition (if any) is open.

In Ref.[3] we have studied the phase structure of the quenched model with D=4, N=2 and $N_f = 0$. Since N plays a role of the volume, we expected to see some signatures of a phase change for finite and even small N³.

To check this we have measured the distribution of the trace of the Polyakov line

$$P = \frac{1}{N} \operatorname{Tr} \left(\prod_{m=1}^{N_t} U_D(m) \right).$$
(4)

which is a very sensitive determinant of the two phases in QCD. Similarly to lattice pure gauge system, symmetric concentration of the trace around zero

 $^{^{2}}$ The full phase structure of the M-theory is much more complex, *op. cit.*

 $^{^{3}}$ It is well known from statistical physics that the singularities (in the temperature) associated with the phase transition, develop gradually and smoothly with increasing the volume and show up (as a broad peaks for example) even for small systems [14].

indicates the low temperature phase with $\langle P \rangle = 0$ (here a "black hole" phase) while clustering around ± 1 (or, for arbitrary N, around the elements of Z_N) is characteristic of the high temperature phase (here the elementary 0-brane phase). It follows from Eq.(3) and $N_t = 1/Ta$ that $\beta = 2NN_t^3T^3$, hence the lattice coupling provides, at fixed N_t , a direct measure of the physical temperature T^4 . It was found that the distribution of (4) is indeed changing from a convex to a concave one at some finite value of $\beta = \beta_c(N_t)$. Moreover the dependence of β_c on N_t is consistent with the canonical scaling. The best fit to Monte Carlo results gave $\beta_c(N_t) = (0.17 \pm 0.05)N_t^{(3.02\pm 0.33)}$ in good agreement with the expected dependence $\beta_c \sim N_t^3$. This means that in the continuum limit the transition occurs at finite temperature T_c , which can be written as $T_c = A_N(g^2N)^{(1/3)}$ with $A_2 = .28 \pm 0.03$.

We have also measured the average size of the system $R^2 = g^2 \sum_a (A_i^a)^2$ for different values of the temperature. It shows entirely different behaviours at low and high temperatures with a narrow transition region. The pseudocritical temperature determined in this way is consistent with the one obtained above from the study of the Polyakov line. Our results also agree qualitatively with the mean field calculation at large N [11]. Obviously there is a long way between N = 2 and the BFSS limit $N \to \infty$ and it was essential to extend this study to the larger groups.

2.2 SU($3 \le N \le 8$)

Recently we have done Monte Carlo simulations with the intermediate size groups, still for $N_f = 0$ and D = 4 [13]. We used the standard Metropolis algorithm employing all independent moves in the SU(N) manifold. For N > 2 the trace P in Eq.(4) is complex. The condition det(U)=1 constraints P to lie inside the "N-star" in the complex plane (*c.f.* Fig.1). New simulations entirely confirm the structure found in the SU(2) case. In the low temperature phase the distribution is symmetric and peaked around P = 0which results in $\langle P \rangle = 0$. At high temperatures P's are expected to concentrate around the elements of Z_N . A sample of our results for N=3,5,8 and at high and low temperatures ($\beta \sim T^3$) is shown in Fig.1. Indeed one clearly sees the existence of both regimes. Moreover, in the high temperature region, the system has a tendency to be stuck in one of the Z_N minima. Correspondingly the thermalization time grows strongly with N. This is a typical indication of the spontaneous symmetry breaking which might occur at $N \to \infty$ leading eventually to a non-zero value of the order

 $^{^{4}}$ Up to the finite *a* corrections.



Figure 1: Distribution of the Polyakov line in the low and high temperature phases (upper and lower rows respectively) of the quenched Yang-Mills quantum mechanics with the SU(3), SU(5) and SU(8) gauge groups.

parameter $\langle P \rangle$.

We cannot determine at present the nature of the transition separating the two phases. In fact it is also possible that this is not the phase transition at all, but just a gentle crossover between the high and low temperature regimes. More extensive simulations are required to answer this question. Nevertheless, our data show unambiguously that the system behaves differently in both regions. For example, the dependence of the size of the system on the temperature is definitely different, and the change in the behaviour occurs at the same T where the distributions in Fig. 1 change their shapes. Moreover, the characteristic temperature separating the two regions (loosely called phases) is physical in the sense that it is finite in the continuum limit. This follows simply from the N_t^3 scaling of the lattice coupling β_c .

With present data one can study, for the first time, the N dependence of the transition temperature. It follows from Eq.(3) and $N_t = 1/Ta$, that the 't Hooft scaling $T_c \approx (g^2 N)^{1/3}$ implies that the lattice coupling $\beta_c \approx N^2$ at fixed N_t . This prediction is tested in Fig.2, where β_c/N^2 is plotted versus



Figure 2: Dependence of the transition lattice coupling on the number of colours N at fixed time extent $N_t = 4$.

N. The values of β_c were determined for each N, analogously to Sect.2.1, as the couplings for which the distributions, Fig.1, change their shape. The errors represent the subjective uncertainty of such an estimate. Indeed a nice saturation of the N dependence is observed in Fig.2 confirming the 't Hooft scaling of T_c . The solid line is a fit of the $1/N^2$ correction, and the flat solid line indicates the fitted asymptotic value. The data for the first two N require yet higher order corrections, while simulations for SU(8) give the asymptotic value with less than 15% accuracy.

It is interesting to compare above results with the N dependence found in the full, spatially extended Yang-Mills theory [15] where the $1/N^2$ correction seems to work well even for N=2,3 in some cases. It would be instructive to repeat their study for the finite temperature phenomena.

3 Noncompact formulation

In this section we propose an alternative discretization of the continuum model (1), which turns out to be easier to simulate numerically. The idea is to leave the D-1 spatial components of the gauge fields noncompact, while keeping the *D*-th component compact. To this end split the continuous

action of the reduced theory into the kinetic and potential terms.

$$S = S_{kin} + S_{pot},$$

$$S_{pot} = \frac{1}{2g^2} \int dt Tr(\mathbf{X}^i \mathbf{X}^k)^2,$$

$$S_{kin} = \frac{1}{g^2} \int dt Tr(D_t X^i)^2.$$
(5)

where the sums over i and k are implied and D_t denotes the covariant derivative along the time direction After discretizing the time we obtain

$$S_{pot} = \frac{a}{2g^2} \sum_m Tr(\mathbf{X}^i \mathbf{X}^k)^2,$$

$$S_{kin} = \frac{1}{ag^2} \sum_m Tr(\Delta \mathbf{X}^i)^2,$$
(6)

with the D-1 noncompact matrix coordinates $\mathbf{X}^{i}(m)$ defined at a discrete time intervals $t_{m} = am$. The covariant finite difference along the time direction

$$\Delta \mathbf{X}^{i}(m+1) = \mathbf{X}^{i}(m+1) - U(m+1,m)\mathbf{X}^{i}(m)U(m,m+1).$$
(7)

takes into account the parallel transport between adjacent lattice cites. The system now has D-1 noncompact coordinates \mathbf{X}^i and one compact degree of freedom corresponding to the timelike link $U(m+1,m) \equiv U_D(m)$. The discretized action is invariant under the same local gauge transformations as the compact version (2)

$$\begin{aligned} \mathbf{X}^{i}(m) &\to V^{-1}(m)\mathbf{X}^{i}(m)V(m), \\ U(m+1,m) &\to V^{-1}(m+1)U(m+1,m)V(m), \end{aligned}$$

$$\end{aligned}$$

$$\end{aligned}$$

As a consequence of the dimensional reduction there is no inhomogeneous term in the gauge transformation of the spatial coordinates \mathbf{X}^i . Also, they are rotated by *the same* element of the gauge group V(m) on both sides, since the space like links on $N_i = 1$ lattice, Eq.(2) close back onto themselves.

With the noncompact action (6) one can readily perform simulations closer to the continuum limit than those done with the fully compact variables. Also higher N's can be easier achieved. This is because the lattice coupling $\beta = 2N/a^3g^2$ diverges strongly at small *a* and this causes very severe critical slowing down in the compact case. The problem is even more difficult for larger *N*. On the other hand the effective coupling in the noncompact formulation (6) diverges only as 1/a which alleviates the above difficulty. We have done some exploratory simulations with the new formulation for a range of N's $(2 \le N \le 9)$ and for different dimensions of unreduced theory $(4 \le D \le 10)$. All simulations in all these dimensions confirm existence of the two different regimes of the model in full analogy with the compact, N = 2 case. We have encountered no difficulty extending these study to higher D. In fact we have found a steady decrease of the average size of a system (defined per one link) with D, in both phases, in agreement with the mean field predictions [11].

4 The future

Lattice simulations provide a new approach to a quantitative study of the Yang-Mills quantum mechanics and possibly to the M-theory. Preliminary results are encouraging, but a lot remains to be done. Simulations of the quenched model work for all interesting values of the dimension D and are feasible for a range of N. Recent results give us a rough idea how the large N limit is approached and where the asymptotics sets it. All quenched simulations performed up do date indicate existence of the two regions at finite temperature. This intriguing correspondence with the predictions of the M-theory should be (and can be) further quantified. Of course, the next step is to include the dynamical fermions. This can be done by a brute force for D=4 and for first few N's at D=10. The one dimensional nature of the system should help considerably. For higher N, at D=10, we face the problem of the complex pfaffian and new ideas are needed. Finally, the full potential of the small volume approach [16, 17] may provide an important insight into the problem and should be applied to these supersymmetric systems.

Acknowledgements

JW thanks the organizers of this Workshop for the invitation and support. JW also thanks the Theory Group of the Max-Planck-Institute in Munich, where this work was completed, for their hospitality and support. This research is supported by the Polish Committee for Scientific Research under the grant 2 P03B 019 17.

References

[1] A. Sen, Nucl. Phys. B(Proc. Suppl.) 94, 35 (2000).

- [2] T. Banks, W. Fishler, S. Shenker and L. Susskind, *Phys. Rev.* D 55, 6189 (1997).
- [3] R. A. Janik and J. Wosiek, Acta. Phys. Polon. B 32, 2143 (2001), hep-th/0003121; Nucl. Phys. B(Proc. Suppl.) 94, 711 (2000).
- [4] M. Claudson and M. B. Halpern, Nucl. Phys. B 250, 689 (1985).
- [5] U. H. Danielsson, G. Ferretti and B. Sundborg, Int. J. Mod. Phys. A 11, 5463 (1996).
- [6] M. B. Halpern and C. Schwartz, Int. J. Mod. Phys. A 13, 4367 (1998).
- [7] N. Ishibashi, H. Kawai, Y. Kitazawa and A. Tsushijya, *Nucl. Phys.* B 498, 467 (1997), hep-th/9612115.
- [8] J. Ambjorn et al., JHEP 9911, 029 (1999); JHEP 0007, 011 (2000).
- [9] A. Strominger and C. Vafa, *Phys. Lett.* B **379**, 99 (1996).
- [10] E. J. Martinec, hep-th/9909049, Lectures given at NATO Advanced Study Institute: TMR Summer School on Progress in String Theory and M-Theory, Cargesse, Corsica, France, May 1999.
- [11] D. Kabat, G. Lifszytz and D. A. Lowe, *Phys. Rev. Lett.* 86, 1426 (2001);
 Int. J. Mod. Phys. A 16, 856 (2001);hep-th/0105171.
- [12] D. J. Gross and E. Witten, *Phys. Rev.* D **21**, 446 (1980).
- [13] P. Bialas and J. Wosiek, hep-lat/0105031, to appear in the Proceedings of the 36th Rencontres de Moriond, Les Arc, France, March 2001.
- [14] L. Stodolsky and J. Wosiek, Nucl. Phys. B 413, 813 (1994).
- [15] M. Teper, *Phys. Lett.* B **397**, 223 (1997); *Phys. Rev.* D **59**, 014512 (1999); B. Lucini and M. Teper, *JHEP* **0106**, 050 (2001); hep-lat/0107007.
- [16] M. Lüscher, Nucl. Phys. B 219, 233 (1983); P. van Baal, Acta. Phys. Polon. B 20, 295 (1989).
- [17] For a recent review see, P. van Baal in Handbook of QCD, Boris Ioffe Festchrift, ed. M. Shifman (World Scientific, Singapore, 2001).