A Numerical Study of the Hierarchical Ising Model: High Temperature Versus Epsilon Expansion

Y. MEURICE, G. ORDAZ, AND V.G.J. RODGERS

Department of Physics and Astronomy, University of Iowa, Iowa City, Iowa 52242, USA

#### ABSTRACT

We study numerically the magnetic susceptibility of the hierarchical model with Ising spins ( $\sigma = \pm 1$ ) above the critical temperature and for two values of the epsilon parameter. The integrations are performed exactly using recursive methods which exploit the symmetries of the model. Lattices with up to  $2^{18}$  sites have been used. Surprisingly, the numerical data can be fitted very well with a simple power law of the form  $(1-\beta/\beta_c)^{-\gamma}$  for the whole temperature range. The numerical values for  $\gamma$  agree within a few percent with the values calculated with a high-temperature expansion but show significant discrepancies with the epsilon-expansion.

#### 1. Introduction

The renormalization group (RG) method<sup>[1]</sup>is a powerful tool to handle critical phenomena and to approach the continuum limit of lattice models. However, its practical implementation usually requires approximations. In his original paper, Wilson made order of magnitude estimates of various terms contributing to the partition function of the Landau-Ginzburg model and derived the so called approximate recursion formula.<sup>[1]</sup>In this approximation, the RG transformation is reduced to a single integral equation which can be studied using numerical methods or functional analysis.

Recursion formulas closely related to the approximate recursion formula hold exactly for the hierarchical models. This is due to the large group of symmetries of the hamiltonians of these models. The RG transformation for these models has been studied in great detail and rigorous results concerning the epsilon expansion of the critical exponents are available in the literature. [3]

The fact that the RG transformation can be handled easily for hierarchical models suggests the use of these models as an approximation [4] for nearest neighbor models. The main technical problems in proceeding this way are: how to derive explicitly the approximate models and how to improve systematically the approximation. Recently, one of us [5] has answered these questions for the gaussian models where everything can be calculated explicitly. In order to extend this method to interacting models, one should be able to calculate the average value of perturbation terms added to the hierarchical hamiltonian. For Ising models, where the spin  $\sigma$  takes only the values  $\pm 1$ , this task can be carried numerically in an efficient way. It has been suggested [6] that such an approach might shed some light on Polyakov's conjecture [7] for the 3D Ising model. In preparation for these calculations, we first checked the agreement between the numerical calculations for the (unperturbed) hierarchical Ising model and analytical results. In doing so, we found surprising results which are reported in the following.

In this paper, we calculate numerically the magnetic susceptibility per site - the

susceptibility for short - of the hierarchical Ising model as a function of the temperature and for two values (0 and 1) of  $\epsilon$ , the parameter used in the  $\epsilon$ -expansion. Calculations have been carried with up to  $2^{18}$  sites. The numerical integration made use of the symmetries of the model in order to cut down the time of computation logarithmically. However, no approximations have been made and the numbers shown below are exact up to round-off errors. These errors were analyzed by changing from simple to double precision. In all the cases considered, this only affected the fifth significant digit of the susceptibility in the worse cases. Our calculations have been mostly restricted to the high temperature region and its boundary. In other words, the parameter  $\beta$ , proportional to the inverse temperature, will run between 0 and a critical value  $\beta_c$ . However, at the beginning, a few calculations will be made in the low temperature region in order to locate  $\beta_c$ .

Surprisingly, we found that the numerical data can be fitted very precisely with a simple power law of the form  $(1-\beta/\beta_c)^{-\gamma}$  in the whole high-temperature region. As a consequence, it is possible to calculate with good accuracy the free entries of this parametrization,  $\gamma$  and  $\beta_c$ , using the two first coefficients of the high-temperature expansion of the susceptibility. On the other hand, the values of  $\gamma$  obtained numerically, differ significantly from the values obtained in the  $\epsilon$ -expansion are incorrect, but only that for the models considered here, we can never get near the region (in the space of theories) where these approximations can be made.

This paper is organized as follows. In section 2, we introduce the hierarchical model and the numerical method of integration. In section 3, we check the scaling laws for the variance of the total spin and we determine  $\beta_c$ . In section 4, we analyze the temperature dependence of the susceptibility and we show that the data can be fitted very well with a simple power law. In section 5, we compare the results with the high temperature expansion and the  $\epsilon$ -expansion. Finally, we discuss our present understanding of the results in the conclusions.

### 2. The Hierarchical Ising Model

In this section we describe the hierarchical Ising model and the basic ideas of the numerical calculation performed. Hierarchical models <sup>[2]</sup> are specified by a non-local hamiltonian bilinear in the spin variables and a local measure of integration. We consider here the case of a Ising measure, where the spin take only the values  $\pm 1$ . The hamiltonian of a hierarchical model with  $2^n$  sites can be written as

$$H = -\frac{1}{2} \sum_{l=1}^{n} \left(\frac{c}{4}\right)^{l} \sum_{i_{n}, \dots, i_{l+1}} \left(\sum_{i_{l}, \dots, i_{1}} \sigma_{(i_{n}, \dots, i_{1})}\right)^{2}$$
(2.1)

For convenience, we have labeled the sites with n indices  $i_n....i_1$ , each index being 1 or 2. In order to visualize the meaning of this notation, one can divide the  $2^n$  sites into two boxes, each containing  $2^{n-1}$  sites. If  $i_n = 1$ , the site is the first box, if  $i_n = 2$ , the site is in the second box. Repeating this procedure n times (for the two boxes, their respective two sub-boxes, etc.), we obtain an unambiguous labeling for each of the sites. The interactions corresponding to a given value of l in (2.1) couple all the sites within each box of size  $2^l$  with the same strength  $(\frac{c}{4})^l$ . The model has a free parameter c for which we shall use the parametrization

$$c = 2^{1 - \frac{2}{D}}. (2.2)$$

The parameter of the epsilon-expansion will be defined as

$$\epsilon = 4 - D \tag{2.3}.$$

This choice has been justified in Ref.[6], but different conventions exist in the literature. In the actual calculations reported below, we have selected the values 0 and 1 for  $\epsilon$ .

We are interested in calculating the magnetic susceptibility of the hierarchical Ising model. This quantity can be calculated easily if we know the probability for the total spin denoted  $P_n(S)$ . This probability is obviously  $\beta$ -dependent even though we shall not write it explicitly. This probability will be calculated recursively using the RG method without a rescaling of the spins. We first integrate the spins inside boxes of size 2 keeping the sum of the spins in each box constant. We then include the terms with l=1 in (2.1) in a new local measure for these sums. We repeat this procedure n times and obtain a measure for the total spin which can be normalized as probability. Note that a choice of indices  $i_n, \ldots, i_{l+1}$  completely specifies a box of size  $2^l$  with the subdivision described above. We call the sum of the spin inside this box  $S_{i_n,\ldots,i_{l+1}}$ . It can take all the even values between  $-2^l$  and  $2^l$ . Obviously,  $S_{i_n,\ldots,i_{l+1}} = S_{i_n,\ldots,i_{l+1},1} + S_{i_n,\ldots,i_{l+1},2}$ . With these notations, the recursion formula reads

$$P_{l+1}(S_{i_{n},\dots,i_{l+1}}) = C_{l+1} Exp(\frac{1}{2}\beta(\frac{c}{4})^{l+1}(S_{i_{n},\dots,i_{l+1}})^{2})$$

$$\sum_{S_{i_{n},\dots,i_{l+1}}=S_{i_{n},\dots,i_{l+1},1}+S_{i_{n},\dots,i_{l+1},2}} P_{l}(S_{i_{n},\dots,i_{l+1},1})P_{l}(S_{i_{n},\dots,i_{l+1},2})$$

$$(2.4)$$

The constant  $C_{l+1}$  is adjusted in such a way that the sum of the probabilities add up to 1. Strictly speaking, it is not necessary to impose such a normalization during the intermediate steps of the calculation, however it keeps the numbers reasonably small. This recursion formula has been implemented with a computer program. In order to calculate  $P_n(S)$ , we have to repeat  $2^n$  times a calculation involving roughly  $2^n$  operations. Consequently, the time necessary to calculate  $P_n(S)$  scales approximately like  $4^n$ . With the fastest computer at our disposal, a DEC alpha 3000/400, it takes about 10 minutes to calculate  $P_{17}(S)$  from  $P_{16}(S)$  when programmed in FORTRAN.

Due to the size of the calculation, it is clearly necessary to check for round-off errors. We have studied the size of these errors by repeating the calculation with double-precision instead of simple precision for a large sample of values of  $\beta$ . For n up to 16, we have found very good agreement between the two calculations, the differences showing up in the sixth significant digit of the susceptibility. Differences

in the fifth digits were observed for n = 17 and 18. Calculations for n = 19 and beyond have shown less stability, require a lot of computer time and will not be reported here.

The main quantity of interest for us is the average of the square of the total spin denoted  $X_n(\beta)$  and defined as

$$X_n(\beta) = \sum_{S} P_n(S)S^2 \tag{2.5}$$

Above the critical temperature, this quantity divided by the number of sites has a finite thermodynamic limit. It will be called the magnetic susceptibility per site and denoted  $\chi_n(\beta)$ 

$$\chi_n(\beta) = \frac{X_n(\beta)}{2^n} \tag{2.6}$$

# 3. The Scaling Laws and the Determination of $\beta_c$

The RG method has definite predictions for the the large n behavior of  $X_n$ . This is explained at length for instance in chapter 7 of Parisi's textbook. In the following we shall just recall the main results and show that our numerical calculations reproduce these results with good precision. The case D=3 will be discussed in complete detail, while the results for D=4 will only be briefly commented upon.

It is convenient to express  $X_n$  as a power of the square of the number of sites, namely

$$X_n(\beta) = 2^{2n\omega(\beta,n)} \tag{3.1}$$

The value of  $\omega(\beta, n)$  can be easily estimated for large or small values of  $\beta$ . For large values of  $\beta$ , all the spins tend to align and  $\omega(\beta, n)$  gets close to 1. For small values of  $\beta$ , the spins at different sites become uncorrelated and the variance of the total spin can be approximated by the sum of the individual variances. In other words  $\omega(\beta, n)$  get close to 1/2. A more detailed analysis [8] shows that for large n,

 $\omega(\beta, n)$  is attracted by 1 if  $\beta$  exceeds a critical value denoted  $\beta_c$ , and by 1/2 if  $\beta$  is less than  $\beta_c$ . When  $\beta$  is exactly  $\beta_c$ ,  $\omega(\beta, n)$  tends to  $\frac{1}{2} + \frac{1}{D}$ . This can be derived from the fact that if we reabsorb a factor  $\sqrt{c/4} = 2^{\frac{1}{2} + \frac{1}{D}}$  in the spin variable after the integration described in the previous section, the Hamiltonian is invariant in the thermodynamic limit.

We shall now illustrate these results and calculate  $\beta_c$  for D=3. In order to get a rough idea of the value of  $\beta_c$ , we have plotted in Fig. 1 the trajectories  $\omega(\beta, n)$ from n=1 to n=16 and for  $\beta=0.1,0.2,...,1.6$ . It appears clearly that the separation between the two domains of attraction occurs for a value of  $\beta$  between 1.1 and 1.2. We have repeated this calculation for  $\beta=1.10,1.11,.....,1.20$  as shown on Fig. 2. This restricts  $\beta_c$  to the interval [1.17, 1.19]. At this point, it is more informative to consider ratios of susceptibilities at two successive values of n, because this quantity depends less sensitively on the constant of proportionality appearing in the scaling laws. A short calculation shows that

$$\lim_{n\to\infty} Log_2(\frac{\chi_{n+1}(\beta_c)}{\chi_n(\beta_c)}) = \frac{2}{D}$$
(3.3)

Fig. 3 displays  $Log_2(\frac{\chi_{n+1}(\beta)}{\chi_n(\beta)})$  for 3 values of  $\beta$  and for n up to 18. For  $\beta = 1.179$ , this quantity stays within one percent of the critical value 2/3 for the last six iterations. If  $\beta$  is increased or decreased by 0.001, the same quantity, departs by about 10 percent from 2/3 when n = 18. From this, we conclude that 1.179 approximates  $\beta_c$  with a precision better than 0.001.

Applying the same procedure for D=4, we found that  $\beta_c$  should be inside the interval [0.66, 0.67]. Fig. 4 shows that 0.665 approximates  $\beta_c$  with a precision better than 0.001.

### 4. Fitting the numerical data with a simple power law

In this section, we report our numerical results concerning the  $\beta$ -dependence of the magnetic susceptibility per site at D=3 and D=4 and for  $\beta < \beta_c$ . Again, we start with the case D=3 and discuss it in detail while the case D=4 will be presented more rapidly later in this section.

We have calculated  $\chi_n(\beta)$  for D=3 and n up to 16 and for values of  $\beta$  between 0 and 1.18 separated by intervals of length 0.01. As expected, the susceptibility rises sharply when  $\beta$  gets close to 1.18. We have displayed the results for n=16 and  $\beta \leq 1.1$  on Fig. 5. Results for n=14 or n=15 would have been hardly distinguishable from n=16 on a graph of this size. On the other hand, the n-dependence becomes more sizable when  $\beta$  is closer to its critical value as shown on Fig. 6 which also includes results for n=17. Except for the difference of scales, the resemblance between Fig. 5 and Fig. 6 is striking. This can be understood from the fact that the numerical data can be fitted very precisely with a simple power law of the form

$$\chi_n(\beta) = (1 - \beta/\beta_c)^{-\gamma} \tag{4.1}$$

in the *whole* interval [0, 1.18]. In order to justify this claim, we first notice that (4.1) implies the inverse logarithmic derivative of the susceptibility is a linear function, namely

$$\left(\frac{d}{d\beta}Log(\chi_n(\beta))\right)^{-1} = (\gamma)^{-1}(\beta_c - \beta) \tag{4.2}$$

We now approximate this inverse logarithmic derivative by

$$\Delta \beta / \Delta Log(\chi_n(\beta + \Delta \beta/2)) = \frac{\Delta \beta}{Log(\chi_n(\beta + \Delta \beta)) - Log(\chi_n(\beta))}$$
(4.3)

with  $\Delta\beta = 0.01$ , the interval used here. This function is plotted in Fig. 7 for n = 16. Remarkably, the numerical data is barely distinguishable from the least

square linear fit

$$\Delta \beta / \Delta Log(\chi_{16}(\beta)) = 0.80839 - 0.67843\beta \tag{4.4}$$

The graphs for n=14 or 15 are almost identical. The linear fits are respectively

$$\Delta \beta / \Delta Log(\chi_{15}(\beta)) = 0.80849 - 0.67712\beta$$
  
$$\Delta \beta / \Delta Log(\chi_{14}(\beta)) = 0.80867 - 0.67508\beta$$
(4.5)

In order to see the corrections to the linear behavior, we have plotted in Fig.8 the difference between the linear fit and the data, denoted  $E(\beta)$ , for n=14, 15 and 16. These difference are not larger than 0.003 in absolute values and have interesting regularities. The reasons why the power law (4.1) is so accurate and the nature of its corrections are being investigated.

We have intentionally used simple precision data to plot Fig. 8 in order to give an idea of the round-off errors of the method. Small irregularities are visible especially in the low  $\beta$  region where their typical size is  $10^{-4}$ . On the other hand,  $E(\beta)$  is smoother for  $\beta > 0.8$ . The size of these numerical errors is compatible with the claim made in section that numerical errors affects only the sixth significant digit of the susceptibility. Indeed, if one of the  $\chi_n$  is replaced by  $\chi_n(1+\delta)$  in Eq. (4.3), this creates an error  $\delta\Delta\beta/(\Delta Log(\chi))^2$ . A simple inspection shows that  $\delta$  can be amplified by a factor of order 100 if  $\beta$  is not too close to  $\beta_c$ .

A similar procedure has been followed for D=4. In Fig. 9 we desplay the quantity  $\Delta\beta/\Delta Log(\chi_{16}(\beta))$ . Again the departures from linearity are small and the changes in the coefficients of the linear fit for n=14 and 15 are of the same order as those for D=3. The difference between the linear fit and the data for n=14, 15 and 16 is displayed on Fig.10.

### 5. Comparisons with High-Temperature and Epsilon Expansions

Approximate values of the critical exponent  $\gamma$  can be extracted from linear fits presented in the last section. For instance, identifying Eqs.(4.2) and (4.4) yields  $\gamma = 1.474$  and  $\beta_c = 1.192$ . Changes of 1 or 2 percent can be obtained if we restrict  $\beta$  to intervals closer to  $\beta_c$ . These results can be compared with analytical results obtained using the high temperature or the  $\epsilon$ -expansion.

The high-temperature expansion of the susceptibility reads

$$\chi_n(\beta) = 1 + b_{(1,n)}\beta + b_{(2,n)}\beta^2 + \dots$$
 (5.1)

A straightforward but tedious calculation yields

$$b_{(1,n)} = \left(1 - \frac{c}{4}\right)^{-1} \left(\frac{c}{4} \left(1 - \left(\frac{c}{2}\right)^n\right) \left(1 - \frac{c}{2}\right)^{-1} - \left(2^n - 1\right) \left(\frac{c}{4}\right)^{n+1}\right)$$
 (5.2.1)

and

$$b_{(2,n)} = (b_{(1,n)})^2 - (1 - \frac{c}{4})^{-2} \left[ \left( \frac{c}{4} \right)^2 \left( 1 - \left( \frac{c^2}{8} \right)^n \right) \left( 1 - \frac{c^2}{8} \right)^{-1} - 2\left( \frac{c}{4} \right)^{n+2} \left( 1 - \left( \frac{c}{2} \right)^n \right) \left( 1 - \frac{c}{2} \right)^{-1} - (2^n - 1)\left( \frac{c}{4} \right)^{2(n+1)} \right]$$

$$(5.2.2)$$

Comparing Eq.(5.1) and Eq.(4.2) expanded about 0, we obtain

$$\gamma = \left(\frac{2b_{(2,n)}}{b_{(1,n)}^2} - 1\right)^{-1} \tag{5.3}$$

On the other hand, the critical exponents of the hierarchical model near Wilson's non-trivial fixed point can be calculated using the  $\epsilon$ - expansion. With the convention of Eq.(2.2), we obtain at first order in  $\epsilon$ .

$$\gamma = 1 + 0.1667\epsilon \tag{5.4}$$

These analytical results are compared with the numerical results in Fig.11 and 12 for D=3 and D=4 respectively. It appears clearly that the data agree within

a few percent with the high-temperature calculation performed above, while a significant departure from the  $\epsilon$ -expansion is shown. This is quite clear in the case D=4 where the  $\epsilon$  expansion predicts  $\gamma=1$ . This is less clear in the case D=3 where we do not know at which order we need to truncate what is presumably an asymptotic series.

#### 6. Conclusions

We have studied the magnetic susceptibility per site of the hierarchical Ising model. We found that the  $\beta$ -dependence of this quantity could be fitted quite accurately with a simple power law. It would be quite interesting to understand the origin of this result. Small corrections to this power law have been observed and it might be possible to handle this problem using Callan-Symanzik's equations. In any case, it is remarkable that one is able to obtain an accurate information about the critical behavior of a model just by calculating two coefficients of a high temperature expansion.

We insist on the fact that we have made no approximations in our numerical calculations and that the results presented are exact up to round-off errors. We have analyzed these errors and found that they do not affect our conclusions. The fact that we were able to reproduce accurately well-understood analytical results such as the scaling laws at the critical temperature and the high temperature behavior of the susceptibility seem to rule out errors in implementing Eq. (2.4) numerically.

The discrepancy with the  $\epsilon$ -expansion is quite surprising. Our limited data cannot rule out the possibility that this discrepancy decreases very slowly with n. Another possibility is that by starting with Ising spins and applying the RG transformation, we never get close enough to the region (in the space of theories) where the  $\epsilon$ -expansion is legitimate. This second possibility seems supported by preliminary results <sup>[9]</sup> obtained in the field theory approach of this problem

Acknowledgements: One of us (Y.M.) would to thank the participants of the Aspen Center for Physics in June and July 1993 for providing a stimulating atmosphere while some initial motivations for this work where found. This work was partially supported by the Carver foundation and NSF grant PHY-9103914.

## REFERENCES

- K. Wilson, Phys. Rev. B 4 (1971) 3185; K. Wilson and J. Kogut, Phys. Rep. 12 (1974) 75.
- F. Dyson, Comm. Math. Phys. 12 (1969) 91; G. Baker, Phys. Rev. B 5 (1972) 2622.
- P. Bleher and Y. Sinai, Comm. Math. Phys. 45 (1975) 247; P. Collet and
   J. P. Eckmann, Comm. Math. Phys. 55 (1977) 67.
- 4. K. Gawedzki and A. Kupiainen, Les Houches 1985, K. Osterwalder and R. Stora, Editors
- 5. Y. Meurice, Univ. of Iowa Preprint 92-12, hep-th 9307128.
- 6. Y. Meurice, Mod. Phys. Lett. **7** (1992) 3331 and Jour. Math. Phys. **35** (1994) 0000 .
- A. Polyakov, Phys. Lett. 82B (1979) 247 and Phys. Lett. 103B (1981) 211
- 8. G. Parisi, *Statistical Field Theory* (Addison-Wesley, New-York, 1988) and references therein.
- 9. Y. Meurice and G. Ordaz, work in progress.

#### FIGURE CAPTIONS

- Fig. 1 :  $\omega(n,\beta)$  versus n for D=3 and values of  $\beta$  going from 0.1 to 1.6 by steps of 0.1
- Fig. 2 :  $\omega(n,\beta)$  versus n for D=3 and values of  $\beta$  going from 1.1 to 1.2 by steps of 0.01
- Fig. 3:  $Log_2$  of the ratio of two successive values of the susceptibility for D=3 and for  $\beta=1.178, 1.179$  and 1.180.
- Fig. 4:  $Log_2$  of the ratio of two successive values of the susceptibility for D=4 and for  $\beta=0.6645,\,0.6655$  and 0.6665.
- Fig. 5: The magnetic susceptibility for D=3, n=16 and values of  $\beta$  going from 0 to 1.18 by steps of 0.01.
- Fig. 6: The magnetic susceptibility for D=3, n=14, 15, 16 and 17, and values of  $\beta$  going from 1.00 to 1.18 by steps of 0.01.
- Fig. 7: A discrete version of the inverse logarithmic derivative of the susceptibility and its linear fit for D = 3 and n = 16.
- Fig. 8: Difference between the discrete version of the inverse logarithmic derivative of the susceptibility and its linear fit for D=3 and n=14, 15 and 16.
- Fig. 9: A discrete version of the inverse logarithmic derivative of the susceptibility and its linear fit for D=4 and n=16.
- Fig. 10: Difference between the discrete version of the inverse logarithmic derivative of the susceptibility and its linear fit for D=4 and n=14, 15 and 16.
- Fig. 11: Comparison between the high temperature calculation and our numerical estimation of the critical exponent  $\gamma$  for D=3 and n between 5 and 16. The solid line is the result for the  $\epsilon$ -expansion at first order.
- Fig. 12: Comparison between the high temperature calculation and our numerical estimation of the critical exponent  $\gamma$  for D=4 and n between 5 and 16. The solid line is the result for the  $\epsilon$ -expansion.

This figure "fig1-1.png" is available in "png" format from:

This figure "fig2-1.png" is available in "png" format from:

This figure "fig3-1.png" is available in "png" format from:

This figure "fig1-2.png" is available in "png" format from:

This figure "fig2-2.png" is available in "png" format from:

This figure "fig3-2.png" is available in "png" format from:

This figure "fig1-3.png" is available in "png" format from:

This figure "fig2-3.png" is available in "png" format from:

This figure "fig3-3.png" is available in "png" format from:

This figure "fig1-4.png" is available in "png" format from:

This figure "fig2-4.png" is available in "png" format from:

This figure "fig3-4.png" is available in "png" format from: