

On the relevance of center vortices to QCD

Philippe de Forcrand¹ and Massimo D'Elia²

¹SCSC, ETH-Zurich, CH-8092 Zurich, Switzerland

²Dipartimento di Fisica dell'Università and INFN., I-56127, Pisa, Italy

(February 5, 2020)

In a numerical experiment, we remove center vortices from an ensemble of lattice SU(2) gauge configurations. This removal adds short-range disorder. Nevertheless, we observe long-range order in the modified ensemble: confinement is lost and chiral symmetry is restored (together with trivial topology), proving that center vortices are responsible for both phenomena. As for the Abelian monopoles, they survive but their percolation properties are lost. These statements hold regardless of the Gribov copy chosen for center projection.

PACS numbers: 11.15.Ha, 12.38.Aw, 12.38.Gc, 11.30.Rd

The essential non-perturbative properties of QCD are confinement and chiral symmetry breaking. It has been observed through numerical lattice simulations that these two properties persist in the quenched theory (where the feedback of quarks on gluons is ignored) until the temperature is raised to a critical value $T_c \approx 220$ MeV, where confinement is lost; chiral symmetry appears to be simultaneously restored [1]. The disorder which leads to the area law for the Wilson loop thus seems to be somehow tied to the existence of a chiral condensate. Effective mechanisms have been proposed to explain confinement or chiral symmetry breaking. But a successful effective model should provide a common explanation for both.

Two effective descriptions of QCD have been receiving a lot of attention: one considers instantons as the effective degrees of freedom, the other chromomagnetic monopoles. Instantons are natural candidates to explain chiral symmetry breaking, because each instanton is associated with a zero mode of the Dirac operator [2], and there must be an accumulation of zero eigenvalues to obtain a quark condensate at low temperature [3]. Above T_c , the vanishing of the quark condensate is accompanied by a qualitative change in the instanton ensemble (see, e.g., [4]). On the other hand, it is unlikely that instantons play a significant role in confinement (see [5] for a recent discussion). An attractive mechanism for confinement is dual superconductivity of the QCD vacuum [6]. Considerable evidence for this dual Meissner effect has been accumulated on the lattice, including a disorder parameter demonstrating the condensation of chromomagnetic monopoles below T_c [7]. This condensation has been observed directly after gauge-fixing to 'Maximal Abelian Gauge' [8], and the idea of 'Abelian dominance' has emerged, according to which the Abelian degrees of freedom of the Yang-Mills field encode all its long-distance

properties. Indeed, chiral symmetry breaking and its restoration have been observed in the Abelian sector [9]. The Abelian dominance scenario has some flaws however: it does not explain the breaking of the adjoint string; and the Abelian string tension differs slightly from the Yang-Mills one [10]. Moreover, degrees of freedom more elementary than the Abelian monopoles, embedded in them and solely responsible for the physics assigned to them, cannot be ruled out.

The idea of center vortices, which initially failed due to their misidentification [11], has been successfully revived, first as an embedded model inside the Abelian sector [12], then without reference to Abelian projection [13]. Center vortices are exposed by gauge-fixing: after a gauge transformation which brings each lattice link as close as possible to a center element of the gauge group, vortices consist of defects in the projected gauge field formed from those center elements alone. The idea of center dominance is again that the center degrees of freedom (d.o.f.) encode all the long-distance physics. This is aesthetically attractive because the d.o.f. are decimated even more than in the Abelian projection. Furthermore, the density of center vortices seems to be a well-defined continuum quantity; the center string tension appears to more or less match the original one; and an explanation for the behaviour of the adjoint potential has been proposed [14]. However, chiral symmetry has not yet been studied in this context.

An additional problem with Abelian and center dominance is that the relevant d.o.f. are identified only after gauge-fixing. Gauge-fixing non-Abelian fields is notoriously ambiguous, and different Gribov copies produce different Abelian monopoles or center vortices, whose properties like the string tension differ slightly. For this reason, we are a priori suspicious of effective models which involve gauge-fixing, and so we designed a simple numerical experiment to disprove the center-dominance scenario. For simplicity, we consider the gauge group SU(2), with center Z_2 . Our starting point is an ensemble of lattice gauge fields representative of the continuum. We identify center vortices in this ensemble, and construct from it a modified ensemble where all center vortices have been removed by flipping the sign of a subset of SU(2) gauge links. As expected, this operation introduces a lot of disorder in the gauge field. Nonetheless, these disordered gauge fields now have a trivial, vortex-free center projection, and so according to the credo of center dominance they should not confine. In other words, our intro-

duction of short-range disorder should at the same time bring long-range order. To our surprise, this is indeed what happens. On this vortex-free ensemble, confinement is lost. Emboldened by this finding, one may ask if the spectral properties of the Dirac operator are not also dominated by the center components of the gauge field. In that case, our modified ensemble should show no sign of chiral symmetry breaking, since its center projection is the trivial (perturbative) vacuum. Indeed, this is what we observe: removal of center vortices causes both loss of confinement and restoration of chiral symmetry.

The next intriguing question regards the fate of Abelian monopoles as center vortices are removed. They do not disappear; on the contrary, the introduction of short-range disorder increases their number. However, we observe the complete disappearance of monopole current loops winding around the periodic lattice: we can thus identify these as the fundamental objects associated with confinement in the Abelian sector, apparently influenced by the underlying center degrees of freedom.

Finally, we investigated the effect of Gribov copies which caused our initial skepticism. We repeated our experiment on the same SU(2) ensemble, but introduced a systematic tolerance in the gauge condition to be satisfied before identifying the center vortices to be removed. Although the location and number of center vortices were removed varied appreciably, the modified SU(2) ensemble was always non-confining and chirally symmetric.

The numerical experiment { We start from an ensemble of SU(2) lattice gauge fields obtained by Monte Carlo using the standard Wilson plaquette action. We want to generate a modified ensemble, free of center vortices, and compare the static potential and the chiral condensate on the original and modified ensembles.

To identify center vortices, we gauge-fix the original configurations in order to bring each SU(2) gauge link $U(x)$ as close as possible to an element of the center $Z_2 = \{ \pm 1 \}$. We therefore try to iteratively maximize

$$Q(\text{fUg}) = \sum_x (\text{Tr} U(x))^2; \quad (1)$$

as in [13], where this gauge is called the "direct maximal center gauge." The gauge-fixed SU(2) links, denoted $U^{GF}(x)$, are then projected to Z_2 elements $Z(x)$ using

$$Z(x) = \text{sign}[\text{Tr} U^{GF}(x)]; \quad (2)$$

Plaquettes in the Z_2 -projected theory with value -1 represent defects of the Z_2 gauge field called P-vortices [12]. Numerical evidence has been presented [12,13] showing that plaquette-like P-vortices signal the presence of macroscopic, physical excitations, called center vortices, in the unprojected original SU(2) configuration.

Consider then the modified SU(2) configuration made of gauge links $U^0(x)$ constructed as

$$U^0(x) = Z(x) U(x); \quad (3)$$

The gauge transformation which maximizes $Q(\text{fUg})$ in (1) also gives the same maximum to $Q(\text{fU}^0\text{g})$, so that the modified gauge-fixed links $U^{GF}(x)$ are simply $Z(x) U^{GF}(x)$. Therefore, we instantly know the center projection $Z^0(x)$ of $U^0(x)$:

$$\begin{aligned} Z^0(x) &= \text{sign}[\text{Tr} U^{GF}(x)] = \text{sign}[\text{Tr} Z(x) U^{GF}(x)] \\ &= \text{sign}[\text{Tr} U^{GF}(x)]^2 = +1; \end{aligned} \quad (4)$$

Every configuration in our modified ensemble thus projects onto the trivial Z_2 vacuum: all center vortices have been removed.

Our ensemble consists of about 1000 SU(2) configurations, on a 16^4 lattice at $\beta = 2.4$. To maximize $Q(\text{fUg})$ (Eq.1) we use standard overrelaxation, stopping when

$$\sum_x (\text{Tr} U(x))^2 < 10^{-6} \quad (5)$$

from one gauge-fixing sweep to the next.




FIG. 1. Normalized plaquette distribution on the original and modified ensembles (SU(2), $\beta = 2.40$). Center-vortex removal increases short-range disorder.

In Fig.1 we show the distribution of SU(2) plaquette values on the original and modified ensembles. It is apparent that under the sign flip Eq.(3), many SU(2) plaquettes acquire a negative value. Removal of the center vortices increases the action: the modified ensemble has more short-range disorder.




FIG. 2. Creutz ratios on the original and modified ensembles. The dashed band is the string tension result of [15].

Results { In Fig2 we present our results for the Creutz ratios $\chi(R;R)$ $\ln[\chi(R;R)/\chi(R-1;R-1)] = \ln[\chi(R;R-1)/\chi(R-1;R-1)]$ constructed from the averages $\chi(R;T)$ of R by T Wilson loops on the original and modified ensembles. For large R , $\chi(R;R)$ tends to the string tension. It is clear that on the modified ensemble the Creutz ratios decrease and tend to zero. In spite of the increased short-range disorder, long-range order has been created and confinement has been lost.

This is even clearer if one looks directly at the Wilson loop values. In Fig3 we show $\ln\chi(R;T)$ as a function of T . For a fixed R , points at successively larger T form a line whose asymptotic slope is the value of the static potential $V(R)$. The lines corresponding to the modified ensemble are parallel, indicating that $V(R)$ does not grow with R : the string tension has vanished.




FIG. 3. Wilson loop values $\ln\chi(R;T)$ on the original and modified ensembles. Note the parallel lines for successive R in the latter: upon center-vortex removal, confinement is lost.

Fig.4 illustrates our study of chiral symmetry on the original and modified ensembles. As is well-known, chiral symmetry breaking cannot occur on a finite lattice. Therefore, we measure $\chi(m_q) = \langle \text{Tr}(\bar{\psi}\psi + m_q) \rangle$ for a range of quark masses m_q where finite-size effects are small, and extrapolate to $m_q \rightarrow 0$. In the original ensemble, χ clearly extrapolates to a non-zero value which signals chiral symmetry breaking. In the modified ensemble, the extrapolated value is zero within errors: the removal of center vortices restores chiral symmetry. We expect then the instanton content of the Yang-Mills field to be modified also. To check this, we use improved cooling [16] to measure the topological charge of the modified field: the striking result is that the removal of center vortices always leads to the trivial topological sector.

We therefore have clear numerical evidence for "center dominance": if the center-projected field is the trivial vacuum with all links equal to 1, as in the modified ensemble, then the Yang-Mills field shows the long-distance

properties of the trivial vacuum, i.e., no confinement, no quark condensate and no topology. The long-distance properties of the Yang-Mills field appear to be determined by its center projection.




FIG. 4. Quark condensate $\chi(m_q)$ on the original and modified ensembles. The dashed line corresponds to "poor" center-vortex identification (see text). In all cases, center-vortex removal restores chiral symmetry.

On the other hand, a large number of studies now support the alternative scenario of "Abelian dominance." We use our approach of center-vortex removal to directly assess the relationship between these two scenarios.

In a first experiment, we construct the Abelian projection of our original SU(2) ensemble by gauging to Maximal Abelian Gauge in the usual way [8], then identify and remove center vortices from the Abelian sector. While the original Abelian-projected ensemble shows confinement, with a string tension similar to the non-Abelian one, the modified Abelian-projected configurations do not confine. Therefore, we find no contradiction between "Abelian dominance" and "center dominance." The latter simply appears more fundamental because of the greater reduction of the number of d.o.f.

In a second experiment, we look at clusters of Abelian monopole currents, whose percolation has been identified as the signal for confinement [17], obtained from the original and modified ensembles. We find that the removal of center vortices changes the distribution of monopole cluster sizes in a crucial way (see Fig.5): whereas in the original ensemble, each configuration contains typically one very large, percolating monopole cluster and many very small ones, the modified ensemble gives a more homogeneous size distribution, with a handful of large clusters per configuration. These large clusters are the remnants of the very large one, broken into pieces by the vortex removal. Some of them still percolate, even though confinement has disappeared. Therefore, we are led to associate confinement with a more specific feature of the monopole clusters: monopole current loops which wind around the periodic lattice. Such loops can be found frequently on the original, confining ensemble, but never on the modified, non-confining one. We conclude that: (i) confinement manifests itself in the Abelian sector by the pres-

ence of monopole current loops with non-trivial topology; (ii) center-vortex removal, which destroys confinement, always finds the "weak links" of these non-trivial loops and breaks them into trivial pieces.




FIG. 5. Size distribution of monopole clusters on the original and modified ensembles.

Now let us consider the issue of gauge-fixing ambiguities, which was the reason for our initial skepticism about the center-vortex idea. These ambiguities come from the structure of Q (Eq.1), which has many local maxima, any of which can be selected by a local iterative maximization algorithm. Each local maximum, or Gribov copy, will have its own set of P-vortices, differing in number and location. The proposal of [12] is that, no matter which Gribov copy one chooses, P-vortices are the traces of physical center vortices and are roughly located at their center. This argument may account for P-vortices differing in location but not in number. To study this question in more detail, we magnified the effect of gauge-fixing ambiguities, by stopping our iterative algorithm early, as soon as $\langle E \rangle < 10^2$. Thus we not only explore a different basin of attraction of Q , but we do not even stop at a local maximum. One effect of this partial gauge-fixing is expected: the density of P-vortices increases from 5.504% to 7.414%, i.e. shows an increase of 1.91%. The string tension we measure in the Z_2 -projected ensemble increases accordingly: whereas the Z_2 string tension after "complete" gauge-fixing ($a^2 = 0.075$) is a little larger than but compatible with the non-Abelian string tension (0.0708–0.0711 from [15], see Fig.2), it jumps to 0.12 after partial gauge-fixing. What is remarkable is that this increase

0.045 is similar to that obtained by placing the surplus of P-vortices at random, uncorrelated locations: $\ln(1.2)$. This does not prove, but makes it plausible, that the center projection always captures the core degrees of freedom relevant for long-distance properties, plus a varying amount of unrelated noise. Indeed, it has been argued that gauge-fixing is not even necessary for center projection [18]. In our description, center-gauge-fixing acts as an UV noise-filtering device, with different Gribov copies letting through different noise components. Further evidence for this is obtained by removing from the original SU(2) ensemble the center

vortices identified after partial gauge-fixing only. Just as for "complete" gauge-fixing, we observe that confinement is lost, chiral symmetry restored, and the topology trivial. The only difference is that the modified ensemble now has much more short-range disorder.

In conclusion, we have shown that removal of center-vortices from an SU(2) Yang-Mills ensemble causes the loss of confinement and the restoration of chiral symmetry. Center vortices appear as effective degrees of freedom in terms of which both phenomena can be successfully described. It would be desirable, of course, to formulate an effective action for the center-projected theory. Ref. [19] considers an extension of the Nambu-Goto action, where the fundamental d.o.f. are 2-dimensional random surfaces dual to the P-vortices. Ref. [20] instead proposes to consider center monopoles and their world-lines. We suggest identifying a "minimum spanning tree" of negative Z_2 links responsible for the P-vortices: perhaps only a subset of them form the essential d.o.f. governing the long-distance properties.

Finally, our vortex-removal procedure can be used to study properties of non-confining non-Abelian fields and effects of center-symmetry breaking. For instance, removing time-like center disorder only would be similar to raising the temperature above T_c .

We thank Adriano Di Giacomo, Maarten Golterman, Je G Reensite and O sam u M iyam ura for discussions.

-
- [1] J. Kogut et al., Phys. Rev. Lett. 50 (1983) 393.
 - [2] G. 't Hooft, Phys. Rev. D 14 (1976) 3432.
 - [3] T. Banks and A. Casher, Nucl. Phys. B 169 (1980) 103.
 - [4] T. Schafer and E. V. Shuryak, hep-ph/9610451.
 - [5] R. C. Brower et al., hep-lat/9809091.
 - [6] G. 't Hooft, Nucl. Phys. B 190 (1981) 455; S. Mandelstam, Phys. Rep. 23C (1976) 245.
 - [7] A. Di Giacomo and G. Pauti, Phys. Rev. D 56 (1997) 6816.
 - [8] H. Shiba and T. Suzuki, Phys. Lett. B 333 (1994) 461.
 - [9] O. M. Yamura, Phys. Lett. B 353 (1995) 91.
 - [10] G. S. Bali et al., Phys. Rev. D 54 (1996) 2863.
 - [11] G. Mack and E. Pietarinen, Nucl. Phys. B 205 [FS5] (1982) 141.
 - [12] L. Del Debbio et al., Phys. Rev. D 55 (1997) 2298.
 - [13] L. Del Debbio et al., Phys. Rev. D 58 (1998) 094501.
 - [14] M. Faber et al., Phys. Rev. D 57 (1998) 2603.
 - [15] C. Michael and M. Teper, Phys. Lett. B 199 (1987) 95.
 - [16] P. de Forcrand et al., Nucl. Phys. B 499 (1997) 409.
 - [17] E. G. T. L. Ivanenko et al., Phys. Lett. B 302 (1993) 458.
 - [18] M. Faber et al., hep-lat/9810008.
 - [19] M. Engelhardt, talk at the RIKEN-BNL workshop on QCD Phase Transitions, Nov. 1998.
 - [20] M. N. Chernodub et al., hep-lat/9809158.