

Constraints of the 4th generation from $g_\mu - 2$

Wu-Jun Huo and Tai-Fu Feng

CCAST (World Lab.), P.O. Box 8730, Beijing 100080

and

Institute of High Energy Physics, Academia Sinica, P.O. Box 918(4),

Beijing 100039, P.R. China

Abstract

We investigate the newly observed muon $g_\mu - 2$ anomaly in the framework of a sequential fourth generation model with a heavy fourth neutrino, ν' . We find that $g_\mu - 2$ can exclude most values of $m_{\nu'}$ and put a very stringent constraint on the existence of the fourth generation. We also obtain bounds of a 4×4 leptonic mixing matrix elements, $V_{2\nu'}$ if it really exists.

At present, Standard model (SM) has to face the experimental difficulties which are all relate to leptons. It seems to indicate the presence of new physics just round the corner will be in the leptonic part. Firstly, there are convincing evidences that neutrinos are massive and oscillate in flavor [1]. Secondly, the recent measurement of the muon anomalous magnetic moment by the experiment E821 [2] at Brookhaven National Laboratory disagrees with the SM expectations at more than 2.6σ level.

Defined as $a_\mu \equiv (g_\mu - 2)/2$, the recent measurement of a_μ is

$$a_\mu^{\text{exp}} = (11\,659\,202 \pm 14 \pm 6) \times 10^{-10}, \quad (1)$$

where the SM prediction is

$$a_\mu^{\text{SM}} = (11\,659\,159.7 \pm 6.7) \times 10^{-10}, \quad (2)$$

Thus, one finds

$$a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (42.6 \pm 16.5) \times 10^{-10}. \quad (3)$$

which gives the old 2.6σ deviation. But the revised difference between experiment and SM is

$$\delta a_\mu^{\text{SM}} \equiv a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = 26(16) \times 10^{-10} \quad (4)$$

which is now only a 1.6σ deviation [3]. Although the deviation drops from 2.6σ to 1.6σ , it might seem that the absolute magnitude of the deviation may be a hint of new physics. There have been a lot of scenarios of new physics proposed to interpret the non-vanishing and positive value of δa_μ . In this note, we consider the sequential fourth generation standard model (SM4) to investigate its contributions to a_μ .

From the point of phenomenology, there is a realistic question what are numbers of the fermions generation or weather there are other additional quarks or leptons. The present experiments tell us there are only three generation fermions with *light* neutrinos which mass are less smaller than $M_Z/2$ [4]. But the experiments don't exclude the existence of other additional generation, such as the fourth generation, with a *heavy* neutrino, i.e.

$m_{\nu_4} \geq 45.5\text{GeV}$ [5]. Many refs. have studied the SM4 [6], which is added an up-like quark t' , a down-like quark b' , a lepton τ' , and a heavy neutrino ν' in the SM. The properties of these new fermions are all the same as their corresponding counterparts of other three generations except their masses and CKM mixing, see Tab.1,

	up-like quark	down-like quark	charged lepton	neutral lepton
SM fermions	u	d	e	ν_e
	c	s	μ	ν_μ
	t	b	τ	ν_τ
new fermions	t'	b'	τ'	ν'

Table 1: The elementary particle spectrum of SM4

If there exists a very heavy fourth neutrino ν' , it can contribute to a_μ through diagram of Fig. 1. This is a electroweak interaction. Similar to that of quarks, the corresponding Lagragian is

$$\mathcal{L} = -\frac{g}{\sqrt{2}}(\bar{\nu}'\gamma_\mu a_L V_{2\nu'}^l \mu)W^\mu + h.c. \quad (5)$$

where $a_L = (1-\gamma_5)/2$, $V_{2\nu'}^l$ is the (2,4) element of the four-generation CKM matrix (4×4),

$$V_{\text{CKM}}^{\text{SM4}} = \begin{pmatrix} V_{1\nu_e} & V_{1\nu_\mu} & V_{1\nu_\tau} & V_{1\nu'} \\ V_{2\nu_e} & V_{2\nu_\mu} & V_{2\nu_\tau} & V_{2\nu'} \\ V_{3\nu_e} & V_{3\nu_\mu} & V_{3\nu_\tau} & V_{3\nu'} \\ V_{4\nu_e} & V_{4\nu_\mu} & V_{4\nu_\tau} & V_{4\nu'} \end{pmatrix} \quad (6)$$

Reverting back to the diagrams of Fig. 1, we see that the fourth neutrino contribution to a_μ is

$$a_\mu^{\text{SM4}} = \alpha \left(\frac{g}{\sqrt{2}} V_{2\nu'}^l \right)^2 \frac{2m_\mu^2}{m_W^2} \cdot f(x), \quad (7)$$

where $x \equiv m_{\nu'}^2/m_W^2$, α is the fine construction constant and

$$f(x) = \frac{-5x^3 - 5x^2 + 4x}{12(x-1)^3} + \frac{(2x^3 - x^2) \log x}{2(x-1)^4}. \quad (8)$$

We suppose that the 1.6σ discrepancy of muon anomalous magnetic moment, δa_μ^{SM} , is induced by the fourth sequential neutrino ν' . We can use the above equation to get parameter space of $f(x)$ and $V_{2\nu'}^l$ to $m_{\nu'}$ (see Fig. 2 and 3),

$$(V_{2\nu'}^l)^2 = \frac{\sqrt{2} \cdot \delta a_\mu^{\text{SM}}}{8G_F \alpha m_\mu^2 \cdot f(x)} \quad (9)$$

From Fig. 2, we can see that $f(x)$ is taken negative values except for a very narrow range of $m_{\nu'}$ which is from 58GeV to 80GeV . In other words, the sign of a_μ^{SM4} is only related to ν mass. Only in this narrow mass range, ν' gives positive contribution to $g_\mu - 2$. The low bound of $m_{\nu'}$ we get from $g_\mu - 2$ is consistent with the present experiments [4]. But the upper bound, $m_{\nu'} < 80\text{GeV}$, seems to conflict with the current experiments statue which there is no any new physics signals upper to several GeVs. The fourth generation particles seems not to be so light. Moreover, from Fig. 3, if we consider the unitarity of the matrix V_{CKM}^{SM4} and tiny values of its elements, the reasonable value range of m'_μ will be more narrow.

In summary, we calculate the contribution of the fourth generation to $g - 2$ and get an interesting result which we can exclude most values of $m_{\nu'}$. Considering the new revised deviation of $g - 2$, we give the parameter space of $m_{\nu'}$ and lepton mixing matrix element $V_{2\mu'}$. We find that $g - 2$ can constrain on the neutrino mass of the fourth generation: i.e. its mass should be heavier than 58 GeV and lighter than 80 GeV . It seems that from the lepton part, the current experiments can impose a stringent constraint on the existence of the fourth generation.


Acknowledgments

This research is supported by the the Chinese Postdoctoral Science Foundation and CAS K.C. Wong Postdoctoral Research Award Fund. We are grateful to prof. C.S. Huang and Prof. X.M Zhang for useful discussions.

References

- [1] Y. Fukuda *et al.*, Phys. Lett. **B436** 33 (1998); Phys. Rev. Lett. **81**, 1562 (1998).
- [2] Muon g-2 Collaboration, H.N. Brown *et al.*, Phys. Rev. Lett. **86**, 2227 (2001).
- [3] U. Chattopadhyay, A. Corille and P. Nath, hep-ph/0202275; M. Knecht and A. Nyffeler, hep-ph/0111058; M. Knecht, A. Nyffeler, M. Perrottet and E. de Pafael, Phys. Rev. Lett. **88**, 071802 (2002); M. Hayakawa and T. Kinsoshita, hep-ph/0112102; I. Blokland, A. Czarnecki and K. Melnikov, Phys. Rev Lett. **88**, 071803 (2002); J. Bijnens, E. Pallante and J. Prades, hep-ph/0112255.
- [4] G.S. Abrams et al., Mark II Collab., Phys. Rev. Lett. **63** (1989) 2173; B. Advera et al., L3 Collab., Phys. Lett. B **231** (1989) 509; I. Decamp et al., OPAL Collab., *ibid.*, **231** (1989) 519; M.Z. Akrawy et al., DELPHI Collab., *ibid.*, **231** (1989) 539.
- [5] Z. Berezhiani and E. Nardi, Phys. Rev. D **52** (1995) 3087; C.T. Hill, E.A. Paschos, Phys. Lett. B **241** (1990) 96.
- [6] J.F. Gunion, Douglas W. McKay, H. Pois, Phys. Lett. B **334** (1994) 339; Phys. Rev. D **51** (1995) 201; H.-J. He, N. Polonsky and S. Su, Phys.Rev. **D64** (2001) 053004.
- [7] Particle Data Group, C. Caso *et al.*, Eur. Phys. J. **C3**, 1 (1998).

Figure 1: Feynmann diagram for a_μ induced by μ' .






Figure 2: Diagram of $f(x)$ to $m_{\mu'}$




Figure 3: Diagram of $V_{2\nu'}$ to $m_{\nu'}$.