

# Prospects for detection of $\Upsilon(1D) \rightarrow \Upsilon(1S)\pi\pi$ via $\Upsilon(3S) \rightarrow \Upsilon(1D) + X$ <sup>1</sup>

Jonathan L. Rosner<sup>2</sup>

*Enrico Fermi Institute and Department of Physics  
University of Chicago, 5640 S. Ellis Avenue, Chicago, IL 60637*

At least one state in the first family of D-wave  $b\bar{b}$  quarkonium levels has been discovered near the predicted mass of 10.16 GeV/ $c^2$ . This state is probably the one with  $J = 2$ . This state and the ones with  $J = 1$  and  $J = 3$  may contribute a detectable amount to the decay  $\Upsilon(1D) \rightarrow \Upsilon(1S)\pi\pi$ , depending on the partial widths for these decays for which predictions vary considerably. The prospects for detection of the chain  $\Upsilon(3S) \rightarrow \Upsilon(1D) + X \rightarrow \Upsilon\pi\pi + X$  are discussed.

PACS Categories: 14.40.Gx, 13.20.Gd, 13.40.Hq, 12.39.Ki

A sample of  $4.73 \times 10^6$   $\Upsilon(3S)$  decays obtained by the CLEO III experiment has permitted the first observation of an  $\Upsilon(1D)$  level [1]. This state, which probably is the  $J = 2$  member of the spin-triplet  $1^3D_{1,2,3}$ , has a mass of  $10162.2 \pm 1.6$  MeV/ $c^2$  and is observed with a combined set of branching ratios for production and decay very close to theoretical predictions [2, 3]. The data exceed the CLEO Collaboration's previously reported sample [4],  $237\,000 \pm 6800$   $\Upsilon(3S)$  decays, by about a factor of 20. They are several times that of the CUSB Collaboration [5],  $1.18 \times 10^6$   $\Upsilon(3S)$  decays with  $\mu^+\mu^-$  detection and  $0.64 \times 10^6$  decays with  $e^+e^-$  detection.

The chain through which the  $\Upsilon(1D)$  state was discovered is illustrated in Fig. 1(a). A cascade involving four photons  $\gamma_1\text{--}\gamma_4$  followed by leptonic decay of the  $\Upsilon(1S)$  leads to a very clean final state in which the major limitations are the total size of the sample and backgrounds from cascades via the photons labeled  $\gamma_2'$  and  $\gamma_3'$  involving the intermediate  $\Upsilon(2S)$  state. The combined branching ratio including the decay  $\Upsilon(1S) \rightarrow (e^+e^- \text{ or } \mu^+\mu^-)$ ,  $(3.3 \pm 0.6 \pm 0.5) \times 10^{-5}$ , is consistent with the prediction [3] of  $3.76 \times 10^{-5}$ .

Theoretical predictions [3] indicate that for the CLEO III  $\Upsilon(3S)$  sample mentioned above, one should have produced a total of (5.7, 14.2, 7.7) thousand  $\Upsilon(1D)$  states with  $J = (1, 2, 3)$ , respectively, via the transitions involving  $\gamma_1$  and  $\gamma_2$  in Fig. 1(a). Yan [6] and Tuan [7] have pointed out that another way to detect these states may be through their decays to  $\Upsilon(1S)\pi\pi$ , as illustrated in Fig. 1(b). The present note examines the likelihood of observing these transitions in the CLEO III data set. It is not necessary to detect the photons  $\gamma_1$  and  $\gamma_2$ , though vetoing charged particles aside from the pions in  $\Upsilon(1D) \rightarrow \Upsilon(1S)\pi\pi$  might be helpful.

The predictions for  $\Upsilon(1D) \rightarrow \Upsilon(1S)\pi\pi$  rates differ substantially from one another. All are independent of the spin of the  $1D$  state [6, 8], aside from possible small effects

<sup>1</sup>Enrico Fermi Institute preprint EFI 03-06, hep-ph/0302122. To be submitted to Physical Review D, Brief Reports.

<sup>2</sup>rosner@hep.uchicago.edu




Figure 1: Cascades from the  $\Upsilon(3S)$  leading to production and detection of the  $\Upsilon(1D)$  levels. In each case the  $\Upsilon(1S)$  is detected via its decay to  $e^+e^-$  or  $\mu^+\mu^-$ . (a) Four-photon cascades; (b) sequence  $\Upsilon(3S) \rightarrow \gamma_1\gamma_2\Upsilon(1D)$  followed by  $\Upsilon(1D) \rightarrow \Upsilon(1S)\pi\pi$ .

on the  $J = 1$  state due to  $^3S_1$ - $^3D_1$  mixing [9]. Billoire *et al.* [10] found  $\Gamma(\Upsilon(1D) \rightarrow \Upsilon(1S)\pi\pi) = 0.07\alpha_s^2$  keV; Kuang and Yan [11] obtained 24 KeV; Moxhay [8] found 0.07 keV (a value employed in Refs. [2] and [3]), and Ko [12] argued on the basis of non-perturbative effects that the Billoire *et al.* and Moxhay results should be enhanced, leading to a prediction of  $0.56 \pm 0.07$  keV. I shall compare the sensitivity of the present CLEO III sample to the last three predictions. Discussions of the reasons for the differences may be found in Refs. [2, 12, 13, 14].

In brief, the present data sample should be able to rule out the prediction of Kuang and Yan or observe the predicted transition with great statistical significance. The sensitivity at the level of Ko's prediction will depend on CLEO III's superior performance as a detector in comparison with the previous CLEO limit [4]. In that work a combined upper limit

$$\sum_{J=1}^3 \mathcal{B}[\Upsilon(3S) \rightarrow \Upsilon(1^3D_J)X] \mathcal{B}[\Upsilon(1^3D_J \rightarrow \Upsilon(1S)\pi\pi] < 0.6\% \text{ (90\% c.l.)} \quad (1)$$

was placed.

I first justify the estimate of the total number of  $\Upsilon(1D)$  states produced from the  $\Upsilon(3S)$  through two-photon cascades via the  $\chi_b'(2P)$  states. The branching ratios for the transitions  $\Upsilon(3S) \rightarrow \chi_b'(2P)\gamma_1$  are quoted in Ref. [15] (central values are used here), while for the decays  $\chi_b'(2P) \rightarrow \Upsilon(1D)\gamma_2$  the branching ratios predicted in Table VIII of Ref. [2] are employed. The combined results are summarized in Table I. It is assumed that these electromagnetic transitions are the only source of the  $\Upsilon(1D)$  levels. The energies of the photons  $\gamma_1$  are shown in the Table, while the energies of  $\gamma_2$  range between about 80 and 120 MeV for the masses favored in Ref. [2], shown in parentheses in MeV/ $c^2$  under the corresponding spins of the  $1^3D_J$  levels.

Table I: Branching ratios for electromagnetic transitions contributing to  $\Upsilon(3S) \rightarrow \Upsilon(1^3D_J)X$ . Branching ratios for emission of  $\gamma_1$  are observed values, taken from Ref. [15]. Table entries in Roman type denote  $\mathcal{B}(\chi_{bJ'} \rightarrow \Upsilon(1^3D_J)\gamma_2)$  predictions in percent [2]; numbers in italics denote thousands of  $\Upsilon(1^3D_J)$  states produced for  $4.73 \times 10^6$   $\Upsilon(3S)$  decays. Blank entries denote transitions forbidden by electric dipole selection rules.

| $J(\chi_{b'})$              | 0             | 1              | 2              | Total             |
|-----------------------------|---------------|----------------|----------------|-------------------|
| $\mathcal{B}(\gamma_1)(\%)$ | $5.4 \pm 0.6$ | $11.3 \pm 0.6$ | $11.4 \pm 0.8$ | events            |
| $E_{\gamma_1}$ (MeV)        | 123           | 100            | 87             | ( $\times 10^3$ ) |
| $^3D_1$                     | 0.153         | 0.97           | 0.023          |                   |
| (10156)                     | <i>0.391</i>  | <i>5.185</i>   | <i>0.124</i>   | <i>5.70</i>       |
| $^3D_2$                     |               | 2.36           | 0.29           |                   |
| (10162)                     |               | <i>12.61</i>   | <i>1.564</i>   | <i>14.18</i>      |
| $^3D_3$                     |               |                | 1.42           |                   |
| (10166)                     |               |                | <i>7.657</i>   | <i>7.66</i>       |

To calculate the branching ratios of each of the  $1^3D_J$  states into  $\Upsilon(1S)\pi\pi$  for the various models of rates, one may use the predictions [2] of decay rates of each state into other modes,  $\Gamma_{\text{other}}(1^3D_{1,2,3}) = (35.5, 28.0, 25.4)$  keV. The results are summarized in Table II, taking Ko's central value of  $\Gamma(1^3D_J \rightarrow \Upsilon(1S)\pi\pi) = 0.56$  keV.

The predicted products of branching ratios corresponding to the left-hand side of Eq. (1) are all below the old CLEO upper bound [4]. However, with 20 times the old data, CLEO III should be able to check the Kuang–Yan prediction. (Our value of 0.27% is to be compared with the estimate made by Tuan [7] of 0.43%). The prediction of Moxhay appears to be too low to be testable. Ko [12] has argued that nonperturbative effects raise Moxhay's prediction by a factor of  $8 \pm 1$ , leading to the prediction

$$\sum_{J=1}^3 \mathcal{B}[\Upsilon(3S) \rightarrow \Upsilon(1^3D_J)X] \mathcal{B}[\Upsilon(1^3D_J \rightarrow \Upsilon(1S)\pi\pi)] = 0.011\% \quad . \quad (2)$$

(Ko obtained a slightly smaller value of 0.009%; Eq. (2) takes account of updated estimates [3] of  $\Upsilon(1^3D_J)$  production.) Although this is a factor of about 50 below the upper bound in Eq. (1), the total number of predicted events looks encouraging. Multiplying 532  $\Upsilon(1S)\pi\pi$  decays by 2/3 (for charged pions) and by  $\mathcal{B}[\Upsilon(1S) \rightarrow (e^+e^- \text{ or } \mu^+\mu^-)] = 4.86 \pm 0.13\%$ , we expect about 17 events with two charged pions and two charged leptons ( $e$  or  $\mu$  pairs). Of these, about half should be due to the  $1^3D_2$  state which is the best candidate for that reported in Ref. [1], with about 2/3 of the remaining events due to the  $1^3D_3$  state. Some additional sensitivity might be gained using neutral pions.

I am grateful to Stephen Godfrey, Yu-Ping Kuang, San Fu Tuan, and Eckhard von Toerne for discussions. This work was supported in part by the United States Department of Energy through Grant No. DE FG02 90ER40560.

Table II: Predicted branching ratios for  $\Upsilon(1^3D_J) \rightarrow \Upsilon(1S)\pi\pi$  (Roman entries, in percent) and numbers of  $\Upsilon(1S)\pi\pi$  final states produced for  $4.73 \times 10^6$   $\Upsilon(3S)$  decays (italics).

| State                        | $N(^3D_J)$<br>( $\times 10^3$ ) | Kuang-Yan<br>[11] | Moxhay<br>[8] | Ko<br>[12] |
|------------------------------|---------------------------------|-------------------|---------------|------------|
| $1^3D_1$                     |                                 | 40.4              | 0.20          | 1.55       |
|                              | 5.70                            | <i>2303</i>       | <i>11</i>     | <i>88</i>  |
| $1^3D_2$                     |                                 | 46.2              | 0.25          | 1.96       |
|                              | 14.18                           | <i>6551</i>       | <i>35</i>     | <i>278</i> |
| $1^3D_3$                     |                                 | 48.6              | 0.27          | 2.16       |
|                              | 7.66                            | <i>3721</i>       | <i>21</i>     | <i>165</i> |
| Total                        | 27.54                           | <i>12575</i>      | <i>67</i>     | <i>532</i> |
| $\mathcal{B}$ [Eq. (1) (%) ] |                                 | 0.27              | 0.0014        | 0.011      |

## References

- [1] CLEO Collaboration, S. E. Csorna *et al.*, presented by T. Skwarnicki at 31st International Conference on High Energy Physics (ICHEP 2002), Amsterdam, The Netherlands, 24–31 Jul 2002, CLEO Report No. CLEO-CONF-02-06, hep-ex/0207060.
- [2] W. Kwong and J. L. Rosner, Phys. Rev. D **38**, 279 (1988).
- [3] S. Godfrey and J. L. Rosner, Phys. Rev. D **64**, 097501 (2001); **66**, 059902(E) (2002).
- [4] CLEO Collaboration, I. C. Brock *et al.*, Phys. Rev. D **43**, 1448 (1991).
- [5] CUSB Collaboration, U. Heintz *et al.*, Phys. Rev. D **46**, 1928 (1992).
- [6] T.-M. Yan, Phys. Rev. D **22**, 1652 (1980).
- [7] S. F. Tuan, Commun. Theor. Phys. **26**, 381 (1996).
- [8] P. Moxhay, Phys. Rev. D **37**, 2557 (1988).
- [9] We thank Y.-P. Kuang for a discussion of this point.
- [10] A. Billoire, R. Lacaze, A. Morel, and H. Navelet, Nucl. Phys. **B155**, 493 (1979).
- [11] Y.-P. Kuang and T.-M. Yan, Phys. Rev. D **24**, 2874 (1981).
- [12] P. Ko, Phys. Rev. D **47**, 208 (1993).
- [13] M. Voloshin, Yad. Fiz. **43**, 1571 (1986) [Sov. J. Nucl. Phys. **43**, 1011 (1986)].
- [14] Y.-P. Kuang and T.-M. Yan, Phys. Rev. D **41**, 155 (1990).
- [15] Particle Data Group, K. Hagiwara *et al.*, Phys. Rev. D **66**, 010001 (2002).