

CALT-68-2407, DCPT/03/02
 IPPP/03/01, LMU 18/02
 TUM-HEP-481/02
 hep-ph/0302174

The lightest Higgs Boson of mSUGRA, mGMSB and mAMSB at Present and Future Colliders: Observability and Precision Analyses

A. Dedes¹, S. Heinemeyer^{2y}, S. Su^{3z} and G. Weiglein^{4x}

¹ Physik Department, Technische Universität München, D-85748 Garching, Germany

² Institut für theoretische Elementarteilchenphysik, LMU München, Theresienstr. 37,
 D-80333 München, Germany

³ California Institute of Technology, Pasadena, CA 91125, USA

⁴ Institute for Particle Physics Phenomenology, University of Durham,
 Durham DH1 3LE, UK

Abstract

We investigate the physics of the lightest CP-even MSSM Higgs boson at the Tevatron, the LHC, a linear e^+e^- collider, a $\mu^+\mu^-$ collider and a $\gamma\gamma$ collider. The analysis is performed in the three most prominent soft SUSY-breaking scenarios, mSUGRA, mGMSB and mAMSB. For all colliders the observability and parameter regions with suppressed production cross sections (compared to a SM Higgs boson with the same mass) are investigated. For the lepton and photon colliders the potential is analyzed of precision measurements of the branching ratios of the light CP-even Higgs boson for obtaining indirect bounds on the mass of the CP-odd Higgs boson and the high-energy parameters of the soft SUSY-breaking scenarios. In regions of the parameter space where the LHC can detect the heavy Higgs bosons, precision measurements of the properties of the light Higgs boson at the linear collider can provide valuable information for distinguishing between the mSUGRA, mGMSB and mAMSB scenarios.

email: dedes@ph.tum.de

^yemail: Sven.Heinemeyer@physik.uni-muenchen.de

^zemail: shufang@theory.caltech.edu

^xemail: Georg.Weiglein@durham.ac.uk

1 Introduction

The search for the light neutral Higgs boson is a crucial test of Supersymmetry (SUSY) that can be performed with the present and the next generation of high-energy colliders. The prediction of a relatively light Higgs boson is common to all supersymmetric models whose couplings remain in the perturbative regime up to a very high energy scale [1]. Finding the Higgs boson and subsequently studying its couplings to fermions and bosons is thus one of the main goals of high-energy physics. The data taken during the final year of LEP running at $\sqrt{s} > 206 \text{ GeV}$ have established a 95% C.L. exclusion limit for the Standard Model (SM) Higgs boson of $M_{H_{SM}} > 114.4 \text{ GeV}$. They showed a slight excess at about the 2 level of signal-like events over the background expectation, which would be compatible with the expectation for the production of a Higgs boson with SM-like ZZH coupling with a mass $M_{H_{SM}} = 116 \pm 1 \text{ GeV}$ [2]. In the Minimal Supersymmetric Standard Model (MSSM) the mass of the lightest CP-even Higgs boson, m_h , is bounded from above by $m_h < 135 \text{ GeV}$ [3,4] (taking into account radiative corrections up to two-loop order [3,4,6,7,8,9,10,5,11,12,13,14]).

In the MSSM no specific assumptions are made about the underlying Supersymmetry-(SUSY)-breaking mechanism, and a parameterization of all possible SUSY-breaking terms is used. This gives rise to the huge number of more than 100 new parameters in addition to the SM, which in principle can be chosen independently of each other. A phenomenological analysis of this model in full generality would clearly be very involved, and one usually restricts to certain benchmark scenarios, see e.g. Refs. [15,16,17]. On the other hand, models in which all the low-energy parameters are determined in terms of a few parameters at the Grand Unification scale (or another high-energy scale), employing a specific soft SUSY-breaking scenario, are much more predictive. The most prominent scenarios in the literature are minimal Supergravity (mSUGRA) [18,19], minimal Gauge Mediated SUSY Breaking (mGMSB) [20] and minimal Anomaly Mediated SUSY Breaking (mAMSB) [21,22,23]. Analyses of the Higgs sector in these scenarios, mostly focusing only on the maximum value of m_h , have been performed in Refs. [24,25,26,27,28,29,30,31]. A detailed comparison of the three soft SUSY-breaking scenarios in terms of exclusion regions in the $M_A - \tan\beta$ plane (where A is the CP-odd Higgs boson and $\tan\beta$ the ratio of the vacuum expectation values of the two Higgs doublets), their compatibility with the slight excess observed at LEP, and their corresponding SUSY particle spectra can be found in Ref. [32].

In the present paper the work of Ref. [32] is extended to an analysis of the lightest CP-even Higgs boson phenomenology at present and future colliders. We relate the input from the three soft SUSY-breaking scenarios in a uniform way to the predictions for the low-energy phenomenology in the Higgs sector, allowing thus a direct comparison of the predictions arising from the different scenarios. The high-energy parameters given in the three scenarios are related to the low-energy SUSY parameters via renormalization group (RG) running, taking into account contributions up to two-loop order [33] (for a recent comparison of different codes and current accuracies, see Ref. [34]). After transforming the parameters obtained in this way into the corresponding on-shell parameters [35,36,37], they are used as input for the program FeynHiggs [38,39,40]. As a result the Higgs boson mass spectrum and the Higgs decay rates and branching ratios [41,42] have been obtained. Further restrictions such as from precision observables and the non-observation of SUSY particles at LEP and the Tevatron are also taken into account. For an analysis within the mSUGRA scenario where also the cold dark matter (CDM) constraints are included, see Ref. [43]. Based on

these predictions for the Higgs sector phenomenology, we analyze the observability of the lightest MSSM Higgs boson at the Tevatron, the LHC, a future e^+e^- linear collider (LC), a $\mu\mu$ collider (C), and a $\tau\tau$ collider (C). Regions of the high-energy parameter space with strongly suppressed Higgs production cross sections are identified. As the next step the branching ratios of the lightest Higgs boson into SM fermions ($h \rightarrow b\bar{b}; h \rightarrow c\bar{c}; h \rightarrow \tau^+\tau^-$) and into W bosons ($h \rightarrow W^+W^-$) are compared for the mSUGRA, mGMSB and mAMSB case. We show that the precise measurement of the various Higgs decay branching ratios can be used to impose bounds on the value of M_A and also the high-energy input parameters. Combined with the information on M_A that could be obtained from the LHC Higgs searches, we further discuss the possibility of distinguishing the three scenarios via the Higgs branching ratio measurements at the LC.

The rest of the paper is organized as follows. In Sect. 2 we briefly review the three soft SUSY-breaking scenarios and the evaluation of the low-energy data. The observability of the lightest MSSM Higgs boson is investigated in Sect. 3. In Sect. 4 the potential is analyzed of precision measurements of the Higgs boson branching ratios for obtaining indirect constraints on M_A and the high-energy parameters of the soft SUSY-breaking scenarios. The possibility of a distinction of the mSUGRA, mGMSB and mAMSB scenarios is discussed. The conclusions can be found in Sect. 5.

2 The low-energy sector and phenomenological constraints

In deriving the low-energy parameters in the three soft SUSY-breaking scenarios (mSUGRA, mGMSB and mAMSB) from the high-energy input parameters we follow Ref. [32]. Thus, in this section only the most relevant facts are briefly summarized.

2.1 The soft SUSY-breaking scenarios

The fact that no SUSY partners of the SM particles have so far been observed means that low-energy SUSY cannot be realized as an unbroken symmetry in nature, and SUSY models thus have to incorporate additional Supersymmetry breaking interactions. This is achieved by adding to the Lagrangian (defined by the $SU(3)_C \times SU(2)_L \times U(1)_Y$ gauge symmetry and the superpotential W) some further interaction terms that respect the gauge symmetry but break Supersymmetry (softly, i.e. no quadratic divergences appear), so called "soft SUSY-breaking" (SSB) terms. Assuming that the R-parity symmetry [44] is conserved, which we do in this paper for all SUSY breaking scenarios, reduces the amount of new soft terms allowed in the Lagrangian. Choosing a particular soft SUSY-breaking pattern allows further reduction of the number of free parameters and the construction of predictive models. The three most prominent scenarios for such models are

mSUGRA (minimal Supergravity scenario) [8, 19]:

Apart from the SM parameters (for the experimental values of the SM input parameters we use Ref. [45]), 4 parameters and a sign are required to define the mSUGRA scenario:

$$\{m_0; m_{1/2}; A_0; \tan\beta; \text{sign}(\mu)\} \quad (1)$$

While m_0 , $m_{1=2}$ and A_0 define the scalar and fermionic masses and the trilinear couplings at the GUT scale ($\sim 10^6$ GeV), $\tan \beta$ (the ratio of the two vacuum expectation values) and the $\text{sign}(\mu)$ (μ is the supersymmetric Higgs mass parameter) are defined at the low-energy scale. For our numerical analysis, see Sects. 3, 4, we have scanned over the following parameter space¹:

$$\begin{aligned} 50 \text{ GeV} & \leq m_0 \leq 1 \text{ TeV} ; \\ 50 \text{ GeV} & \leq m_{1=2} \leq 1 \text{ TeV} ; \\ 3 \text{ TeV} & \leq A_0 \leq 3 \text{ TeV} ; \\ 1.5 & \leq \tan \beta \leq 60 ; \\ \text{sign}(\mu) & = +1: \end{aligned} \quad (2)$$

The low-energy spectrum has been evaluated with the programs `SUSIT Y`/`FeynSSG` [48, 49].

m G M S B (minimal Gauge Mediated SUSY-Breaking) [20]:

A very promising alternative to m SUGRA is based on the hypothesis that the soft SUSY-breaking occurs at relatively low energy scales and is mediated mainly by gauge interactions through the so-called "messenger sector" [51, 50, 20]. Also in this scenario, the low-energy parameters depend on 4 parameters and a sign,

$$\{M_{\text{mess}}; N_{\text{mess}}; \mu; \tan \beta; \text{sign}(\mu)\} ; \quad (3)$$

where M_{mess} is the overall messenger mass scale; N_{mess} is a number called the messenger index, parameterizing the structure of the messenger sector; μ is the universal soft SUSY-breaking mass scale felt by the low-energy sector. The phenomenology of m G M S B is characterized by the presence of a very light gravitino $\tilde{G}$ with mass given by $m_{3=2} = m_{\tilde{G}} = \frac{F}{3M_{\text{P}}^2}$, $\frac{F}{100 \text{ TeV}^2} \sim 2.37 \text{ eV}$ [52], where F is the fundamental scale of SSB and $M_{\text{P}}^2 = 2.44 \times 10^8 \text{ GeV}^2$ is the reduced Planck mass. Since F is typically of order 100 TeV, the $\tilde{G}$ is always the LSP in these theories. The numerical analysis in Sects. 3, 4 is based on the following scatter ranges:

$$\begin{aligned} 10^4 \text{ GeV} & \leq M_{\text{mess}} \leq 10^5 \text{ GeV} ; \\ 1.01 & \leq N_{\text{mess}} \leq 10 ; \\ 1 & \leq \tan \beta \leq 60 ; \\ \text{sign}(\mu) & = +1: \end{aligned} \quad (4)$$

The low-energy parameter sets for this scenario have been calculated by using the program `SUSY FIRE` [53] and adopting the phenomenological approach of Refs. [54, 55, 56, 29].

m A M S B (minimal Anomaly Mediated SUSY-Breaking) [21, 22, 23]:

In this model, SUSY breaking happens on a separate brane and is communicated to

¹The sign of μ has been fixed to (+) (for all three soft SUSY-breaking scenarios), since this sign is favored by the $g_{\mu\mu}^2$ [46] and the BR ($b \rightarrow s$) [47] constraints, see Sect. 2.3.

the visible world via the super-Weyl anomaly. The particle spectrum is determined by 3 parameters and a sign:

$$m_{\text{aux}}; m_0; \tan \beta; \text{sign}(\mu): \quad (5)$$

The overall scale of SUSY particle masses is set by m_{aux} , which is the VEV of the auxiliary field in the supergravity multiplet. m_0 is introduced as a phenomenological parameter to avoid negative slepton mass squares, for other approaches to this problem see Refs. [21,57,58,59,60]. The scatter parameter space for the numerical analysis in Sects. 3, 4 is chosen to be

$$\begin{aligned} 20 \text{ TeV} & \leq m_{\text{aux}} \leq 100 \text{ TeV}; \\ 0 & \leq m_0 \leq 2 \text{ TeV}; \\ 1.5 & \leq \tan \beta \leq 60; \\ \text{sign}(\mu) & = +1: \end{aligned} \quad (6)$$

The low-energy spectrum has been derived with the code described in Ref. [26].

2.2 Evaluation of predictions in the Higgs boson sector of the MSSM

The most relevant parameters for Higgs boson phenomenology in the MSSM are the mass of the CP-odd Higgs boson, M_A , the ratio of the two vacuum expectation values, $\tan \beta$, the scalar top masses and mixing angle, $m_{t_1}; m_{t_2}; \epsilon_t$, for large $\tan \beta$ also the scalar bottom masses and mixing angle, $m_{b_1}; m_{b_2}; \epsilon_b$, the supersymmetric Higgs mass parameter, μ , the U(1) and SU(2) gaugino masses, M_1 and M_2 , and the gluino mass, m_g . These low-energy parameters are derived from the high-energy parameters of the three soft SUSY-breaking scenarios via RG running, see Ref. [32]. Since the RG running employed in the three scenarios is based on the $\overline{\text{DR}}$ scheme, the corresponding low-energy parameters are $\overline{\text{DR}}$ parameters. In order to derive predictions for observables, i.e. particle masses and mixing angles, these parameters in general have to be converted into on-shell parameters.

For the predictions in the MSSM Higgs sector we use results obtained in the Feynman diagrammatic (FD) approach within the on-shell renormalization scheme as incorporated in the Fortran code FeynHiggs [38,39,40] based on Refs. [5,4,3].

Our analysis is concerned with the main Higgs production and decay channels at different colliders. To this end the predictions for the Higgs boson masses and effective couplings (especially the effective mixing angle, α_e , in the neutral CP-even Higgs boson sector that includes higher-order corrections) as well as for the branching ratios of the lightest MSSM Higgs boson (and for a SM Higgs boson with the same mass) have been evaluated. While the predictions for the decays of $h \rightarrow b\bar{b}; c\bar{c}; \tau^+\tau^-$ are based on Ref. [41], including the higher-order corrections described in Ref. [61,62], the other decay channels have been derived with the code Hdecay [42], which has been implemented as a subroutine in the latest version of FeynHiggs [40]. The proper transition from on-shell parameters in FeynHiggs to $\overline{\text{DR}}$ parameters in Hdecay has been taken into account [35,36,37].

In order to derive the relative difference between a MSSM production or decay rate and the corresponding SM rate (for the same Higgs boson mass), the following ratios have been

calculated (below the notation $tt \rightarrow tth$ refers to the processes $q\bar{q}; gg \rightarrow tth$, and $VV \rightarrow h$ refers to the vector boson fusion processes at the LHC and the LC, $q\bar{q} \rightarrow q^0 q^0 h, e^+ e^- \rightarrow h$, respectively):

$q\bar{q}; q\bar{q} \rightarrow V \rightarrow Vh$ ($V = W, Z$):

$$\frac{\frac{SU(SY)}{hV}}{\frac{SM}{hV}} \sin^2(\alpha_e) \quad (7)$$

$gg \rightarrow h$:

$$\frac{\frac{SU(SY)}{SM} (gg \rightarrow h)}{\frac{SM}{SM} (gg \rightarrow h)} = \frac{\frac{SU(SY)}{SM} (h \rightarrow gg)}{\frac{SM}{SM} (h \rightarrow gg)} \quad (8)$$

$q\bar{q}; gg \rightarrow t\bar{t} \rightarrow tth$:

$$\frac{\frac{SU(SY)}{SM} (t\bar{t} \rightarrow tth)}{\frac{SM}{SM} (t\bar{t} \rightarrow tth)} = \frac{\cos^2 \alpha_e}{\sin^2 \alpha_e} \quad (9)$$

$VV \rightarrow h$, ($V = W, Z$):

$$\frac{\frac{SU(SY)}{VVh}}{\frac{SM}{VVh}} \sin^2(\alpha_e) \quad (10)$$

$e^+ e^- \rightarrow Z \rightarrow Zh$:

$$\frac{\frac{SU(SY)}{hZ}}{\frac{SM}{hZ}} \sin^2(\alpha_e) \quad (11)$$

$\gamma \gamma \rightarrow h$:

$$\frac{\frac{SU(SY)}{SM} (\gamma \gamma \rightarrow h)}{\frac{SM}{SM} (\gamma \gamma \rightarrow h)} = \frac{\frac{SU(SY)}{SM} (h \rightarrow \gamma \gamma)}{\frac{SM}{SM} (h \rightarrow \gamma \gamma)} \quad (12)$$

$g g \rightarrow h$:

$$\frac{\frac{SU(SY)}{SM} (g g \rightarrow h)}{\frac{SM}{SM} (g g \rightarrow h)} = \frac{\frac{SU(SY)}{SM} (h \rightarrow g g)}{\frac{SM}{SM} (h \rightarrow g g)} \quad (13)$$

Higgs boson decays:

$$\begin{aligned} & \frac{BR^{SU(SY)}(h \rightarrow b\bar{b})}{BR^{SM}(h \rightarrow b\bar{b})}; \frac{BR^{SU(SY)}(h \rightarrow c\bar{c})}{BR^{SM}(h \rightarrow c\bar{c})}; \frac{BR^{SU(SY)}(h \rightarrow \tau^+ \tau^-)}{BR^{SM}(h \rightarrow \tau^+ \tau^-)}; \\ & \frac{BR^{SU(SY)}(h \rightarrow \gamma \gamma)}{BR^{SM}(h \rightarrow \gamma \gamma)}; \frac{BR^{SU(SY)}(h \rightarrow W^+ W^-)}{BR^{SM}(h \rightarrow W^+ W^-)}; \frac{BR^{SU(SY)}(h \rightarrow gg)}{BR^{SM}(h \rightarrow gg)} : \end{aligned} \quad (14)$$

For some of the cross sections in Eqs. (7) – (12) more complete results for the MSSM exist in the literature than those used in our analysis, see e.g. Refs. [63,64,65,66]. For the qualitative analysis below, however, the approximations used here should be sufficient, see e.g. the comparison in [63].

For the numerical analysis the above production and decay modes have been combined to the most relevant channels for each collider, i.e. we have calculated the product

$$\frac{\sigma^{\text{SUSY}}(\text{Higgs prod.})}{\sigma^{\text{SM}}(\text{Higgs prod.})} \frac{\text{BR}^{\text{SUSY}}(\text{Higgs decay})}{\text{BR}^{\text{SM}}(\text{Higgs decay})} : \quad (15)$$

The following channels have been considered:

Tevatron:

$$q\bar{q} \rightarrow V \rightarrow Vh \rightarrow Vbb; (V = W, Z) \quad (16)$$

LHC:

$$gg \rightarrow h \quad (17)$$

$$q\bar{q}; gg \rightarrow t\bar{t} \rightarrow t\bar{t}h \rightarrow t\bar{t}bb \quad (18)$$

$$VV \rightarrow h \rightarrow \gamma\gamma, W^+W^- \quad (19)$$

LC:

$$e^+e^- \rightarrow Z \rightarrow Zh \rightarrow Zbb; Zcc; Z\gamma\gamma; ZWW; Zgg \quad (20)$$

$$WW \rightarrow h \rightarrow bb; cc; \gamma\gamma; W^+W^-; gg \quad (21)$$

Due to our approximations for the production processes the relative results of $\frac{\sigma^{\text{SUSY}}}{\sigma^{\text{SM}}} \frac{\text{BR}^{\text{SUSY}}}{\text{BR}^{\text{SM}}}$ for the same final states in the two chains numerically agree.

C:

$$\gamma\gamma \rightarrow h \rightarrow bb; W^+W^-; \quad (22)$$

C:

$$\gamma\gamma \rightarrow h \rightarrow bb; \gamma\gamma; W^+W^- \quad (23)$$

2.3 Phenomenological constraints

While our main focus in this paper is on the physics in the Higgs sector, we also take into account some further (relatively mild) constraints when determining the allowed parameter space. These constraints are briefly summarized here. (A more detailed discussion can be found in Ref. [32].)

LEP Higgs bounds:

The results from the Higgs search at LEP have excluded a considerable part of the MSSM parameter space [67]. The results of the search for the MSSM Higgs bosons are usually interpreted in three different benchmark scenarios [15]. The 95% C.L. exclusion limit for the SM Higgs boson of $M_{H_{SM}} > 114.4 \text{ GeV}$ [2] applies also for the lightest CP-even Higgs boson of the MSSM except for the parameter region with small M_A and large $\tan\beta$. In the unconstrained MSSM this bound is reduced to $m_h > 91.0 \text{ GeV}$ [67] for $M_A < 150 \text{ GeV}$ and $\tan\beta > 8$ as a consequence of a reduced coupling of the Higgs to the Z boson. For the CP-odd Higgs boson a lower bound of $M_A > 91.9 \text{ GeV}$ has been obtained [67]. In order to correctly interpolate between the parameter regions where the SM lower bound² of $M_{H_{SM}} > 113 \text{ GeV}$ and the bound $m_h > 91 \text{ GeV}$ apply, we use the result for the Higgs-mass exclusion given with respect to the reduced ZZh coupling squared (i.e. $\sin^2(\alpha_e)$) [68]. We have compared the excluded region with the theoretical prediction obtained at the two-loop level for m_h and $\sin^2(\alpha_e)$ for each parameter set (using $m_t = 175 \text{ GeV}$).

Precision observables:

The electroweak precision observables are affected by the whole spectrum of SUSY particles. The main SUSY contributions enter via the β -parameter [69]. In our analysis we take into account the corrections arising from $\bar{t}b$ loops up to two-loop order [70]. A value of β outside the experimentally preferred region of $\beta^{SUSY} < 3 \cdot 10^3$ [45] indicates experimentally disfavored $\bar{t}$ and b masses. The evaluation of β^{SUSY} is implemented in FeynHiggs.

Experimental bounds on SUSY particle masses

In order to restrict the allowed parameter space in the three soft SUSY-breaking scenarios we employed the current experimental constraints on their low-energy mass spectrum [45]. The precise values of the bounds that we have applied can be found in Ref. [32].

Other restrictions

We just briefly list here the further restrictions that we have taken into account. For a detailed discussion see Ref. [32].

- { The top quark mass is fixed to $m_t = 175 \text{ GeV}$.
- { The GUT or high-energy scale parameters are taken to be real, no SUSY CP-violating phases are assumed.
- { In all models under consideration the R-parity symmetry [44] is taken to be conserved.

² Instead of the actual experimental lower bound, $M_{H_{SM}} > 114.4 \text{ GeV}$ [2], we use the value of 113 GeV in order to allow for some uncertainty in the theoretical evaluation of m_h from unknown higher-order corrections, which is currently estimated to be $\sim 3 \text{ GeV}$ [4].

- { Parameter sets that do not fulfil the condition³ of radiative electroweak symmetry breaking (REWSB) are discarded (already at the level of model generation).
- { Parameter sets that do not fulfil the constraints that there should be no charge or color breaking minima are discarded (already at the level of model generation).
- { Contrary to Ref. [32] we did not apply a "naturalness bound" on the stau and gluino mass, but just restricted the scanned parameter space as indicated in Sect. 2.1. This is in fact not an important restriction for the BR calculation, since very heavy SUSY particles tend to decouple from the observables we are considering here, i.e. the quantity in Eq. (15) approaches 1.
- { We demand that the lightest SUSY particle (LSP) is uncolored and uncharged. In the mGMSB scenario the LSP is always the gravitino, so this condition is automatically fulfilled. Within the mSUGRA and mAMSB scenario, the LSP is required to be the lightest neutralino. Parameter sets that result in a different LSP are excluded.
- { We do not apply any further cosmological constraints, i.e. we do not demand a relic density in the region favored by dark matter constraints [72].
- { Although we do not apply constraints from BR($b \rightarrow s$) [47] or $g_{\mu\mu}^2$ of the muon [46], we restrict ourselves to the case where the Higgsino mixing parameter is positive, $\mu > 0$. This choice is favored by the current data. The results with negative μ can differ significantly from the case we consider here and would require an additional analysis.

2.4 Bounds on m_h and $\tan\beta$

Scanning over the parameter space of mSUGRA, mGMSB and mAMSB as described in Sect. 2.1 and applying the constraints as described in Sect. 2.3 results in upper maximal values of m_h and lower bounds on $\tan\beta$ (for the general MSSM case, see Refs. [67, 73]). This analysis has been performed in Ref. [32]. However, due to the progress in the m_h evaluation, see Ref. [4] for a review, these bounds have changed as compared to our earlier analysis. Tab. 1 gives an update of the obtained m_h and $\tan\beta$ bounds.

	mSUGRA	mGMSB	mAMSB
$m_h^{\text{max}} [\text{GeV}]$	126.6	123.2	124.5
$\tan\beta^{\text{min}}$	2.9	3.1	3.8

Table 1: Upper bound on m_h , $m_h < m_h^{\text{max}}$, and lower bound on $\tan\beta$, $\tan\beta > \tan\beta^{\text{min}}$, in the three soft SUSY-breaking scenarios (for $m_t = 175 \text{ GeV}$ and taking into account the phenomenological constraints of Sect. 2.3; no theoretical uncertainties from unknown higher-order corrections are included in m_h^{max}).

³We use here the one-loop minimization conditions. Analytical two-loop expressions of $O(\alpha_s + \alpha_t^2)$ for the minimization conditions have been recently given in Ref. [71]. The full two-loop corrections can be derived numerically from the work of Ref. [14].

3 Observability of the lightest MSSM Higgs boson

In this section the observability of the lightest MSSM Higgs boson at the different colliders is analyzed. Especially at the hadron colliders a reduced BR for certain channels compared to the SM value could make it more difficult to establish a Higgs signal over the background.

Tevatron:

In all three soft SUSY-breaking scenarios the Tevatron search channels are not significantly suppressed compared to the SM rates. The channel $V \rightarrow Vh \rightarrow Vbb$ is suppressed by not more than 10% as compared to the SM value. Therefore the prospects at the Tevatron for the discovery of the lightest MSSM Higgs boson are as good as for the SM Higgs boson. A similar observation has already been made in Ref. [43] for the mSUGRA scenario, where also other phenomenological constraints have been taken into account.

LHC:

We start our discussion with the channel $gg \rightarrow h \rightarrow$ that is of particular importance for a light SM-like Higgs boson with $m_h < 130$ GeV [74]. As has been shown in Ref. [75,76,16], for certain values of the SUSY parameters this production channel can be heavily suppressed for a wide region of the M_A – $\tan\beta$ parameter space of the unconstrained MSSM (see also Ref. [77] for an analysis in the mSUGRA scenario). Since the event rate for this channel is relatively low ($h \rightarrow$ being a rare decay), a suppression of 50% or more would certainly pose a challenge to the experiment.

The situation within the three soft SUSY-breaking scenarios can be read off from Fig. 1⁴. Within the mSUGRA scenario a suppression of up to 10{20% is found for $M_A < 700$ GeV for all $\tan\beta$ values. For very large $\tan\beta$, $\tan\beta > 50$, a reduction even larger than 20% can be found. This is in agreement with the result obtained in Ref. [43], where no substantial reduction has been found, after other phenomenological constraints like $BR(b \rightarrow s)$, $g_{\mu\mu}^2$ and especially the CDM restrictions had been applied. Concerning the parameter space of the GUT scale parameters m_0 and $m_{1=2}$ (which is not shown here explicitly), the reduction is found for all m_0 and $m_{1=2} < 350$ GeV.

The suppression can be stronger in the other two scenarios. Nearly all model points with $M_A < 400$ GeV for mGMSB and mAMSB, which correspond to $\tan\beta$ values of $\tan\beta > 20$, show a suppression of 20–50% (or even more). For $M_A < 200$ GeV (300 GeV) and $\tan\beta > 50$ ($30 < \tan\beta < 40$) the reduction is even larger than 50% in mGMSB (mAMSB). For larger M_A values, $M_A > 600$ GeV, the SM value of BR is approached. Concerning the high-energy parameters, the largest reduction in the mGMSB scenario is found all over the parameter space for M_{mess} and N_{mess} (with $\tan\beta > 50$ and > 30 TeV), whereas a reduction by 10–20% is mostly found for the lowest values for all M_{mess} values. Within mAMSB the largest reduction is found for $m_0 < 700$ GeV and $m_{\text{aux}} < 4 \cdot 10^4$ GeV. Small values of the high-energy parameters correspond to relatively small values of the low-energy SUSY parameters, which are required for a sizable suppression of the $gg \rightarrow h \rightarrow$ channel [75,76,16].

⁴For a similar analysis for the charged Higgs bosons, see Ref. [78].




Figure 1: The rate for the LHC channel $gg \rightarrow h \rightarrow \gamma\gamma$, normalized to the SM case with the same Higgs boson mass, $(\text{BR})^{\text{SUSY}} = (\text{BR})^{\text{SM}}$, is shown in the M_A ($\tan\beta$) plane for the mSUGRA, mGMSB and mAMSB scenarios (from top to bottom). The unconnected dots appearing in this and the following figures reflect the fact that only a few points with very low density appear in this area. However, increasing the density of the scatter data would cover the whole area in which the dots are located.

We now consider the associated production channel at the LHC, $q\bar{q}; g\bar{g} \rightarrow t\bar{t} \rightarrow t\bar{t}h \rightarrow t\bar{t}b\bar{b}$.⁵ In all three scenarios the rate is not suppressed compared to the SM rate by more than 10% or even an enhancement by up to 10–20% occurs. Therefore this channel, which is most relevant in the region $100 \text{ GeV} < m_h < 120 \text{ GeV}$ [74] does not suffer from suppression due to SUSY corrections in the three soft SUSY-breaking scenarios.

More recently also the Higgs boson production via W boson fusion, $W^+W^- \rightarrow h$, with a subsequent decay to $\tau^+\tau^-$ pairs, W bosons or photons has been discussed, see Refs. [79,80] for a SM analyses and Refs. [81,82] for the mode $h \rightarrow \tau^+\tau^-$ in the MSSM case. These channels can be relevant for the whole allowed m_h range in the MSSM. The mSUGRA scenario offers very good prospects for the decay into $\tau^+\tau^-$, which is characterized by the coupling $\sin \theta_e = \cos \theta$. Over almost the whole parameter space the rate for $\tau^+\tau^-$ BR differs by less than 10% from the SM rate. Only for very small M_A and very high $\tan \beta$ a suppression is possible. This corresponds to the parameter space where the heavy Higgs boson can have SM-like couplings, see Fig. 1 in Ref. [32]. The other channels ($h \rightarrow W^+W^-$; $\gamma\gamma$) show the following pattern: due to the increased decay rates to fermions, at relatively small M_A , $250 \text{ GeV} < M_A < 500 \text{ GeV}$, a suppression of the decays into W bosons or photons is possible, while the value of the SM rate is approached for larger M_A .

The situation is similar in the mGMSB and mAMSB scenarios, besides that the suppression of the $\tau^+\tau^-$ channel for very small M_A and very large $\tan \beta$ is not found. For the smallest possible values of M_A and the largest possible values of $\tan \beta$ a strong enhancement of these channels can be observed. This agrees with the results of Ref. [32], where only in the mSUGRA scenario but not in the mGMSB and mAMSB scenario a region of the parameter space has been found where the heavy Higgs boson is SM-like. Thus the $h \rightarrow \tau^+\tau^-$ channel is enhanced everywhere in mGMSB and mAMSB. Correspondingly the two other channels, $h \rightarrow W^+W^-$ and $h \rightarrow \gamma\gamma$ are suppressed everywhere. As an example, in Fig. 2 we show the channel $W^+W^- \rightarrow h \rightarrow W^+W^-$ in the mGMSB and the mAMSB scenario. For not too large M_A , $M_A < 550 \text{ GeV}$ (700 GeV), in mGMSB (mAMSB) a reduction larger than 10% can be observed. For larger M_A the results in the mGMSB and mAMSB scenarios approach the one in the SM.

LC :

Due to its clean experimental environment, Higgs boson production should be easily observable at the LC, i.e. in various channels even a significant suppression compared to the SM rate would not be harmful [84,85,86]. Therefore the production channels $Z \rightarrow Zh$ and $WW \rightarrow h$ (which yield the same numerical result in our analysis, see Sect. 2.2) and all decay channels, $h \rightarrow b\bar{b}, c\bar{c}, \tau^+\tau^-, W^+W^-, \gamma\gamma$, are observable in the three soft SUSY-breaking scenarios. This applies also in the region of small M_A , $M_A < 200 \text{ GeV}$, where a suppression of more than 50% can occur in all three scenarios. We will therefore present a detailed analysis of the LC production and decay channels only in the context of precision measurements, see Sect. 4.

⁵Also some other channels for a SM Higgs boson have been studied, e.g. $t\bar{t} \rightarrow t\bar{t}h \rightarrow t\bar{t}W^+W^-$ [83]. They could be easily included in our analysis as outlined in Sect. 2. Since including these channels does not change our qualitative results, we do not discuss them here explicitly.




Figure 2: The rate for the LHC channel $WW \rightarrow h \rightarrow WW^*$, normalized to the SM case with the same Higgs boson mass, is shown in the M_A - $\tan\beta$ plane for the mGMSB (upper plot) and the mAMSB scenario (lower plot).

C :

Also the C, due to the Higgs boson production in the s-channel, offers very good prospects for the Higgs boson observation [87,88]. Only the decay $h \rightarrow \gamma\gamma$ could become problematic if it is strongly suppressed compared to the SM value (e.g. due to an enhanced hbb coupling).

In the mSUGRA scenario the $h \rightarrow \gamma\gamma$ channel and the $h \rightarrow WW^*$ channel are very similar. They can be suppressed by more than 20% only for very large $\tan\beta$, $\tan\beta > 50$, or for $\tan\beta \approx 10$ with the smallest allowed M_A values. On the other hand, the $h \rightarrow b\bar{b}$ and $h \rightarrow \tau^+\tau^-$ modes are unproblematic within mSUGRA and show either only a very small suppression or even some enhancement, with the only exception of a possible suppression at very large $\tan\beta$.

In the mGMSB scenario the $h \rightarrow b\bar{b}$ and $h \rightarrow \tau^+\tau^-$ channels are always enhanced. The $h \rightarrow \gamma\gamma$ and $h \rightarrow WW^*$ channels show a suppression of more than 50% for




Figure 3: The rate for the C channel $h \rightarrow \gamma\gamma$, normalized to the SM case with the same Higgs boson mass, is shown in the M_A - $\tan\beta$ plane for the mGMSB (upper plot) and the mAMSB scenario (lower plot).

$M_A < 100$ GeV, while for $M_A < 600$ GeV still a suppression of more than 10% occurs, see Fig. 3 for the $h \rightarrow \gamma\gamma$ channel. The situation is quite similar in the mAMSB scenario, with an exception in the intermediate M_A region, $600-1300$ GeV, where there is an enhancement of the $h \rightarrow \gamma\gamma$ mode compared to the one obtained in the mGMSB scenario, see Fig. 3.

C :

Finally, to complete our analysis, we also briefly look at the C . This collider offers good prospects since the Higgs boson can be produced in the s -channel without a loop suppression like at the C , however with the relatively small $t \rightarrow h$ Yukawa coupling [89]. However, in the unconstrained MSSM it is possible that the $t \rightarrow h$ coupling, being $\sin\theta_e = \cos\theta$, can become very small if $\theta_e \neq 0$ because of loop corrections [89, 90]. In this parameter region, on the other hand, H, A production at the C happens with an enhanced rate and offers good prospects for resolving H

and A as separate resonances [89,90].

The feature of a suppressed $h \rightarrow \gamma\gamma$ coupling can also be realized in the mSUGRA scenario when the heavy (and not the light) CP-even Higgs boson is SM-like. This is possible for very high $\tan\beta$ and small M_A , $M_A < 300$ GeV (note that here no CDM constraints are taken into account, in contrast to the analysis of Ref. [89]). In this parameter region a strong suppression is possible for all C channels of the light CP-even Higgs boson (while H, A production happens at enhanced rates). In the rest of the parameter space the $h \rightarrow b\bar{b}$ and $h \rightarrow \tau\tau$ channel are strongly enhanced for $M_A < 700$ GeV. For very large M_A an enhancement of up to 10% occurs. Correspondingly, the $h \rightarrow W^+W^-$ channel is not enhanced, but still within 10% of the SM value.

Within mGMSB and mAMSB the suppression of the $h \rightarrow \gamma\gamma$ coupling is not present. Because of the coupling factor $\sin\alpha = \cos\beta$ the $h \rightarrow b\bar{b}$ and $h \rightarrow \tau\tau$ channels are strongly enhanced for small M_A , while the SM value is approached for large values of M_A . The $h \rightarrow W^+W^-$ channel, being enhanced with the $\sin\alpha = \cos\beta$ factor only at the production vertex, is less enhanced, but should be unproblematic for the whole parameter space.

The results of this section are summarized in Tab. 2. The modes $gg \rightarrow h \rightarrow \gamma\gamma, \tau\tau, t\bar{t}$ and $W^+W^- \rightarrow h \rightarrow W^+W^-; \gamma\gamma; \tau\tau$ allow the detection of the lightest MSSM Higgs boson in all three scenarios over the whole indicated parameter space. Possible exceptions occur for the $gg \rightarrow h \rightarrow \gamma\gamma$ channel for the very small M_A region in the mGMSB and mAMSB scenarios, where a strong suppression of more than 50% could happen. Because of the clean experimental environment at a LC, detection of the light Higgs is ensured for all the three scenarios. For a C , the region of very small M_A values in the mGMSB and mAMSB scenarios might be difficult for the $h \rightarrow W^+W^-$ and $h \rightarrow \tau\tau$ mode because of a strong suppression of more than 50%. At the C , the region $M_A < 150$ GeV and $\tan\beta > 50$ in the mSUGRA scenario exhibits a significant suppression of the production of the lightest CP-even MSSM Higgs boson. Besides the "difficult" regions mentioned above, the main search modes at the C and the C are not affected by strong suppression for all three scenarios, and the lightest MSSM Higgs boson in these scenarios will clearly be detectable at all possible future colliders.

4 Precision analyses of the Higgs masses and branching ratios

We now investigate the potential of Higgs branching ratio measurements at future colliders for testing the underlying SUSY model. We concentrate our analysis on the LC and the C , since the anticipated precisions of at the LHC will in general be much worse, while on the other hand branching ratio measurements of the light Higgs boson at the C are not expected to yield substantially better results than at the LC.

Over most of the parameter space of the scenarios discussed here one would of course also expect to observe direct production of SUSY particles at the next generation of colliders.

Channel	Collider	m_h	mSUGRA	mGMSB	mAMSB
$V^* \rightarrow Vh \rightarrow Vb\bar{b}$	Tev LC	$\lesssim 135$ $\lesssim 0.7\sqrt{s}$	$\pm 10\%$	$\pm 10\%$	$M_A \lesssim 300 : +50\%$
$V^* \rightarrow Vh \rightarrow Vc\bar{c}$	LC	$\lesssim 0.7\sqrt{s}$	$M_A \lesssim 500 : \pm 20\%$	$400 \lesssim M_A \lesssim 600 : -20\%$ $200 \lesssim M_A \lesssim 400 : -50\%$ $M_A \lesssim 200 : \lesssim -50\%$	$500 \lesssim M_A \lesssim 650 : -20\%$ $300 \lesssim M_A \lesssim 500 : -50\%$ $M_A \lesssim 300 : \lesssim -50\%$
$VV \rightarrow h \rightarrow WW^*$ $V^* \rightarrow Vh \rightarrow VWW^*$	LHC LC	$\gtrsim 110$ $\lesssim 0.7\sqrt{s}$	$M_A \lesssim 150 : \pm 20\%$, for $\tan\beta \lesssim 30$ or $\gtrsim 50$	like $Vc\bar{c}$	like $Vc\bar{c}$
$VV \rightarrow h \rightarrow \tau^+\tau^-$ $V^* \rightarrow Vh \rightarrow V\tau^+\tau^-$	LHC LC	$\lesssim 150$ $\lesssim 0.7\sqrt{s}$	$M_A \lesssim 150 : \pm 50\%$, for $\tan\beta \gtrsim 50$	$\tan\beta \gtrsim 50 : \gtrsim 50\%$	$M_A \lesssim 500 : +50\%$
$V^* \rightarrow Vh \rightarrow Vgg$	LC	$\lesssim 130$	$M_A \lesssim 200 : \pm 20\%$	like $Vc\bar{c}$	$550 \lesssim M_A \lesssim 650 : -20\%$ $250 \lesssim M_A \lesssim 550 : -50\%$ $M_A \lesssim 250 : \lesssim -50\%$
$gg \rightarrow h \rightarrow \gamma\gamma$	LHC	$\lesssim 150$	$M_A \lesssim 150 : \pm 20\%$ $\tan\beta \lesssim 30$ or $\gtrsim 50$	like $Vc\bar{c}$	$550 \lesssim M_A \lesssim 750 : -20\%$ $250 \lesssim M_A \lesssim 550 : -50\%$ $M_A \lesssim 250 : \lesssim -50\%$
$t\bar{t} \rightarrow t\bar{t}h \rightarrow t\bar{t}b\bar{b}$	LHC	$\lesssim 120$	SM like	SM like $+50\%$ for $\tan\beta \gtrsim 50$	$M_A \lesssim 300 : +50\%$
$\gamma\gamma \rightarrow h \rightarrow b\bar{b}$	γC	n.f.e.	$M_A \lesssim 150 : +50\%$	SM like	$M_A \lesssim 300 : +50\%$
$\gamma\gamma \rightarrow h \rightarrow WW^*$	γC	n.f.e.	$M_A \lesssim 200 :$ -20% up to $+50\%$	$400 \lesssim M_A \lesssim 600 : -20\%$ $200 \lesssim M_A \lesssim 400 : -50\%$ $M_A \lesssim 200 : \lesssim -50\%$	$500 \lesssim M_A \lesssim 650 : -20\%$ $250 \lesssim M_A \lesssim 500 : -50\%$ $M_A \lesssim 250 : \lesssim -50\%$
$\gamma\gamma \rightarrow h \rightarrow \tau^+\tau^-$	γC	n.f.e.	$\tan\beta \sim 50 : \pm 50\%$ $\tan\beta < 50 : +50\%$ for $M_A \lesssim 150$	$\tan\beta \gtrsim 50 : \gtrsim 50\%$	$M_A \lesssim 600 : +50\%$
$\gamma\gamma \rightarrow h \rightarrow \gamma\gamma$	γC	n.f.e.	$M_A \lesssim 200 :$ -20% up to $+50\%$	like $\gamma\gamma \rightarrow h \rightarrow WW^*$	$500 \lesssim M_A \lesssim 650 : -20\%$ $250 \lesssim M_A \lesssim 500 : -50\%$ $M_A \lesssim 250 : \lesssim -50\%$
$\mu^+\mu^- \rightarrow h \rightarrow b\bar{b}$	μC	n.f.e.	$M_A \lesssim 250 : +50\%$ $M_A \lesssim 150, \tan\beta > 50 : -50\%$	$200 \lesssim M_A \lesssim 600 : +50\%$ $M_A \lesssim 200 : \gtrsim 50\%$	$450 \lesssim M_A \lesssim 900 : +50\%$ $M_A \lesssim 450 : \gtrsim 50\%$
$\mu^+\mu^- \rightarrow h \rightarrow WW^*$	μC	n.f.e.	$\pm 10\%$	$\tan\beta \simeq 50 : +50\%$	$M_A \lesssim 600 : +50\%$
$\mu^+\mu^- \rightarrow h \rightarrow \tau^+\tau^-$	μC	n.f.e.	$M_A \lesssim 250 : +50\%$ $M_A \lesssim 150, \tan\beta > 50 : -50\%$	like $\mu^+\mu^- \rightarrow h \rightarrow b\bar{b}$	like $\mu^+\mu^- \rightarrow h \rightarrow b\bar{b}$

Table 2: Behavior of the production and decay modes of the lightest CP-even Higgs boson of the MSSM, see Eq. (15), for the most relevant channels at present and future colliders in three different SUSY breaking scenarios. When there is suppression or enhancement we indicate its maximum magnitude together with the parameters where this happens. If not stated explicitly, for the rest of the parameter space the mode behaves roughly like in the SM. The Higgs mass range where detection of a statistically significant signal is possible is also shown [91,92]. The phrase "n.f.e." stands for "not fully explored" and refers to channels where the studies so far have been performed for some fixed m_h values only.

However, we concentrate our analysis on information obtainable from the Higgs sector without assuming further knowledge of the SUSY spectrum. In a realistic situation one would of course confront the model under study with all available experimental information.

Tab. 3 lists the anticipated accuracies in different channels at the LC [84,93] and the C [87,88]. The values given in Tab. 3 correspond to a SM-like Higgs boson with a mass compatible with the allowed mass range of the lightest CP-even Higgs boson in the three soft SUSY-breaking scenarios according to the upper bounds given in Tab. 1. There is of course some variation in the accuracy with which the branching ratios can be measured over the allowed range of m_h . For simplicity, we assume a constant precision over the allowed mass range for each channel with a value referring to the middle of the allowed range. In parameter regions where the MSSM rate differs drastically from the SM rate the prospective precision will of course be different than in the SM. While in extreme cases like this it will be easy to infer properties of the SUSY model from Higgs sector measurements, we will focus in our analysis below on moderate deviations between the MSSM and the SM, for which the values given in Tab. 3 can be applied in good approximation. We will indicate the deviation between the MSSM and the SM in terms of the accuracies given in Tab. 3, i.e. a $\pm n\%$ deviation means that the calculated MSSM value of BR deviates from the corresponding SM value (with $M_{H_{SM}} = m_h$) by ($\pm n\%$ precision).

collider	decay mode	precision
LC	$h \rightarrow b\bar{b}$	1.5%
LC	$h \rightarrow \tau^+\tau^-$	4.5%
LC	$h \rightarrow c\bar{c}$	6%
LC	$h \rightarrow g\bar{g}$	4%
LC	$h \rightarrow W^+W^-$	3%
C	$h \rightarrow b\bar{b}$	2%
C	$h \rightarrow W^+W^-$	5%
C	$h \rightarrow \gamma\gamma$	11%

Table 3: Anticipated precisions from measurements of Higgs branching ratios at the LC [84,93] and the C [87,88]. The values are given for a SM-like Higgs boson with a mass compatible with the allowed mass range of the lightest CP-even Higgs boson in the three soft SUSY-breaking scenarios, see text.

4.1 Sensitivity to M_A and $\tan\beta$

While within the MSSM the prospects for the detection of the lightest CP-even Higgs boson at the next generation of colliders are very good, the situation is quite different for direct observation of the CP-odd A boson. At the LHC the detection of this particle can be very

difficult over sizable fractions of the MSSM parameter space (see e.g. Ref. [74,94]), while it may be outside the kinematical reach of the LC (see Refs. [64,95] for a recent account of this subject). Thus, it is of interest to study the potential for obtaining indirect bounds on M_A from precision measurements. Exploiting the sensitivity to M_A can be done in a similar fashion as nowadays for the SM Higgs, where indirect bounds are derived from electroweak precision tests. Since in the decoupling limit, $M_A \rightarrow M_Z$, the Higgs sector of the MSSM becomes SM-like, deviations in the production and decay of the lightest CP-even Higgs boson of the MSSM can in principle be translated into an upper bound on M_A . If direct information on M_A is available, the indirect sensitivity to M_A allows a stringent test of the model.

Several analyses of the sensitivity to M_A at the LC or the C have been carried out in the literature [43,88,96,97,98] (for an analysis focusing on the measurements with a GigaZ option of the LC see Ref. [99]). While in many of these analyses particular "benchmark" values of the SUSY parameters have been chosen, we perform a detailed scan over the parameter space of the three soft SUSY-breaking scenarios. Our study differs considerably especially from previous studies of the Higgs branching ratios in Ref. [88,96,97]. In these previous analyses, all parameters except for the one under investigation have been kept fixed and the effect of an assumed deviation between the MSSM and the SM has solely been attributed to this single free parameter. This would correspond to a situation with a complete knowledge of all SUSY parameters without any experimental or theoretical uncertainty, which obviously leads to an unrealistic enhancement of the sensitivity to the investigated parameter. Allowing the other SUSY (and SM) parameters to vary within reasonable ranges would result in reduced sensitivities as compared to the ones reported in these studies.

Since assumptions about which part of the SUSY spectrum might be accessible at the next generation of colliders are necessarily very speculative, we do not assume any further information beyond the Higgs sector at all and perform a full scan over the parameter space of the three soft SUSY-breaking scenarios. The resulting sensitivity to M_A (which effectively covers also possible theoretical uncertainties⁶) can thus be interpreted as a "worst case" scenario within mSUGRA, mGMSB and mAMSB, which could be improved by incorporating further information from other sectors of the model.

In Fig. 4 the indirect sensitivity to M_A within the mSUGRA scenario is investigated for the channels $h \rightarrow b\bar{b}$; $h \rightarrow \tau\bar{\tau}$; $h \rightarrow W^+W^-$ at the LC. The figure shows the regions in the M_A – $\tan\beta$ plane where the result in the mSUGRA scenario differs from the SM prediction by 1, 2 or 3%, according to the prospective accuracy at the LC as given in Tab. 3. The corresponding sensitivities at the C (which are not shown here) turn out to be usually worse than at the LC for the mSUGRA scenario.

If a 2 or 3% deviation of the Higgs branching ratios from the corresponding SM values is found at the LC, an upper bound on M_A can be inferred within the mSUGRA scenario according to Fig. 4. In particular, the $h \rightarrow W^+W^-$ channel yields an upper bound on M_A of 500–600 GeV (depending on $\tan\beta$) for a more than 3% deviation, 600–700 GeV for a deviation in excess of 2%, while deviations of more than 1% occur for M_A up to 800–1000 GeV within the mSUGRA scenario. On the other hand, measuring a suppression in the $h \rightarrow b\bar{b}$ and/or $h \rightarrow \tau\bar{\tau}$ channel (left column of Fig. 4) or an enhancement in the $h \rightarrow \tau\bar{\tau}$

⁶Note that the presently largest theoretical uncertainty in the MSSM Higgs sector, which arises from the experimental error of the top-quark mass, will be drastically reduced by the precise measurement of m_t at the LC.




Figure 4: Indirect sensitivity to M_A in the $mSUGRA$ scenario: for the channels $h \rightarrow b\bar{b}$, $h \rightarrow c\bar{c}$, $h \rightarrow \tau^+\tau^-$ and $h \rightarrow W^+W^-$ (from top to bottom) the regions in the M_A ($\tan\beta$) plane are shown where the result in the $mSUGRA$ scenario differs from the SM prediction by 1, 2 or 3 σ , assuming the prospective accuracy at the LC according to Tab. 3.

and/or $h \rightarrow W W$ channel (right column of Fig. 4) would determine $\tan \beta$ to lie within $35 < \tan \beta < 55$ in the mSUGRA scenario. The mSUGRA scenario is the only of the three soft SUSY-breaking scenarios that could accommodate a suppression of the $h \rightarrow b\bar{b}$ and/or $h \rightarrow \tau^+\tau^-$ channel⁷. Thus these measurements can help to distinguish the soft SUSY-breaking scenarios, see Sect. 4.3.




Figure 5: Indirect sensitivity to M_A in the mGMSB scenario: for the channels $h \rightarrow b\bar{b}$ (top left), $h \rightarrow c\bar{c}$ (top right), $h \rightarrow \tau^+\tau^-$ (bottom left) and $h \rightarrow W W$ (bottom right) the regions in the M_A $\{\tan \beta$ plane are shown where the result in the mGMSB scenario differs from the SM prediction by 1, 2 or 3, assuming the prospective accuracy at the LC according to Tab. 3.

In Fig. 5 we show the sensitivity to M_A within the mGMSB scenario for the channels $h \rightarrow b\bar{b}; c\bar{c}; \tau^+\tau^-; W W$ at the LC. The corresponding results for $h \rightarrow b\bar{b}; W W$ at the C are displayed in Fig. 6 (which yields comparable sensitivities in this scenario). As for mSUGRA, the observation of a 2 or 3 deviation compared to the SM prediction will allow to establish an upper bound on M_A within the mGMSB scenario. Also in this case the $h \rightarrow W W$ channel shows significant deviations from the SM prediction over a wider range of the M_A $\{\tan \beta$ plane than the other channels. For a more than 2 deviation in this channel an upper bound on M_A of about 700 GeV (depending somewhat on $\tan \beta$) can be inferred.

⁷In our O(50000) mGMSB scatter points we have found two points with $M_A \approx 100$ GeV and $\tan \beta \approx 55$ that exhibit a very strong suppression of the $h \rightarrow b\bar{b}$ and $h \rightarrow \tau^+\tau^-$ channel by more than 50%. However, these points appear to be rather fine-tuned and we did not include them into our analysis.




Figure 6: Indirect sensitivity to M_A in the mGMSB scenario: for the channels $h \rightarrow b\bar{b}$ (left) and $h \rightarrow W W$ (right) the regions in the M_A $\{\tan \beta$ plane are shown where the result in the mGMSB scenario differs from the SM prediction by 1, 2 or 3, assuming the prospective accuracy at the CLC according to Tab. 3.




Figure 7: Indirect sensitivity to M_A in the mAMSB scenario: for the channels $h \rightarrow b\bar{b}$ (top left), $h \rightarrow c\bar{c}$ (top right), $h \rightarrow \tau^+\tau^-$ (bottom left) and $h \rightarrow W W$ (bottom right) the regions in the M_A $\{\tan \beta$ plane are shown where the result in the mAMSB scenario differs from the SM prediction by 1, 2 or 3, assuming the prospective accuracy at the LC according to Tab. 3.

Bigger deviations result in correspondingly lower upper bounds on M_A .

In Fig. 7 the sensitivity to M_A within the mAMSB scenario is displayed for the channels $h \rightarrow b\bar{b}; c\bar{c}; \tau^+\tau^-; W^+W^-$ at the LC. The sensitivities at the C are usually worse in this scenario. As for the other two scenarios, in general an upper bound on M_A can be established if a 2 or 3 deviation from the SM result is observed. Again in particular the $h \rightarrow W^+W^-$ channel offers good prospects for observing sizable deviations. It allows to set an upper bound on M_A of 800–900 GeV (depending on $\tan\beta$) if a deviation of more than 2 is observed. Higher deviations result in correspondingly lower upper bounds on M_A . Comparing the results for the $h \rightarrow W^+W^-$ channel in the mAMSB scenario with the other scenarios, in the mAMSB scenario sizable deviations from the SM prediction occur over a wider parameter space in the $M_A\{\tan\beta$ plane than in the other scenarios. Thus, the prospects for experimentally establishing a deviation from the SM prediction and in this way inferring an upper bound on M_A appear to be particularly good in the mAMSB scenario.

4.2 Sensitivity to high-energy parameters

Besides providing sensitivity to M_A , precise measurements of Higgs branching ratios at the LC can also yield indirect information on the high-energy parameters of the different soft SUSY-breaking scenarios. In Fig. 8 the results for the channels $h \rightarrow b\bar{b}; c\bar{c}; \tau^+\tau^-; W^+W^-$ are shown in the $m_{1=2}\{m_0$ plane for the mSUGRA scenario. While the indirect constraints that can be obtained with a 2 or 3 deviation on m_0 are rather mild, stronger bounds can be obtained for $m_{1=2}$. This reflects the fact that M_A and the squark masses are strongly correlated with the $m_{1=2}$ value. Combining the channels, an upper bound of 350 GeV on $m_{1=2}$ can be set if a deviation of more than 3 from the SM prediction is observed, a 2 deviation constrains $m_{1=2}$ to be smaller than 450 GeV, while deviations of more than 1 occur for $m_{1=2} < 650$ GeV.

Concerning the mGMSB scenario (which is not displayed here) the indirect constraints are weaker. Results deviating from the SM prediction for the $h \rightarrow W^+W^-$ channel by 3, for instance, are distributed over nearly the whole $M_{\text{mess}}\{$ plane. Thus, establishing a non SM-like behavior in the Higgs sector alone is not sufficient to derive indirect bounds on M_{mess} and μ , i.e. further experimental information is necessary to constrain these parameters. On the other hand, weak lower limits on M_{mess} , could be set, which can cut out the lower edge of the mGMSB allowed $M_{\text{mess}}\{$ area, if the deviation from the SM value is found to be small.

In Fig. 9 the results for the channels $h \rightarrow b\bar{b}; c\bar{c}$ are shown in the $m_0\{m_{\text{aux}}$ plane for the mAMSB scenario. Like in the mGMSB scenario, deviations of 3 or 2 with respect to the SM prediction occur over a rather wide range of m_0 and m_{aux} values. An observed deviation of 3 would constrain m_0 to be smaller than 1100 GeV, while m_{aux} would have to be smaller than 6–10 GeV. Observation of a 2 deviation would allow to set an upper bound on m_0 of $m_0 < 1400$ GeV, while restricting the parameter space to the one compatible with a 2 deviation does not significantly reduce the range of possible values of m_{aux} in Fig. 9.




Figure 8: Indirect sensitivity to $m_0, m_{1/2}$ in the mSUGRA scenario: for the channels $h \rightarrow b\bar{b}$, $h \rightarrow c\bar{c}$, $h \rightarrow \tau^+\tau^-$ and $h \rightarrow W^+W^-$ (from top to bottom) the regions in the $m_{1/2}\{m_0\}$ plane are shown where the result in the mSUGRA scenario differs from the SM prediction by 1, 2 or 3 σ , assuming the prospective accuracy at the LC according to Tab. 3.




Figure 9: Indirect sensitivity to m_0, m_{aux} in the $m\text{AMSB}$ scenario: for the channels $h \rightarrow b\bar{b}$ (left) and $h \rightarrow c\bar{c}$ (right) the regions in the m_0 - m_{aux} plane are shown where the result in the $m\text{AMSB}$ scenario differs from the SM prediction by 1, 2 or 3, assuming the prospective accuracy at the LC according to Tab. 3.

4.3 Discrimination between soft SUSY-breaking scenarios

We now investigate the potential of precise measurements of Higgs branching ratios at a LC for distinguishing between the three soft SUSY-breaking scenarios. The main experimental test of different SUSY-breaking scenarios will of course be to confront their predictions for the SUSY spectrum with the results obtained from the direct observation of SUSY particles. The analysis of the Higgs sector, however, may contribute further valuable information. Since the different soft SUSY-breaking scenarios predict different mass patterns for the SUSY particles, variations in the results for the Higgs sector observables for the same value of M_A can be expected.

As above, we do not assume experimental input from other sectors of the MSSM, but concentrate on the Higgs sector. Clearly, resolving differences between the predictions of the three scenarios via measurements in the Higgs sector will require some experimental information on the tree-level parameters of the Higgs sector, M_A and $\tan\beta$. Therefore we will focus on a scenario where the LHC can detect the heavy MSSM Higgs bosons via their decays $H \rightarrow A \rightarrow \tau^+\tau^-$ (with the main production channel $b\bar{b} \rightarrow b\bar{b} H \rightarrow A$), which can be realized for large $\tan\beta$ and not too large M_A [74,100]. As a specific example we assume that the LHC provides a measurement of M_A as well as a lower bound on $\tan\beta$,

$$500 \text{ GeV} < M_A < 600 \text{ GeV}; \quad \tan\beta > 30 : \quad (24)$$

The results of the analysis below would improve for smaller values of M_A , while for a larger M_A and smaller $\tan\beta$ observation of the heavy Higgs bosons at the LHC would become increasingly difficult.

Restricting the data set of our scan, see Sect. 2.1, to those parameter points fulfilling Eq. (24) we compare the predictions for the different branching ratios arising from the three scenarios. As above, we indicate the deviations from the SM prediction in terms of the prospective accuracy at the LC according to Tab. 3.

In Fig. 10 we show the results for the channels $h \rightarrow b\bar{b}$ and $h \rightarrow gg$. The results for these channels, as for the others that are not shown ($h \rightarrow \tau^+\tau^-$, $h \rightarrow c\bar{c}$ and $h \rightarrow WW$), are similar: for the M_A values corresponding to Eq. (24) the $m\text{AMSB}$ scenario gives rise to larger





Figure 10: Comparison of $\text{BR}(h \rightarrow b\bar{b})$ (top) and $\text{BR}(h \rightarrow gg)$ (bottom) in the three soft SUSY-breaking scenarios via LC measurements.

deviations in the branching ratios from the SM values than the mGMSB scenario. Thus, if in the situation of Eq. (24) a 3 deviation from the SM value were found in $BR(h \rightarrow bb)$ and a 4 deviation in $BR(h \rightarrow gg)$, this would be better compatible with an AMSB scenario than with a mGMSB scenario. If, on the other hand, the branching ratios were found to agree well with the SM prediction, this would be best compatible with a SUSY-breaking scenario of mSUGRA type.

As a consequence, precision measurements at the LC of the branching ratios of the light CP-even Higgs boson of the MSSM may indicate a preference among the three soft SUSY-breaking scenarios at the 1-2 level. This information will be complementary to the information from the direct observation of SUSY particles.

5 Conclusions

We have investigated the relevant production and decay channels of the lightest CP-even MSSM Higgs boson at the Tevatron, the LHC, an e^+e^- LC, a C and a C within the mSUGRA, mGMSB and mAMSB scenarios. The values of BR have been compared with the corresponding SM values with the same Higgs boson mass, $M_{H_{SM}} = m_h$. In this context we have also updated earlier results on the upper bound on m_h within the three scenarios and on the lower bounds on $\tan\beta$ that can be inferred by confronting the theoretical predictions with the LEP exclusion limit.

We have first analyzed the observability of the lightest MSSM Higgs boson at the different colliders. The modes $gg \rightarrow h$, $tt \rightarrow tth$ and $WW \rightarrow h$ allow the detection of the lightest MSSM Higgs boson in all three scenarios over the whole corresponding parameter space. Possible exceptions occur for the very small M_A region in the mGMSB and mAMSB scenarios, where a strong suppression of more than 50% could happen for $gg \rightarrow h$. Within the clean experimental environment of the LC the observation of the light Higgs will be ensured for all three scenarios. For a C, the very small M_A region in mGMSB and mAMSB can be problematic for the $h \rightarrow bb$ and $h \rightarrow \tau\tau$ mode. At the C, $M_A < 150$ GeV and $\tan\beta > 50$ for mSUGRA exhibits a strong suppression for the $h \rightarrow bb$ and $h \rightarrow \tau\tau$ mode, while on the other hand in this parameter region the production of the heavy MSSM Higgs bosons H, A happens with an enhanced rate. Besides these difficult regions, the main search modes at a C and a C do not suffer from severe suppressions with respect to the SM case in all three scenarios. The results of this analysis are summarized in Tab. 2. Thus, all possible future colliders offer very good prospects for detecting the lightest CP-even Higgs boson of the MSSM in an mSUGRA, mGMSB or mAMSB scenario.

We then investigated the potential of precision measurements of Higgs branching ratios at the LC and the C for establishing indirect constraints on M_A and $\tan\beta$. For this analysis we have not assumed any further experimental information on the SUSY spectrum, i.e. a full scan over the parameter space (restricting to the case $\mu > 0$) has been performed. If deviations of the Higgs branching ratios from their SM values will be found at the 2-3 level, it will be possible to establish an upper bound for M_A significantly below 1 TeV in all three scenarios. The biggest sensitivity will come from the $h \rightarrow WW$ and $h \rightarrow \tau\tau$ channels. Within the mSUGRA scenario, furthermore a bound on $\tan\beta$ of $35 < \tan\beta < 55$ can be obtained if a suppression of the $h \rightarrow bb$ and/or $h \rightarrow \tau\tau$ channel or an enhancement in the $h \rightarrow \tau\tau$ and/or $h \rightarrow WW$ channel with respect to the SM values is observed. If this would

be the case, this could be independently constrained by Higgs mediated B-physics observables like $B^0 \rightarrow \pi^+ \pi^-$ or $B^0 \rightarrow B^0$ mixing.

Similarly, precise measurements of $\Gamma(B \rightarrow \pi)$ BR at the LC can also provide indirect information on the high-energy parameters of the three soft SUSY-breaking scenarios. While within the mGMSB scenario the experimental determination of the Higgs branching ratios will allow to set only very weak bounds on the high-energy parameters, within mSUGRA relatively strong bounds on $m_{1=2}$ and in mAMSB moderate bounds on m_0 could be set.

Finally we have investigated the potential of precise measurements of $\Gamma(B \rightarrow \pi)$ BR at a LC to distinguish between the three soft SUSY-breaking scenarios. For this analysis we have assumed a situation where experimental information on M_A (and to a lesser extent on $\tan \beta$) obtained at the LHC can be combined with precision measurements of the properties of the light Higgs boson at the LC, see also Ref. [101]. If a significant suppression of the $h \rightarrow b\bar{b}$ and/or $h \rightarrow \tau^+ \tau^-$ channel with respect to its SM value were found, this would point towards the mSUGRA scenario, irrespectively of the actual value of M_A (with $M_A < 1$ TeV). Otherwise, assuming in our example M_A to be restricted to $500 \text{ GeV} < M_A < 600 \text{ GeV}$, precise measurements of $\Gamma(B \rightarrow \pi)$ BR in particular in the $h \rightarrow b\bar{b}$ and $h \rightarrow \tau^+ \tau^-$ channels may indicate a preference among the three soft SUSY-breaking scenarios at the 1-2 level. This information might be valuable as it complements the one about the SUSY spectrum from the direct observation of SUSY particles.

Acknowledgements

We thank S. Ambrosanio for helpful discussion on the GMSB scenario as well as for his Fortran code SUSYFIRE. We thank T. Plehn for discussions. A.D. thanks P. Slavich for interesting discussions. S.S. has been supported by the DOE grant DE-FG 03-92-ER-40701. A.D. would like to acknowledge financial support from the Network RTN European Program HPRN-CT-2000-00148 "Physics Across the Present Energy Frontier: Probing the Origin of Mass". This work has been supported by the European Community's Human Potential Programme under contract HPRN-CT-2000-00149 "Physics at Colliders".

References

- [1] G. Kane, C. Kolda and J. Wells, Phys. Rev. Lett. 70 (1993) 2686, hep-ph/9210242;
J. Espinosa and M. Quiros, Phys. Lett. B 302 (1993) 51, hep-ph/9212305.
- [2] [LEP Higgs working group], LHWG Note/2002-01, see:
lephiggs.web.cern.ch/LEPHIGGS/papers/.
- [3] S. Heinemeyer, W. Hollik and G. Weiglein, Eur. Phys. Jour. C 9 (1999) 343,
hep-ph/9812472.
- [4] G. Degrandi, S. Heinemeyer, W. Hollik, P. Slavich and G. Weiglein, to appear in Eur.
Phys. Jour. C, hep-ph/0212020.
- [5] S. Heinemeyer, W. Hollik and G. Weiglein, Phys. Rev. D 58 (1998) 091701,
hep-ph/9803277; Phys. Lett. B 440 (1998) 296, hep-ph/9807423.

- [6] M . Carena, J. Espinosa, M . Quiros and C . W agner, Phys. Lett. B 355 (1995) 209, hep-ph/9504316.
- [7] M . Carena, M . Quiros and C . W agner, Nucl. Phys. B 461 (1996) 407, hep-ph/9508343.
- [8] H . Haber, R . H em p ing and A . Hoang, Z . Phys. C 75 (1997) 539, hep-ph/9609331.
- [9] R . H em p ing and A . Hoang, Phys. Lett. B 331 (1994) 99, hep-ph/9401219.
- [10] R . Zhang, Phys. Lett. B 447 (1999) 89, hep-ph/9808299.
- [11] J. Espinosa and R . Zhang, Nucl. Phys. B 586 (2000) 3, hep-ph/0003246.
- [12] J. Espinosa and I. Navarro, Nucl. Phys. B 615 (2001) 82, hep-ph/0104047.
- [13] G . D egrassi, P . Slavich and F . Zw imer, Nucl. Phys. B 611 (2001) 403, hep-ph/0105096;
A . Brignole, G . D egrassi, P . Slavich and F . Zw imer, Nucl. Phys. B 631 (2002) 195, hep-ph/0112177; Nucl. Phys. B 643 (2002) 79, hep-ph/0206101.
- [14] S. M artin, Phys. Rev. D 65 (2002) 116003, hep-ph/0111209; Phys. Rev. D 66 (2002) 096001, hep-ph/0206136; hep-ph/0211366.
- [15] M . Carena, S. Heinem eyer, C . W agner and G . W eiglein, hep-ph/9912223.
- [16] M . Carena, S. Heinem eyer, C . W agner and G . W eiglein, Eur. Phys. Jour. C 26 (2003) 601, hep-ph/0202167.
- [17] B . A llanach et al., Eur. Phys. J. C 25 (2002) 113, hep-ph/0202233.
- [18] H . N illes, Phys. Lett. B 115 (1982) 193; Nucl. Phys. B 217 (1983) 366.
A . Cham seddine, R . A mow itt and P . Nath, Phys. Rev. Lett. 49 (1982) 970;
R . Barbieri, S. Ferrara and C . Savoy, Phys. Lett. B 119 (1982) 343;
H . N illes, M . Srednicki and D . W yler, Phys. Lett. B 120 (1983) 346.
E . C rem m er, P . Fayet and L . G irardello, Phys. Lett. B 122 (1983) 41.
S. Ferrara, L . G irardello and H . N illes, Phys. Lett. B 125 (1983) 457. L . H all, J. Lykken
and S. W einberg, Phys. Rev. D 27 (1983) 2359;
S. Soni and H . W eldon, Phys. Lett. B 126 (1983) 215.
R . A mow itt, A . Cham seddine and P . Nath, Nucl. Phys. B 227 (1983) 121.
For more details see, S. W einberg, "The quantum theory of elds. Vol 3: Supersym m e-
try," C am bridge U niversity P ress (2000).
- [19] For review s see also:
H . N illes, Phys. Rept. 110 (1984) 1;
H . Haber and G . K ane, Phys. Rept. 117 (1985) 75;
A . Lahanas and D . N anopoulos, Phys. Rept. 145 (1987) 1;
S. M artin, in "Perspectives on supersym m etry", ed. G . K ane, hep-ph/9709356, see:
zippy.physics.niu.edu/primer.shtml.
- [20] For a review , see: G F . Giudice and R . Rattazzi, Phys. Rept. 322 (1999) 419, hep-ph/9801271.

- [21] L. Randall and R. Sundrum, Nucl. Phys. B 557 (1999) 79, hep-th/9810155.
- [22] G.F. Giudice, M.A. Luty, H. Murayama and R. Rattazzi, JHEP 9812 (1998) 027, hep-ph/9810442.
- [23] T. Gherghetta, G.F. Giudice, J.D. Wells, Nucl. Phys. B 559 (1999) 27, hep-ph/9904378.
- [24] K. Matchev and D. Pierce, Phys. Lett. B 445 (1999) 331, hep-ph/9805275; W. de Boer, hep-ph/9808448.
- [25] T. Kaeding and S. Nandi, hep-ph/9906342.
- [26] S. Su, Nucl. Phys. B 573 (2000) 87, hep-ph/9910481.
- [27] B. Allanach and A. Dedes, JHEP 0006 (2000) 017, hep-ph/0003222.
- [28] A. Dedes, S. Heinemeyer, P. Teixeira-Dias and G. Weiglein, Jour. Phys. G 26 (2000) 582, hep-ph/9912249.
- [29] S. Ambrosanio, S. Heinemeyer and G. Weiglein, in hep-ph/0002191 and hep-ph/0005142.
- [30] J. Ellis, G. Gani, D. Nanopoulos and K. Olive, Phys. Lett. B 502 (2001) 171, hep-ph/0009355.
- [31] A. Djouadi, M. Drees and J. Kneur, JHEP 0108 (2001) 055, hep-ph/0107316.
- [32] S. Ambrosanio, A. Dedes, S. Heinemeyer, S. Su and G. Weiglein, Nucl. Phys. B 624 (2001) 3, hep-ph/0106255; A. Dedes, S. Heinemeyer, S. Su and G. Weiglein, hep-ph/0110219.
- [33] S. Martin and M. Vaughn, Phys. Rev. D 50 (1994) 2282, hep-ph/9311340; Y. Yamada, Phys. Rev. D 50 (1994) 3537, hep-ph/9401241; I. Jack and D. Jones, Phys. Lett. B 333 (1994) 372, hep-ph/9405233.
- [34] B. Allanach, S. Kramer and W. Porod, hep-ph/0302102.
- [35] M. Carena, H. Haber, S. Heinemeyer, W. Hollik, C. Wagner and G. Weiglein, Nucl. Phys. B 580 (2000) 29, hep-ph/0001002.
- [36] S. Heinemeyer, W. Hollik and G. Weiglein, hep-ph/9910283.
- [37] J. Espinosa and R. Zhang, JHEP 0003 (2000) 026, hep-ph/9912236.
- [38] S. Heinemeyer, W. Hollik and G. Weiglein, Comp. Phys. Comm. 124 2000 76, hep-ph/9812320; hep-ph/0002213.
The codes are accessible via www.feynhiggs.de.
- [39] M. Frank, S. Heinemeyer, W. Hollik and G. Weiglein, hep-ph/0202166.
- [40] An advanced (and so far unpublished) version of FeynHiggs contains also the evaluation of the Higgs boson decay rates, combining the results from Refs. [41,42].

- [41] S. Heinemeyer, W. Hollik and G. Weiglein, Eur. Phys. Jour. C 16 (2000) 139, hep-ph/0003022.
- [42] A. Djouadi, J. Kalinowski and M. Spira, Comput. Phys. Commun. 108 (1998) 56, hep-ph/9704448.
- [43] J. Ellis, S. Heinemeyer, K. Olive and G. Weiglein, Phys. Lett. B 515 (2001) 348, hep-ph/0105061; JHEP 0301 (2003) 006, hep-ph/0211206.
- [44] H. Dreiner, hep-ph/9707435;
G. Bhattacharyya, hep-ph/9709395;
B. Allanach, A. Dedes and H. Dreiner, Phys. Rev. D 60 (1999) 075014, hep-ph/9906209.
- [45] Particle Data Group, Phys. Rev. D 66 (2002) 010001.
- [46] H. Brown et al. [Muon g-2 Collaboration], Phys. Rev. Lett. 86 (2001) 2227, hep-ex/0102017; updated as hep-ex/0208001;
A. Czarnecki and W. Marciano, Phys. Rev. D 64 (2001) 013014, hep-ph/0102122;
M. Davier, S. Eidelman, A. Hocker and Z. Zhang, hep-ph/0208177.
- [47] S. Chen et al. [CLEO Collaboration], Phys. Rev. Lett. 87 (2001) 251807, hep-ex/0108032;
Y. Ushiroda [Belle Collaboration], hep-ex/0104045;
B. Aubert et al. [BaBar Collaboration], hep-ex/0207076.
- [48] A. Dedes, A. Lahanas and K. Tamvakis, Phys. Rev. D 53, 3793 (1996), hep-ph/9504239.
- [49] A. Dedes, S. Heinemeyer and G. Weiglein, "FeynSSG: A program for the minimal supergravity and Higgs spectrum", in
G. Azeels et al., [Les Houches Beyond the SM working group], Summary report, hep-ph/0204031.
- [50] M. Dine, A. Nelson, Phys. Rev. D 48 (1993) 1277, hep-ph/9303230;
M. Dine, A. Nelson, Y. Shiman, Phys. Rev. D 51 (1995) 1362, hep-ph/9408384;
M. Dine, A. Nelson, Y. Nir, Y. Shiman, Phys. Rev. D 53 (1996) 2658, hep-ph/9507378.
- [51] M. Dine, W. Fischler, M. Srednicki, Nucl. Phys. B 189 (1981) 575;
S. Dimopoulos, S. Raby, Nucl. Phys. B 192 (1981) 353;
M. Dine, W. Fischler, Phys. Lett. B 110 (1982) 227;
M. Dine, M. Srednicki, Nucl. Phys. B 202 (1982) 238;
M. Dine, W. Fischler, Nucl. Phys. B 204 (1982) 346;
L. Alvarez-Gaume, M. Claudson, M. Wise, Nucl. Phys. B 207 (1982) 96;
C. Nappi, B. Ovrut, Phys. Lett. B 113 (1982) 175;
S. Dimopoulos, S. Raby, Nucl. Phys. B 219 (1983) 479.
- [52] P. Fayet, Phys. Lett. B 70 (1977) 461; Phys. Lett. B 86 (1979) 272; Phys. Lett. B 175 (1986) 471; and in "Unification of the fundamental particle interactions", eds. S. Ferrara, J. Ellis, P. van Nieuwenhuizen (Plenum, New York, 1980) p. 587.

- [53] An updated, generalized and Fortran-linked version of the program used in Ref. [54]. It generates minimal and non-minimal GMSB and SUGRA models. For inquiries about this software package, please send e-mail to Sandro.Ambrosanio@bancaroma.it.
- [54] S. Ambrosanio, G. Kribs, S. Martin, Phys. Rev. D 56 (1997) 1761, hep-ph/9703211.
- [55] S. Ambrosanio, G. Blair, Eur. Phys. Jour. C 12 (2000) 287, hep-ph/9905403.
- [56] S. Ambrosanio et al., JHEP 0101 (2001) 014, hep-ph/0010081.
- [57] A. Pomarol and R. Rattazzi, JHEP 9905 (1999) 013, hep-ph/9903448.
- [58] Z. Chacko, M. Luty and E. Ponton, JHEP 0004 (2000) 001, hep-ph/9905390.
- [59] E. Katz, Y. Shadmi and Y. Shiman, JHEP 9908 (1999) 015, hep-ph/9906296.
- [60] I. Jack, D. Jones and R. Ward, Phys. Lett. B 535, 193 (2002), hep-ph/0202101.
- [61] L. J. Hall, R. Rattazzi and U. Sarid, Phys. Rev. D 50 (1994) 7048, hep-ph/9306309.
- [62] M. Carena, S. Mrenna and C. Wagner, Phys. Rev. D 60 (1999) 075010, hep-ph/9808312; Phys. Rev. D 62 (2000) 055008, hep-ph/9907422; H. Eberl, K. Hidaka, S. Kramer, W. Majerotto and Y. Yamada, Phys. Rev. D 62 (2000) 055006, hep-ph/9912463; M. Carena, D. Garcia, U. Nierste and C. Wagner, Nucl. Phys. B 577 (2000) 88, hep-ph/9912516.
- [63] S. Heinemeyer, W. Hollik, J. Rosiek and G. Weiglein, Eur. Phys. Jour. C 19 (2001) 535, hep-ph/0102081.
- [64] T. Hahn, S. Heinemeyer and G. Weiglein, Nucl. Phys. B 652 (2003) 229, hep-ph/0211204.
- [65] H. Eberl, W. Majerotto, and V. Spanos, Phys. Lett. B 538 (2002) 353, hep-ph/0204280.
- [66] S. Dawson, A. Djouadi and M. Spira, Phys. Rev. Lett. 77 (1996) 16, hep-ph/9603423.
- [67] [LEP Higgs Working Group Collaboration], hep-ex/0107030; LHWG Note 2001-4, see: lephiggs.web.cern.ch/LEPHIGGS/papers/.
- [68] R. Barate et al. [ALEPH Collaboration], Phys. Lett. B 499 (2001) 53, hep-ex/0010062.
- [69] M. Veltman, Nucl. Phys. B 123 (1977) 89.
- [70] A. Djouadi, P. Gambino, S. Heinemeyer, W. Hollik, C. Junger and G. Weiglein, Phys. Rev. Lett. 78 (1997) 3626, hep-ph/9612363; Phys. Rev. D 57 (1998) 4179, hep-ph/9710438; S. Heinemeyer and G. Weiglein, in Proc. of the 5th International Symposium on Radiative Corrections (RADCOR 2000) ed. H. Haber, hep-ph/0102317. For a recent update, see also: S. Heinemeyer and G. Weiglein, JHEP 0210 (2002) 072, hep-ph/0209305; hep-ph/0301062.

- [71] A. D edes and P. Slavich, hep-ph/0212132.
- [72] J. Ellis, J. Hagelin, D. Nanopoulos, K. Olive and M. Srednicki, Nucl. Phys. B 238 (1984) 453;
H. Goldberg, Phys. Rev. Lett. 50 (1983) 1419;
J. Ellis, T. Falk, G. Gani, K. Olive and M. Srednicki, Phys. Lett. B 510 (2001) 236, hep-ph/0102098.
- [73] S. Heinemeyer, W. Hollik and G. Weiglein, JHEP 0006 (2000) 009, hep-ph/9909540.
- [74] ATLAS Collaboration, Detector and Physics Performance Technical Design Report, CERN/LHCC/99-15 (1999), see:
atlasinfo.cern.ch/Atlas/GROUPS/PHYSICS/TDR/access.html.
- [75] A. Djouadi, Phys. Lett. B 435 (1998) 101, hep-ph/9806315.
- [76] G. Belanger, F. Boudjema and K. Sridhar, Nucl. Phys. B 568 (2000) 3, hep-ph/9904348;
G. Belanger, F. Boudjema, F. Donato, R. Godbole and S. Rosier-Lees, Nucl. Phys. B 581 (2000) 3, hep-ph/0002039.
- [77] E. Richter-Was and D. Froidevaux, ATL-COM-PHYS-98-012.
- [78] A. Belyaev, D. Garcia, J. Guasch and J. Sola, Phys. Rev. D 65 (2002) 031701, hep-ph/0105053.
- [79] D. Zeppenfeld, R. Kinnunen, A. Nikitenko and E. Richter-Was, Phys. Rev. D 62 (2000) 013009, hep-ph/0002036.
- [80] D. Rainwater and D. Zeppenfeld, Phys. Rev. D 60 (1999) 113004, E: ibid D 61 (2000) 099901, hep-ph/9906218;
D. Rainwater, hep-ph/9908378;
N. Kauer, T. Plehn, D. Rainwater and D. Zeppenfeld, Phys. Lett. B 503 (2001) 113, hep-ph/0012351.
- [81] T. Plehn, D. Rainwater and D. Zeppenfeld, Phys. Lett. B 454 (1999) 297, hep-ph/9902434.
- [82] T. Plehn, D. Rainwater and D. Zeppenfeld, Phys. Rev. D 61 (2000) 093005, hep-ph/9911385.
- [83] F. Maltoni, D. Rainwater and S. Willenbrock, Phys. Rev. D 66 (2002) 034022, hep-ph/0202205.
- [84] TESLA TDR Part 3: "Physics at an e^+e^- Linear Collider", eds. R. Heuer, D. Miller, F. Richard and P. Zerwas, hep-ph/0106315, see: tesla.desy.de/tdr.
- [85] T. Abe et al. [American Linear Collider Working Group Collaboration], Resource book for Snowmass 2001, hep-ex/0106055, hep-ex/0106056, hep-ex/0106057, hep-ex/0106058.
- [86] K. Abe et al. [ACFA Linear Collider Working Group Collaboration], hep-ph/0109166.

- [87] TESLA TDR Part 6, Chapter 1: "Photon collider at TESLA", [hep-ex/0108012](#).
- [88] D. Asner et al., to appear in Eur. Phys. Jour. C , [hep-ex/0111056](#).
- [89] C. Blochinger et al. [Higgs factory working group of the ECFA-CERN study on Neutrino Factory & Muon Storage Rings at CERN], "Physics opportunities at e^+e^- Higgs factories", [hep-ph/0202199](#).
- [90] S. Heinemeyer and G. Weiglein, BNL-22-01.
- [91] For a summary see: D. Zeppenfeld, in Proc. of the APS/DPF/DPB Summer Study on the Future of Particle Physics (Snowmass 2001) ed. R. Davidson and C. Quigg, [hep-ph/0203123](#).
- [92] For a summary see: H. Haber, [hep-ph/9703381](#).
- [93] J. Brient, talk at the Linear Collider Workshop, Cracow, Poland, September 2001, see: [webnt.physics.ox.ac.uk/lc/ecfadesy](#)
- [94] CMS Collaboration, see: [cmsinfo.cern.ch/Welcome.html/CMSdocuments/CMSplots/](#).
- [95] T. Hahn, S. Heinemeyer and G. Weiglein, [hep-ph/0211384](#).
- [96] J. Guasch, W. Hollik and S. Penaranda, Phys. Lett. B 515 (2001) 367, [hep-ph/0106027](#).
- [97] M. Carena, H. Haber, H. Logan and S. Mrenna, Phys. Rev. D 65 (2002) 055005, E: *ibid* D 65 (2002) 099902, [hep-ph/0106116](#).
- [98] S. Dawson and S. Heinemeyer, Phys. Rev. D 66 (2002) 055002, [hep-ph/0203067](#).
- [99] J. Erler, S. Heinemeyer, W. Hollik, G. Weiglein and P. Zerwas, Phys. Lett. B 486 (2000) 125, [hep-ph/0005024](#);
J. Erler and S. Heinemeyer, in Proc. of the 5th International Symposium on Radiative Corrections (RADCOR 2000) ed. H. Haber, [hep-ph/0102083](#).
- [100] D. Cavalliet al., [Les Houches Higgs working group], Summary report, [hep-ph/0203056](#).
- [101] G. Weiglein, [hep-ph/0301111](#), see also: [www.ippp.dur.ac.uk/georg/lhclc/](#).