



Determining weak phases using $B \rightarrow D^* V$ Decays*

David London^a, Nita Sinha^b and Rahul Sinha^b

^a Laboratoire René J.-A. L  vesque, Universit   de Montr  al,
C.P. 6128, succ. centre-ville, Montr  al, QC, Canada H3C 3J7

^bInstitute of Mathematical Sciences, Taramani, Chennai 600113, India

We describe how the angular analysis of vector-vector final states in B decays provides theoretically clean techniques for determination of CP violating phases. The quantity $\sin^2(2\beta + \gamma)$ can be cleanly obtained from the time dependent study of decays such as $B_d^0(t) \rightarrow D^{*\pm} \rho^\mp$, $D^{*\pm} a_1^\mp$ etc. Similarly, one can use $B_s^0(t) \rightarrow D_s^{*\pm} K^{*\mp}$ to extract $\sin^2 \gamma$. A time independent study of the charged decay modes $B^\pm \rightarrow D^{(*)0} K^{*\pm}$ can also be used to extract γ .

1 Why look at VV modes for determining CPV Phases?

A precise determination of all CP violating angles [1] is one of the major goals of the current and future B Physics experiments. The measured values of these angles may be consistent with the standard model (SM) predictions, or they may indicate the presence of physics beyond the standard model. Early indications are that new physics, if present, is likely to have rather small effects on the angles of the unitarity triangle. Hence, to uncover any new physics, it is extremely important that all the CP violating angles be determined without theoretical uncertainties.

In the early days of the field, it was thought that the CP angles could be easily measured in $B_d^0(t) \rightarrow \pi^+ \pi^-$ (α), $B_d^0(t) \rightarrow \Psi K_s$ (β), and $B_s^0(t) \rightarrow \rho K_s$ (γ). However, it soon became clear that things would not be so easy: the presence of penguin amplitudes [2] makes the extraction of α from $B_d^0(t) \rightarrow \pi^+ \pi^-$ quite difficult, and completely spoil the measurement of γ in $B_s^0(t) \rightarrow \rho K_s$. In fact, determining γ through techniques which are theoretically clean as well as experimentally feasible has been a challenge.

The problem of penguin pollution can be avoided by considering decay modes that involve only tree amplitudes. In this talk we first indicate the practical problems encountered by most of the clean methods to determine γ , using pseudoscalar-pseudoscalar (PP) and pseudoscalar-vector (PV) final states and involving only tree amplitudes. We next demonstrate, how the use of the corresponding vector-vector (VV) final state modes, resolves these problems. The rich kinematics of the VV modes accessible via an angular analysis, provides a large number of observables which allows clean extraction of CP phases.

As a first example, we consider the PP or PV final states, $f \equiv D^- \pi^+ (D^{*-} \pi^+)$. Both B^0 and \bar{B}^0 can decay to these final

states and only one weak amplitude contributes. Hence, the amplitudes for these decay modes may be written as,

$$A \equiv \text{Amp}(B^0 \rightarrow f) = a e^{i\delta_a} e^{i\phi_a}, \quad (1)$$

$$A' \equiv \text{Amp}(\bar{B}^0 \rightarrow f) = b e^{i\delta_b} e^{i\phi_b}. \quad (2)$$

where, $\phi_a = 0$ and $\phi_b = -\gamma$. Because of B^0 - \bar{B}^0 mixing, CP violation comes about due to an interference between the amplitudes $B^0 \rightarrow f$ and $\bar{B}^0 \rightarrow f$. Note that since both B^0 and \bar{B}^0 can also decay to \bar{f} , one can measure the four time dependent decay rates, $\Gamma(B_d^0(t) \rightarrow f)$, $\Gamma(\bar{B}_d^0(t) \rightarrow f)$, $\Gamma(B_d^0(t) \rightarrow \bar{f})$ and $\Gamma(\bar{B}_d^0(t) \rightarrow \bar{f})$. It is therefore possible to determine the weak phase $\phi = (2\beta + \gamma)$ [3, 4]. However, the decay amplitude, $b \ll a$ and hence, the decay rate for $\Gamma(\bar{B}_d^0 \rightarrow D^- \pi^+)$ is expected to be small.

The ratio, $\Gamma(\bar{B}_d^0 \rightarrow D^- \pi^+)/\Gamma(B_d^0 \rightarrow D^- \pi^+)$, is essentially $|V_{ub} V_{cd}^* / V_{cb}^* V_{ud}|^2 \simeq 4 \times 10^{-4}$. Obviously, it will be very difficult to measure this tiny quantity with any precision, and therefore it would not be viable to carry out this method in practice. On the other hand, we will demonstrate that with the corresponding VV final states ($D^{*\pm} \rho^\mp$), one can extract $\sin^2(2\beta + \gamma)$ using an angular analysis, without the knowledge of the smaller amplitude [5].

The second example that we examine, is that of direct CP violation, in the modes: $B^\pm \rightarrow \bar{D}^0 K^\pm$, $B^\pm \rightarrow \bar{D}^0 \pi^\pm$. In this method [6], γ is obtained from an interference of the mode $B \rightarrow D^0 K$ with $B \rightarrow \bar{D}^0 K$, which occurs if and only if, both D^0 and \bar{D}^0 decay to a common final state f ; in particular, f is taken to be a CP eigenstate. This technique of extracting γ requires a measurement of the branching ratio for $B^+ \rightarrow D^0 K^+$ which is not experimentally feasible as pointed out in [7]. Moreover, the CP violating asymmetries tend to be small as the interfering amplitudes are not comparable. The use of non-CP eigenstates ' f ' has also been considered [8] in literature. Atwood, Dunitz and Soni (ADS) [7] extended this proposal by considering

*Talk presented by Nita Sinha

' f ' to be non- CP eigenstates that are also doubly Cabbibo suppressed modes of D . The two interfering amplitudes then are of the same magnitude resulting in large asymmetries. Their proposal is to use two final states f_1 and f_2 with at least one being a non- CP eigenstate. The use of more than one final state enables not only the determination of γ , but also of all the strong phases involved and the difficult to measure branching ratio $Br(B^+ \rightarrow D^0 K^+)$. However, an input into the determination of γ is the branching ratio of the doubly Cabbibo suppressed mode of D . Here again, the VV final states, provide an alternative [9]. (Some other plausible methods have also been recently proposed [10].) The VV modes $\bar{D}^{*0} K^{*\pm}, \bar{D}^{*0} \rho^\pm$, enable extraction of γ and all the unknowns involved, including the BR for Doubly Cabibbo-suppressed mode of D .

2 Vector-Vector final state decay amplitudes

The most general covariant amplitude for a B meson decaying to a pair of vector mesons has the form [11]

$$A(B(p) \rightarrow V_1(k, \epsilon_1) V_2(q, \epsilon_2)) = \epsilon_1^{*\mu} \epsilon_2^{*\nu} \times \left(a g_{\mu\nu} + \frac{b}{m_1 m_2} p_\mu p_\nu + i \frac{c}{m_1 m_2} \epsilon_{\mu\nu\alpha\beta} k^\alpha q^\beta \right), \quad (3)$$

where, ϵ_1, ϵ_2 and m_1, m_2 represent the polarization vectors and the masses of the vector mesons V_1 and V_2 respectively. The coefficients a, b , and c can be expressed in terms of the linear polarization basis A_\parallel, A_\perp and A_0 as follows:

$$\begin{aligned} A_0 &= -xa - (x^2 - 1)b, \\ A_\parallel &= \sqrt{2}a, \\ A_\perp &= \sqrt{2(x^2 - 1)}c, \end{aligned} \quad (4)$$

where $x = k \cdot q / (m_1 m_2)$. If both mesons subsequently decay into two $J^P = 0^-$ mesons, *i.e.* $V_1 \rightarrow P_1 P'_1$ and $V_2 \rightarrow P_2 P'_2$, the amplitude can be expressed as [9, 12]

$$A(B \rightarrow V_1 V_2) \propto (A^0 \cos \theta_1 \cos \theta_2 + \frac{A^\perp}{\sqrt{2}} \sin \theta_1 \sin \theta_2 \cos \phi - i \frac{A^\perp}{\sqrt{2}} \sin \theta_1 \sin \theta_2 \sin \phi), \quad (5)$$

where $\theta_1(\theta_2)$ is the angle between the $P_1(P_2)$ three-momentum vector, $\vec{k}_1(\vec{q}_1)$ in the $V_1(V_2)$ rest frame and the direction of total V_1 (V_2) three-momentum vector defined in the B rest frame. ϕ is the angle between the normals to the planes defined by $P_1 P'_1$ and $P_2 P'_2$, in the B rest frame.

3 Time dependent analysis in $B \rightarrow VV$

We consider a final state f , consisting of two vector mesons, to which both B^0 and \bar{B}^0 can decay. If only one

weak amplitude contributes to $B^0 \rightarrow f$ and $\bar{B}^0 \rightarrow f$, we can write the helicity amplitudes as follows:

$$A_\lambda \equiv \text{Amp}(B^0 \rightarrow f)_\lambda = a_\lambda e^{i\delta_\lambda^a} e^{i\phi_a}, \quad (6)$$

$$A'_\lambda \equiv \text{Amp}(\bar{B}^0 \rightarrow f)_\lambda = b_\lambda e^{i\delta_\lambda^b} e^{i\phi_b}, \quad (7)$$

$$\bar{A}'_\lambda \equiv \text{Amp}(B^0 \rightarrow \bar{f})_\lambda = b_\lambda e^{i\delta_\lambda^b} e^{-i\phi_b}, \quad (8)$$

$$\bar{A}_\lambda \equiv \text{Amp}(\bar{B}^0 \rightarrow \bar{f})_\lambda = a_\lambda e^{i\delta_\lambda^a} e^{-i\phi_a}, \quad (9)$$

where the helicity index λ takes the values $\{0, \parallel, \perp\}$. In the above, $\phi_{a,b}$ and $\delta_\lambda^{a,b}$ are the weak and strong phases, respectively. Using CPT invariance, the total decay amplitudes can be written as

$$\mathcal{A} = \text{Amp}(B^0 \rightarrow f) = A_0 g_0 + A_\parallel g_\parallel + i A_\perp g_\perp, \quad (10)$$

$$\bar{\mathcal{A}} = \text{Amp}(\bar{B}^0 \rightarrow \bar{f}) = \bar{A}_0 g_0 + \bar{A}_\parallel g_\parallel - i \bar{A}_\perp g_\perp, \quad (11)$$

$$\mathcal{A}' = \text{Amp}(\bar{B}^0 \rightarrow f) = A'_0 g_0 + A'_\parallel g_\parallel - i A'_\perp g_\perp, \quad (12)$$

$$\bar{\mathcal{A}}' = \text{Amp}(B^0 \rightarrow \bar{f}) = \bar{A}'_0 g_0 + \bar{A}'_\parallel g_\parallel + i \bar{A}'_\perp g_\perp, \quad (13)$$

where the g_λ are the coefficients of the helicity amplitudes, defined using Eq. (5) and depend only on the angles describing the kinematics. With the above equations, the time-dependent decay rate for a B^0 decaying into the two vector-meson final state, *i.e.* $B^0(t) \rightarrow f$, can be written as

$$\Gamma(B^0(t) \rightarrow f) = e^{-\Gamma t} \sum_{\lambda \leq \sigma} \left(\Lambda_{\lambda\sigma} + \Sigma_{\lambda\sigma} \cos(\Delta M t) - \rho_{\lambda\sigma} \sin(\Delta M t) \right) g_\lambda g_\sigma. \quad (14)$$

By performing a time-dependent study and angular analysis of the decay $B^0(t) \rightarrow f$, one can measure the 18 observables $\Lambda_{\lambda\sigma}, \Sigma_{\lambda\sigma}$ and $\rho_{\lambda\sigma}$. In terms of the helicity amplitudes $A_0, A_\parallel, A_\perp$, these can be expressed as:

$$\begin{aligned} \Lambda_{\lambda\lambda} &= \frac{|A_\lambda|^2 + |A'_\lambda|^2}{2}, & \Sigma_{\lambda\lambda} &= \frac{|A_\lambda|^2 - |A'_\lambda|^2}{2}, \\ \Lambda_{\perp i} &= -\text{Im}(A_\perp A_i^* - A'_\perp A_i'^*), & \Lambda_{\parallel 0} &= \text{Re}(A_\parallel A_0^* + A'_\parallel A_0'^*), \\ \Sigma_{\perp i} &= -\text{Im}(A_\perp A_i^* + A'_\perp A_i'^*), & \Sigma_{\parallel 0} &= \text{Re}(A_\parallel A_0^* - A'_\parallel A_0'^*), \\ \rho_{\perp i} &= -\text{Re}\left(\frac{q}{p} [A_\perp^* A_i' + A_i^* A'_\perp]\right), & \rho_{\perp\perp} &= -\text{Im}\left(\frac{q}{p} A_\perp^* A'_\perp\right), \\ \rho_{\parallel 0} &= \text{Im}\left(\frac{q}{p} [A_\parallel^* A_0' + A_0^* A'_\parallel]\right), & \rho_{ii} &= \text{Im}\left(\frac{q}{p} A_i^* A_i'\right), \end{aligned} \quad (15)$$

where $i = \{0, \parallel\}$. In the above, $q/p = \exp(-2i\phi_M)$, where ϕ_M is the weak phase present in B^0 - \bar{B}^0 mixing.

Similarly, the decay rate for $B^0(t) \rightarrow \bar{f}$ is given by

$$\Gamma(B^0(t) \rightarrow \bar{f}) = e^{-\Gamma t} \sum_{\lambda \leq \sigma} \left(\bar{\Lambda}_{\lambda\sigma} + \bar{\Sigma}_{\lambda\sigma} \cos(\Delta M t) - \bar{\rho}_{\lambda\sigma} \sin(\Delta M t) \right) g_\lambda g_\sigma. \quad (16)$$

The expressions for these another 18 observables, $\bar{\Lambda}_{\lambda\sigma}$, $\bar{\Sigma}_{\lambda\sigma}$ and $\bar{\rho}_{\lambda\sigma}$ are similar to those given in Eq. (15), with the replacements $A_\lambda \rightarrow \bar{A}'_\lambda$ and $A'_\lambda \rightarrow \bar{A}_\lambda$.

Angular analysis is more powerful than previously realized. Due to the interference between the different helicity states, there are enough independent measurement that one can obtain weak phase information as we now show. First, we note that

$$\Lambda_{\lambda\lambda} = \bar{\Lambda}_{\lambda\lambda} = \frac{(a_\lambda^2 + b_\lambda^2)}{2}, \quad \Sigma_{\lambda\lambda} = -\bar{\Sigma}_{\lambda\lambda} = \frac{(a_\lambda^2 - b_\lambda^2)}{2}. \quad (17)$$

Thus, one can determine the magnitudes of the amplitudes appearing in Eqs. (6)–(9), a_λ^2 and b_λ^2 . However, it must be stressed that *the knowledge of b_λ^2 will not be necessary within our method.*

Next, we have

$$\begin{aligned} \Lambda_{\perp i} &= -\bar{\Lambda}_{\perp i} = b_\perp b_i \sin(\delta_\perp - \delta_i + \Delta_i) - a_\perp a_i \sin(\Delta_i), \\ \Sigma_{\perp i} &= \bar{\Sigma}_{\perp i} = -b_\perp b_i \sin(\delta_\perp - \delta_i + \Delta_i) - a_\perp a_i \sin(\Delta_i), \end{aligned} \quad (18)$$

where $\Delta_i \equiv \delta_\perp^a - \delta_i^a$ and $\delta_\lambda \equiv \delta_\lambda^b - \delta_\lambda^a$. Using Eq. (18) one can solve for $a_\perp a_i \sin \Delta_i$. We will see that this is the only combination needed to cleanly extract weak phase information.

The coefficients of the $\sin(\Delta mt)$ term, which can be obtained in a time-dependent study, can be written as

$$\rho_{\lambda\lambda} = \pm a_\lambda b_\lambda \sin(\phi + \delta_\lambda), \quad \bar{\rho}_{\lambda\lambda} = \pm a_\lambda b_\lambda \sin(\phi - \delta_\lambda), \quad (19)$$

where the sign on the right hand side is positive for $\lambda = \parallel, 0$, and negative for $\lambda = \perp$. In the above, we have defined the CP phase $\phi \equiv -2\phi_M + \phi_b - \phi_a$. These quantities can be used to determine

$$2b_\lambda \cos \delta_\lambda = \pm \frac{\rho_{\lambda\lambda} + \bar{\rho}_{\lambda\lambda}}{a_\lambda \sin \phi}, \quad 2b_\lambda \sin \delta_\lambda = \pm \frac{\rho_{\lambda\lambda} - \bar{\rho}_{\lambda\lambda}}{a_\lambda \cos \phi}. \quad (20)$$

Similarly, the terms involving interference of different helicities are given as

$$\begin{aligned} \rho_{\perp i} &= -a_\perp b_i \cos(\phi + \delta_i - \Delta_i) - a_i b_\perp \cos(\phi + \delta_\perp + \Delta_i), \\ \bar{\rho}_{\perp i} &= -a_\perp b_i \cos(\phi - \delta_i + \Delta_i) - a_i b_\perp \cos(\phi - \delta_\perp - \Delta_i). \end{aligned} \quad (21)$$

Putting all the above information together, we are now in a position to extract the weak phase ϕ . Using Eq. (20), the expressions in Eq. (21) can be used to yield

$$\begin{aligned} \rho_{\perp i} + \bar{\rho}_{\perp i} &= -\cot \phi a_i a_\perp \cos \Delta_i \left[\frac{\rho_{ii} + \bar{\rho}_{ii}}{a_i^2} - \frac{\rho_{\perp\perp} + \bar{\rho}_{\perp\perp}}{a_\perp^2} \right] \\ &\quad - a_i a_\perp \sin \Delta_i \left[\frac{\rho_{ii} - \bar{\rho}_{ii}}{a_i^2} + \frac{\rho_{\perp\perp} - \bar{\rho}_{\perp\perp}}{a_\perp^2} \right], \end{aligned} \quad (22)$$

$$\begin{aligned} \rho_{\perp i} - \bar{\rho}_{\perp i} &= \tan \phi a_i a_\perp \cos \Delta_i \left[\frac{\rho_{ii} - \bar{\rho}_{ii}}{a_i^2} - \frac{\rho_{\perp\perp} - \bar{\rho}_{\perp\perp}}{a_\perp^2} \right] \\ &\quad - a_i a_\perp \sin \Delta_i \left[\frac{\rho_{ii} + \bar{\rho}_{ii}}{a_i^2} + \frac{\rho_{\perp\perp} + \bar{\rho}_{\perp\perp}}{a_\perp^2} \right]. \end{aligned} \quad (23)$$

In the above two equations: (i) $\rho_{\lambda\sigma}$ and $\bar{\rho}_{\lambda\sigma}$ are measured quantities, (ii) the a_λ^2 are determined from the relations in Eq. (17), and (iii) $a_i a_\perp \sin \Delta_i$ is obtained from Eq. (18). Thus, the above two equations involve only two unknown quantities, $\tan \phi$ and $a_i a_\perp \cos \Delta_i$, which can easily be solved for (up to a sign ambiguity in each of these quantities). Hence, $\tan^2 \phi$ (or, equivalently, $\sin^2 \phi$) can be obtained from the angular analysis.

Note that this method relies on the measurement of the interference terms between different helicities. However, we do not actually require that all three helicity components of the amplitude be used. In fact, one can use observables involving any two of largest helicity amplitudes. In the above description, one could have chosen ‘ $\parallel 0$ ’ instead of ‘ $\perp \parallel$ ’ or ‘ $\perp 0$ ’.

We now turn to specific applications of this method. Consider first the final states where, $f = \pm \bar{f}$. In this case, the parameters of Eqs. (6)–(9) satisfy $a_\lambda = b_\lambda$, $\delta_\lambda^a = \delta_\lambda^b$ (which implies that $\delta_\lambda = 0$), and $\phi_a = -\phi_b$ (so that $\phi \equiv -2\phi_M + 2\phi_b$). As described above, a_λ^2 can be obtained from Eq. (17). But now the measurement of $\rho_{\lambda\lambda}$ [Eq. (19)] directly yields $\sin \phi$. In fact, this is the conventional way [13] of using the angular analysis to measure the weak phases: each helicity state separately gives clean CP-phase information. Thus, for such states, nothing is gained by including the interference terms.

In order to have final states with only one weak amplitude, we consider states that do not receive penguin contributions. The only such Cabibbo-allowed quark-level decays are $\bar{b} \rightarrow \bar{c}u\bar{d}$, $\bar{u}c\bar{d}$. The meson level examples of these are $B_d^0/\bar{B}_d^0 \rightarrow D^{*-}\rho^+, D^{*+}\rho^-$. These are the VV counterparts of the PP/PV modes described in first example of section 1. As we have already emphasized in the discussion following Eq. (17), *none of the observables or combinations required for the analysis to extract $\sin^2(2\beta + \gamma)$ are proportional to b_λ^2 .* Thus, we avoid the practical problems present in Duni-etz’s method [3].

The two decay amplitudes for the final states $D^{*\pm}\rho^\mp$ have very different sizes, *i.e.* $b_\lambda \ll a_\lambda$. This results in a very small CP-violating asymmetry whose size is approximately $|V_{ub}V_{cd}^*/V_{cb}V_{ud}| \approx 2\%$. Thankfully, the situation is alleviated by the large branching ratio for the decay $B_d^0 \rightarrow D^{*-}\rho^+$, roughly 1%.

The Cabibbo-suppressed quark-level decays which do not receive penguin contributions are $\bar{b} \rightarrow \bar{c}u\bar{s}$, $\bar{u}c\bar{s}$, at meson level these would correspond to $B_d^0 \rightarrow \bar{D}^{*0}K^{*0}$, $D^{*0}K^{*0}$ and $\bar{B}_d^0 \rightarrow D^{*0}\bar{K}^{*0}$, $\bar{D}^{*0}\bar{K}^{*0}$, with K^{*0} and \bar{K}^{*0} decaying to $K_s\pi^0$. However, in these modes one has the old problem of tagging the neutral $D's$ [7].

One can also consider B_s^0 and \bar{B}_s^0 decays. corresponding to the quark-level decays $\bar{b} \rightarrow \bar{c}u\bar{d}$, $\bar{u}c\bar{d}$, or $\bar{b} \rightarrow \bar{c}u\bar{s}$, $\bar{u}c\bar{s}$. The most promising processes are the Cabibbo-suppressed

decay modes $B_s^0/\bar{B}_s^0 \rightarrow D_s^{*\pm} K^{*\mp}$. Here the $B_s^0 - \bar{B}_s^0$ mixing phase is almost 0, so that the quantity $\sin^2 \gamma$ can be extracted from the angular analysis of $B_s^0(t) \rightarrow D_s^{*\pm} K^{*\mp}$. Other methods for obtaining the CP phase γ using similar final states have also been studied [14].

4 Direct Asymmetry in $B \rightarrow VV$

It is also possible to cleanly extract the weak phase γ using only charged B^\pm decays, by studying the angular distribution [9]. The decays $B^+ \rightarrow D^{*0} V^+$, $B^+ \rightarrow \bar{D}^{*0} V^+$ and $B^- \rightarrow D^{*0} V^-$, $B^- \rightarrow \bar{D}^{*0} V^-$ can be related by CPT. Consider D^{*0}/\bar{D}^{*0} decaying into $D^0 \pi^0/\bar{D}^0 \pi^0$, with D^0/\bar{D}^0 meson further decaying to a final state ‘ f ’ that is common to both D^0 and \bar{D}^0 . f is chosen to be a Cabibbo-allowed mode of D^0 or a doubly-suppressed mode of \bar{D}^0 . The amplitudes for the decays of B^+ and B^- to a final state involving f will be a sum of the contributions from D^{*0} and \bar{D}^{*0} , and similarly for the CP-conjugate processes. In this case one can experimentally measure the magnitudes of the 12 helicity amplitudes, as well as the interference terms, leading to a total of 24 independent observables. However, there are just 15 unknowns involved in the amplitudes: $a_\lambda, b_\lambda, \delta_\lambda^a, \delta_\lambda^b, \gamma, \Delta$ and \mathcal{R} ; where, $\mathcal{R}^2 = \frac{Br(\bar{D}^0 \rightarrow f)}{Br(D^0 \rightarrow f)}$, and Δ is the strong phase difference between $D^0 \rightarrow f$ and $\bar{D}^0 \rightarrow \bar{f}$. Hence, the weak phase γ may be cleanly extracted.

In addition, this technique has some interesting features:

- The signals of CP violation can be obtained by adding B and \bar{B} events, i.e, flavour tagging is not required.
- The CP violation signals are not diluted by the sine of strong phases.

5 Conclusions

We have presented new methods, that use angular analysis in $B \rightarrow VV$ decay modes, to cleanly extract weak phases. Using modes which do not receive penguin contributions, the weak phases $(2\beta + \gamma)$ and γ can be determined. We have shown that the quantity $\sin^2(2\beta + \gamma)$ can be cleanly obtained from the time dependent angular analysis study of the decays $B_d^0(t) \rightarrow D^{*\pm} \rho^\mp$, $D^{*\pm} a_1^\mp$ etc. Similarly, $\sin^2 \gamma$ can be cleanly extracted from $B_s^0(t) \rightarrow D_s^{*\pm} K^{*\mp}$, or simply from an angular analysis of the decay mode $B^\pm \rightarrow \bar{D}^{*0} K^{*\pm}$. The large number of data samples already collected for the $B_d^0(t) \rightarrow D^{*-} \rho^+$ mode [15], may enable, $\sin^2(2\beta + \gamma)$ to be the second clean measurement, after $\sin 2\beta$, to be performed at B -factories.

Acknowledgements

N.S. and R.S. would like to thank Prof. Patricia Ball and the local organizers of the CKM Workshop for financial support. We thank the organizers for an exciting conference. The work of D.L. was financially supported by NSERC of Canada. The work of N.S. was supported by the Department of Science and Technology, India.

References

1. For a review, see H. Quinn and A. I. Sanda, *Eur. Phys. J.* **C15**, 626 (2000), in The Review of Particle Physics, (Particle Data Group), D.E. Groom *et. al.*, *Eur. Phys. J.* **C15**, 1 (2000).
2. D. London and R. Peccei, *Phys. Lett.* **223B**, 257 (1989); M. Gronau, *Phys. Rev. Lett.* **63**, 1451 (1989), *Phys. Lett.* **300B**, 163 (1993); B. Grinstein, *Phys. Lett.* **229B**, 280 (1989).
3. I. Dunietz, *Phys. Lett.* **427B**, 179 (1998).
4. R. Aleksan, I. Dunietz, B. Kayser and F. Le Diberder, *Nucl. Phys.* **B361**, 141 (1991).
5. David London, Nita Sinha & Rahul Sinha, *Phys. Rev. Lett.* **85**, 1807 (1990).
6. M. Gronau and D. Wyler, *Phys. Lett.* **265B**, 172 (1991); M. Gronau and D. London., *Phys. Lett. B* **253**, 483 (1991).
7. D. Atwood, I. Dunietz and A. Soni, *Phys. Rev. Lett.* **78**, 3257 (1997).
8. I. Dunietz, *Z. Phys.* **C56**, 129 (1992).
9. N. Sinha and R. Sinha, *Phys. Rev. Lett.* **80**, 3706 (1998).
10. D. Atwood and A. Soni, hep-ph/0206045; hep-ph/0304085; R. Aleksan, T. C. Petersen and A. Soffer, *Phys. Rev. D* **67**, 096002 (2003); A. Giri, Y. Grossman, A. Soffer and J. Zupan, hep-ph/0303187; Y. Grossman, Z. Ligeti and A. Soffer, *Phys. Rev. D* **67**, 071301 (2003).
11. G. Valencia, *Phys. Rev. D* **39**, 3389 (1989); G. Kramer and W. F. Palmer, *Phys. Rev. D.* **45**, 193 (1992).
12. A. S. Dighe I. Dunietz and R. Fleischer, *Eur. Phys. J.* **C6**, 647 (1999).
13. I. Dunietz, H.R. Quinn, A. Snyder, W. Toki and H.J. Lipkin, *Phys. Rev.* **D43**, 2193 (1991).
14. R. Aleksan, I. Dunietz and B. Kayser, *Zeit. Phys.* **C54**, 653 (1992); I. Dunietz and R. Fleischer, *Phys. Lett.* **387B**, 361 (1996).
15. B. Aubert *et. al.*, BABAR Collaboration, *Phys. Rev.* **D67** (2003) 091101.