The anomalous Higgs-top couplings in the MSSM

Tai-Fu Feng^a Xue-Qian $\mathrm{Li}^{b,c}$ Jukka Maalampi a

^aDepartment of Physics, 40014 University of Jyväskylä, Finland
 ^b CCAST (World Laboratory), P.O. Box 8730, Beijing 100080, China
 ^cDepartment of Physics, Nankai University, Tianjin 300070, China

Abstract

The anomalous couplings of the top quark and the Higgs boson have been studied in an effective theory deduced from the minimal supersymmetric extension of the standard model (MSSM) as the heavy fields are integrated out. Constraints on the parameters of the model from the experimental data of $R_b = \Gamma(Z \to b\bar{b})/\Gamma(Z \to hadrons)$ are obtained.

PACS numbers: 12.38.Bx, 12.60.Jv, 14.65.Ha, 14.80.Cp

1 Introduction

The standard model (SM) has been very successfull phenomenologically, but nevertheless it should be considered just as an effective theory valid for physics at the electroweak (EW) scale. In higher-energy regimes new physics beyond the SM must exist. Irrespective to what this new physics might be, it should be able to give a satisfactory answer to the most fundamental open question of the electroweak physics, that is, it must explain the origin of the electroweak gauge symmetry breaking [1, 2]. In the SM this is arranged through the spontaneous symmetry breaking mechanism by introducing a doublet of scalars with a nonzero vacuum expectation value (VEV). This mechanism, despite its simplisity and economy, has well known problems, which has enforced theorists and experimenters to look for new physics beyond the SM. Among the possible ways of extending the SM, supersymmetry is considered as a particularly attractive one. The minimal supersymmetry extension of the standard model (MSSM) provides an appealing solution to the gauge hierarchy problem by guaranteeing the perturbative stability of the theory from the electroweak scale to the Planck scale.

The MSSM contains two complex Higgs doublets, denoted by H_u , H_d and assigned with opposite hypercharges $Y_B(H_u) = -Y_B(H_d) = 1$. There are altogether four neutral scalar degrees of freedom, three of which correspond to physical scalar fields. In the case where CP is conserved one can define two CP-even neutral Higgs fields, H, h, and one CP-odd neutral Higgs field, A. The present experimental bounds

on the Higgs boson masses set strong restrictions on the parameter space of the CP-conserving MSSM [3]. Radiative corrections to the lightest CP-even Higgs boson mass have been computed by using the renormalization group equation (RGE) method and diagram technique [4], and the resulting upper bound is 135 GeV, which is not much above the present experimental lower bound of 95GeV (95% CL).

The possibility of CP violation makes the situation drastically different. There are three main sources of the CP violation in the MSSM Lagrangian. The first one is the well known μ parameter of the superpotential, which is in general complex. The second source is constituted by the soft mass terms of the $SU(3) \times SU(2) \times U(1)$ gauginos. The third source are the phases of the soft supersymmetry-breaking mass terms of scalar fermions and of the soft trilinear couplings, which are presented by the matrices $\mathbf{m}_{Q,U,D,L,R}^2$ and $\mathbf{A}_{U,D,E}$, respectively. Actually, only the off-diagonal elements of the soft mass matrices can be complex due to the hermiticity of these matrices. The matrices $\mathbf{A}_{U,D,E}$ in contrast can have complex phases also in their diagonal elements [5]. Not all the phases of these soft SUSY breaking parameters are physical and lead to the violation of CP parity. The physical CP phases are restricted by experimental observations, the most rigorous constraints originating from the measurements of the electron and neutron electric dipole moments (edm). The present upper limits for these edms are $d_e < 4.3 \times 10^{-27} \mathrm{e} \cdot \mathrm{cm}$ [6] and $d_n < 6.5 \times 10^{-26} \mathrm{e} \cdot \mathrm{cm}$ [7], respectively. Also the edm of H_g^{199} is quite accurately measured, the present upper limit being $d_{H_g^{199}} < 9. \times 10^{-28} \mathrm{e} \cdot \mathrm{cm}$ [8].

It has been demonstrated that the MSSM can be consistent with these constraints in some regions of the parameter space when suitable cancellations between different contributions occur [9] or when CP violation effects are associated with the third generation of squarks only [10]. The mixing of neutral Higgs bosons in the latter scenario is analyzed in [11, 12, 13, 14]. It is found that the CP-violating phases and large Yukawa couplings of the third generation fermions can lead to large mixings among the neutral Higgs bosons as a consequence of radiative effects. These mixings can drastically change the couplings between the neutral Higgs bosons and gauge bosons, as well as the self-couplings of the Higgs fields. One consequence of this is that the experimental lower bound on the lightest neutral Higgs mass is relaxed to 60 GeV, while the predicted upper limit for the lightest Higgs boson mass remains about 135 GeV.

If the new physics scale is much higher than the EW scale, one would have at the EW scale a great number of higher-dimensional operators \mathcal{O}_i ($dim(\mathcal{O}_i) > 4$) induced by the beyond-the-SM physics [15, 16, 17, 18]. The resulting effective Lagrangian is of the general form

$$\mathcal{L}_{eff} = \mathcal{L}_0 + \frac{1}{\mu_{NP}^2} \sum_i C_i \mathcal{O}_i + \mathcal{O}(\frac{1}{\mu_{NP}^4}) . \tag{1}$$

Here \mathcal{L}_0 is the SM Lagrangian, C_i are Wilson coefficients, and μ_{NP} the energy scale of new physics. The Wilson coefficients are in general dependent on the new energy scale, but in addition to this all the higher-dimensional operators in \mathcal{L}_{eff} have a common suppression factor $1/\mu_{NP}^2$.

In this paper we shall study anomalous couplings (to use the terminology of [19, 20]), i.e. the couplings not present in the SM Lagrangian \mathcal{L}_0 , between the lightest neutral Higgs scalar (h) and the top quark induced by the new physics of MSSM. We assume that the other Higgs bosons, as well as all supersymmetric particles, are much heavier than the lightest neutral Higgs particle, so that the corresponding fields can be integrated out. A well known fact is that the masses of the other two neutral Higgs bosons are approximately equal to that of the charged Higgs boson (H^+) under the condition $m_{H^+} \gg m_h$, and hence one can consider the lighter Higgs doublet as the SM Higgs field and integrate out the heavier Higgs doublet.

Our presentation is organized as follows. In Section II, the notations adopted in this work are introduced. In Section III we shall describe the method of obtaining the Wilson coefficients by integrating out the heavy degrees of freedom in the full theory. The numerical analysis of the constraints on the parameter space from the present experiments, especially by the R_b data, is given in Section IV. Section V summarizes

our results. Some lengthy formulae, such as the expressions for the Wilson coefficients and the loop integral functions, are collected in Appendices.

2 Preliminaries

The most general gauge invariant superpotential, which retains all the conservation laws of the SM, is given by

$$\mathcal{W} = \mu \epsilon_{ij} \hat{H}_u^i \hat{H}_d^j + \epsilon_{ij} h_L^{JI} \hat{H}_d^i \hat{L}_j^{II} \hat{R}^J + \epsilon_{ij} h_D^{JI} \hat{H}_d^i \hat{Q}_j^{II} \hat{D}^J + \epsilon_{ij} h_U^{JI} \hat{H}_u^i \hat{Q}_j^{II} \hat{U}^J. \tag{2}$$

Here \hat{H}_u , \hat{H}_d are the two Higgs superfield doublets, \hat{Q}^I and \hat{L}^I are the doublets of quark and lepton superfields, and \hat{U}^I , \hat{D}^I and \hat{R}^I are the singlet superfields of u- and d-type quarks and charged leptons, respectively (I=1, 2, 3 is generation index, i,j=1,2 are SU(2) indices). Yukawa coupling constants are denoted by h_L , $h_{U,D}$. The breaking of supersymmetry happens through the so-called soft terms, which are in the most general case given by

$$\mathcal{L}_{soft} = -m_{H_{u}}^{2} H_{u}^{i*} H_{u}^{i} - m_{H_{d}}^{2} H_{d}^{i*} H_{d}^{i} - m_{L^{IJ}}^{2} \tilde{L}_{i}^{I*} \tilde{L}_{i}^{J} - m_{R^{IJ}}^{2} \tilde{R}^{I*} \tilde{R}^{J} - m_{Q^{IJ}}^{2} \tilde{Q}_{i}^{I*} \tilde{Q}_{i}^{J} - m_{U^{IJ}}^{2} \tilde{U}^{I*} \tilde{U}^{J}
- m_{D^{IJ}}^{2} \tilde{D}^{I*} \tilde{D}^{J} + (m_{1} \lambda_{B} \lambda_{B} + m_{2} \lambda_{A}^{i} \lambda_{A}^{i} + m_{3} \lambda_{G}^{a} \lambda_{G}^{a} + h.c.) + \left[\epsilon_{ij} m_{H_{12}}^{2} H_{u}^{i} H_{d}^{j} + \epsilon_{ij} A_{L}^{JI} H_{d}^{i} \tilde{L}_{j}^{I} \tilde{R}^{J} \right]
+ \epsilon_{ij} A_{D}^{JI} H_{d}^{i} \tilde{Q}_{j}^{I} \tilde{D}^{J} + \epsilon_{ij} A_{U}^{JI} H_{u}^{i} \tilde{Q}_{j}^{I} \tilde{U}^{J} + h.c. \right].$$
(3)

Here λ_G^a $(a=1,\ 2,\ \cdots\ 8),\ \lambda_A^i$ $(i=1,\ 2,\ 3)$ and λ_B denote the $SU(3),\ SU(2)$ and U(1) gauginos, respectively, and $A_{U,D,L}^{IJ}$ are coupling constants of the unit of mass.

Let us define scalar doublets Φ and Φ_H as follows:

$$\begin{pmatrix} \Phi \\ \Phi_H \end{pmatrix} = \begin{pmatrix} c_{\beta} & -s_{\beta} \\ s_{\beta} & c_{\beta} \end{pmatrix} \begin{pmatrix} \tilde{H}_d \\ H_u \end{pmatrix} , \tag{4}$$

where $\tilde{H}_d = i\sigma_2 H_d^*$ and $c_\beta = \cos\beta$, $s_\beta = \sin\beta$ with $\tan\beta = v_u/v_d$, the ratio of the VEVs of H_u , H_d . With this definition Φ is identified as the SM Higgs doublet, consisting of Goldstone bosons and a physical neutral Higgs field. More explicitly, one can write the two Higgs doublets as

$$\Phi = \begin{pmatrix} G^{+} \\ \frac{1}{\sqrt{2}} \left(v + H_{1}^{0} + iG^{0} \right) \end{pmatrix} , \quad \Phi_{H} = \begin{pmatrix} H^{+} \\ \frac{1}{\sqrt{2}} \left(H_{2}^{0} + iA \right) \end{pmatrix} , \tag{5}$$

where G^0 , G^+ denote the Goldstone bosons, H_1^0 and H_2^0 are the neutral Higgs fields, H^+ and A are the physical charged Higgs and CP-odd neutral Higgs bosons, respectively, and $v = \sqrt{v_u^2 + v_d^2} = 246$ GeV. At the electro-weak scale, the two physical CP-even neutral Higgs fields are obtained through the mixing between the fields H_1^0 and H_2^0 . The masses of the physical Higgs bosons are given by

$$\mathbf{m}_{even}^{2} = \begin{pmatrix} m_{\rm Z}^{2} \left(s_{\beta}^{2} - c_{\beta}^{2}\right)^{2} & 2m_{\rm Z}^{2} s_{\beta} c_{\beta} \left(c_{\beta}^{2} - s_{\beta}^{2}\right) \\ 2m_{\rm Z}^{2} s_{\beta} c_{\beta} \left(c_{\beta}^{2} - s_{\beta}^{2}\right) & \frac{m_{H_{12}}^{2}}{s_{\beta} c_{\beta}} + 4m_{\rm Z}^{2} s_{\beta}^{2} c_{\beta}^{2} \end{pmatrix} \quad \text{(in the basis } (\mathbf{H}_{1}^{0}, \mathbf{H}_{2}^{0})^{\mathrm{T}}),$$

$$m_{A}^{2} = \frac{m_{H_{12}}^{2}}{s_{\beta} c_{\beta}},$$

$$m_{H^{+}} = \frac{m_{H_{12}}^{2}}{s_{\beta} c_{\beta}} + m_{\mathrm{W}}^{2}. \quad (6)$$

In the limit $m_H^2 \gg m_{\rm W}^2$ the two doublets Φ and Φ_H decouple, the former remaining light and the latter being associated with a large mass $m_H^2 = m_{H_{12}}^2/s_\beta c_\beta$. In the following, we will use the four-component spinor representation for fermions. From the two-

In the following, we will use the four-component spinor representation for fermions. From the two-component Wyel spinors ψ_{Q_I} , ψ_{U_I} , ψ_{D_I} , ψ_{H_u} and ψ_{H_d} , we form the following four-component Dirac fermions:

$$q_{L}^{I} = \begin{pmatrix} \psi_{Q_{I}} \\ 0 \end{pmatrix}, \quad u_{R}^{I} = \begin{pmatrix} 0 \\ \bar{\psi}_{U_{I}} \end{pmatrix},$$

$$d_{R}^{I} = \begin{pmatrix} 0 \\ \bar{\psi}_{D_{I}} \end{pmatrix}, \quad \psi_{H} = \begin{pmatrix} \frac{\psi_{H_{u}}}{\tilde{\psi}_{H_{d}}} \end{pmatrix},$$

$$(7)$$

with $\tilde{\psi}_{H_d} = (i\sigma^2)\psi_{H_d}$. Similarly, for the $SU(3)\times SU(2)\times U(1)$ gauginos λ_G^a , λ_A^i , λ_B can we define the following four-component Majorana spinors:

$$\psi_G^a = \begin{pmatrix} i\lambda_G^a \\ -i\bar{\lambda}_G^a \end{pmatrix} , \quad \psi_A^i = \begin{pmatrix} i\lambda_A^i \\ -i\bar{\lambda}_A^i \end{pmatrix}$$

$$\psi_B = \begin{pmatrix} i\lambda_B \\ -i\bar{\lambda}_B \end{pmatrix} . \tag{8}$$

Diagonalizing the soft mass terms is done with help of the sfermion mixing matrices $\mathcal{Z}_{O,U,D}$ defined as

$$\mathcal{Z}_{Q}^{\dagger} \mathbf{m}_{Q}^{2} \mathcal{Z}_{Q} = \hat{\mathbf{m}}_{Q}^{2} ,
\mathcal{Z}_{U}^{\dagger} \mathbf{m}_{U}^{2} \mathcal{Z}_{U} = \hat{\mathbf{m}}_{U}^{2} ,
\mathcal{Z}_{D}^{\dagger} \mathbf{m}_{D}^{2} \mathcal{Z}_{D} = \hat{\mathbf{m}}_{D}^{2} ,$$
(9)

where the matrices $\hat{\mathbf{m}}_{Q,U,D}^2$ on the right-handed side are diagonal.

Finally, we will benefit in our calculations from the following rearrangement identities of the SU(2) group indices:

$$\mathbf{1}_{\alpha\alpha'} \otimes \mathbf{1}_{\beta\beta'} = \frac{1}{2} \left\{ \mathbf{1}_{\alpha\beta} \otimes \mathbf{1}_{\beta'\alpha'} + \sum_{a} \sigma_{\alpha\beta}^{a} \otimes \sigma_{\beta'\alpha'}^{a} \right\},$$

$$\sigma_{\alpha\alpha'}^{a} \otimes \sigma_{\beta\beta'}^{a} = \frac{1}{2} \left\{ 3\mathbf{1}_{\alpha\beta} \otimes \mathbf{1}_{\beta'\alpha'} - \sum_{a} \sigma_{\alpha\beta}^{a} \otimes \sigma_{\beta'\alpha'}^{a} \right\},$$

$$\sum_{a,b} \epsilon_{abc} \epsilon_{abd} = 2\delta_{cd}.$$
(10)

3 The Higgs-top anomalous couplings

In this section we shall discuss the anomalous couplings of the top quark and Higgs bosons. Considering the suppression of the new physics energy scale, we just keep operators up to dimension-six in the effective Lagrangian Eq. (1). The top-Higgs anomalous couplings of interest can be classified into three types: the anomalous couplings involving a left-handed quark, the right-handed top quark and Higgs boson ($\mathcal{O}_{tq\Phi}$), the couplings between the Higgs boson and a left-handed quark ($\mathcal{O}_{q\Phi}$), and the couplings between the Higgs boson and the right-handed top quark ($\mathcal{O}_{t\Phi}$). After the EW symmetry breaking, these operators produce

not only corrections to the effective couplings $Wt\bar{b}$, $Xt\bar{t}$, $Xb\bar{b}$ ($X=\gamma,\ Z,\ H$), but also induce anomalous couplings such as $\gamma Ht\bar{t}$. All those effects may be detectable at the Next Linear Collider (NLC) and at the Tevatron.

In the following subsections we will give the explicit expressions for the contributions of supersymmetric particles and the heavy Higgs boson doublet to the effective operators mentioned above by deriving the relevant Wilson coefficients. We will give our results in terms of the following loop integral functions:

$$\mathcal{B}_{j,k}^{i}(x_{a}, x_{b}) = \frac{(4\pi)^{2}}{i} \int \frac{d^{4}q}{(2\pi)^{4}} \frac{\left(q^{2}\right)^{i}}{(q^{2} - x_{a})^{j}(q^{2} - x_{b})^{k}} ,$$

$$\mathcal{C}_{jkl}^{i}(x_{a}, x_{b}, x_{c}) = \frac{(4\pi)^{2}}{i} \int \frac{d^{4}q}{(2\pi)^{4}} \frac{\left(q^{2}\right)^{i}}{(q^{2} - x_{a})^{j}(q^{2} - x_{b})^{k}(q^{2} - x_{c})^{l}} ,$$

$$\mathcal{D}_{jklm}^{i}(x_{a}, x_{b}, x_{c}, x_{d}) = \frac{(4\pi)^{2}}{i} \int \frac{d^{4}q}{(2\pi)^{4}} \frac{\left(q^{2}\right)^{i}}{(q^{2} - x_{a})^{j}(q^{2} - x_{c})^{k}(q^{2} - x_{c})^{l}(q^{2} - x_{d})^{m}} .$$
(11)

The explicit expressions of these are given in Ref. [21].

3.1 The anomalous couplings $\mathcal{O}_{_{tq\Phi}}$

This class of operators includes the CP-even operators

$$\mathcal{O}_{tq\Phi1} = \left(\Phi^{\dagger}\Phi\right) \left(\bar{q}_{L} t_{R} \tilde{\Phi} + \tilde{\Phi}^{\dagger} \bar{t}_{R} q_{L}\right),
\mathcal{O}_{tq\Phi2} = \bar{q}_{L} \left(D_{\mu} t_{R}\right) D^{\mu} \tilde{\Phi} + \left(D_{\mu} \tilde{\Phi}\right)^{\dagger} \overline{\left(D^{\mu} t_{R}\right)} q_{L},
\mathcal{O}_{tq\Phi3} = \overline{\left(D_{\mu} q_{L}\right)} \left(D^{\mu} t_{R}\right) \tilde{\Phi} + \tilde{\Phi}^{\dagger} \overline{\left(D_{\mu} t_{R}\right)} \left(D^{\mu} q_{L}\right),
\mathcal{O}_{tq\Phi4} = \overline{\left(D_{\mu} q_{L}\right)} t_{R} \left(D^{\mu} \tilde{\Phi}\right) + \left(D_{\mu} \tilde{\Phi}\right)^{\dagger} \bar{t}_{R} \left(D^{\mu} q_{L}\right),
\mathcal{O}_{tq\Phi5} = i \overline{\left(D_{\mu} q_{L}\right)} \sigma^{\mu\nu} t_{R} \left(D_{\nu} \tilde{\Phi}\right) + i \left(D_{\nu} \tilde{\Phi}\right)^{\dagger} \bar{t}_{R} \sigma^{\mu\nu} \left(D_{\mu} q_{L}\right),$$
(12)

where the covariant derivative D_{μ} is given by $D_{\mu} = \partial_{\mu} - \frac{i}{2}g_3 T^A G_{\mu}^A - \frac{i}{2}g_2 \sigma^a W_{\mu}^a - \frac{i}{2}g_1 Y_B B_{\mu}$. The CP-odd counterparts of these operators are

$$\mathcal{O}_{tq\Phi6} = \left(\Phi^{\dagger}\Phi\right) \left(\bar{q}_{L} t_{R} \tilde{\Phi} - \tilde{\Phi}^{\dagger} \bar{t}_{R} q_{L}\right),$$

$$\mathcal{O}_{tq\Phi7} = \bar{q}_{L} \left(D_{\mu} t_{R}\right) D^{\mu} \tilde{\Phi} - \left(D_{\mu} \tilde{\Phi}\right)^{\dagger} \overline{\left(D^{\mu} t_{R}\right)} q_{L},$$

$$\mathcal{O}_{tq\Phi8} = \overline{\left(D_{\mu} q_{L}\right)} \left(D^{\mu} t_{R}\right) \tilde{\Phi} - \tilde{\Phi}^{\dagger} \overline{\left(D_{\mu} t_{R}\right)} \left(D^{\mu} q_{L}\right),$$

$$\mathcal{O}_{tq\Phi9} = \overline{\left(D_{\mu} q_{L}\right)} t_{R} \left(D^{\mu} \tilde{\Phi}\right) - \left(D_{\mu} \tilde{\Phi}\right)^{\dagger} \bar{t}_{R} \left(D^{\mu} q_{L}\right),$$

$$\mathcal{O}_{tq\Phi10} = i \overline{\left(D_{\mu} q_{L}\right)} \sigma^{\mu\nu} t_{R} \left(D_{\nu} \tilde{\Phi}\right) - i \left(D_{\nu} \tilde{\Phi}\right)^{\dagger} \bar{t}_{R} \sigma^{\mu\nu} \left(D_{\mu} q_{L}\right).$$
(13)

For the CP-even operator $\mathcal{O}_{tq\Phi 1}$ and for the corresponding CP-odd operator $\mathcal{O}_{tq\Phi 6}$, nonzero contributions to the Wilson coefficients originate from the Feynman diagrams shown in Fig. 1, and they are given by

$$C_{tq\Phi 1} = \frac{1}{2} \left(g_1^2 + g_2^2 \right) \mathbf{Re}(\mathbf{h}_U^{33}) \frac{s_{\beta} c_{\beta}^2 (c_{\beta}^2 - s_{\beta}^2)}{x_H} ,$$

$$C_{tq\Phi 6} = i \frac{1}{2} \left(g_1^2 + g_2^2 \right) \mathbf{Im}(\mathbf{h}_U^{33}) \frac{s_{\beta} c_{\beta}^2 (c_{\beta}^2 - s_{\beta}^2)}{x_H} ,$$
(14)

where $x_H = m_H^2/\mu_{NP}^2$. In the full theory, the Feynman diagrams that induce the anomalous couplings $\mathcal{O}_{tq\Phi 1}$, $\mathcal{O}_{tq\Phi 6}$ should also include diagrams involving virtual SM fields. However, these diagrams have no contribution to the Wilson coefficients after matching the effective Lagrangian Eq. (1) with the Lagrangian of the full theory (MSSM) (see below for more details).

For the CP-even anomalous operators $\mathcal{O}_{tq\Phi i}$ (i=2,3,4,5) and the CP-odd anomalous operators $\mathcal{O}_{tq\Phi i}$ (i=7,8,9,10), the derivation of the Wilson coefficients leads to relatively tedious calculations. In Fig. 2 we show the Feynman diagrams, which induce nontrival contributions to the Wilson coefficients after matching the amplitude of the effective theory with that of the MSSM. In these diagrams, the black blobs represent the self-energy diagrams of $\bar{q}_L q_L$, $\bar{t}_R t_R$, and $\Phi^{\dagger}\Phi$ (Φ_H) (Fig. 4).

The matching procedure, to which we refer above, is extensively applied in the derivation of the effective Lagrangian in the hadron physics, especially in the application of the effective Lagrangian to the rare B decay [22]. The main idea of this procedure is the following. We derive the amplitude corresponding to the relevant Feynman diagrams both in the full theory and in the effective theory. In both derivations we only keep the momenta p_i of external particles to the second order. Through a comparison of the amplitudes of the full theory and the effective theory we then obtain the Wilson coefficients of interest.

For a demonstration, let us consider the first diagram of Fig. 2. In the full theory we can write the amplitude corresponding to this diagram as

$$A_{2(1)}^{FT}(p,q) = -\frac{i}{(4\pi)^2} s_{\beta} c_{\beta}^2 \left(\mathbf{h}_D^{\dagger} \mathbf{h}_D \mathbf{h}_U^{\dagger}\right)_{33} \left\{ \left[\Delta + 1 + \ln \frac{\mu_{NP}^2}{m_H^2} \right] \left(\bar{q}_L \tilde{\Phi}\right) t_R - \frac{1}{2m_H^2} \left(1 + \ln \frac{m_q^2}{m_H^2}\right) \left(\bar{q}_L \tilde{\Phi}\right) q^2 t_R + \frac{1}{2m_H^2} \left(\bar{q}_L \tilde{\Phi}\right) (p+q)^2 t_R - \frac{1}{2m_H^2} \left(\bar{q}_L \tilde{\Phi}\right) q \cdot (p+q) t_R + \frac{1}{4m_H^2} \left(\bar{q}_L \tilde{\Phi}\right) [\not p, \not q] t_R \right\}.$$

$$(15)$$

Here $\Delta = \frac{1}{\epsilon} - \gamma_E + \ln 4\pi$ denotes the ultraviolet divergence $(D = 4 - 2\epsilon)$ is the time-space dimension in the dimensional regularization scheme), μ_{NP} is the scale of new physics, and p and q denote the four-momenta of the external particles t_R and Φ , respectively. In the full theory the light fields and the heavy fields co-exist in the Lagrangian. When the heavy degrees of freedom are integrated out and the light fields are treated as massless, infrared divergences are encountered. They are regulated by the parameter m_q .

The amplitude of the corresponding Feynman diagram in the effective theory, presented in Fig. 3, is given by

$$A_{2(1)}^{ET}(p,q) = -\frac{i}{2(4\pi)^2 m_H^2} s_\beta c_\beta^2 \left(\mathbf{h}_D^\dagger \mathbf{h}_D \mathbf{h}_U^\dagger\right)_{33} \left(\Delta - \frac{1}{2} + \ln \frac{\mu_{NP}^2}{m_g^2}\right) \left(\bar{q}_L \tilde{\Phi}\right) q^2 t_R \ . \tag{16}$$

In the operators of the effective theory, only the light fields exist, and the Wilson coefficients do not depend on their masses. As in the full theory, an infrared divergence emerges here, and it is also regularized by the parameter m_q . As expected, the infrared divergence appearing in the effective theory is the same as that appearing in the full-theory. By matching the amplitudes Eq. (16) and Eq. (15), one gets rid of the infrared divergence. After this matching step, we can present the amplitude in its final form:

$$A_{2(1)}(p,q) = -\frac{i}{(4\pi)^2} s_{\beta} c_{\beta}^2 \left(\mathbf{h}_{D}^{\dagger} \mathbf{h}_{D} \mathbf{h}_{U}^{\dagger}\right)_{33} \left\{ \left[\Delta + 1 + \ln \frac{\mu_{NP}^2}{m_H^2} \right] \left(\bar{q}_L \tilde{\Phi} \right) t_R + \frac{1}{2m_H^2} \left(1 + \ln \frac{\mu_{NP}^2}{m_H^2} \right) \left(\bar{q}_L \tilde{\Phi} \right) p \cdot q t_R \right.$$

$$\left. - \frac{1}{2m_H^2} \left(1 + \ln \frac{\mu_{NP}^2}{m_H^2} \right) \left(\bar{q}_L \tilde{\Phi} \right) q \cdot (p+q) t_R + \frac{1}{2m_H^2} \left(\bar{q}_L \tilde{\Phi} \right) p \cdot (p+q) t_R \right.$$

$$\left. - \frac{1}{4m_H^2} \left(\bar{q}_L \tilde{\Phi} \right) [\mathbf{q}, \mathbf{p}] t_R \right\}. \tag{17}$$

The first term in the parenthesis of Eq. (17) contributes to the renormalization of the Yukawa couplings \mathbf{h}_U , and it is irrelevant to our present discussion, taking into account the approximation level we work on. For those diagrams where the inner lines are supersymmetry particles, the Wilson coefficients of the anomalous couplings can be directly read from the amplitudes, because we integrate out all the supersymmetry fields in the effective theory.

Now we will turn to show how to obtain the contributions of the self-energy diagrams to the anomalous couplings. As mentioned above, there are three possible self-energy diagrams that contribute to the coefficients indirectly, namely the self-energy corrections to the $\bar{q}_L q_L$, $\bar{t}_R t_R$, and Higgs doublet currents. For a fermion, the renormalized fields are defined by

$$f_{L,i}^{0} = Z_{L,ij}^{\frac{1}{2}} f_{L,j} ,$$

$$f_{R,i}^{0} = Z_{R,ij}^{\frac{1}{2}} f_{R,j} ,$$
(18)

where i,j are generation indices, $f_{L,i}^0$, $f_{R,i}^0$ are the left- and right-handed bare fields, respectively, $f_{L,i}$, $f_{R,i}$ are the corresponding renormalized fields, and $Z_{L,R}$ are the wave function renormalization constants. Ignoring the fermion masses, we can write down the counter terms for the fermions in Eq. (18) as follows:

$$\Sigma_{ij}^{L,c}(p) = \left(Z_{L,iI}^{\dagger \frac{1}{2}} Z_{L,Ij}^{\frac{1}{2}} - \delta_{ij}\right) \not p = \frac{1}{2} \left(\delta Z_{L,ij}^{\dagger} + \delta Z_{L,ij}\right) \not p ,$$

$$\Sigma_{ij}^{R,c}(p) = \left(Z_{R,iI}^{\dagger \frac{1}{2}} Z_{R,Ij}^{\frac{1}{2}} - \delta_{ij}\right) \not p = \frac{1}{2} \left(\delta Z_{R,ij}^{\dagger} + \delta Z_{R,ij}\right) \not p ,$$
(19)

where p denotes the external momentum of the fermion. In the full theory, we express the bare self-energy of the fermions as

$$\Sigma_{ij}^{L,0} = \left[\delta_{ij} + A_{ij}^L + B_{ij}^L p^2 \right] \not p ,$$

$$\Sigma_{ij}^{R,0} = \left[\delta_{ij} + A_{ij}^R + B_{ij}^R p^2 \right] \not p$$
(20)

where the first term δ_{ij} represents the Born approximation part and $A^{L,R}$, $B^{L,R}$ originate from raditive corrections. From Eq. (19) and Eq. (20) one finds the following form for the renormalized self-energies:

$$\hat{\Sigma}_{ij}^{L} = \left[\delta_{ij} + \frac{1}{2} \left(\delta Z_{L,ij}^{\dagger} + \delta Z_{L,ij} \right) + A_{ij}^{L} + B_{ij}^{L} p^{2} \right] \not p ,
\hat{\Sigma}_{ij}^{R} = \left[\delta_{ij} + \frac{1}{2} \left(\delta Z_{R,ij}^{\dagger} + \delta Z_{R,ij} \right) + A_{ij}^{R} + B_{ij}^{R} p^{2} \right] \not p .$$
(21)

The explicit expressions of the renormalization constant $\delta Z_{L,R}$ depend upon the renormalization scheme, i.e., the renormalization conditions. Instead of the often-used renormalization schemes, i.e. the minimal

subtraction scheme (MS) or the modified minimal subtraction scheme (\overline{MS}) , we adopt here the physical renormalization conditions

$$\hat{\Sigma}_{ij}^{L} f_{L,j}|_{\not p=0} = 0 ,$$

$$\hat{\Sigma}_{ij}^{R} f_{R,j}|_{\not p=0} = 0 ,$$

$$\frac{1}{\not p} \hat{\Sigma}_{ij}^{L} f_{L,j}|_{\not p=0} = f_{L,i} ,$$

$$\frac{1}{\not p} \hat{\Sigma}_{ij}^{R} f_{R,j}|_{\not p=0} = f_{R,i} .$$
(22)

The first two conditions mean that the renormalized fields satisfy the equations of motion of free particles (for massless fermions this is a trivial constraint), and the last two conditions set the residue of the propagators at the pole equal to unity. In fact, this scheme is just the on-shell renormalization scheme often used when calculating radiative corrections to electroweak processes [23]. Of course, for high energy processes we can ignore the fermion mass in our approximation. Using the condition Eq. (22), we achieve the renormalized fermion self-energies:

$$\hat{\Sigma}_{ij}^{L} = B_{ij}^{L} p^{2} \not p ,$$

$$\hat{\Sigma}_{ij}^{R} = B_{ij}^{R} p^{2} \not p .$$
(23)

We can attribute those terms to the contributions of the high dimension operators $\bar{q}_L(\mathcal{D})^3 q_L$, $\bar{t}_R(\mathcal{D})^3 t_R$. After the matching of the full and effective theories, there is no contribution to the operators of our interesting given in Eq. (12) and Eq. (13) from the fermion self-energy diagrams.

For the Higgs sector, the bare self-energies are given as

$$\begin{split} \Sigma^0_{\Phi\Phi}(p^2) &= D_{\Phi\Phi} + \left(1 + E_{\Phi\Phi}\right) p^2 + F_{\Phi\Phi} p^4 \;, \\ \Sigma^0_{\Phi\Phi_H}(p^2) &= D_{\Phi\Phi_H} + E_{\Phi\Phi_H} p^2 + F_{\Phi\Phi_H} p^4 \;, \end{split} \tag{24}$$

where p denotes the momentum of the external particle. In Eq. (24), D, E, F are standard integral functions that appear in radiative corrections. For the renormalization of the Higgs boson wave function and mass, we request the renormalized boson self-energy to satisfy the conditions

$$\begin{split} \hat{\Sigma}_{\Phi\Phi}(p^2)|_{p^2=0} &= 0 ,\\ \frac{1}{p^2} \hat{\Sigma}_{\Phi\Phi}(p^2)|_{p^2=0} &= 0 ,\\ \hat{\Sigma}_{\Phi\Phi_H}(p^2)|_{p^2=0} &= 0 ,\\ \hat{\Sigma}_{\Phi\Phi_H}(p^2)|_{p^2=m_H^2} &= 0 . \end{split} \tag{25}$$

It is easy to find the renormalized Higgs field self-energies which meet the conditions of Eq. (25):

$$\hat{\Sigma}_{\Phi\Phi}(p^2) = F_{\Phi\Phi}p^4 ,$$

$$\hat{\Sigma}_{\Phi\Phi_H}(p^2) = F_{\Phi\Phi_H}p^2(p^2 - m_H^2) .$$
(26)

The function $F_{\Phi\Phi}$ is attributed to the contribution of the high dimensional operator $\Phi^{\dagger} \left(D_{\mu} D^{\mu} \right)^2 \Phi$. After the matching procedure, this piece will not contribute to the operators in (12) and (13) which we are interested in. In fact, after the matching the only nonvanishing contributions from the self-energy diagrams to these operators originate from the integral function $F_{\Phi\Phi_H}$, because we integrate the heavy Higgs doublet out in the effective theory.

After these preparations, we can now derive the Wilson coefficients of the operators $\mathcal{O}_{tq\Phi i}$ $(i=2,\ 3,\ 4,\ 5)$ and $\mathcal{O}_{tq\Phi i}$ $(i=7,\ 8,\ 9,\ 10)$. For clarity, we present their lengthy expressions in Appendix A.

3.2 The anomalous couplings $\mathcal{O}_{t\Phi}$

This class of anomalous couplings includes the effective operators

$$\mathcal{O}_{t\Phi 1} = i \Big(\Phi^{\dagger} D_{\mu} \Phi - (D_{\mu} \Phi)^{\dagger} \Phi \Big) \bar{t}_{R} \gamma_{\mu} t_{R} ,$$

$$\mathcal{O}_{t\Phi 2} = i \Big(\Phi^{\dagger} \Phi \Big) \Big(\bar{t}_{R} \gamma^{\mu} (D_{\mu} t_{R}) - (\overline{D_{\mu} t_{R}}) \gamma^{\mu} t_{R} \Big) ,$$

$$\mathcal{O}_{t\Phi 3} = i \Big(\Phi^{\dagger} D_{\mu} \Phi + (D_{\mu} \Phi)^{\dagger} \Phi \Big) \bar{t}_{R} \gamma_{\mu} t_{R} ,$$
(27)

where the operators $\mathcal{O}_{t\Phi 1}$, $\mathcal{O}_{t\Phi 2}$ have even CP-parity and $\mathcal{O}_{t\Phi 3}$ has odd CP-parity. In Fig. 5, we present those Feynman diagrams, which induce nontrivial contributions to the Wilson coefficients when matching the amplitude obtained in the effective theory with that in the full theory (MSSM). The ensuing Wilson coefficients are collected in Appendix B.

In the full theory, we also include the 1PI diagrams depicted in Fig. 6, where the gray blobs represent the corresponding diagrams of Fig. 2. However, the contributions from these diagrams disappear as a result of the matching of the effective theory and full theory amplitudes. In order to demonstrate this, let us consider an example. From the subsection 3.1, we find that the contributions of the subdiagram (framed by the dashed lines) in Fig. 7(a) induce the following term to the effective Lagrangian:

$$\mathcal{L}'_{eff} = \frac{1}{2\mu_{NP}^2} \sum_{\alpha=2}^{5} \left(C'_{tq\Phi\alpha} + C'_{tq\Phi(5+\alpha)} \right) \left(\mathcal{O}_{tq\Phi\alpha} + \mathcal{O}_{tq\Phi(5+\alpha)} \right), \tag{28}$$

where

$$\begin{split} C'_{tq\Phi2} &= -\frac{1}{48\pi^2} g_1^2 \sum_I \left[\Lambda_{U,I}^T \Big(\mathcal{C}_{121}^1(x_\mu, x_1, x_{U_I}) - 2s_\beta x_\mu \mathcal{C}_{131}^1(x_\mu, x_1, x_{U_I}) \Big) \right. \\ &\quad + 2c_\beta \Lambda_{U,I}^{R,1} \mathcal{C}_{131}^0(x_\mu, x_1, x_{U_I}) \right] \,, \\ C'_{tq\Phi3} &= \frac{1}{24\pi^2} g_1^2 \sum_I x_{U_I} \Big[s_\beta \Lambda_{U,I}^T \mathcal{C}_{113}^1(x_\mu, x_1, x_{U_I}) - c_\beta \Lambda_{U,I}^{R,1} \mathcal{C}_{113}^0(x_\mu, x_1, x_{U_I}) \Big] \,, \\ C'_{tq\Phi4} &= -\frac{1}{48\pi^2} g_1^2 \sum_I \Big[\Lambda_{U,I}^T \Big(Q_9(x_1, x_\mu, x_{U_I}) - 2s_\beta Q_4(x_\mu, x_1, x_{U_I}) \Big) + 2c_\beta \Lambda_{U,I}^{R,1} Q_3(x_\mu, x_1, x_{Q_I}) \Big] \,, \\ C'_{tq\Phi5} &= -\frac{1}{48\pi^2} g_1^2 \sum_I \Lambda_{U,I}^T \mathcal{C}_{112}^1(x_\mu, x_1, x_{U_I}) \,, \\ C'_{tq\Phi7} &= -\frac{i}{48\pi^2} g_1^2 \sum_I \Big[\Lambda_{U,I}^C \Big(\mathcal{C}_{121}^1(x_\mu, x_1, x_{U_I}) - 2s_\beta x_\mu \mathcal{C}_{131}^1(x_\mu, x_1, x_{U_I}) \Big) \end{split}$$

$$+2c_{\beta}\Lambda_{U,I}^{A,1}C_{131}^{0}(x_{\mu},x_{1},x_{U_{I}})\right],$$

$$C'_{tq\Phi8} = \frac{i}{24\pi^{2}}g_{1}^{2}\sum_{I}x_{U_{I}}\left[s_{\beta}\Lambda_{U,I}^{C}C_{113}^{1}(x_{\mu},x_{1},x_{U_{I}}) - c_{\beta}\Lambda_{U,I}^{A,1}C_{113}^{0}(x_{\mu},x_{1},x_{U_{I}})\right],$$

$$C'_{tq\Phi9} = \frac{i}{48\pi^{2}}g_{1}^{2}\sum_{I}\left[\Lambda_{U,I}^{C}\left(Q_{9}(x_{1},x_{\mu},x_{U_{I}}) - 2s_{\beta}Q_{4}(x_{\mu},x_{1},x_{U_{I}})\right) + 2c_{\beta}\Lambda_{U,I}^{A,1}Q_{3}(x_{\mu},x_{1},x_{Q_{I}})\right],$$

$$C'_{tq\Phi10} = -\frac{i}{48\pi^{2}}g_{1}^{2}\sum_{I}\Lambda_{U,I}^{C}C_{112}^{1}(x_{\mu},x_{1},x_{U_{I}}),$$

$$(29)$$

and $x_{\mu}=|\mu|^2/\mu_{NP}^2$, $x_{U_I}=m_{U_I}^2/\mu_{NP}^2$ and $x_i=|m_i|^2/\mu_{NP}^2$ $(i=1,\ 2,\ 3)$. The definition of the coupling constants $\Lambda_{U,I}$ and functions $Q_i(x,y,z)$ can be found in Appendix D.

In the effective theory, the amplitude of Fig. 7(b) is written as

$$A^{ET}(p_{1}, q_{1}, q_{2}) = \frac{i}{2} s_{\beta} \mathbf{h}_{U}^{3K} \left(\Phi^{\dagger} \Phi \right) \left\{ \left(C'_{tq\Phi 2} + C'_{tq\Phi 7} \right) \bar{t}_{R} \frac{1}{\not p_{1} + \not q_{1}} q_{1} \cdot p_{1} t_{R} \right.$$

$$\left. + \left(C'_{tq\Phi 3} + C'_{tq\Phi 8} \right) \bar{t}_{R} \frac{1}{\not p_{1} + \not q_{1}} p_{1} \cdot (q_{1} + p_{1}) t_{R} \right.$$

$$\left. + \left(C'_{tq\Phi 4} + C'_{tq\Phi 9} \right) \bar{t}_{R} \frac{1}{\not p_{1} + \not q_{1}} q_{1} \cdot (q_{1} + p_{1}) t_{R} \right.$$

$$\left. - \frac{1}{2} \left(C'_{tq\Phi 5} + C'_{tq\Phi 10} \right) \bar{t}_{R} \frac{1}{\not p_{1} + \not q_{1}} [\not q_{1}, \not p_{1}] t_{R} \right\}$$

$$(30)$$

where p_1 , q_1 , q_2 denote the four-momenta of the initial right-handed top quark and the Higgs bosons, respectively. In the full theory, the corresponding amplitude obtains the form

$$\begin{split} A^{FT}(p_{1},q_{1},q_{2}) &= -\frac{i}{96\pi^{2}\mu_{NP}^{2}}s_{\beta}g_{1}^{2}\Big(\mathbf{h}_{U}\mathbf{h}_{U}^{\dagger}\mathcal{Z}_{U}^{\dagger}\Big)^{3I}\mathcal{Z}_{U}^{I3}\Big(\Phi^{\dagger}\Phi\Big)\Big\{4\Big[c_{\beta}\mu_{NP}^{2}\sqrt{x_{\mu}x_{1}}e^{i(\varphi_{1}+\varphi_{\mu})}\mathcal{C}_{111}^{0}(x_{\mu},x_{1},x_{U_{I}})\\ &-s_{\beta}\mathcal{C}_{111}^{1}(x_{\mu},x_{1},x_{U_{I}})\Big]\bar{t}_{R}\frac{1}{\not{p}_{1}+\not{q}_{1}}t_{R}\\ &+\Big[2\mathcal{C}_{121}^{1}(x_{\mu},x_{1},x_{U_{I}})+2\mathcal{C}_{112}^{1}(x_{\mu},x_{1},x_{U_{I}})\\ &+4c_{\beta}\sqrt{x_{\mu}x_{1}}e^{i(\varphi_{1}+\varphi_{\mu})}Q_{3}(x_{\mu},x_{1},x_{U_{I}})\\ &-4s_{\beta}Q_{4}(x_{\mu},x_{1},x_{U_{I}})\Big]\bar{t}_{R}\frac{1}{\not{p}_{1}+\not{q}_{1}}q_{1}\cdot(q_{1}+p_{1})t_{R}\\ &-\Big[2\mathcal{C}_{121}^{1}(x_{\mu},x_{1},x_{U_{I}})+4c_{\beta}x_{\mu}\sqrt{x_{\mu}x_{1}}e^{i(\varphi_{1}+\varphi_{\mu})}\mathcal{C}_{131}^{0}(x_{\mu},x_{1},x_{U_{I}})\\ &-4s_{\beta}x_{\mu}\mathcal{C}_{131}^{1}(x_{\mu},x_{1},x_{U_{I}})\Big]\bar{t}_{R}\frac{1}{\not{p}_{1}+\not{q}_{1}}q_{1}\cdot p_{1}t_{R}\\ &+4x_{U_{I}}\Big[c_{\beta}x_{\mu}\sqrt{x_{\mu}x_{1}}e^{i(\varphi_{1}+\varphi_{\mu})}\mathcal{C}_{113}^{0}(x_{\mu},x_{1},x_{U_{I}})\\ &-s_{\beta}\mathcal{C}_{113}^{1}(x_{\mu},x_{1},x_{U_{I}})\Big]\bar{t}_{R}\frac{1}{\not{p}_{1}+\not{q}_{1}}p_{1}\cdot(q_{1}+p_{1})t_{R}\\ &-\mathcal{C}_{112}^{1}(x_{\mu},x_{1},x_{U_{I}})\bar{t}_{R}\frac{1}{\not{p}_{1}+\not{q}_{1}}\Big[\not{q}_{1},\not{p}_{1}\Big]t_{R}\Big\}, \end{split}$$

where φ_{μ} and φ_{i} (i=1,2,3) denote the *CP*-phases of the parameter μ and m_{i} , respectively. As already mentioned before, the first term of Eq. (31) is related to the Yukawa coupling renormalization in the full theory and it does not affect our computation. While matching Eq. (31) with Eq. (30), we find that the diagram does not contribute to the Wilson coefficients of the operators $\mathcal{O}_{t\Phi}$. A similar conclusion is true also for the other 1PI diagrams in Fig. 6.

3.3 The anomalous couplings $\mathcal{O}_{_{q\Phi}}$

This class of anomalous couplings includes the effective operators

$$\mathcal{O}_{q\Phi 1} = i \Big(\Phi^{\dagger} D_{\mu} \Phi - (D_{\mu} \Phi)^{\dagger} \Phi \Big) \bar{q}_{L} \gamma^{\mu} q_{L} ,$$

$$\mathcal{O}_{q\Phi 2} = i \Big(\Phi^{\dagger} \sigma^{a} D_{\mu} \Phi - (D_{\mu} \Phi)^{\dagger} \sigma^{a} \Phi \Big) \bar{q}_{L} \sigma^{a} \gamma^{\mu} q_{L} ,$$

$$\mathcal{O}_{q\Phi 3} = i \Big(\Phi^{\dagger} \Phi \Big) \Big(\bar{q}_{L} \gamma^{\mu} (D_{\mu} q_{L}) - \overline{(D_{\mu} q_{L})} \gamma^{\mu} q_{L} \Big) ,$$

$$\mathcal{O}_{q\Phi 4} = i \Big(\Phi^{\dagger} \sigma^{a} \Phi \Big) \Big(\bar{q}_{L} \sigma^{a} \gamma^{\mu} (D_{\mu} q_{L}) - \overline{(D_{\mu} q_{L})} \sigma^{a} \gamma^{\mu} q_{L} \Big) ,$$

$$\mathcal{O}_{q\Phi 5} = i \Big(\Phi^{\dagger} D_{\mu} \Phi + (D_{\mu} \Phi)^{\dagger} \Phi \Big) \bar{q}_{L} \gamma^{\mu} q_{L} ,$$

$$\mathcal{O}_{q\Phi 6} = i \Big(\Phi^{\dagger} \sigma^{a} D_{\mu} \Phi + (D_{\mu} \Phi)^{\dagger} \sigma^{a} \Phi \Big) \bar{q}_{L} \sigma^{a} \gamma^{\mu} q_{L} ,$$
(32)

where the last two operators are CP-odd and the others are CP-even. The Feynman diagrams, which induce nontrivial contributions to the Wilson coefficients, are presented in Fig. 8. We collect the expressions for the Wilson coefficients of the corresponding operators in Appendix C.

4 Experimental bounds on the Wilson coefficients

At present, the most rigorous constraint on the Wilson coefficients considered in this work comes from the decay $Z \to b\bar{b}$. For an on-shell Z, one can write the general effective vertex $Zb\bar{b}$ as [19]

$$\Gamma_{\mu}^{Zbb} = -i \frac{e}{4s_{W}c_{W}} \left[V_{b}^{Z} \gamma_{\mu} - A_{b}^{Z} \gamma_{\mu} \gamma_{5} + \frac{1}{2m_{b}} S_{b}^{Z} (p_{b} - p_{\bar{b}}) \right], \tag{33}$$

where $s_{\rm W}\equiv \sin\theta_{\rm W},\ c_{\rm W}\equiv \cos\theta_{\rm W},$ and $p_b,\ p_{\bar b}$ are the momenta of the outgoing quark and antiquark, respectively. For the operators listed in Eqs. (12), (13), (27) and (32), $S_b^Z=0$. The vector and axial-vector couplings can be written as

$$V_b^Z = V_b^{Z,0} + \delta V_b^Z ,$$

$$A_b^Z = A_b^{Z,0} + \delta A_b^Z ,$$
(34)

where $V_b^{Z,0}$, $A_b^{Z,0}$ represent the SM couplings and δV_b^Z , δA_b^Z are the new physics contributions. Ignoring the bottom quark mass, the lowest order theoretical prediction on the observable R_b at the Z pole is given as

$$R_b = \frac{\Gamma(Z \to b\bar{b})}{\Gamma(Z \to hadrons)} = R_b^{SM} \left\{ 1 + 2 \frac{V_b^{Z,0} \delta V_b^Z + A_b^{Z,0} \delta A_b^Z}{\left(V_b^{Z,0}\right)^2 + \left(A_b^{Z,0}\right)^2} (1 - R_b^{SM}) \right\}. \tag{35}$$

With the Born approximation, we can obtain modifications to the couplings V_b^Z , A_b^Z induced by the new physics operators $\mathcal{O}_{q\Phi 1}$ and $\mathcal{O}_{q\Phi 2}$. Provided that there is no accidental cancellation between these contributions, the corrections are given by [20]

$$\delta V_b^Z = \delta A_b^Z = \frac{2s_W m_W v}{e\mu_{NP}^2} \left[C_{q\Phi 1} + C_{q\Phi 2} \right], \tag{36}$$

where v denotes the VEV of the SM Higgs field doublet and $m_{\rm W}$ is the W-boson mass. From Eq. (35), we have

$$\delta V_b^Z = \delta A_b^Z = \frac{R_b^{exp} - R_b^{SM}}{(1 - R_b^{SM})R_b^{SM}} \frac{\left(V_b^{Z,0}\right)^2 + \left(A_b^{Z,0}\right)^2}{2(V_b^{Z,0} + A_b^{Z,0})} \,. \tag{37}$$

The SM prediction on R_b and the most recent experimental value are, respectively, given by [24]:

$$R_b^{SM} = 0.21572 \pm 0.00015, \quad R_b^{exp} = 0.21664 \pm 0.00065$$
 (38)

If we attribute the difference of these two values to the new physics effects, we get a bound for the new physics corrections on the R_b . At the 1σ tolerance we obtain:

$$0.00012 \le \Delta R_b \le 0.00172 \ . \tag{39}$$

Correspondingly, the bound for the Wilson coefficients is

$$3.1 \times 10^{-4} \le \frac{v^2}{\mu_{NP}^2} C_{q\Phi(1+2)} \le 4.5 \times 10^{-3}$$
(40)

with $C_{q\Phi(1+2)} = C_{q\Phi1} + C_{q\Phi2}$. Using the same method, we can also analyze the forward-backward asymmetry, A_{FB}^b , of the decay $Z \to b\bar{b}$. However, our theoretical result indicates that the present experiment data on this quantity set a weaker bound on the Wilson coefficients than R_b .

The other Wilson coefficients of the operators appearing in the Lagrangian are not constrained by R_b on the Born approximation level. With higher-order approximations, those operators contribute to the gauge boson self-energies, and thus we can get for them only a rather loose bound with a significant uncertainty. We can also have loose bounds from the argument of partial wave unitarity [25]:

$$|C_{tq\Phi 1}| \le \frac{16\pi}{3\sqrt{2}} \left(\frac{\mu_{NP}}{v}\right), \quad |C_{t\Phi 1}| \le 8\pi\sqrt{3},$$

$$-6.4 \le C_{tq\Phi 2} \le 10.4. \tag{41}$$

At present, there are no strong experimental constraints on the CP-odd couplings involving the top quark. It is well known that the MSSM contains in its general form unfortunately many 'new' free parameters in addition to the SM parameters. In order to simplify our discussion, we take the following assumption to restrict the MSSM parameter space:

- All possible CP phases are taken to be zero or π . A direct consequence of this choice is that there are no CP-odd operators in the effective Lagrangian Eq. (1)
- All Yukawa couplings and the soft breaking parameters are flavor conserving, *i.e.*, the mixing matrices $\mathcal{Z}_O = \mathcal{Z}_U = \mathcal{Z}_D = \mathbf{1}$.

Under these assumptions, the parameters relevant to our discussion are the gauge coupling constants g_1, g_2, g_3 , the higgsino and gaugino masses μ , m_1 , m_2 , m_3 , the Yukawa couplings of the third generation quarks and the corresponding soft breaking parameters $h_t = h_U^{33}$, $h_b = h_D^{33}$, $A_t = A_U^{33}$, $A_b = A_D^{33}$, and the square masses of the heavy Higgs boson doublet and the third generation squarks m_H^2 , $m_{Q_I}^2$, $m_{U_I}^2$, $m_{D_I}^2$ (I=3). In our numerical analysis, we will disregard the loose bounds from partial wave unitarity on the Wilson coefficients $C_{tq\Phi 1}$, $C_{tq\Phi 2}$ due to the large uncertainties mentioned above.

Without losing generality, we assume $m_Q=m_U=m_D$, $A_t=A_b$, $m_1=m_2=m_3$ in our numerical computations. Setting $\mu_{NP}=1000$ GeV, $m_H=500$ GeV, $m_1=m_2=m_3=500$ GeV, $A_t=A_b=1000$ GeV, $M_t=1000$ GeV, $M_t=10000$ GeV 100 GeV, we obtain constraints set by Eq. (40) on the soft breaking parameters. In Fig. 9, we plot the values of $m_Q = m_U = m_D$ versus the parameter μ with (a) $\tan \beta = 2$, and (b) $\tan \beta = 40$, where the gray regions are allowed by the condition for $v^2C_{q\Phi(1+2)}/\mu_{NP}^2$ set by Eq. (40). From this plot we observe that the restriction set on the parameter space with $\tan \beta = 40$ is more rigorous than that with $\tan \beta = 2$. As $\tan \beta = 2$, the contribution from the supersymmetric box diagrams varies from negative to positive gradually, then tends to zero after its maximum as $m_Q = m_U = m_D$ increase from 200 GeV. When $m_Q=m_U=m_D\geq 1.3$ TeV, the contribution is definitely less than 10^{-4} . Beside those box diagrams, $C_{q\Phi 1}$ also receives a contribution from the heavy Higgs doublet. Under our choice about the parameter space, the Higgs contribution to the Wilson coefficient $v^2 C_{q\Phi 1}/\mu_{\rm NP}^2$ is proportional to $\left|1/\tan^2\beta - \left(m_b/m_t\right)^4\tan^2\beta\right|$. Taking the bottom quark mass $m_b = 4.5 \text{ GeV}$ and top mass $m_t = 174 \text{ GeV}$, this contribution is about 5×10^{-4} for $\tan \beta = 2$. As $\tan \beta$ increasing, the contribution of the heavy Higgs doublet is strongly suppressed and less than 10^{-7} for $\tan \beta = 40$. The fact can help us understanding why very massive supersymmetry particles are allowed by the experimental bound for $\tan \beta = 2$ (Fig. 9(a)), whereas the most part of the parameter space is excluded by the bound except a narrow band at the neighborhood of $\mu = 0 \text{ for } \tan \beta = 40 \text{ (Fig. 9(b))}.$

In the figures discussed above, we have considered the 1σ tolerance for the experimental data. Since the central value of the experimentally measured R_b is only about one standard deviation away from the SM prediction, this sets a lower bound on the $C_{q\Phi(1+2)}$ which is positive and very close to zero as shown in Eq.(40). Certainly, very massive supersymmetry particles are excluded by this condition in the large $\tan\beta$ case. In fact, considering the practical situation of the experiments, we may relax the lower bound on the $C_{q\Phi(1+2)}$ to -5×10^{-5} , while the upper bound remains unchanged (this is just only slightly beyond the standard deviation). In Fig. 10 we plot $m_Q = m_U = m_D$ versus μ by using the constraint $-5\times 10^{-5} \le v^2 C_{q\Phi(1+2)}/\mu_{NP}^2 \le 4.5\times 10^{-3}$. One can see that the allowed parameter region is drastically enlarged in comparison with the case of the strict 1σ tolerance for the large $\tan\beta$.

Now, we discuss the operators $\mathcal{O}_{q\Phi_{1},2}$ corrections to R_b in the MSSM. Taking $\mu=m_i=500~{\rm GeV}$ (i=1,2,3), we plot ΔR_b versus squark masses $m_Q=m_U=m_D$ with $\tan\beta=2,40$ in Fig. 11. The gray band is the experimentally allowed region at the 1σ tolerance. When the scalar quark mass is less than 700 GeV, the supersymmetric box diagrams determine the leading contribution and results in a negative ΔR_b , the corresponding parameter space is excluded by Eq. (39). As the parameters $m_Q=m_U=m_D$ increase, the supersymmetric contribution turns to be positive, then tends to zero after the maximum. With $\tan\beta=2$, the correction of the heavy Higgs doublet to the ΔR_b is about 2.3×10^{-4} , and plays the leading role when $m_Q=m_U=m_D\geq 1.3~{\rm TeV}$. For $\tan\beta=40$, the total corrections from the Higgs and supersymmetric sectors to R_b do not satisfy Eq. (39), because the contribution of the heavy Higgs doublet is strongly suppressed. For $-\mu=m_i=500~{\rm GeV}$ (i=1,2,3), $m_Q=m_U=m_D=500$, 1000 GeV, and $\tan\beta=2$, 40, we present ΔR_b versus the parameter μ in Fig. 12. For $m_Q=m_U=m_D=500~{\rm GeV}$ (dot- and dot-dashed-lines), the corresponding parameter space is excluded by the condition Eq. (39) due to the negative supersymmetry contribution. With $m_Q=m_U=m_D=1~{\rm TeV}$ and $\tan\beta=2$ (solid-line), ΔR_b satisfies the condition Eq. (39) when $\mu\geq -700~{\rm GeV}$. As for the case $m_Q=m_U=m_D=1~{\rm TeV}$ and

 $\tan \beta = 40$ (dashed-line), the new physics correction to ΔR_b is excluded by the 1σ tolerance experimental bound except the region neighboring $\mu = 0$ GeV. Finally, we investigate the new physics prediction on ΔR_b with the assumption $m_Q = m_U = m_D = |\mu|$. Choosing $m_i = 500$ GeV (i = 1, 2, 3), $\tan \beta = 2, 40$, we plot ΔR_b versus the parameter $m_Q = m_U = m_D = |\mu|$ in Fig. 13. For the case $\tan \beta = 40$, the correction to ΔR_b exceeds the 1σ tolerance experimental bound. As $\tan \beta = 2$ and $\mu > 800$ GeV, the new physics prediction on ΔR_b satisfies this bound because of the relatively large contribution from the heavy Higgs. In those analyses the experimental bound with 1σ standard deviation are adopted. After we relax the condition (Eq. 39) slightly, the more massive supersymmetry particles are also permitted by the corresponding experimental bound.

Since the experimental data constrain the coefficients $C_{q\Phi1,2}$ strongly, the operators $\mathcal{O}_{q\Phi1,2}$ have only negligible effects on the measurements at the proposed future colliders [19]. Other operators will produce the observable effects in the next generation colliders. In the associated production of the Higgs boson and top quark pair $e^+e^- \to t\bar{t}$ h, the CP-even operators will affect the energy and angular distributions of the final state particles [20]. Through the measurements of various distributions, such as $d\sigma/dE_t$, $d\sigma/dE_h$ and $d\sigma/d\cos\theta_h$ (E_t , E_h denote the outgoing energy of the top quark and Higgs boson respectively, θ_h is the angle of three-momentum of the outgoing Higgs boson with respect to the electron beam direction), we can obtain useful information about the operators. The constraints on the CP-odd operators can be obtained through measuring various CP violation observables in this process. In the process $e^+e^- \to t\bar{t}$, we can analyze the effects of the operators on various polarized top-quark production cross sections. On the other hand, more strict constraints on the supersymmetry parameter space will be set by more precise measurements on the widths of $Z \to b\bar{b}$ and the top quark decays. All of these will provide valuable information for the search of supersymmetry particles on the future colliders.

It should be stressed that the above numerical analysis is performed under special assumptions about the MSSM parameter space. For example, we assume that all the parameters are real and flavor-conserving, the universal soft parameters are: $m_Q = m_U = m_D$, $A_t = A_b$, $m_1 = m_2 = m_3$. In a practical phenomenology analysis, those priori conditions should be dismissed. Nevertheless, it is quite obvious that the experimental data on R_b set significant bounds on the parameter space even in a more general case than that we have considered here.

5 The Summary

We have considered in this work the anomalous couplings between top quark and Higgs boson induced by the MSSM when the heavy Higgs doublet and all supersymmetry fields are integrated out. An essential assumption made here is that there is only one neutral Higgs boson with the electroweak mass, the other Higgs particles are much heavier. We have derived the Wilson coefficients of the relevant higher dimensional operators in the ensuing effective theory. We have also studied numerically the constraints set by the experimental results for $R_b = \Gamma(Z \to b\bar{b})/\Gamma(Z \to hadrons)$ on the parameters of the MSSM.

Acknowledgement

The work has been supported by the Academy of Finland under the contract no. 40677 and by the National Natural Science Foundation of China.

A The Wilson coefficients for the operators $\mathcal{O}_{tq\Phi i}~(i=2,~\cdots,5,~7,~\cdots,~10)$

$$\begin{split} C_{\iota_{q42}} &= -\frac{c_s}{(4\pi)^2} \bigg\{ \sum_{i=1,2} F_{i} g_{i}^{2} \xi_{\mathrm{R}_{i}}^{R} P_{H}(x_{\mu}, x_{i}) - s_{\mu} c_{\mu} \sum_{q=U,D} \Big(-1 \Big)^{\frac{1}{2} - T_{\mathrm{Z}}^{2}} \mathbf{Re} \Big[\mathbf{h}_{0}^{33} \Big(Z_{i} \hat{\mathbf{A}}_{a} Z_{Q} \Big)_{IJ} \Big] P_{H}(x_{\iota_{J}}, x_{q_{I}}) + s_{\sigma} c_{\mu} \mathbf{Re} \Big[\mathbf{h}_{0}^{33} \Big(Z_{h} \hat{\mathbf{A}}_{L} Z_{L} \Big)_{IJ} \Big(Z_{h} \hat{\mathbf{A}}_{h}^{L} Z_{L} \Big)_{JI} \Big] P_{H}(x_{\iota_{J}}, x_{n_{I}}) \bigg\} \\ &+ \frac{1}{32\pi^{2} x_{H}} s_{\mu} c_{\mu}^{2} \mathbf{Re} \Big(\mathbf{h}_{0}^{\dagger} \mathbf{h}_{0} \mathbf{h}_{0}^{\dagger} \Big)_{33} \Big(1 - \mathbf{ln} x_{H} \Big) + \frac{g_{\sigma}^{2}}{6\pi^{2}} s_{\mu} \mathbf{F}_{\iota_{JJ}}^{\mathrm{Ra}} x_{U_{J}} \mathcal{C}_{\iota_{13}}^{2} (x_{\mu}, x_{U_{J}}, x_{U_{J}}) \\ &+ \frac{g_{1}^{2}}{72\pi^{2}} s_{\mu} \mathbf{F}_{\iota_{IJ}}^{\mathrm{Ra}} x_{U_{J}} \mathcal{C}_{013}^{0} (x_{\mu}, x_{U_{J}}, x_{U_{J}}) + \frac{1}{16\pi^{2}} c_{\mu} \mathbf{F}_{\iota_{JJ}}^{\mathrm{Ra}} x_{U_{J}} \mathcal{C}_{133}^{0} (x_{\mu}, x_{U_{J}}, x_{U_{J}}) \\ &- \frac{1}{192\pi^{2}} \sum_{i=1,2} F_{i}^{2} g_{i}^{2} \Big[\mathbf{A}_{IJ}^{T} \Big(\mathcal{C}_{211}^{1} (x_{\mu}, x_{i}, x_{U_{J}}) + 2 s_{\mu} x_{\mu} \mathcal{C}_{311}^{1} (x_{\mu}, x_{i}, x_{U_{J}}) \Big) \\ &+ 2 c_{\mu} x_{\mu} \mathbf{A}_{iJ}^{\mathrm{RL}} \mathcal{C}_{031}^{0} (x_{\mu}, x_{i}, x_{U_{J}}) \Big] - \frac{1}{48\pi^{2}} g_{1}^{2} \Big[\mathbf{A}_{IJ}^{T} \Big(\mathcal{C}_{121}^{1} (x_{\mu}, x_{i}, x_{U_{J}}) \Big) \\ &+ 2 c_{\mu} x_{\mu} \mathbf{A}_{iJ}^{\mathrm{RL}} \mathcal{C}_{031}^{0} (x_{\mu}, x_{i}, x_{U_{J}}) \Big] - \frac{1}{48\pi^{2}} g_{1}^{2} \Big[\mathbf{A}_{IJ}^{T} \Big(\mathcal{C}_{121}^{1} (x_{\mu}, x_{i}, x_{U_{J}}) \Big) \\ &+ 2 c_{\mu} x_{\mu} \mathbf{A}_{iJ}^{\mathrm{RL}} \mathcal{C}_{031}^{0} (x_{\mu}, x_{\mu}, x_{U_{J}}) \Big] - \frac{1}{32\pi^{2} x_{\mu}} s_{\mu} \mathbf{A}_{iJ}^{\mathrm{RL}} \mathcal{C}_{031}^{0} (x_{\mu}, x_{\mu}, x_{U_{J}}) \Big] \\ &+ 2 c_{\mu} x_{\mu} \mathbf{A}_{iJ}^{\mathrm{RL}} \mathcal{C}_{031}^{0} (x_{\mu}, x_{\mu}, x_{U_{J}}) \Big] - \frac{1}{16\pi^{2}} s_{\mu} \mathbf{B}_{iJ}^{\mathrm{RL}} \mathcal{C}_{031}^{\mathrm{RL}} (x_{\mu}, x_{\mu}, x_{U_{J}}) \Big] \\ &+ 2 c_{\mu} x_{\mu} \mathbf{A}_{\mu} \mathbf{A}_{iJ}^{\mathrm{RL}} \mathcal{C}_{031}^{\mathrm{RL}} \Big[\mathbf{A}_{iJ}^{\mathrm{RL}} \mathbf{A}_{iJ}^{\mathrm{RL}} \mathbf{A}_{iJ}^{\mathrm{RL}} \Big] \Big[\mathbf{A}_{iJ}^{\mathrm{RL}} \mathcal{C}_{031}^{\mathrm{RL}} \mathbf{A}_{iJ}^{\mathrm{RL}} \mathbf{A}_{iJ}^{\mathrm{RL}} \mathbf{A}_{iJ}^{\mathrm{RL}} \Big] \Big] \\ &+ \frac{1}{96\pi^{2}} s_{\mu} \mathbf{A}_{iJ}^{\mathrm{RL}} \mathbf{A}_{iJ}^{\mathrm{RL}} \mathbf{A}_{iJ}^{\mathrm{RL}} \mathbf{A}_{iJ}^{\mathrm{RL}} \mathbf{A}_{iJ}^{\mathrm{RL}} \mathbf{A}_{iJ}^{\mathrm{RL}} \mathbf{A}_{iJ}$$

$$\begin{split} C_{l_0 \neq 5} &= -\frac{1}{32\pi^2 x_n} s_\beta c_\beta^2 \mathbf{Re} \Big(\mathbf{h}_D^\dagger \mathbf{h}_D \mathbf{h}_D^\dagger \Big)_{33} - \frac{1}{192\pi^2} \sum_{i=1,2} F_i^2 g_i^2 \lambda_{Q,I}^T \mathcal{C}_{112}^I (x_\mu, x_i, x_{Q_I}) \\ &- \frac{1}{48\pi^2} g_i^2 \lambda_{U,I}^T \mathcal{C}_{112}^I (x_\mu, x_i, x_{U_I}) \,, \\ C_{t_0 \neq \tau} &= i \frac{c_\beta}{(4\pi)^2} \Big\{ \sum_{i=1,2} F_{ij}^2 \xi_{R_I}^4 P_H(x_\mu, x_i) - s_\beta c_\beta \sum_{q=C,D} \Big(-1 \Big)^{\frac{1}{2} - T_2^2} \mathbf{Im} \Big[\mathbf{h}_0^{33} \Big(\mathcal{Z}_q \hat{\mathbf{A}}_\sigma \mathcal{Z}_Q \Big)_{IJ} \Big] P_H(x_{Q_J}, x_{q_I}) + s_\beta c_\beta \mathbf{Im} \Big[\mathbf{h}_0^{33} \Big(\mathcal{Z}_R \hat{\mathbf{A}}_K \mathcal{Z}_L \Big)_{IJ} \Big(\mathcal{Z}_R \hat{\mathbf{A}}_S^I \mathcal{Z}_L \Big)_{IJ}^\dagger P_H(x_{L_J}, x_{R_I}) \Big\} \\ &+ \frac{i}{32\pi^2 x_H} s_\beta c_\beta^2 \mathbf{Im} \Big(\mathbf{h}_D^\dagger \mathbf{h}_D \mathbf{h}_0^\dagger \Big)_{33} \Big(1 - \mathbf{ln} x_H \Big) + i \frac{g_3^2}{6\pi^2} s_\beta \Gamma_{0,IJ}^{*,13} x_{U_J} \mathcal{C}_{113}^{\dagger} \Big(x_\mu, x_{U_J} + x_{R_I} \Big) \Big\} \\ &+ i \frac{g_1^2}{92\pi^2} s_\beta \Gamma_{I,IJ}^{*,14} x_{U_J} \mathcal{C}_{113}^{\dagger} \Big(x_\mu, x_{U_J}, x_{U_J} \Big) + \frac{i}{16\pi^2} c_\beta \Gamma_{U,IJ}^B x_{U_J} \mathcal{C}_{113}^{\dagger} \Big(x_\mu, x_{U_J}, x_{U_J} \Big) \\ &+ i \frac{g_1^2}{192\pi^2} \sum_{i=1,2} F_i^2 g_i^2 \Big[\Lambda_{Q,I}^C \Big(\mathcal{C}_{211}^\dagger (x_\mu, x_\iota, x_{Q_I}) + 2 s_\beta x_\mu \mathcal{C}_{211}^\dagger (x_\mu, x_\iota, x_{Q_I}) \Big) \\ &+ 2 c_\beta x_\mu \mathcal{C}_{113}^{*,14} \Big(x_\mu, x_\iota, x_{Q_I} \Big) \Big] - \frac{i}{48\pi^2} g_1^2 \Big[\Lambda_{C,I}^C \Big(\mathcal{C}_{121}^\dagger (x_\mu, x_\iota, x_{Q_I}) \Big) \\ &+ 2 c_\beta x_\mu \mathcal{C}_{121}^A \Big(x_\mu, x_\iota, x_{Q_I} \Big) \Big] + 2 c_\beta x_\mu \Lambda_{U,I}^A \mathcal{C}_{03}^0 \Big(x_\mu, x_\iota, x_{Q_I} \Big) \Big] \\ &+ 2 c_\beta x_\mu \mathcal{C}_{121}^A \Big(x_\mu, x_\iota, x_{Q_I} \Big) \Big] + 2 c_\beta x_\mu \Lambda_{U,I}^A \mathcal{C}_{03}^0 \Big(x_\mu, x_\iota, x_{Q_I} \Big) \Big] \\ &+ 2 c_\beta x_\mu \mathcal{C}_{121}^A \Big(x_\mu, x_\iota, x_{Q_I} \Big) \Big] + 2 c_\beta x_\mu \Lambda_{U,I}^A \mathcal{C}_{03}^0 \Big(x_\mu, x_{Q_I}, x_{Q_I} \Big) \Big] \\ &+ 2 c_\beta x_\mu \mathcal{C}_{121}^A \Big(x_\mu, x_\iota, x_{Q_I} \Big) \Big] + 2 c_\beta x_\mu \Lambda_{U,I}^A \mathcal{C}_{03}^0 \Big(x_\mu, x_{Q_I}, x_{Q_I} \Big) \Big] \\ &+ 2 c_\beta \mathcal{C}_{121}^A \mathcal{C}_{121}^A \Big(x_\mu, x_\iota, x_{Q_I} \Big) + 2 c_\beta \mathcal{C}_{121}^A \mathcal{C}_{121}^0 \Big(x_\mu, x_{Q_I} \Big) \Big] \\ &+ 2 c_\beta \mathcal{C}_{121}^A \mathcal{C}_{121}^2 \Big(x_\mu, x_\mu, x_{Q_I} \Big) + 2 c_\beta \mathcal{C}_{121}^A \mathcal{C}_{121}^1 \Big(x_\mu, x_\iota, x_{Q_I} \Big) + c_\beta \mathcal{C}_{121}^A \mathcal{C}_{121}^0 \Big(x_\mu, x_\mu, x_{Q_I} \Big) \Big] \\ &+ \frac{i}{16\pi^2} \mathcal{C}_{121}^A \mathcal{C}_{121}^A \Big(x_\mu, x_\iota, x_{Q_I} \Big) +$$

$$C_{tq\Phi10} = -\frac{i}{32\pi^2 x_H} s_{\beta} c_{\beta}^2 \mathbf{Im} \left(\mathbf{h}_{D}^{\dagger} \mathbf{h}_{D} \mathbf{h}_{U}^{\dagger} \right)_{33} - \frac{i}{192\pi^2} \sum_{i=1,2} F_i^2 g_i^2 \Lambda_{Q,I}^C \mathcal{C}_{112}^1(x_{\mu}, x_i, x_{Q_I})$$

$$-\frac{i}{48\pi^2} g_1^2 \Lambda_{U,I}^C \mathcal{C}_{112}^1(x_{\mu}, x_1, x_{U_I})$$

$$(42)$$

with $F_1=1,~F_2=3,~T_Z^U=-T_Z^D=\frac{1}{2},~Y_U^B=\frac{4}{3},~Y_D^B=-\frac{2}{3}$ and $\hat{\mathbf{A}}_U=\frac{1}{\mu_{NP}}\Big(\mathbf{A}_U-\frac{\mu^*}{\tan\beta}\mathbf{h}_U\Big),~\hat{\mathbf{A}}_D=\frac{1}{\mu_{NP}}\Big(\mathbf{A}_D-\mu^*\tan\beta\mathbf{h}_D\Big),~\hat{\mathbf{A}}_E=\frac{1}{\mu_{NP}}\Big(\mathbf{A}_E-\mu^*\tan\beta\mathbf{h}_E\Big)$ and $\hat{\mathbf{A}}_F'=\hat{\mathbf{A}}_F+\frac{\mu^*}{s_\beta c_\beta \mu_{NP}}\mathbf{h}_F~(F=U,~D,~E).$ In the above expression, the sum with the generation indices I,~J is implied.

B The Wilson coefficients for operators $\mathcal{O}_{t\Phi i}$ $(i=1,\ 2,\ 3)$

The corresponding Wilson coefficients are

$$\begin{split} C_{t \bullet 1} &= -\frac{1}{32 \pi^2 x_H} c_{\beta}^2 \Big[\mathbf{h}_{U} \Big(s_{\beta}^2 \mathbf{h}_{U}^{\dagger} \, \mathbf{h}_{U} - c_{\beta}^2 \mathbf{h}_{D}^{\dagger} \, \mathbf{h}_{D} \Big) \mathbf{h}_{U}^{\dagger} \Big]_{33} \Big(1 - \ln x_H \Big) \\ &+ \frac{\mathcal{X}_{U,IJK}^n}{72 \pi^2} \Big[g_{1}^2 \mathcal{D}_{1121}^1 (x_{1}, x_{U_{I}}, x_{Q_{J}}, x_{U_{K}}) + 3 g_{3}^2 \mathcal{D}_{1121}^1 (x_{3}, x_{U_{I}}, x_{Q_{J}}, x_{U_{K}}) \Big] \\ &+ \frac{1}{64 \pi^2} \sum_{q = U, D} \mathcal{X}_{q,IJK}^S \mathcal{D}_{1121}^1 (x_{\mu}, x_{Q_{I}}, x_{q_{J}}, x_{Q_{K}}) + \frac{1}{144 \pi^2} g_{1}^4 \mathcal{Z}_{U}^{43I} \mathcal{Z}_{U}^{I3} \Big[c_{\beta}^2 Q_{5}(x_{\mu}, x_{1}, x_{U_{I}}) \\ &+ 2 \xi_{H1}^S Q_{6}(x_{\mu}, x_{1}, x_{U_{I}}) + s_{\beta}^2 x_{1} Q_{7}(x_{\mu}, x_{1}, x_{U_{I}}) \Big] \\ &- \sum_{i = 1, 2} \frac{F_{i} g_{i}^2}{128 \pi^2} \Big(\mathbf{h}_{U} \mathcal{Z}_{Q} \Big)_{3I} \Big(\mathcal{Z}_{Q}^{\dagger} \mathbf{h}_{U}^{\dagger} \Big)_{I3} \Big[2 \xi_{H}^S Q_{6}(x_{i}, x_{\mu}, x_{Q_{I}}) \\ &+ s_{\beta}^2 Q_{5}(x_{i}, x_{\mu}, x_{Q_{I}}) + c_{\beta}^2 x_{\mu} Q_{7}(x_{i}, x_{\mu}, x_{Q_{I}}) \Big] \\ &+ \frac{1}{48 \pi^2} g_{1}^2 s_{\beta} \Big[2 \Gamma_{U,IJ}^U \mathcal{D}_{1111}^0 (x_{\mu}, x_{1}, x_{Q_{I}}, x_{U_{J}}) - \Gamma_{U,IJ}^T U_{5}(x_{\mu}, x_{Q_{I}}, x_{1}, x_{U_{J}}) \Big] , \\ C_{t \bullet 2} &= \frac{\left(\mathbf{h}_{U} \mathbf{h}_{U}^{\dagger} \right)_{33}}{256 \pi^2 x_{H}} c_{\beta}^2 \Big[4 s_{\beta}^2 (s_{\beta}^2 - c_{\beta}^2) (2 g_{1}^2 + 3 g_{2}^2) \ln x_{H} - 12 s_{\beta}^4 g_{2}^2 + (c_{\beta}^4 - 7 s_{\beta}^4) g_{1}^2 \Big] \\ &+ \frac{g_{1}^2}{108 \pi^2} (s_{\beta}^2 - c_{\beta}^2) \mathcal{Z}_{U}^{\dagger 3I} \mathcal{Z}_{U}^{\dagger 3I} \Big[g_{1}^2 \mathcal{B}_{3,1}^1 (x_{U_{I}}, x_{1}) + 3 g_{3}^2 \mathcal{B}_{3,1}^1 (x_{U_{I}}, x_{3}) \Big] \\ &+ \frac{c_{\beta}^2}{72 \pi^2} \mathbf{Re} \Big[\mathcal{Z}_{U}^{\dagger 3I} \Big(\mathcal{Z}_{U} \mathbf{h}_{U} \mathbf{h}_{U}^{\dagger} \mathcal{Z}_{U}^{\dagger} \Big)^{IJ} \mathcal{Z}_{U}^{J3} \Big] \Big[g_{1}^2 Q_{9}(x_{U_{I}}, x_{1}, x_{U_{J}}) + 3 g_{3}^2 Q_{9}(x_{U_{I}}, x_{3}, x_{U_{J}}) \Big] \\ &- \frac{1}{192 \pi^2} g_{1}^2 (s_{\beta}^2 - c_{\beta}^2) \Big(\mathbf{h}_{U} \mathcal{Z}_{Q} \Big)_{3I} \Big(\mathcal{Z}_{Q}^{\dagger} \mathbf{h}_{U}^{\dagger} \Big)_{I3} \mathcal{B}_{3,1}^3 (x_{Q_{I}}, x_{\mu}) \\ &+ \frac{1}{16 \pi^2} \mathbf{Re} \Big[\Big(\mathbf{h}_{U} \mathcal{Z}_{Q} \Big)^{3I} \Big(\mathcal{Z}_{Q}^{\dagger} (c_{\beta}^2 \mathbf{h}_{U}^{\dagger} \mathbf{h}_{U} + s_{\beta}^2 \mathbf{h}_{D}^{\dagger} \mathbf{h}_{D} \Big) \mathcal{Z}_{Q} \Big)^{IJ} \Big(\mathcal{Z}_{Q}^{\dagger} \mathbf{h}_{U}^{\dagger} \Big)^{J3} \Big] Q_{9}(x_{Q_{I}}, x_{\mu}, x_{Q_{I}}) \\ &- \frac{c_{\beta}^2}{64 \pi^2 x_{H}} \Big[\mathbf{h}_{U} \Big(s_{\beta}^2 \mathbf{h}_{U}$$

$$\begin{split} &+\frac{1}{144\pi^{2}}g_{1}^{4}\mathcal{Z}_{U}^{\dagger 3I}\mathcal{Z}_{U}^{I3}\left[c_{\beta}^{2}\mathcal{C}_{122}^{2}(x_{\mu},x_{1},x_{U_{I}})+s_{\beta}^{2}\mathcal{C}_{122}^{1}(x_{\mu},x_{1},x_{U_{I}})+2\xi_{HI}^{S}\mathcal{C}_{122}^{1}(x_{\mu},x_{1},x_{U_{I}})\right]\\ &+\sum_{i=1,2}\frac{F_{i}g_{i}^{2}}{128\pi^{2}}\left(h_{U}\mathcal{Z}_{Q}\right)_{3I}\left(\mathcal{Z}_{Q}^{\dagger}h_{U}^{\dagger}\right)_{I3}\left[c_{\beta}^{2}x_{\mu}\mathcal{C}_{122}^{1}(x_{\mu},x_{i},x_{Q_{I}})+s_{\beta}^{2}\mathcal{C}_{212}^{2}(x_{\mu},x_{i},x_{Q_{I}})\right]\\ &+2\xi_{Hi}^{S}\mathcal{C}_{212}^{1}(x_{\mu},x_{i},x_{Q_{I}})\right]-\frac{1}{48\pi^{2}}g_{1}^{2}s_{\beta}\Gamma_{U,IJ}^{T}U_{5}(x_{\mu},x_{1},x_{Q_{I}},x_{U_{J}})\;,\\ &\mathcal{C}_{t\oplus 3}=i\frac{c_{\beta}^{2}}{72\pi^{2}}\mathbf{Im}\left[\mathcal{Z}_{U}^{\dagger 3I}\left(\mathcal{Z}_{U}\mathbf{h}_{U}\mathbf{h}_{U}^{\dagger}\mathcal{Z}_{U}^{\dagger}\right)^{IJ}\mathcal{Z}_{U}^{J3}\right]\left[g_{1}^{2}\mathcal{C}_{211}^{1}(x_{U_{I}},x_{1},x_{U_{J}})+3g_{3}^{2}\mathcal{C}_{211}^{1}(x_{U_{I}},x_{3},x_{U_{J}})\right]\\ &+\frac{i}{16\pi^{2}}\mathbf{Im}\left[\left(\mathbf{h}_{U}\mathcal{Z}_{Q}\right)^{3I}\left(\mathcal{Z}_{Q}^{\dagger}(c_{\beta}^{2}\mathbf{h}_{U}^{\dagger}\mathbf{h}_{U}+s_{\beta}^{2}\mathbf{h}_{D}^{\dagger}\mathbf{h}_{D})\mathcal{Z}_{Q}\right)^{IJ}\left(\mathcal{Z}_{Q}^{\dagger}\mathbf{h}_{U}^{\dagger}\right)^{J3}\right]\mathcal{C}_{211}^{1}(x_{Q_{I}},x_{\mu},x_{Q_{J}})\\ &+i\frac{1}{36\pi^{2}}\mathcal{X}_{U,IJK}^{A}\left[g_{1}^{2}\mathcal{D}_{1211}^{1}(x_{1},x_{U_{I}},x_{Q_{J}},x_{U_{K}})+3g_{3}^{2}\mathcal{D}_{1211}^{1}(x_{3},x_{U_{I}},x_{Q_{J}},x_{U_{K}})\right]\\ &+i\frac{1}{32\pi^{2}}\sum_{q=U,D}\mathcal{X}_{q,IJK}^{B}\mathcal{D}_{1211}^{1}(x_{\mu},x_{Q_{I}},x_{Q_{I}},x_{Q_{K}})+\frac{i}{36\pi^{2}}g_{1}^{4}\xi_{Hi}^{B}\mathcal{Z}_{U}^{\dagger 3I}\mathcal{Z}_{U}^{I3}\mathcal{C}_{211}^{0}(x_{\mu},x_{1},x_{U_{I}})\\ &+i\sum_{i=1,2}\frac{F_{i}g_{i}^{2}}{32\pi^{2}}\xi_{Hi}^{B}\left(h_{U}\mathcal{Z}_{Q}\right)_{3I}\left(\mathcal{Z}_{Q}^{\dagger}h_{U}^{\dagger}\right)_{I3}\mathcal{C}_{211}^{0}(x_{\mu},x_{i},x_{Q_{I}})\\ &+\frac{i}{48\pi^{2}}g_{1}^{2}s_{\beta}\left[2\Gamma_{U,IJ}^{D}\mathcal{D}_{1111}^{0}(x_{\mu},x_{1},x_{Q_{I}},x_{U_{J}})-\Gamma_{U,IJ}^{C}U_{5}(x_{\mu},x_{U_{J}},x_{U_{J}},x_{Q_{I}},x_{1})\right]. \end{split}$$

Just as in the appendix A, the sum with the generation indices I, J, K is implied.

C The Wilson coefficients for operators $\mathcal{O}_{q\Phi i}$ $(i=1,\ \cdots\ 6)$

The Wilson coefficients for those operators are written as

$$\begin{split} C_{q^{\Phi 1}} &= \frac{1}{32\pi^2 x_H} s_{\beta}^2 \mathcal{C}_{\beta}^2 \Big[\Big(\mathbf{h}_{U}^{\dagger} \mathbf{h}_{U} \mathbf{h}_{U}^{\dagger} \mathbf{h}_{U} \Big)_{33} - \Big(\mathbf{h}_{D}^{\dagger} \mathbf{h}_{D} \mathbf{h}_{D}^{\dagger} \mathbf{h}_{D} \Big)_{33} \Big] \Big(1 - \ln x_H \Big) \\ &+ \sum_{q = U, D} (-1)^{\frac{1}{2} - T_{Z}^{q}} \mathcal{X}_{q, IJK}^T \Big[\sum_{i = 1, 2} \frac{F_{i}^2 g_{i}^2}{2304\pi^2} \mathcal{D}_{1121}^1 (x_{i}, x_{Q_{I}}, x_{q_{J}}, x_{Q_{K}}) \\ &+ \frac{g_{3}^2}{48\pi^2} \mathcal{D}_{1121}^1 (x_{3}, x_{Q_{I}}, x_{q_{J}}, x_{Q_{K}}) \Big] - \frac{1}{64\pi^2} \sum_{q = U, D} (-1)^{\frac{1}{2} - T_{Z}^{q}} \mathcal{X}_{q, IJK}^U \mathcal{D}_{1121}^1 (x_{\mu}, x_{q_{I}}, x_{Q_{J}}, x_{q_{K}}) \\ &+ \frac{g_{1}^4}{2304\pi^2} (c_{\beta}^2 - s_{\beta}^2) \mathcal{Z}_{Q}^{3I} \mathcal{Z}_{Q}^{\dagger I3} Q_{8}(x_{\mu}, x_{1}, x_{Q_{I}}) \\ &+ \sum_{i = 1, 2} \frac{F_{i} g_{i}^2}{256\pi^2} \Big(\mathbf{h}_{U}^{\dagger} \mathcal{Z}_{U}^{\dagger} \Big)_{3I} \Big(\mathcal{Z}_{U} \mathbf{h}_{U} \Big)_{I3} \Big[s_{\beta}^2 Q_{10}(x_{\mu}, x_{i}, x_{U_{I}}) - 4 \xi_{Hi}^S \mathcal{C}_{211}^0 (x_{\mu}, x_{i}, x_{U_{I}}) \\ &+ \eta_{Hi}^R Q_{7}(x_{i}, x_{\mu}, x_{U_{I}}) \Big] - \sum_{i = 1, 2} \frac{F_{i} g_{i}^2}{256\pi^2} \Big(\mathbf{h}_{D}^{\dagger} \mathcal{Z}_{D}^{\dagger} \Big)_{3I} \Big(\mathcal{Z}_{D} \mathbf{h}_{D} \Big)_{I3} \Big[c_{\beta}^2 Q_{10}(x_{\mu}, x_{i}, x_{D_{I}}) \\ &- 4 \xi_{Hi}^S \mathcal{C}_{211}^0 (x_{\mu}, x_{i}, x_{D_{I}}) + \eta_{Hi}^S Q_{7}(x_{i}, x_{\mu}, x_{D_{I}}) \Big] \end{split}$$

$$\begin{split} &+\frac{g_{1}^{2}}{256\pi^{2}}\sum_{q=U,D}Y_{q}^{B}\left[2\Gamma_{\eta,IJ}^{W}\mathcal{D}_{1111}^{0}(x_{\mu},x_{1},x_{q_{I}},x_{Q_{J}})-\Gamma_{\eta,IJ}^{V,1}U_{5}(x_{\mu},x_{q_{I}},x_{1},x_{Q_{J}})\right]\\ &-\frac{3g_{2}^{2}}{128\pi^{2}}\sum_{q=U,D}\left[2\Gamma_{\eta,IJ}^{W}\mathcal{D}_{1111}^{0}(x_{\mu},x_{2},x_{q_{I}},x_{Q_{J}})-\Gamma_{\eta,IJ}^{V,2}U_{5}(x_{\mu},x_{q_{I}}^{2},x_{2},x_{Q_{J}})\right],\\ &C_{g^{2}2}=\sum_{q=U,D}\mathcal{X}_{q,IJK}^{F}\left[\sum_{i=1,2}(-1)^{i}\frac{F_{i}^{2}g_{i}^{2}}{2304\pi^{2}}\mathcal{D}_{1121}^{1}(x_{i},x_{Q_{I}},x_{q_{I}},x_{Q_{K}})\\ &-\frac{g_{3}^{2}}{48\pi^{2}}\mathcal{D}_{1121}^{1}(x_{3},x_{Q_{I}},x_{q_{I}},x_{Q_{K}})\right]+\frac{g_{2}^{4}}{128\pi^{2}}(s_{g}^{2}-c_{g}^{2})\mathcal{Z}_{q}^{3I}\mathcal{Z}_{q}^{\dagger I3}Q_{8}(x_{\mu},x_{2},x_{Q_{I}})\\ &-\frac{g_{1}^{2}g_{2}^{2}}{384\pi^{2}}(s_{g}^{2}-c_{g}^{2})\mathcal{Z}_{q}^{3I}\mathcal{Z}_{q}^{\dagger I3}\left[U_{2}(x_{\mu},x_{1},x_{2},x_{Q_{I}})+\sqrt{x_{1}x_{2}}\cos(\varphi_{2}-\varphi_{1})U_{3}(x_{\mu},x_{1},x_{2},x_{Q_{I}})\right]\\ &+\sum_{i=1,2}(-1)^{i}\frac{g_{1}^{2}}{256\pi^{2}}\left(\mathbf{h}_{v}^{1}\mathcal{Z}_{v}^{1}\right)_{3I}\left(\mathcal{Z}_{v}\mathbf{h}_{v}\right)_{I3}\left[s_{g}^{2}Q_{10}(x_{\mu},x_{i},x_{u_{I}})-4\xi_{H}^{S}\mathcal{C}_{211}^{0}(x_{\mu},x_{i},x_{u_{I}})\right]\\ &+\mu_{H_{i}}^{R}Q_{7}(x_{i},x_{\mu},x_{u_{I}})\right]+\sum_{i=1,2}(-1)^{i}\frac{g_{i}^{2}}{256\pi^{2}}\left(\mathbf{h}_{v}^{1}\mathcal{Z}_{v}^{1}\right)_{3I}\left(\mathcal{Z}_{v}\mathbf{h}_{v}\right)_{I3}\left[c_{g}^{2}Q_{10}(x_{\mu},x_{i},x_{u_{I}},x_{u_{I}})\right]\\ &-4\xi_{H_{i}}^{S}\mathcal{C}_{211}^{O}(x_{\mu},x_{i},x_{u_{I}})+\eta_{H_{i}}^{S}Q_{7}(x_{i},x_{\mu},x_{u_{I}})\right]\\ &-\frac{g_{1}^{2}}{256\pi^{2}}\sum_{q=U,D}(-1)^{\frac{1}{2}-T_{u}^{2}}\mathcal{Y}_{q}^{B}\left[2\Gamma_{u_{i}D}^{W}\mathcal{D}_{1111}^{O}(x_{\mu},x_{2},x_{q_{I}},x_{q_{I}})-\Gamma_{u,I}^{V,1}U_{5}(x_{\mu},x_{q_{I}},x_{1},x_{Q_{I}})\right]\\ &-\frac{g_{2}^{2}}{128\pi^{2}}\sum_{q=U,D}(-1)^{\frac{1}{2}-T_{u}^{2}}\mathcal{Y}_{q}^{B}\left[2\Gamma_{u_{i}D}^{W}\mathcal{D}_{1111}^{O}(x_{\mu},x_{2},x_{q_{I}},x_{Q_{I}})-\Gamma_{u,I}^{V,1}U_{5}(x_{\mu},x_{q_{I}},x_{2},x_{Q_{I}})\right],\\ &C_{g^{2}3}&=3\frac{g_{1}^{2}+g_{2}^{2}}{2}s_{p}^{2}c_{g}^{2}(s_{g}^{2}-c_{g}^{2})\left[\left(\mathbf{h}_{v}^{1}\mathbf{h}_{v}\mathbf{h}_{v}\right)_{33}-\left(\mathbf{h}_{v}^{1}\mathbf{h}_{v}\right)_{33}\right]+\frac{1}{1}\frac{1}{128\pi^{2}}\sum_{q=U,D}\left[Y_{q}^{B}\mathcal{G}_{1}^{2}(s_{g}^{2}-c_{g}^{2})\left(\mathbf{h}_{v}^{1}\mathcal{F}_{v}^{1}\right)_{3}\left(\mathcal{Z}_{q}^{1}\mathbf{h}_{v}\right)_{I3}\mathcal{B}_{s,1}^{2}(x_{q_{I}},x_{s},x_{q_{I}})\right]\\ &+\frac{1}{128\pi^{2}}\sum_$$

$$\begin{split} & + \frac{g_1^4}{2304\pi^2} Z_q^{g_1} Z_q^{IT3} \Big[C_{122}^2(x_n, x_1, x_{Q_I}) + \eta_{II}^T C_{122}^\dagger(x_\mu, x_1, x_{Q_I}) \Big] \\ & + \sum_{i=1,2} \frac{F_{ij}^2}{256\pi^2} \Big(\mathbf{h}_{i}^\dagger Z_{i}^\dagger \Big)_{3I} \Big(Z_{i}^\dagger \mathbf{h}_{i} \Big)_{13} \Big[\eta_{Ri}^R C_{12}^\dagger(x_\mu, x_1, x_{Q_I}) + s_s^2 C_{212}^2(x_\mu, x_1, x_{Q_I}) \Big] \\ & + \sum_{i=1,2} \frac{F_{ij}^2}{256\pi^2} \Big(\mathbf{h}_{i}^\dagger Z_{i}^\dagger \Big)_{3I} \Big(Z_{D}^\dagger \mathbf{h}_{D} \Big)_{13} \Big[\eta_{Ri}^R C_{12}^\dagger(x_\mu, x_1, x_{D_I}) + s_s^2 C_{212}^2(x_\mu, x_1, x_{D_I}) \Big] \\ & + \sum_{i=1,2} \frac{F_{ij}^2}{256\pi^2} \Big(\mathbf{h}_{i}^\dagger Z_{i}^\dagger \Big)_{3I} \Big(Z_{D}^\dagger \mathbf{h}_{D} \Big)_{3I} \Big[\eta_{Ri}^R C_{12}^\dagger(x_\mu, x_1, x_{D_I}) + s_s^2 C_{212}^2(x_\mu, x_1, x_{D_I}) \Big] \\ & + \frac{1}{64\pi^2} \sum_{q=U,D} (-1)^{\frac{1}{2}-T_2^2} \Gamma_{ri}^{YZ} U_1(x_\mu, x_{Q_I}, x_{Q_I}) + \frac{g_1^2}{256\pi^2} \sum_{q=U,D} (-1)^{\frac{1}{2}-T_2^2} Y_q^B \Gamma_{u,IJ}^{YI} U_5(x_\mu, x_1, x_{Q_I}, x_{Q_I}) \Big] \\ & - \frac{3g_2^2}{128\pi^2} \sum_{q=U,D} (-1)^{\frac{1}{2}-T_2^2} \Gamma_{ri}^{YZ} U_5(x_\mu, x_2, x_{Q_I}, x_{Q_I}) \Big] \\ & - \frac{3g_2^2}{128\pi^2} \sum_{s_i^2} C_s^2 (s_s^2 - c_s^2) \Big[\Big(\mathbf{h}_i^\dagger \mathbf{h}_u \Big)_{33} + \Big(\mathbf{h}_i^\dagger \mathbf{h}_u \Big)_{33} \Big] \frac{1 + \ln x_H}{x_H} \\ & - \frac{1}{256\pi^2} x_H \Big(g_2^2 - 2s_s^2 c_g^2 (g_1^2 + g_2^2) \Big) \Big[c_x^2 \Big(\mathbf{h}_b^\dagger \mathbf{h}_u \Big)_{33} \Big] \frac{1 + \ln x_H}{x_H} \\ & - \frac{g_2^2}{2304\pi^2} \Big(s_s^2 - c_s^2 \Big) Z_q^3 Z_q^{2I_3} \Big[\sum_{i=1,2} (-1)^{i} F_{i}^2 g_1^2 B_{3,1} (x_{Q_I}, x_i) - 48g_3^2 B_{3,1}^4 (x_{Q_I}, x_3) \Big] \\ & + \frac{1}{2304\pi^2} \lambda_{IJ}^8 \Big[\sum_{i=1,2} (-1)^{i-1} F_i^2 g_1^2 Q_9 (x_{Q_I}, x_i, x_{Q_I}) + 48g_3^2 Q_9 (x_{Q_I}, x_3, x_{Q_I}) \Big] \\ & + \frac{1}{2304\pi^2} \lambda_{IJ}^8 \Big[\sum_{i=1,2} (-1)^{\frac{1}{2}-T_2^2} X_{q,IK}^2 \Big[\sum_{i=1,2} (-1)^{i} F_i^2 g_1^2 U_1 (x_i, x_{Q_I}, x_{Q_I}, x_{Q_I}) + \eta_{II}^T C_{12}^4 (x_\mu, x_2, x_{Q_I}) \Big] \\ & + \frac{1}{2304\pi^2} \Big[D_{1112}^2 (x_\mu, x_1, x_{Q_I}, x_{Q_I}) + \Big(\sqrt{x_1 x_2} \cos(\varphi_2 - \varphi_1) + 2 \Big[\xi_{Ri}^2 + \xi_{Ri}^2 \Big] D_{1112}^2 (x_\mu, x_1, x_2, x_{Q_I}) \Big] \Big] \\ & + \sum_{i=1,2} \Big[-1 \Big[\frac{g_1^2}{256\pi^2} \Big(\mathbf{h}_b^\dagger Z_b^\dagger \Big)_{3I} \Big(Z_U \mathbf{h}_U \Big)_{3I} \Big[\eta_{Ri}^R C_{212}^2 (x_\mu, x_i, x_{Q_I}) + s_3^2 C_{212}^2 (x_\mu, x_i, x_{Q_I}) \Big] \Big] \\ & - \sum_{i=1,2} \Big[-1 \Big[\frac{g_$$

$$\begin{split} &+\frac{i}{1152\pi^{2}}\sum_{q=U,D}\sum_{i=1,2}\mathcal{X}_{q,IJK}^{C}\left[F_{i}^{3}g_{i}^{2}\mathcal{D}_{1211}^{1}(x_{i},x_{Q_{I}},x_{q_{J}},x_{Q_{K}})\right.\\ &+48g_{3}^{2}\mathcal{D}_{1211}^{1}(x_{3},x_{Q_{I}},x_{q_{J}},x_{Q_{K}})\right]+\frac{i}{32\pi^{2}}\sum_{q=U,D}\mathcal{X}_{q,IJK}^{D}\mathcal{D}_{1211}^{1}(x_{\mu},x_{q_{I}},x_{Q_{J}},x_{q_{K}})\\ &-i\frac{g_{1}^{4}}{288\pi^{2}}\xi_{HI}^{B}\mathcal{Z}_{q}^{3I}\mathcal{Z}_{Q1}^{1/3}\mathcal{C}_{021}^{0}(x_{\mu},x_{1},x_{Q_{I}})\\ &-\frac{i}{64\pi^{2}}\sum_{q=U,D}\sum_{i=1,2}F_{i}g_{i}^{2}\xi_{Hi}^{B}\left(\mathbf{h}_{q}^{\dagger}\mathcal{Z}_{q}^{\dagger}\right)_{3I}\left(\mathcal{Z}_{q}\mathbf{h}_{q}\right)_{I3}\mathcal{C}_{211}^{0}(x_{\mu},x_{i},x_{q_{I}})\\ &+i\frac{g_{1}^{2}}{256\pi^{2}}\sum_{q=U,D}\left(-1\right)^{\frac{1}{2}-T_{2}^{q}}Y_{q}^{B}\left[\Gamma_{q,IJ}^{E,1}U_{5}(x_{\mu},x_{Q_{J}},x_{q_{I}},x_{1})-2\Gamma_{q,IJ}^{r}\mathcal{D}_{1111}^{0}(x_{\mu},x_{1},x_{q_{I}},x_{Q_{J}})\right]\\ &-i\frac{3g_{2}^{2}}{128\pi^{2}}\sum_{q=U,D}\left(-1\right)^{\frac{1}{2}-T_{2}^{q}}\left[\Gamma_{q,IJ}^{E,2}U_{5}(x_{\mu},x_{Q_{J}},x_{q_{I}},x_{2})-2\Gamma_{q,IJ}^{r}\mathcal{D}_{1111}^{0}(x_{\mu},x_{2},x_{q_{I}},x_{Q_{J}})\right],\\ &C_{q^{46}} &=\frac{\Lambda_{IJ}^{B}}{2304\pi^{2}}\left(\sum_{i=1,2}\left(-1\right)^{i-1}g_{i}^{2}F_{i}^{3}\mathcal{C}_{112}^{1}(x_{Q_{I}},x_{i},x_{Q_{J}})+48g_{3}^{2}\mathcal{C}_{112}^{1}(x_{Q_{I}},x_{3},x_{Q_{J}})\right)\\ &+\frac{i}{1152\pi^{2}}\sum_{q=U,D}\left(-1\right)^{\frac{1}{2}-T_{2}^{q}}\left[\sum_{i=1,2}\left(-1\right)^{i}F_{i}^{2}g_{i}^{2}\mathcal{X}_{q,IJK}^{c}\mathcal{D}_{1211}^{1}(x_{1},x_{Q_{I}},x_{q_{J}},x_{Q_{J}})\right)\\ &+\frac{i}{384\pi^{2}}\mathcal{Z}_{q,IJK}^{3}\mathcal{D}_{1211}^{1}(x_{3},x_{Q_{I}},x_{q_{J}},x_{Q_{K}})\right]\\ &-i\frac{g_{1}^{2}g_{2}^{2}}{384\pi^{2}}\mathcal{Z}_{3}^{3I}\mathcal{Z}_{1}^{\dagger I3}\left[4\left(\xi_{Hi}^{B}+\xi_{H2}^{B}\right)\mathcal{D}_{1111}^{0}(x_{\mu},x_{1},x_{2},x_{Q_{I}})\right.\\ &-\left.\left.\left(c_{g}^{2}-s_{g}^{2}\right)\sqrt{x_{1}x_{2}}\sin(\varphi_{2}-\varphi_{1}\right)-2\left[\xi_{Hi}^{B}-\xi_{H2}^{B}\right]\right)U_{4}(x_{\mu},x_{1},x_{2},x_{Q_{I}})\right]\\ &-\frac{1}{64\pi^{2}}\sum_{q=U,D}\left(-1\right)^{\frac{1}{2}-T_{2}^{q}}\sum_{i=1,2}\left(-1\right)^{i}g_{i}^{2}\xi_{Hi}^{B}\left(\mathbf{h}_{U}^{\dagger}\mathcal{Z}_{U}^{\dagger}\right)_{3I}\left(\mathcal{Z}_{U}\mathbf{h}_{U}\right)_{I3}\mathcal{C}_{211}^{0}(x_{\mu},x_{1},x_{Q_{I}},x_{Q_{I}})\right]\\ &-i\frac{g_{1}^{2}}{256\pi^{2}}\sum_{q=U,D}\left[\Gamma_{q,IJ}^{E,IJ}U_{5}(x_{\mu},x_{Q_{J}},x_{q_{I}},x_{1})-2\Gamma_{q,IJ}^{p}\mathcal{D}_{1111}^{0}(x_{\mu},x_{2},x_{q_{I}},x_{Q_{I}})\right]. \end{split}$$

(44)

D The coupling constants and loop functions

The loop functions are defined as

$$\begin{split} P_{H}(x,y) &= -\mathcal{B}^{0}_{1,3}(x,y) + 3\mathcal{B}^{1}_{1,4}(x,y) - \frac{2}{3}\mathcal{B}^{2}_{1,5}(x,y) \;, \\ Q_{1}(x,y,z) &= \mathcal{C}^{1}_{122}(x,y,z) + y\mathcal{C}^{0}_{131}(x,y,z) + z\mathcal{C}^{0}_{113}(x,y,z) \;, \end{split}$$

$$\begin{aligned} Q_{2}(x,y,z) &= \mathcal{C}_{122}^{2}(x,y,z) + y\mathcal{C}_{131}^{1}(x,y,z) + z\mathcal{C}_{113}^{1}(x,y,z) \,, \\ Q_{3}(x,y,z) &= \mathcal{C}_{122}^{1}(x,y,z) + x\mathcal{C}_{131}^{0}(x,y,z) + z\mathcal{C}_{113}^{0}(x,y,z) \,, \\ Q_{4}(x,y,z) &= \mathcal{C}_{122}^{2}(x,y,z) + x\mathcal{C}_{131}^{1}(x,y,z) + z\mathcal{C}_{113}^{1}(x,y,z) \,, \\ Q_{5}(x,y,z) &= \mathcal{C}_{122}^{2}(x,y,z) + 2\mathcal{C}_{131}^{2}(x,y,z) - 4\mathcal{C}_{121}^{1}(x,y,z) \,, \\ Q_{6}(x,y,z) &= \mathcal{C}_{122}^{1}(x,y,z) + 2\mathcal{C}_{131}^{1}(x,y,z) + 2\mathcal{C}_{121}^{0}(x,y,z) \,, \\ Q_{7}(x,y,z) &= \mathcal{C}_{122}^{1}(x,y,z) + 2\mathcal{C}_{131}^{1}(x,y,z) \,, \\ Q_{8}(x,y,z) &= 4\mathcal{C}_{121}^{1}(x,y,z) - 2\mathcal{C}_{131}^{2}(x,y,z) - \mathcal{C}_{122}^{2}(x,y,z) + y\left(2\mathcal{C}_{131}^{1}(x,y,z) + \mathcal{C}_{122}^{1}(x,y,z)\right) \\ Q_{9}(x,y,z) &= \mathcal{C}_{11}^{1}(x,y,z) + \mathcal{C}_{112}^{1}(x,y,z) \\ Q_{10}(x,y,z) &= \mathcal{C}_{212}^{2}(x,y,z) + 2\mathcal{C}_{311}^{2}(x,y,z) - 4\mathcal{C}_{211}^{1}(x,y,z) \\ U_{1}(x,y,z,w) &= 2\mathcal{D}_{1211}^{1}(x,y,z,w) + \mathcal{D}_{1121}^{1}(x,y,z,w) \,, \\ U_{2}(x,y,z,w) &= 4\mathcal{D}_{1111}^{1}(x,y,z,w) - \mathcal{D}_{1211}^{2}(x,y,z,w) - \mathcal{D}_{1211}^{2}(x,y,z,w) - \mathcal{D}_{1121}^{2}(x,y,z,w) \\ U_{3}(x,y,z,w) &= \mathcal{D}_{1211}^{1}(x,y,z,w) + \mathcal{D}_{1121}^{1}(x,y,z,w) + \mathcal{D}_{1112}^{1}(x,y,z,w) \\ U_{4}(x,y,z,w) &= \mathcal{D}_{1211}^{1}(x,y,z,w) + \mathcal{D}_{1121}^{1}(x,y,z,w) \\ U_{5}(x,y,z,w) &= \mathcal{D}_{1121}^{1}(x,y,z,w) + \mathcal{D}_{1121}^{1}(x,y,z,w) \end{aligned}$$

$$(45)$$

The coupling constants are

$$\begin{split} \xi_{Hi}^{R} &= \sqrt{x_{\mu}x_{i}} \Big((c_{\beta}^{2} - s_{\beta}^{2}) \mathbf{Re}(\mathbf{h}_{U})_{33} \cos(\varphi_{\mu} + \varphi_{i}) - \mathbf{Im}(\mathbf{h}_{U})_{33} \sin(\varphi_{\mu} + \varphi_{i}) \Big) \;, \; (i = 1, \; 2) \;, \\ \xi_{Hi}^{A} &= \sqrt{x_{\mu}x_{i}} \Big(\mathbf{Im}(\mathbf{h}_{U})_{33} (c_{\beta}^{2} - s_{\beta}^{2}) \cos(\varphi_{\mu} + \varphi_{i}) + \mathbf{Re}(\mathbf{h}_{U})_{33} \sin(\varphi_{\mu} + \varphi_{i}) \Big) \;, \; (i = 1, \; 2) \;, \\ \xi_{Hi}^{S} &= s_{\beta}c_{\beta}\sqrt{x_{\mu}x_{i}} \cos(\varphi_{\mu} + \varphi_{i}) \;, \; (i = 1, \; 2) \;, \\ \xi_{Hi}^{B} &= s_{\beta}c_{\beta}\sqrt{x_{\mu}x_{i}} \sin(\varphi_{\mu} + \varphi_{i}) \;, \; (i = 1, \; 2) \;, \\ \eta_{Hi}^{R} &= c_{\beta}^{2}x_{\mu} + 2s_{\beta}c_{\beta}\sqrt{x_{\mu}x_{i}} \cos(\varphi_{\mu} + \varphi_{i}) \;, \; (i = 1, \; 2) \;, \\ \eta_{Hi}^{T} &= s_{\beta}^{2}x_{\mu} + 2s_{\beta}c_{\beta}\sqrt{x_{\mu}x_{i}} \cos(\varphi_{\mu} + \varphi_{i}) \;, \; (i = 1, \; 2) \;, \\ \eta_{Hi}^{T} &= x_{i} + 4s_{\beta}c_{\beta}\sqrt{x_{\mu}x_{i}} \cos(\varphi_{\mu} + \varphi_{i}) \;, \; (i = 1, \; 2) \;, \\ \Lambda_{U,I}^{R,i} &= \sqrt{x_{\mu}x_{i}} \Big(\mathbf{Re} \Big[\mathcal{Z}_{Q}^{3I} \Big(\mathcal{Z}_{Q}^{\dagger} \mathbf{h}_{U}^{\dagger} \Big)^{I3} \Big] \cos(\varphi_{\mu} + \varphi_{i}) - \mathbf{Im} \Big[\mathcal{Z}_{Q}^{3I} \Big(\mathcal{Z}_{Q}^{\dagger} \mathbf{h}_{U}^{\dagger} \Big)^{I3} \Big] \sin(\varphi_{\mu} + \varphi_{i}) \Big) \;, \; (i = 1, \; or \; 2) \\ \Lambda_{U,I}^{A,i} &= \sqrt{x_{\mu}x_{i}} \Big(\mathbf{Re} \Big[\mathcal{Z}_{Q}^{3I} \Big(\mathcal{Z}_{Q}^{\dagger} \mathbf{h}_{U}^{\dagger} \Big)^{I3} \Big] \sin(\varphi_{\mu} + \varphi_{i}) + \mathbf{Im} \Big[\mathcal{Z}_{Q}^{3I} \Big(\mathcal{Z}_{Q}^{\dagger} \mathbf{h}_{U}^{\dagger} \Big)^{I3} \Big] \cos(\varphi_{\mu} + \varphi_{i}) \Big) \;, \; (i = 1, \; or \; 2) \\ \Lambda_{IJ}^{S} &= \mathbf{Re} \Big[\mathcal{Z}_{Q}^{3I} \Big(\mathcal{Z}_{Q}^{\dagger} (\mathbf{h}_{U}^{\dagger} \mathbf{h}_{U} \Big)_{33} + s_{\beta}^{2} \Big(\mathbf{h}_{D}^{\dagger} \mathbf{h}_{D} \Big) \mathcal{Z}_{Q} \Big)^{IJ} \mathcal{Z}_{Q}^{\dagger J3} \Big] \;, \\ \Lambda_{IJ}^{B} &= \mathbf{Im} \Big[\mathcal{Z}_{Q}^{3I} \Big(\mathcal{Z}_{Q}^{\dagger} (c_{\beta}^{2} \Big(\mathbf{h}_{U}^{\dagger} \mathbf{h}_{U} \Big)_{33} + s_{\beta}^{2} \Big(\mathbf{h}_{D}^{\dagger} \mathbf{h}_{D} \Big) \mathcal{Z}_{Q} \Big)^{IJ} \mathcal{Z}_{Q}^{\dagger J3} \Big] \;, \\ \Lambda_{U,I}^{T} &= \mathbf{Re} \Big((\mathbf{h}_{U}^{\dagger} \mathcal{Z}_{U}^{\dagger})^{3I} \mathcal{Z}_{U}^{J3} \Big) \;, \; \Lambda_{U,I}^{C} &= \mathbf{Im} \Big((\mathbf{h}_{U}^{\dagger} \mathcal{Z}_{U}^{\dagger})^{3I} \mathcal{Z}_{U}^{J3} \Big) \;, \end{cases}$$

$$\begin{split} & \Lambda_{Q,I}^{T} = \mathbf{Re} \Big(\mathcal{Z}_{Q}^{3I} (\mathcal{Z}_{Q}^{\dagger} \mathbf{h}_{Q}^{\dagger})^{IS} \Big), \quad \Lambda_{Q,I}^{C} = \mathbf{Im} \Big(\mathcal{Z}_{Q}^{3I} (\mathcal{Z}_{Q}^{\dagger} \mathbf{h}_{U}^{\dagger})^{IS} \Big), \\ & \Lambda_{U,IJ}^{U} = c_{\rho}^{2} \mathbf{Re} \Big[\Big(\mathcal{Z}_{U} \mathbf{h}_{U} \mathbf{h}_{U}^{\dagger} \mathcal{Z}_{U}^{\dagger} \mathbf{h}_{U}^{\dagger} \mathcal{Z}_{U}^{\dagger} \Big)^{IJ} \Big(\mathcal{Z}_{U} \mathbf{h}_{U} \Big)^{JS} \Big], \\ & \Lambda_{D,IJ}^{U} = s_{\rho}^{2} \mathbf{Re} \Big[\Big(\mathcal{Z}_{U} \mathbf{h}_{U} \mathbf{h}_{U}^{\dagger} \mathcal{Z}_{U}^{\dagger} \mathbf{h}_{U}^{\dagger} \mathcal{Z}_{U}^{\dagger} \Big)^{IJ} \Big(\mathcal{Z}_{U} \mathbf{h}_{U} \Big)^{JS} \Big], \\ & \Lambda_{D,IJ}^{U} = s_{\rho}^{2} \mathbf{Re} \Big[\Big(\mathcal{Z}_{D} \mathbf{h}_{D} \Big)^{1SI} \Big(\mathcal{Z}_{D} \mathbf{h}_{D} \mathbf{h}_{D}^{\dagger} \mathcal{Z}_{D}^{\dagger} \Big)^{IJ} \Big(\mathcal{Z}_{D} \mathbf{h}_{D} \Big)^{JS} \Big], \\ & \Lambda_{D,IJ}^{D} = s_{\rho}^{2} \mathbf{Re} \Big[\Big(\mathcal{Z}_{D} \mathbf{h}_{D} \Big)^{1SI} \Big(\mathcal{Z}_{D} \mathbf{h}_{D} \mathbf{h}_{D}^{\dagger} \mathcal{Z}_{D}^{\dagger} \Big)^{IJ} \Big(\mathcal{Z}_{D} \mathbf{h}_{D} \Big)^{JS} \Big], \\ & \Lambda_{U,IJ}^{D} = \sqrt{s_{\nu}} \Big(\mathbf{Re} \Big[\mathcal{Z}_{Q}^{3I} \Big(\mathcal{Z}_{U} \hat{\mathbf{A}}_{U} \mathcal{Z}_{Q} \Big)^{\dagger IJ} \mathcal{Z}_{U}^{JS} \Big] \cos \varphi_{\ell} - \mathbf{Im} \Big[\mathcal{Z}_{Q}^{3I} \Big(\mathcal{Z}_{U} \hat{\mathbf{A}}_{U} \mathcal{Z}_{Q} \Big)^{\dagger IJ} \mathcal{Z}_{U}^{JS} \Big] \sin \varphi_{\ell} \Big], \\ & \Lambda_{U,IJ}^{D} = \sqrt{s_{\nu}} \Big(\mathbf{Re} \Big[\mathcal{Z}_{Q}^{3I} \Big(\mathcal{Z}_{U} \hat{\mathbf{A}}_{U} \mathcal{Z}_{Q} \Big)^{\dagger IJ} \mathcal{Z}_{U}^{JS} \Big] \sin \varphi_{\ell} + \mathbf{Im} \Big[\mathcal{Z}_{Q}^{3I} \Big(\mathcal{Z}_{U} \hat{\mathbf{A}}_{U} \mathcal{Z}_{Q} \Big)^{\dagger IJ} \mathcal{Z}_{U}^{JS} \Big] \sin \varphi_{\ell} \Big), \\ & (i = 1, \text{ or } 3) \Big[\mathcal{Z}_{U}^{2} \hat{\mathbf{A}}_{U} \mathcal{Z}_{Q} \Big)^{\dagger IJ} \mathcal{Z}_{U}^{JS} \Big] \sin \varphi_{\ell} + \mathbf{Im} \Big[\mathbf{A}_{U}^{\dagger} \mathcal{Z}_{U}^{\dagger} \mathbf{A}_{U}^{\dagger} \mathcal{Z}_{U}^{\dagger} \Big] \Big[\mathcal{Z}_{Q}^{\dagger} \mathbf{h}_{U}^{\dagger} \Big]^{JS} \Big] \cos \varphi_{\mu} \\ & - \mathbf{Im} \Big[\Big(\mathbf{h}_{U}^{\dagger} \mathcal{Z}_{D}^{\dagger} \mathcal{Z}_{D}^{\dagger} \mathcal{Z}_{U}^{\dagger} \mathcal{Z}_{U}^{\dagger} \mathcal{Z}_{U}^{\dagger} \Big] \Big[\mathcal{Z}_{Q}^{\dagger} \mathbf{h}_{U}^{\dagger} \Big]^{JS} \Big] \sin \varphi_{\mu} \Big), \\ & \Gamma_{U,IJ}^{T} = \mathbf{Re} \Big[\Big(\mathbf{h}_{U} \mathcal{Z}_{Q} \Big)_{3I} \Big(\mathcal{Z}_{D} \hat{\mathbf{A}}_{D} \mathcal{Z}_{Q} \Big)^{IJ} \Big(\mathcal{Z}_{Q}^{\dagger} \mathbf{h}_{U}^{\dagger} \Big)^{JS} \Big] \sin \varphi_{\mu} \Big), \\ & \Gamma_{U,IJ}^{D} = \mathbf{Re} \Big[\Big(\mathbf{h}_{U} \mathcal{Z}_{Q} \Big)_{3I} \Big(\mathcal{Z}_{U} \hat{\mathbf{A}}_{U} \mathcal{Z}_{Q} \Big)^{IJ} \mathcal{Z}_{U}^{JS} \Big] \Big[\mathcal{Z}_{Q}^{J} \mathbf{h}_{U}^{J} \mathcal{Z}_{U}^{JS} \Big] \cos \varphi_{\mu} + \mathbf{Im} \Big[\Big(\mathbf{h}_{U} \mathcal{Z}_{Q} \mathcal{A}_{U} \mathcal{Z}_{Q} \Big)^{J} \mathcal{Z}_{U}^{JS} \Big] \sin \varphi_{\mu} \Big\}, \\ & \Gamma_{U,IJ}^{D} = \mathbf{Re} \Big[\Big(\mathbf{h}_{U} \mathcal{Z}_{Q} \Big)_{3I} \Big(\mathcal{Z}_{U} \hat{$$

$$\begin{split} \Gamma^{W}_{q,IJ} &= s_{\beta} c_{\beta} \sqrt{x_{\mu}} \bigg\{ \mathbf{Re} \Big[\Big(\mathbf{h}_{q}^{\dagger} \mathcal{Z}_{q}^{\dagger} \Big)_{3I} \Big(\mathcal{Z}_{q} \hat{\mathbf{A}}_{q} \mathcal{Z}_{Q} \Big)_{IJ} \mathcal{Z}_{Q}^{\dagger I3} \Big] \cos \varphi_{\mu} \\ &- \mathbf{Im} \Big[\Big(\mathbf{h}_{q}^{\dagger} \mathcal{Z}_{q}^{\dagger} \Big)_{3I} \Big(\mathcal{Z}_{q} \hat{\mathbf{A}}_{q} \mathcal{Z}_{Q} \Big)_{IJ} \mathcal{Z}_{Q}^{\dagger I3} \Big] \sin \varphi_{\mu} \bigg\} \,, \\ \Gamma^{F}_{q,IJ} &= s_{\beta} c_{\beta} \sqrt{x_{\mu}} \bigg\{ \mathbf{Re} \Big[\Big(\mathbf{h}_{q}^{\dagger} \mathcal{Z}_{q}^{\dagger} \Big)_{3I} \Big(\mathcal{Z}_{q} \hat{\mathbf{A}}_{q} \mathcal{Z}_{Q} \Big)_{IJ} \mathcal{Z}_{Q}^{\dagger I3} \Big] \sin \varphi_{\mu} \\ &+ \mathbf{Im} \Big[\Big(\mathbf{h}_{q}^{\dagger} \mathcal{Z}_{q}^{\dagger} \Big)_{3I} \Big(\mathcal{Z}_{q} \hat{\mathbf{A}}_{u} \mathcal{Z}_{Q} \Big)_{IJ} \mathcal{Z}_{Q}^{\dagger I3} \Big] \cos \varphi_{\mu} \bigg\} \,, \\ \Gamma^{V,i}_{D,IJ} &= - \mathbf{Re} \Big[\Big(\mathbf{h}_{D}^{\dagger} \mathcal{Z}_{D}^{\dagger} \Big)_{3I} \Big(\mathcal{Z}_{D} \hat{\mathbf{A}}_{D} \mathcal{Z}_{Q} \Big)_{IJ} \mathcal{Z}_{Q}^{\dagger I3} \Big] \cos \varphi_{\mu} \bigg\} \,, \\ \Gamma^{V,i}_{D,IJ} &= - \mathbf{Re} \Big[\Big(\mathbf{h}_{D}^{\dagger} \mathcal{Z}_{D}^{\dagger} \Big)_{3I} \Big(\mathcal{Z}_{D} \hat{\mathbf{A}}_{D} \mathcal{Z}_{Q} \Big)_{IJ} \mathcal{Z}_{Q}^{\dagger I3} \Big] c_{\beta} \Big(s_{\beta} \sqrt{x_{\mu}} \cos \varphi_{\mu} + c_{\beta} \sqrt{x_{i}} \cos \varphi_{i} \Big) \,, \\ \Gamma^{E,i}_{D,IJ} &= - \mathbf{Re} \Big[\Big(\mathbf{h}_{D}^{\dagger} \mathcal{Z}_{D}^{\dagger} \Big)_{3I} \Big(\mathcal{Z}_{D} \hat{\mathbf{A}}_{D} \mathcal{Z}_{Q} \Big)_{IJ} \mathcal{Z}_{Q}^{\dagger I3} \Big] c_{\beta} \Big(s_{\beta} \sqrt{x_{\mu}} \sin \varphi_{\mu} - c_{\beta} \sqrt{x_{i}} \sin \varphi_{i} \Big) \,, \\ - \mathbf{Im} \Big[\Big(\mathbf{h}_{D}^{\dagger} \mathcal{Z}_{D}^{\dagger} \Big)_{3I} \Big(\mathcal{Z}_{D} \hat{\mathbf{A}}_{D} \mathcal{Z}_{Q} \Big)_{IJ} \mathcal{Z}_{Q}^{\dagger I3} \Big] c_{\beta} \Big(s_{\beta} \sqrt{x_{\mu}} \sin \varphi_{\mu} - c_{\beta} \sqrt{x_{i}} \sin \varphi_{i} \Big) \,, \\ \mathcal{Z}^{R}_{U,IJK} &= s_{\beta}^{2} \mathbf{Re} \Big(\mathcal{Z}_{D}^{\dagger I3} \Big(\mathcal{Z}_{D} \hat{\mathbf{A}}_{D} \mathcal{Z}_{Q} \Big)_{IJ} \mathcal{Z}_{Q}^{\dagger I3} \Big] c_{\beta} \Big(s_{\beta} \sqrt{x_{\mu}} \cos \varphi_{\mu} + c_{\beta} \sqrt{x_{i}} \cos \varphi_{i} \Big) \,, \\ \mathcal{X}^{R}_{U,IJK} &= s_{\beta}^{2} \mathbf{Re} \Big(\mathcal{Z}_{D}^{\dagger I3} \Big(\mathcal{Z}_{D} \hat{\mathbf{A}}_{D} \mathcal{Z}_{Q} \Big)_{IJ} \mathcal{Z}_{Q}^{\dagger I3} \Big) c_{\beta} \Big(s_{\beta} \sqrt{x_{\mu}} \cos \varphi_{\mu} + c_{\beta} \sqrt{x_{i}} \cos \varphi_{i} \Big) \,, \\ \mathcal{X}^{R}_{U,IJK} &= s_{\beta}^{2} \mathbf{Re} \Big(\mathcal{Z}_{D}^{\dagger I3} \Big(\mathcal{Z}_{D} \hat{\mathbf{A}}_{U} \mathcal{Z}_{Q} \Big)^{IJ} \Big(\mathcal{Z}_{U} \hat{\mathbf{A}}_{U} \mathcal{Z}_{Q} \Big)^{\dagger JK} \mathcal{Z}_{U}^{\dagger K3} \Big) \,, \\ \mathcal{X}^{R}_{U,IJK} &= s_{\beta}^{2} \mathbf{Re} \Big(\Big(\mathbf{h}_{U} \mathcal{Z}_{Q} \Big)^{3I} \Big(\mathcal{Z}_{Q} \hat{\mathbf{A}}_{U} \mathcal{Z}_{Q} \Big)^{IJ} \Big(\mathcal{Z}_{Q} \hat{\mathbf{A}}_{U} \mathcal{Z}_{Q} \Big)^{JK} \Big(\mathcal{Z}_{Q}^{\dagger h}_{U} \Big)^{3I} \Big) \,, \\ (q = U, D; \chi_{U} = s_{\beta}^{2}, \chi_{D} = c_{\beta}^{2} \Big) \,, \\ \mathcal{X}^{R}_{u,IJK}$$

References

- S. Dawson, ICTP Summer School in High-Energy Physics and Cosmology, Miramare, Trieste, 29 June - 17 July, 1998.
- [2] M. Carena and H. E. Haber, Prog. Part. Nucl. Phys. **50**, 63(2003); see also the Report of the Higgs working group of the Physics at Run II Supersymmetry/Higgs workshop, hep-ph/0010338.
- [3] J. Ellis, T. Falk, K. Olive, M. Schmitt, Phys. Lett. B388, 97(1996); B413, 355(1997); S. A. Abel, B. C. Allanach, Phys. Lett. B431, 339(1998); J. A. Casas, J. R. Espinosa, H. E. Haber, Nucl. Phys. B526, 3(1998); M. Carena, P. Chankowski, S. Pokorski, C. E. M. Wagner, Phys. Lett. B441, 205(1998);
- [4] M. S. Berger, Phys. Rev. D41, 225(1990); Y. Okada, M. Yamaguchi, T. Yanagida, Prog. Theor.
 Phys. 85, 1(1991); Phys. Lett. B262, 54(1991); H. E. Haber and R. Hempfling, Phys. Rev. Lett. 66,

- 1815(1991); Phys. Rev. **D48**, 4280(1993); J. Ellis, G. Ridolfi, Z. Zwirner, Phys. Lett. **B257**, 83(1991); R. Barbieri, M. Frigeni, F. Caravaglios, Phys. Lett. **B258**, 167(1991); J. R. Espinosa, M. Quiros, Phys. Lett. **B266**, 389(1991); J. L. Lopez, D. V. Nanopoulos, Phys. Lett. **B266**, 397(1991); M. Carena, K. Sasaki, C. E. M. Wagner, Nucl. Phys. **B381**, 66(1992); P. H. Chankowski, S. Pokorski, J. Rosiek, Phys. Lett. **B281**, 100(1992); Nucl. Phys. **B423**, 437(1994); D. M. Pierce, A. Papadopoulos, S. B. Johnson, Phys. Rev. Lett. **B68**, 3678(1992); A. Brignole, Phys. Lett. **B281**, 284(1991); V. Barger, M. S. Berger, Ohmann, Phys. Rev. **D49**, 4908(1994); G. L. Kane, C. Kolda, L. Roszkowksi, J. D. Wells, Phys. Rev. **D49**, 6173(1994); R. Hempfling, A. H. Hoang, Phys. Lett. **B331**, 99(1994); P. Langacker, N. Polonsky, Phys. Rev. **D50**, 2199(1994); M. Carena, J. R. Espinosa, M. Quirós, C. E. M. Wagner, Phys. Lett. **B355**, 209(1995).
- M. Brhlik, G. J. Good, G. L. Kane, Phys. Rev. D59, 115004(1999).
- [6] E. Commins et.al., Phys. Rev. **A50**, 2960(1994).
- [7] P. G. Harris *et.al.*, Phys. Rev. Lett. **82**, 904(1999).
- [8] S. K. Lamoreaux et. al., Phys. Rev. Lett. 57, 3125(1986).
- [9] T. Ibrahim, P. Nath, Phys. Lett. B418, 98(1998); Phys. Rev. D58, 111301(1998); ibid. 61, 093004(2000); A. Bartl, T. Gajdosik, W. Porod, P. Stockinger, H. Stremnitzer, ibid. 60, 073003(1999); S. Pokorski, J. Rosiek, C. A. Savoy, Nucl. Phys. B570, 8(2000); E. Accomando, R. Arnowitt, B. Dutta, Phys. Rev. D61, 115003(2000); U. Chattopadhyay, T. ibrahim, D. P. Roy, ibid. 64, 013004(2001); M. Brhlik, L. Everett, G. Kane, J. Lykken, Phys. Rev. Lett. 83, 2124(1999); Phys. Rev. D62, 035005(2000); E. Accomando, R. Arnowitt, B. Dutta, ibid. 61, 075010(2000); A. Bartl, W. Majerotto, W. Porod, D. Wyler, hep-ph/0306050.
- [10] D. Chang, W.-Y. Keung, A. Pilaftsis, Phys. Rev. Lett. 82, 900(1999); Phys. Rev. Lett. 83, 3972(1999)(E); A.Pilaftsis, Phys. Lett. B471, 174(1999); D. Chang, W.-F. Chang, W.-Y. Keung, Phys. Lett. B478, 239(2000); A.Pilaftsis, Nucl. Phys. B644, 263(2002).
- [11] A. Pilaftsis, Phys. Rev. **D58**, 096010(1998); Phys. Lett. **B435**, 88(1998).
- [12] A. Pilaftsis, C. E. M. Wagner, Nucl. Phys. **B533**, 3(1999).
- [13] M. Carena, J. Ellis, A. Pilaftsis, C. E. M. Wagner, Nucl. Phys. **B586**, 92(2000).
- [14] M. Carena, J. Ellis, A. Pilaftsis, C. E. M. Wagner, Nucl. Phys. **B625**, 345(2002).
- [15] C. J. C. Burgess, H. J. Schnitzer, Nucl. Phys. **B228**, 464(1983).
- [16] C. N. Leung, S. T. Love, S. Rao, Z. Phys. C31, 433(1986).
- [17] B. Buchmuller, D. Wyler, Nucl. Phys. **B268**, 621(1986).
- [18] K. Hagiwara, S. Ishihara, R. Szalapski, D. Zeppenfeld, Phys. Rev. D48, 2182(1993); B. Grzadkowski, Z. Hioki, K. Ohkuma, J. Wudka, hep-ph/0310159; V. Barger, T. Han, P. Langacker, B. McElrath, P. Zerwas, Phys. Rev. D67, 115001(2003).
- [19] K. Whisnant, J.-M. Yang, B.-L. Young, X. Zhang, Phys. Rev. D56, 467(1997); J.-M. Yang, B.-L. Young, ibid. 56, 5907(1997).
- [20] T. Han, T. Huang, Z.-H. Lin, J. -X. Wang, X. Zhang, Phys. Rev. D61, 015006(1999); Z.-H. Lin, T. Han, T. Huang, J. -X. Wang, X. Zhang, Phys. Rev. D65, 014008(2001).

- [21] T. Feng, T. Huang, X. Li, X. Zhang, S. Zhao, Phys. Rev. **D68**, 016004(2003).
- [22] G. Buchalla, A. J. Buras, M. E. Lautenbacher, Rev. Mod. Phys. 68, 1125(1996).
- [23] For the review, see, for example: K. Aoki, Z. Hioki, R. Kawabe, M. Konuma, T. Muta, Prog. Theor. Phys. Suppl. 73, 1(1982); M. Bohm, H. Spiesberger, W. Hollik, Fortsch. Phys. 34, 687(1986); W. Hollik, *ibid.* 38, 165(1990); A. Denner, *ibid.* 41, 307(1993).
- [24] P. Langacker, hep-ph/0308145; P. Gambino, hep-ph/0311257.
- [25] J. Gounaris, D. T. Papadamou, F. M. Renard, Z. Phys. C76, 333(1997).




Figure 1: The Feynman diagrams inducing nontrivial contributions to the Wilson coefficients of the operators $\mathcal{O}_{tq\Phi 1}$ and $\mathcal{O}_{tq\Phi 6}$




Figure 2: The Feynman diagrams inducing nontrivial contribution to the Wilson coefficients of the operators $\mathcal{O}_{tq\Phi i}~(i=2,~\cdots,~5,~7~\cdots,~10)$ in the full theory.




Figure 3: The Feynman diagram corresponds to the first diagram of Fig. 2 in the effective theory.




Figure 4: The Higgs self-energy diagrams which induce nonzero contributions to the Wilson coefficients of operators Eq. (12) and Eq. (13).




Figure 5: The Feynman diagrams which induce nontrivial contributions to the Wilson coefficients of the operators $\mathcal{O}_{t\Phi i}$ $(i=1,\ 2,\ 3).$




Figure 6: The one-particle-irreducible (1PI) Feynman diagrams which are related to the Wilson coefficients of the operators $\mathcal{O}_{t\Phi i}$ $(i=1,\ 2,\ 3)$ in the full theory, the gray bulbs represent the diagrams of Fig. 2.




Figure 7: The 1PI Feynman diagrams related to the Wilson coefficients of the operators $\mathcal{O}_{t\Phi i}$ $(i=1,\ 2,\ 3)$ in (a) the full theory, (b) the effective theory.





Figure 8: The Feynman diagrams which induce nontrivial contributions to the Wilson coefficients of the operators $\mathcal{O}_{q\Phi i}$ $(i=1,\ \cdots,\ 6)$.






Figure 9: The constraint from the anomalous coupling $\mathcal{O}_{q\Phi1,2}$ set by the R_b experimental data with 1σ tolerance,on the soft breaking parameters $m_Q=m_U=m_D$ versus the μ parameter in the superpotential with $m_1=m_2=m_3=500$ GeV, $A_t=A_b=100$ GeV, $m_H=500$ GeV and (a) $\tan\beta=2$; (b) $\tan\beta=40$.




Figure 10: Relaxing the lower bound to -5×10^{-5} and keeping the upper bound unchanged as in Eq.(40), the constraint from the anomalous coupling $\mathcal{O}_{q\Phi1,2}$ on the soft breaking parameters $m_Q=m_U=m_D$ versus the μ parameter in the superpotential with $m_1=m_2=m_3=500$ GeV, $A_t=A_b=100$ GeV, $m_H=500$ GeV and (a) $\tan\beta=2$; (b) $\tan\beta=40$.




Figure 11: Taking $\mu=m_i=500$ GeV $(i=1,~2,~3),~A_t=A_b=100$ GeV, $m_H=500$ GeV, ΔR_b versus squark masses $m_Q=m_U=m_D$ with $\tan\beta=2$ (solid line) or $\tan\beta=40$ (dashed line).




Figure 12: ΔR_b versus the parameter μ with $m_i=500$ GeV ($i=1,\ 2,\ 3$), $A_t=A_b=100$ GeV, $m_H=500$ GeV, and (a) solid-line: $m_Q=m_U=m_D=1$ TeV, $\tan\beta=2$, (b) dashed-line: $m_Q=m_U=m_D=1$ TeV, $\tan\beta=40$, (c) dot-line: $m_Q=m_U=m_D=500$ GeV, $\tan\beta=2$, (d) dot-dashed-line: $m_Q=m_U=m_D=500$ GeV, $\tan\beta=40$.




Figure 13: Under the assumption $m_Q=m_U=m_D=|\mu|,~\Delta R_b$ versus the parameter μ . The other parameters are taken as $m_i=500~{\rm GeV}$ (i=1,~2,~3), $A_t=A_b=100~{\rm GeV},~m_H=500~{\rm GeV}$ and $\tan\beta=2$ (solid-line) $\tan\beta=40$ (dashed-line).