International Journal of Modern Physics A © World Scientific Publishing Company

NUCLEAR SHADOWING AND DIFFRACTION

J. L. ALBACETE

Departamento de Física, Módulo C2, Planta baja, Campus de Rabanales, Universidad de Córdoba, 14071 Córdoba, Spain

N. ARMESTO AND C. A. SALGADO

Department of Physics, CERN, Theory Division, CH-1211 Genève 23, Switzerland

A. CAPELLA

Laboratoire de Physique Théorique, Université de Paris XI, Bâtiment 210, F-91405 Orsay Cedex, France

A. B. KAIDALOV

Institute of Theoretical and Experimental Physics, B. Cheremushkinskaya 25, Moscow 117259, Russia

> Received (Day Month Year) Revised (Day Month Year)

The relation between diffraction in lepton-proton collisions and shadowing of nuclear structure functions which arises from Gribov inelastic shadowing, is described. A model realizing such relation, which produces a parameter-free description of experimental data on nuclear structure functions at small x, is presented. The application to the description of multiplicities in nuclear collisions is discussed and related to other approaches.

Keywords: Nuclear structure functions; small x; diffraction.

1. Introduction and formalism

The modification of parton densities inside nuclei 1,2 is one the most interesting nuclear effects. It has strong practical implications on particle production in nuclear collisions and also offers the possibility to constrain theoretical models. We will be interested here 3 in the region of small x, x < 0.01, relevant for high energy collisions. The observed feature in this region where isospin corrections are negligible is that $F_{2A} < AF_{2p}$ i.e. the shadowing phenomenon. Explanations of this shadowing range from leading twist to all twist effects. Here (see 3 for longer discussions, model description, a more extensive comparison with experimental data and other models, and full references) we examine the possibility of describing the small x data on nuclear structure functions by the relation between diffraction and nuclear shadowing which comes from inelastic shadowing 4,5,6 .

2 J. L. Albacete, N. Armesto, A. Capella, A. B. Kaidalov and C. A. Salgado

Reggeon Field Theory 7 and its extension nuclei 4,5 are pre-QCD models in the form of field theories, which have been very successful in describing experimental data on hadronic collisions for the last 40 years. In Fig. 1 we show the relation between the diagrams describing diffraction in γ^* -nucleon collisions and double rescattering of the γ^* on a nucleus. In such relation the cut through an intermediate state of the hadronic component of the γ^* with arbitrary mass M^2 occurs, the so-called Gribov inelastic shadowing 4,5 . Through the AGK cutting rules 6 the contribution of the cut between the amplitudes exchanged with the nucleus is minus the total 2-scattering contribution,

$$\sigma_A^{(2)} = -4\pi A(A-1) \int d^2b \ T_A^2(b) \int_{M_{min}^2}^{M_{max}^2} dM^2 \ \frac{d\sigma^{\mathcal{D}}}{dM^2 dt} \bigg|_{t=0} F_A^2(t_{min}), \tag{1}$$

with $M_{min}^2 = 4m_{\pi}^2 = 0.08 \text{ GeV}^2$, M_{max}^2 fixed by $x_P < 0.1$, $T_A(b)$ the nuclear profile function normalized to 1 and coherence effects, important for not so small $x > (R_A m_N)^{-1}$, included in

$$F_A(t_{min}) = \int d^2b J_0(b\sqrt{-t_{min}})T_A(b), \quad t_{min} = -m_N^2 x_P^2.$$
 (2)




Fig. 1. Link between the diffractive cut and double scattering which produces the first correction (nuclear shadowing) to the additivity of cross section on a nucleus.

Higher order rescatterings are model-dependent 8 . To estimate such uncertainty we use two models:

$$\frac{\sigma_{\gamma^*A}^{Schw}(b)}{A\sigma_{\gamma^*N}} = \frac{T_A(b)}{1 + (A-1)f(x,Q^2)T_A(b)} > \frac{\sigma_{\gamma^*A}^{eik}(b)}{A\sigma_{\gamma^*N}} = \frac{1 - e^{-2(A-1)f(x,Q^2)T_A(b)}}{2(A-1)f(x,Q^2)}, (3)$$

with $f(x,Q^2)$ chosen to match the two scattering result.

For the diffractive cross section on the nucleon $d\sigma^{\mathcal{D}}/dM^2dt|_{t=0}$, we use a model 9 which includes as main ingredients a separation between short distance (modeled in the dipole model) and long distance (modeled by a pomeron + 1 lower Regge trajectory) contributions, with separation $r_0 = 0.2$ fm, and a triple pomeron contribution required to describe diffraction. With 9 fitted parameters it provides a

good description of F_2 and F_{2D} for the proton at x < 0.01, $Q^2 < 10$ GeV². The t-dependence of the diffractive cross section is taken $\propto \exp(Bt)$ with $B=6~{\rm GeV^{-2}}$.

2. Nuclear structure functions

Within the explained formalism we obtain a parameter-free description of shadowing in nuclear structure functions. In Fig. 2 the results of the model for the ratios $R(A/B) = (B\sigma_{\gamma^*A})/(A\sigma_{\gamma^*B})$ are compared with experimental data for C/D, Ca/D, Pb/D ¹⁰ and Xe/D ¹¹. Both results, joined by straight lines, and data have been obtained for a different Q^2 at each value of x. Taking into account the absence of free parameters, the agreement can be considered as quite good. In Fig. 3 results




Fig. 2. Results of the model using Schwimmer (Schw, solid lines) and eikonal (eik, dashed lines) unitarization compared with experimental data versus x, for the ratios C/D, Ca/D, Pb/D and Xe/D by the E665 Collaboration (filled circles correspond to the analysis with hadron requirement and open circles to that with electromagnetic cuts, see the experimental paper for more details).

of the model are compared with data 12 on the Q^2 evolution of the ratio Sn/C at fixed x. The model shows too flat a behavior, which may point to the lack of perturbative (DGLAP) evolution to be eventually applied for Q^2 of the order or larger than a few GeV^2 . Predictions for smaller x which may be relevant for future lepton-ion colliders can be found in ³ together with a comparison among available models, which could be verified through a measurement of R(A/B) with $\sim 10\%$ precision at $x \sim 10^{-4}$, $Q^2 \sim 2 \text{ GeV}^2$.

4 J. L. Albacete, N. Armesto, A. Capella, A. B. Kaidalov and C. A. Salgado




Fig. 3. Results of the model using Schwimmer (solid lines) and eikonal (dashed lines) unitarization compared with experimental data versus Q^2 , for the ratio Sn/C by the NMC Collaboration at two fixed small values of x.

3. Multiplicities

Using the AGK rules 6 it can be shown (Fig. 4) within this kind of models that multiplicities are proportional to the number of nucleon-nucleon collisions $ABT_{AB}(b)/\sigma_{AB}(b)$ times a reduction factor 13,14




Fig. 4. Diagram showing the contribution to particle production in AB collisions.

$$R_{AB}(b, y^*) = \int d^2s \frac{R_A(\vec{s}, y^*) R_B(\vec{s} - \vec{b}, y^*)}{T_{AB}(b)}, \tag{4}$$

with $T_{AB}(b) = \int d^2s \, T_A(\vec{s}) T_B(\vec{s} - \vec{b}), \, R_A(s, y^*) = T_A(s)/[1 + (A-1)f(y^*, Q^2)T_A(s)], \, R_A \propto A^{-1/3}$. Function $f(y^*, Q^2)$ can be computed in several ways 3 : It can be taken from the model described in the previous section but with integration limits inspired by the parton model: for projectile A (target B), $x_{A(B)} = m_T e^{\pm y^*}/\sqrt{s}$ with $y^* > 0$ for the projectile hemisphere and $y^* < 0$ for the target one, $M_{max}^{2(A(B))} = Q^2 \left(x_{Pmax}/x_{A(B)} - 1\right) = Q^2 \left(x_{Pmax}\sqrt{s}\,e^{\mp y^*}/m_T - 1\right), \, M_{min}^2$ fixed and equal to $0.08 \, \text{GeV}^2$, and $Q^2 = m_T^2 = 0.4 \, \text{GeV}^2$. On the other hand, the reduction factor can be computed as in 14 :

$$f(s, y^*) = 4\pi \int_{y_{min}}^{y_{max}} dy \, \frac{1}{\sigma_P(s)} \, \frac{d\sigma^{PPP}}{dydt} \bigg|_{t=0} F_A^2(t_{min}),$$
 (5)

with $y_{min}^{(A(B))} = \ln{(s/M_{max}^{2(A(B))})}$ and $y_{max}^{(A(B))} = \ln{(s/m_T^2)}/2 \mp y^*$. $\sigma_P(s)$ $(d\sigma^{PPP}/dydt)$ is the single (triple) Pomeron cross section with parameters ³ taken from ¹⁵. Fig. 5 shows predictions for different symmetric nucleus-nucleus collisions at $y^* = 0$. The reduction factor (4) is needed ^{16,17,18} to agree with experimental data.




Fig. 5. Results of the model for the multiplicity reduction factor versus impact parameter b at $y^* = 0$, for AuAu collisions at $\sqrt{s} = 19$, 130 and 200 GeV per nucleon, and for PbPb collisions at $\sqrt{s} = 5500$ GeV per nucleon, in the parton model-like realization (solid lines) and for a triple pomeron contribution alone (dashed lines), see the text for explanations.

6 J. L. Albacete, N. Armesto, A. Capella, A. B. Kaidalov and C. A. Salgado

4. Comments

In this contribution we have explored how lepton-nucleon scattering can be related to lepton-nucleus and nucleus-nucleus through Gribov inelastic shadowing within the framework of Glauber-Gribov theory. Other relations have been found ¹⁹ within saturation physics. The model we have presented shows some similarities to saturation ideas. Shadowing is taken into account by similar diagrams. A factorization formula analogous to (4) has been shown ²⁰ to hold in saturation for gluon production in proton-nucleus collisions. It is also usually employed for nucleus-nucleus e.g. in ²¹, but in this case corrections to factorization have been found ^{22,23}. Studies at higher energies on the centrality dependence of multiplicities at fixed and integrated transverse momentum will help to discriminate between available approaches ²⁴.

Acknowledgments

N.A. thanks the organizers for their invitation to present this talk in such a nice meeting.

References

- 1. M. Arneodo, Phys. Rept. 240, 301 (1994).
- D. F. Geesaman, K. Saito and A. W. Thomas, Ann. Rev. Nucl. Part. Sci. 45, 337 (1995).
- N. Armesto, A. Capella, A. B. Kaidalov, J. Lopez-Albacete and C. A. Salgado, Eur. Phys. J. C29, 531 (2003).
- 4. V. N. Gribov, Sov. Phys. JETP 29, 483 (1969) [Zh. Eksp. Teor. Fiz. 56, 892 (1969)].
- V. N. Gribov, Sov. Phys. JETP 30, 709 (1970) [Zh. Eksp. Teor. Fiz. 57, 1306 (1969)].
- V. A. Abramovsky, V. N. Gribov and O. V. Kancheli, Yad. Fiz. 18, 595 (1973) [Sov. J. Nucl. Phys. 18, 308 (1974)].
- 7. V. N. Gribov, Sov. Phys. JETP 26, 414 (1968) [Zh. Eksp. Teor. Fiz. 53, 654 (1967)].
- 8. B. Z. Kopeliovich, J. Raufeisen and A. V. Tarasov, Phys. Rev. C62, 035204 (2000).
- A. Capella, E. G. Ferreiro, C. A. Salgado and A. B. Kaidalov, *Phys. Rev.* D63, 054010 (2001).
- 10. M. R. Adams et al. [E665 Collaboration], Z. Phys. C67, 403 (1995).
- 11. M. R. Adams et al. [E665 Collaboration], Phys. Rev. Lett. 68, 3266 (1992).
- 12. M. Arneodo et al. [New Muon Collaboration], Nucl. Phys. B481, 23 (1996).
- 13. L. Caneschi, A. Schwimmer and R. Jengo, Nucl. Phys. B108, 82 (1976).
- 14. A. Capella, A. Kaidalov and J. Tran Thanh Van, Heavy Ion Phys. 9, 169 (1999).
- A. Capella, A. Kaidalov, C. Merino, D. Pertermann and J. Tran Thanh Van, Eur. Phys. J. C5, 111 (1998).
- 16. A. Capella and D. Sousa, Phys. Lett. **B511**, 185 (2001).
- 17. A. Capella, these proceedings.
- 18. E. G. Ferreiro, arXiv:hep-ph/0409287.
- 19. N. Armesto, C. A. Salgado and U. A. Wiedemann, arXiv:hep-ph/0407018.
- 20. Y. V. Kovchegov and K. Tuchin, Phys. Rev. D65, 074026 (2002).
- 21. D. Kharzeev and E. Levin, Phys. Lett. **B523**, 79 (2001).
- 22. A. Krasnitz and R. Venugopalan, Nucl. Phys. **B557**, 237 (1999).
- 23. I. Balitsky, arXiv:hep-ph/0409314.
- 24. N. Armesto and C. Pajares, Int. J. Mod. Phys. A15, 2019 (2000).

