The Updated MSW Analysis and the Standard Solar Model Uncertainties *

Naoya Hata and Paul Langacker

Department of Physics, University of Pennsylvania, Philadelphia, PA 19104 (August 6, 1993, UPR-0581T)

Abstract

We update the analysis of the MSW and general astrophysical solutions to the combined solar neutrino observations by including the GALLEX II result. We also show that our parametrized flux uncertainties are equivalent to the Monte-Carlo results of Bahcall and Ulrich.

Typeset using REVT_EX

^{*}This is an update of a talk presented at SUSY93, Northeastern University, Boston, MA, April 1993.

I. GLOBAL ANALYSIS UPDATED

A. MSW

By including the GALLEX II data [1], we update our previous MSW analysis [2,3] which includes a joint χ^2 analysis of all experiments, a full treatment of theoretical uncertainties including correlations between experiments, and the Earth effect. (See also [4–8].) The experimental data used in the analysis are summarized in Table I. The combined result of the Homestake [9], gallium [10,11,1], and Kamiokande (including the day-night data) [12] experiments are shown in Fig. 1. The allowed large-angle region for $\Delta m^2 \sim 10^{-5} \text{eV}^2$ has been reduced significantly by including the new GALLEX result. The allowed regions ¹ at 90, 95, and 99% C.L. are shown in Fig. 2; there is a third allowed solution in the large-angle region at 99% C.L for $\Delta m^2 \sim 10^{-7} \text{eV}^2$. In Fig. 3 we show the combined result including the Earth effect, but without the Kamiokande day-night data.

B. General Astrophysical Solution

When the solar neutrino data are fit to a general solar model [14], using the pp, CNO, ⁷Be, and ⁸B fluxes as free parameters with the luminosity constraint imposed, the best fit value requires $\phi(\text{Be})/\phi(\text{Be})_{SSM} < 0.08$ and $\phi(\text{B})/\phi(\text{B})_{SSM} = 0.37 \pm 0.04$, but with a large χ^2 (= 5.1 for 1 degrees of freedom); this solution is excluded at 98% C.L.

¹ We define the 90, 95, and 99% C.L. regions by increases of 4.6, 6.0, and 9.2 in the overall χ^2 with respect to the global minimum. This prescription is only rigorously valid when the probability distribution in the parameter space is gaussian around the global minimum. Improved treatments will be discussed in a subsequent paper [13].

II. THE PARAMETRIZED SSM UNCERTAINTIES

In the MSW analysis it is important to include the Standard Solar Model (SSM) flux uncertainties properly, since their magnitudes are comparable to the experimental uncertainties and also strongly correlated from experiment to experiment. One way to incorporate those SSM uncertainties is to carry out Monte Carlo simulations, as done by Bahcall and Haxton [4], which is robust but time consuming. Alternatively, we parameterize the flux uncertainties with the uncertainties of the central temperature and of the the relevant nuclear reaction cross sections. This method was, however, questioned by Bahcall on the grounds that this simplification might fail in estimating the uncertainties accurately [15]. We have numerically compared the two methods and conclude that they yields almost identical results [13].

First we have compared the uncertainties of the major fluxes $(pp, {}^{7}\text{Be}, \text{ and }{}^{8}\text{B})$ and their correlations. The results are listed in Table II for the magnitudes and Table III for the correlations. The results are also graphically shown in Fig. 6. The numerical results show that the parameterization method accurately reproduces the Monte-Carlo flux uncertainties.

We have also carried out the comparison for the uncertainties of the rate predictions for the experiments. The magnitudes and the correlations of the uncertainties are compared in Table IV and Table V, respectively. Again the agreement is excellent.

REFERENCES

- P. Anselmann *et al.*, GALLEX collaboration Report No. GX 27a–1993 (to be published in Phys. Lett. B).
- [2] S. A. Bludman, N. Hata, D. C. Kennedy, and P. G. Langacker, Phys. Rev. D 47, 2220 (1993).
- [3] N. Hata and P. Langacker, University of Pennsylvania preprint UPR-0570T (1993).
- [4] J. N. Bahcall and W. C. Haxton, Phys. Rev. D 40, 931 (1989).
- [5] X. Shi, D. N. Schramm, and J. N. Bahcall, Phys. Rev. Lett. 69, 717 (1992); X. Shi and
 D. N. Schramm, Phys. Lett. B 283,305 (1992); Fermilab preprint 92/322-A.
- [6] J. M. Gelb, W. Kwong, and S. P. Rosen, Phys. Rev. Lett. 69, 1864 (1992).
- [7] P. I. Krastev and S. T. Petcov, Phys. Lett. B **299**, 99 (1993).
- [8] L. Krauss, E. Gates, and M. White, Phys. Lett. B **299**, 94 (1993).
- [9] R. Davis, Jr., et al., in Proceedings of the 21th International Cosmic Ray Conference, Vol. 12, edited by R. J. Protheroe (University of Adelaide Press, Adelaide, 1990), p. 143; R. Davis, Jr., in International Symposium on Neutrino Astrophysics, Takayama, Kamioka, 1992 (unpublished).
- [10] A. I. Abazov, et al., Phys. Rev. Lett. 67, 3332 (1991); V. N. Gavlin, in XXVI International Conference on High Energy Physics, Dallas, 1992, edited by J. Sanford (AIP, New York, 1993).
- [11] GALLEX Collaboration, P. Anselmann *et al.*, Phys. Lett. B 285, 376 (1992); 285, 390 (1992).
- [12] K. S. Hirata *et al.*, Phys. Rev. Lett. **65**, 1297 (1990); **65**, 1301(1990); **66**, 9 (1991);
 Phys. Rev. D **44**, 2241 (1991). Y. Suzuki, in *International Symposium on Neutrino Astrophysics*, Takayama, Kamioka, 1992 (unpublished).

- [13] N. Hata and P. Langacker, to be published.
- [14] S. Bludman, N. Hata, and P. Langacker, University of Pennsylvania preprint, UPR-0572T (1993).
- [15] J. N. Bahcall, Institute of Advanced Study preprint IASSNS 92/54 (1992) (To be published in the Proceedings of the XXVI International Conference on High Energy Physics, Dallas, Texas (1992).)
- [16] J. N. Bahcall and M. H. Pinsonneault, Rev. Mod. Phys. 64, 885 (1992).
- [17] S. Turck-Chièze and I. Lopes, Astrophys. J. 408, 347 (1993). S. Turck-Chièze, S. Cahen,
 M. Cassé, and C. Doom, Astrophys. J. 335, 415 (1988).
- [18] J. N. Bahcall and R. N. Ulrich, Rev. Mod. Phys. 60, 297 (1988); J. N. Bahcall, Neutrino Astrophysics, (Cambridge University Press, Cambridge, England, 1989).

TABLES

TABLE I. The standard solar model predictions of Bahcall and Pinsonneault [16] and of Turck-Chiéze and Lopes [17], along with the results of the solar neutrino experiments. The gallium experiment is the combined result of SAGE (58^{+22}_{-27} SNU) and GALLEX I and II (87 ± 16 SNU).

	BP SSM	TCL SSM	Experiments
Kamiokande	1 ± 0.14	$0.77 {\pm} 0.19$	0.50 ± 0.07 BP-SSM
Homestake (Cl)	8 ± 1 SNU	$6.4{\pm}1.4$ SNU	2.23 ± 0.23 SNU (0.28 ± 0.03 BP-SSM)
SAGE & GALLEX (Ga)	131.5^{+7}_{-6} SNU	122.5 ± 7 SNU	$77{\pm}13$ SNU (0.59{\pm}0.10 BP-SSM)

TABLE II. The magnitudes of flux uncertainties $(\Delta \phi / \phi \text{ at } 1\sigma)$ of the Bahcall-Ulrich SSM [18], Bahcall-Ulrich Monte Carlo SSMs, and the parametrized SSM using the central temperature and the nuclear reaction cross sections. Also listed are the uncertainties of the Bahcall-Pinsonneault SSM and of its parametrized fluxes. (The Monte Carlo study of the Bahcall-Pinsonneault model is not available.)

	pp	$^{7}\mathrm{Be}$	⁸ B	
Bahcall-Ulrich SSM	0.0059	0.050	0.12	
Bahcall-Ulrich SSM (Monte Carlo)	0.0067	0.05	0.17	
Parametrized ($\Delta T_C = 0.0053$)	0.0068	0.05	0.12	
Bahcall-Pinsonneault SSM	0.0067	0.06	0.14	
Parametrized ($\Delta T_C = 0.0057$)	0.007	0.06	0.14	

TABLE III. The correlation matrices of flux uncertainties obtained from the Bahcall-Ulrich Monte Carlo SSMs and of the parameterization method. The agreement between the two methods is excellent except the $pp - {}^{7}$ Be element. This can be improved by anticorrelating the uncertainties from the S_{11} for the two fluxes, which is a consequence of the luminosity constraint. The effect of anticorrelating S_{11} is completely negligible in the MSW fits, and we present the MSW results without the anticorrelation.

	pp	⁸ B	$^{7}\mathrm{Be}$
Bahcall-Ulrich SSM (Monte C	'arlo)		
pp	1		
⁸ B	-0.726	1	
⁷ Be	-0.917	0.742	1
Parametrized with ΔT_C and Δ	Δs		
pp	1		
⁸ B	-0.679	1	
⁷ Be	-0.677	0.773	1
Parametrized with ΔT_C and Δ	Δs (S_{11} is anticorrelated	between pp and ^{7}Be)	
pp	1		
⁸ B	-0.679	1	
⁷ Be	-0.916	0.773	1

TABLE IV. The comparison of the magnitudes of rate uncertainties for the Bahcall-Ulrich SSM, Monte Carlo SSMs, and the parametrized SSM. Also listed are the uncertainties in the Bahcall-Pinsonneault SSM and its parametrized SSM.

	Kamiokande	Cl	Ga
Bahcall-Ulrich SSM	0.12	0.11	+0.05 - 0.04
Monte Carlo	0.12	0.11	0.05
Parametrized ($\Delta T_C = 0.0053$)	0.12	0.12	0.05
Bahcall-Pinsonneault SSM	0.14	0.13	0.05
Parametrized ($\Delta T_C = 0.0057$)	0.14	0.13	0.05

TABLE V. The comparison of the correlations of rate uncertainties for Monte Carlo SSMs and the parametrized SSM. Also listed is the parametrized SSM with the S_{11} uncertainty anticorrelated between the pp and ⁷Be fluxes.

Kamiokande		Cl	Ga
Bahcall-Ulrich	h SSM (Monte Carlo)		
Kamiokande	1		
Cl	-0.997	1	
Ga	-0.920	0.947	1
Parametrized	with ΔT_C and Δs		
Kamiokande	1		
Cl	-0.995	1	
Ga	-0.888	0.928	1
Parametrized	with ΔT_C and Δs , with	S_{11} anticorrelated between pp and ⁷ Be	
Kamiokande	1		
Cl	-0.995	1	
Ga	-0.898	0.936	1

FIGURES

FIG. 1. The allowed regions of the combined Homestake, Kamiokande, and gallium observations including the Kamiokande day-night data. The region excluded by the day-night data is also shown (dotted line).

FIG. 2. The allowed regions of the combined fit at 90, 95, and 99% C.L. There is a third solution in the large-angle region for $\Delta m^2 \sim 10^{-7} \text{eV}^2$ at 99% C.L.

FIG. 3. The allowed region with the Earth effect, but without the Kamiokande day-night data.

FIG. 4. The distribution of the pp, ⁷Be, and ⁸B flux of the Bahcall-Ulrich Monte Carlo SSMs (histograms) and the parametrized method (solid curves) that assumes Gaussian distributions of the central temperatures and the nuclear reaction cross sections around their central values.

FIG. 5. The distributions of the expected rate for the Kamiokande, chlorine, and gallium experiments obtained from Bahcall-Ulrich Monte Carlo fluxes (histograms) and the parametrized fluxes (solid curve). In both cases the detector cross section uncertainties are not included. (They are included in the analysis of the MSW and of general astrophysical solutions.)

FIG. 6. The distributions of the ⁷Be and ⁸B flux of the Bahcall-Ulrich SSMs (dots) and the 90% C.L. contour of the parametrized SSM (solid curve). The magnitudes and the correlations are the same for the two methods.









BA