SOFT FIELDS AND HARD PARTICLES IN HOT GAUGE PLASMAS

Edmond IANCU Service de Physique Théorique ¹, CE-Saclay 91191 Gif-sur-Yvette, France

October 1993

Abstract

It is shown that the long wavelength excitations of a quark-gluon plasma may be described as collective oscillations of self-consistent average fields to which the plasma particles couple. Their properties are obtained from a set of coupled mean field and kinetic equations, derived from the general Dyson-Schwinger equations, as the leading order in a systematic expansion in powers of the coupling. By solving the kinetic equations, one obtains in closed form the generating functionals for all the leading order amplitudes between soft quasiparticles, the so-called "hard thermal loops".

Talk given at the 3rd Workshop on Thermal Field Theories and their Applications, August 15–27, 1993, Banff, Alberta, Canada

¹Laboratoire de la Direction des Sciences de la Matière du Commissariat à l'Energie Atomique

SOFT FIELDS AND HARD PARTICLES IN HOT GAUGE PLASMAS

Edmond IANCU Service de Physique Théorique †, CE-Saclay 91191 Gif-sur-Yvette, France

ABSTRACT

It is shown that the long wavelength excitations of a quark-gluon plasma may be described as collective oscillations of self-consistent average fields to which the plasma particles couple. Their properties are obtained from a set of coupled mean field and kinetic equations, derived from the general Dyson-Schwinger equations, as the leading order in a systematic expansion in powers of the coupling. By solving the kinetic equations, one obtains in closed form the generating functionals for all the leading order amplitudes between soft quasiparticles, the so-called "hard thermal loops".

1. Introduction

The present talk is concerned with some recent progress in the understanding of the collective modes propagating in a gauge plasma at high temperature. The results to be reported here offer a consistent and gauge covariant description for the off-equilibrium dynamics of a quark-gluon (or electron-photon) plasma, valid at leading order in the gauge coupling constant g. (I consider QCD at a temperature T well into its deconfined phase and assume a perturbative regime, $g \ll 1$.) Such a description may be viewed as a first step towards a more complete treatement of the plasma transport and relaxation properties. Besides, as we shall see, it is also relevant for a physical understanding of the newly developed resummation schemes, allowing for consistent computations in hot gauge theories¹.

As it is clear by now, in high-T field theory the naive loop expansion fails. As shown by Braaten and Pisarski¹, one-loop corrections to amplitudes with soft external momenta $P_1, ..., P_n$, (i.e., such that $P_j \lesssim gT$ for any j) are as important as the corresponding tree level amplitudes and, correspondingly, have to be resummed in consistent higher order computations¹. The dominant one-loop contributions are called "hard thermal loops" (HTL), as they arise from integrating over hard internal momenta k, (i.e., over k of order T). Isolated first in the 2-point amplitudes², and, later on, in general n-point amplitudes^{3,1} ($n \ge 2$), the HTL's show some outstanding properties, which are left largely unexplained by their original derivation in terms of Feynman diagrams. I summarize here some of them:

- (a) There are some systematic "selection" rules in the appearance of the HTL's:
 - no HTL with more then one pair of fermion lines;
 - in QED: no HTL with more then 2 external photons;

[†]Laboratoire de la Direction des Sciences de la Matière du Commissariat à l'Energie Atomique

- in QCD: HTL's in all multi-gluon amplitudes;
 no HTL with ghost external lines.
- (b) The HTL's present a simple structure. E.g., the HTL for the gluon polarization tensor is identical to the dielectric tensor for a *classical* ultrarelativistic plasma, as already computed by Silin in 1960 from kinetic theory⁷.
- (c) HTL's are generally momentum dependent through "Vlasov-like" (or "eikonal-like") denominators, of the type $1/(v \cdot P)$, where $v^{\mu} = (1, \vec{v})$ is the loop velocity, while P is a linear combination of the external momenta.
 - (d) HTL's are independent of the gauge fixing condition used in their evaluation.
- (e) HTL's obey "abelian-like" Ward identities even in the case of QCD. This is equivalent to a statement of gauge invariance and suggested the construction of an effective action generating all the HTL's^{4,5}. This is a nonlocal functional of the soft fields, whose properties are further investigated in various recent works⁶.
- (f) As already notified, the HTL's have to be resummed, for consistency, into (non-local) effective propagators and vertices. By using them in perturbative computations, one has obtained gauge independent results for the damping of the collective modes¹, as well as sensible values for some transport coefficients⁸.

The need for resummation in the perturbative expansion, as well as the simple properties above, suggest some *simple underlying physics*, whose understanding was the main motivation of the present work. To this aim, a new approach to the problem has been initiated by Jean-Paul Blaizot and myself, by working directly at the level of the equations of motion rather than on the Feynman diagrams⁹. Our results, to be discussed below, may be summarized in the following *physical picture*:

The HTL's describe polarization properties related to the "semiclassical" collective dynamics of the hard particles in selfconsistent mean fields. Resummation is necessary to properly account for the collective character of the soft excitations in higher orders of perturbation theory.

In order to see how this picture emerges, let me start by recalling some typical energy and length scales in the ultrarelativistic plasma. In thermal equilibrium, at high T, quarks and gluons have typical energies and momenta of order T, and will be referred to as "hard" particles. Their number density is proportional to T^3 , and therefore the typical interparticle distance r_0 is $\sim 1/T$. When coupled to slowly varying disturbances, the plasma may develop a collective behaviour on a typical length/time scale of order 1/gT. A familiar example is Debye's screening. Collectivity arises because any motion taking place over a distance scale $\lambda \sim 1/gT$ may involve coherently a large number of hard particles. This gives rise to long range fluctuations in the densities of the associated quantum numbers (like charge, color, spin, baryonic nb.), which are conveniently described in terms of long wavelength oscillations in the selfconsistent mean fields carrying the appropriate quantum nbs.: $\langle A_{\mu}(X) \rangle$, $\langle \psi(X) \rangle$, $\langle \bar{\psi}(X) \rangle$, ... Such fields, to be referred as "soft" in what follows $(\lambda \sim 1/gT)$, represent the average motion of the hard particles over distances large compared to their mean separation $(\lambda \gg r_0)$.

Our main objective is to characterize the properties of the collective modes (i.e., their propagation and mutual interactions) at leading order in g. As we shall see

shortly, these properties are essentially determined by the interactions between the soft fields and the hard thermal particles, which explains the title of this talk. In particular, the HTL's play a decisive role in this description, and all their remarkable properties alluded to before find in this context a natural explanation.

2. An approximation scheme with three facets

I shall consider the response of the plasma to weak and slowly varying external disturbances. When studying off-equilibrium perturbations, one is usually facing three kinds of approximations, involving, respectively, the strength of the coupling, the amplitude of the field oscillations, and the wavelength or period of the modes that one is studying. However, for the ultrarelativistic plasma, it turns out that all three approximations are controlled by the single small parameter, g. They should be viewed, then, as three facets of the same approximation, namely an expansion in powers of g. The naive loop expansion fails because it takes care only of the first facet, namely the expansion in powers of the interaction vertices. To understand how the coupling constant g controls the mean fields aspects of this problem, recall that

- (a) the mean fields are slowly varying, $\lambda \sim 1/gT$, i.e. they carry a typical momentum $P \sim gT$. Then any soft derivative carries a power of $g: i\partial \langle \phi(x) \rangle \sim gT \langle \phi(x) \rangle$, where $\langle \phi(x) \rangle$ denotes any of the average fields $\langle A_{\mu}(x) \rangle$, $\langle \psi(x) \rangle$, ...
- (b) the mean fields are weak (plasma close to equilibrium). This puts constraints on the strength of the mean fields which also depend on g. Indeed, the plasma will be weakly perturbed if, for instance, the color force $\sim g\langle F\rangle$ acting on a typical particle with momentum $k \sim T$, produces a small change in its momentum, i.e. $\Delta k = g\langle F\rangle\Delta t \ll k$; (here, $\langle F_{\mu\nu}\rangle$ is the gauge field strength tensor). One sees that, for $\Delta t \sim \lambda \sim 1/gT$ and $\Delta k \lesssim gT$, we have $\langle F\rangle \lesssim gT^2$. More generally, there exists a limit in which all terms in our final equations of motion (1)–(7) are of the same order of magnitude. This is achieved when $\langle F\rangle \sim gT^2$ and $\langle \bar{\psi}\rangle \langle \psi\rangle \sim gT^3$, i.e., g times a power of T equal to their canonical dimension. In this limit, $g\langle A_{\mu}\rangle \sim gT$ is of the same order as the derivative of a "slowly varying" quantity, $i\partial_{\mu}\langle\phi\rangle \sim g\langle A_{\mu}\rangle \langle\phi\rangle$, which insures the consistency of the soft covariant derivatives $D_{\mu} = \partial_{\mu} + ig\langle A_{\mu}\rangle$. In what follows, I shall consider this limit, where the equations keep their full nonabelian structure.

Our strategy is by now clear: we have to expand the exact (real-time) Dyson-Schwinger equations of motion in powers of g and to preserve consistently the leading, nontrivial, terms. In applying this procedure, we find that the dominant interactions which determine the response of the plasma are those which take place between the hard particles and the soft mean fields. Collisions between hard particles may be ignored in this order, which allows us to truncate the hierarchy at the level of the 2-point functions. That is, we end up with a coupled system of equations for the soft mean fields and for the hard particles 2-point functions, to be presented now.

3. Mean field and kinetic equations

The equations for the soft mean fields are

$$i \not \! D \psi(X) = \eta(X) + \eta^{ind}(X), \tag{1}$$

$$[D^{\nu}, F_{\nu\mu}(X)]^{a} - g\bar{\psi}(X)\gamma_{\mu}t^{a}\psi(X) = j_{\mu}^{a}(X) + j_{\mu}^{ind\,a}(X), \tag{2}$$

where the average gauge and fermionic fields are denoted, respectively, by A(x) and $\psi(x)$, omitting the brackets. The covariant derivative is $D_{\mu} \equiv \partial_{\mu} + igt^a A_{\mu}^a$ and $F_{\mu\nu} \equiv [D_{\mu}, D_{\nu}]/(ig)$. In the r.h.s., $\eta(X)$ and $j_{\mu}^a(X)$ are external perturbations; in particular, they may vanish, in which case Eqs. (1)-(2) describe the soft normal modes of the plasma. The "induced sources" show how the collective motion of the hard particles affects – through the associated polarization effects – the properties of the mean fields. They are generally expressed in terms of connected, off-equilibrium, 2-point functions, and have to be considered as functionals of the fields themselves. For example, $\eta^{ind} = g\gamma^{\nu}t_a\langle A_{\nu}^a(x)\psi(x)\rangle_c$ involves an abnormal quark-gluon propagator, which is nonvanishing only in the presence of the fermionic mean field ψ . The induced color current j_{μ}^{ind} receives contributions from various species of hard particles (quarks and gluons). The fermion piece involves the quark propagator in the presence of the fields: $j_{\mu}^{\mu} = g t^a \langle \bar{\psi}(x) \gamma^{\mu} t^a \psi(x) \rangle_c$. In a covariant gauge, the bosonic piece j_{ν}^{μ} involve both gluons and ghosts⁹.

The 2-point functions which enter the induced sources are obtained in terms of the mean fields by solving the appropriate eqs. of motion at the level of the present approximation (see below Eqs. (5)–(7)). It is remarkable that the corresponding solutions may be expressed in terms of on-shell generalized distribution functions for the hard thermal particles (quarks and transverse gluons). To be specific, we found that, in leading order, the induced sources can be written as

$$j_{\mu}^{ind\,a}(X) = g \int \frac{d^3k}{(2\pi)^3} \frac{k_{\mu}}{k} \left\{ N_{\rm f} \left[\delta n_+^a(\vec{k}, X) - \delta n_-^a(\vec{k}, X) \right] + 2N \, \delta N^a(\vec{k}, X) \right\}, \quad (3)$$

$$\eta^{ind}(x) = g \int \frac{d^3k}{(2\pi)^3} \frac{1}{k} \not\Lambda(\vec{k}, X), \tag{4}$$

where $N_{\rm f}$ is the number of quark flavors. Here, $\delta n_{\pm}^a(\vec{k},X)$ and $\delta N^a(\vec{k},X)$ denote oscillations induced by the soft mean fields in the color densities carried by the hard quarks, antiquarks and, respectively, gluons with momentum \vec{k} . Less familiar, $\not A(\vec{k},X)$ represents a generalized one-body density matrix mixing fermionic and bosonic degrees of freedom. It carries fermionic quantum number and describes fluctuations where, under the action of a soft fermionic mean field, hard quarks are converted into hard gluons and vice-versa.

All the fluctuations above are determined by the following kinetic equations⁹:

$$(v \cdot D_X) \Lambda(\vec{k}, X) = -i C_f (N_k + n_k) \psi \psi(X), \tag{5}$$

$$\left[v \cdot D_X, \, \delta n_{\pm}(\vec{k}, X)\right] = \mp g \, \vec{v} \cdot \vec{E}(X) \, \frac{dn_k}{dk},\tag{6}$$

$$\left[v \cdot D_X, \, \delta N(\vec{k}, X)\right] = -g \, \vec{v} \cdot \vec{E}(X) \frac{dN_k}{dk}.\tag{7}$$

Here, $v^{\mu} \equiv (1, \vec{v})$, where $\vec{v} \equiv \vec{k}/\epsilon_k$ is the velocity of the hard particle, C_f is the quark Casimir, $E^i \equiv E_a^i t^a$ is the chromoelectric field, while $\delta N \equiv \delta N^a t^a$, etc. In the r.h.s., N_k and n_k denote, respectively, equilibrium boson and fermion occupation factors. These eqs. determine the collective dynamics of the hard particles in the presence of soft background fields. They exhibit many features of classical dynamics, and, in fact, they generalize the Vlasov eq. to nonabelian plasmas (the covariant line derivative in the l.h.s. is the familiar drift term of elementary kinetic equations). Also note that i) Eqs. (5)–(7) are independent of the gauge fixing parameter which enters calculations in general covariant gauges⁹; ii) they transform covariantly under a local gauge transformation of the mean fields A_{μ} , ψ and $\bar{\psi}$; iii) they are nonlinear in A_{μ} , because of the covariant derivatives; iv) there is a remarkable symmetry between the response of the hard quarks and that of the hard gluons to soft fields.

4. From kinetic equations to HTL's

Eqs. (5)–(7) may be exactly solved for specified boundary conditions. Then, the induced sources result after a supplimentary integration over \vec{k} (recall Eqs. (3)–(4)). Consider, e.g., retarded conditions, which is appropriate for a plasma being initially in equilibrium. The corresponding solutions, which express the response of the plasma to "soft" perturbations vanishing as $X_0 \to -\infty$, may be written as

$$\eta^{ind}(X) = -i\omega_0^2 \int \frac{d\Omega}{4\pi} \psi \int_0^\infty dt \, U(X, X - vt) \psi(X - vt), \tag{8}$$

$$j_{\mu}^{ind}(X) = 3 \omega_p^2 \int \frac{d\Omega}{4\pi} v_{\mu} \int_0^{\infty} dt \, U(X, X - vt) \, \vec{v} \cdot \vec{E}(X - vt) \, U(X - vt, X), \tag{9}$$

in terms of the parallel transporter U(x,y) along a straight line γ joining x and y, $U(x,y) = P \exp\{-ig \int_{\gamma} dz^{\mu} A_{\mu}(z)\}$. The plasma frequencies are $\omega_0^2 \equiv C_f g^2 T^2/8$ and $\omega_p^2 \equiv (2N+N_{\rm f})g^2 T^2/18$; the angular integral runs over all the directions of the unit vector \vec{v} .

The induced sources (8)–(9) show how the mean fields are renormalized by their interaction with the hard particles. Accordingly, they act as generating functionals

for all (retarded) amplitudes between soft quasiparticles. E.g., the soft quark self-energy in the presence of a background gauge field A_{μ} is $\Sigma(X,Y) = \delta \eta^{ind}(X)/\delta \psi(Y)$, which for $A_{\mu} = 0$ gives precisely the corresponding HTL:

$$\Sigma(P) = \omega_0^2 \int \frac{d\Omega}{4\pi} \frac{\psi}{v \cdot P + in}.$$
 (10)

By further differentiations with respect to A_{μ} in (8), we generate all the HTL's between a quark pair and any number of soft gluons⁹. Note that η^{ind} is linear in ψ , so there is no amplitude with more than one pair of soft external fermions. Similarly, all multi-gluon HTL's arise when differentiating $j_{\mu}^{ind}(X)$. For instance, $\Pi_{\mu\nu}(X,Y) = \delta j_{\mu}^{ind}(X)/\delta A^{\nu}(Y)$ is the well-known polarization tensor for soft gluons:

$$\Pi_{\mu\nu}^{ab}(P) = 3\,\omega_p^2\,\delta^{ab} \left\{ -\delta_\mu^0 \delta_\nu^0 + P^0 \int \frac{d\Omega}{4\pi} \frac{v_\mu \, v_\nu}{v \cdot P + i\eta} \right\}. \tag{11}$$

Remark that the denominator $1/(v \cdot P)$ typical for the HTL's reflects simply the drift operator in the l.h.s. of Eqs. (5)–(7).

Gauge symmetry implies a covariant conservation law for the induced current

$$\left[D^{\mu}, j_{\mu}^{ind}(X)\right] = 0, \tag{12}$$

which leads, by differentiation with respect to the mean fields, to the simple Ward identities relating the HTL's, alluded to in the introductory section. For example,

$$\Pi_{\mu\nu} P^{\nu} = P^{\mu} \Pi_{\mu\nu} = 0,$$

$$P_1^{\mu} \Gamma_{\mu\nu\lambda}(P_1, P_2, P_3) = \Pi_{\nu\lambda}(P_3) - \Pi_{\nu\lambda}(P_2), \dots$$
(13)

where $\Gamma_{\mu\nu\lambda}$ is the HTL for the 3-gluon vertex.

5. Effective action for soft fields

By eliminating the induced sources from Eqs. (1)–(2), using their explicit expressions (8)–(9), one obtains nonlinear and nonlocal eqs. of motion which generalize the Maxwell equations in a polarizable medium. Note that the quark and gluon modes mix, as a consequence of gauge covariance.

In general, these eqs. also include dissipative effects, related to the Landau damping of the mean fields. This explains the nonvanishing imaginary parts of the self-energies (10)-(11), occurring for space-like momenta P_{μ} (i.e., when $P_0 = \vec{v} \cdot \vec{P}$). However, at the level of the present approximation, there is a well defined transparency range where dissipation is absent: this happens for time-like momenta P_{μ} , where the Landau damping is kinematically forbidden. In this range, the mean field eqs. of motion are generated by the minimal action principle applied to an effective action $S_{eff} = S_0 + S_{ind}$. Here, S_0 is the classical QCD action, while S_{ind}

contains the effects of the interactions between the soft fields and the hard particles: $S_{ind} = S_f + S_b$, with⁹

$$S_{\rm f} = \omega_0^2 \int \frac{d\Omega}{4\pi} \int d^4X \int d^4Y \,\bar{\psi}(X) \langle X | \frac{\psi}{i(v \cdot D)} | Y \rangle \psi(Y), \tag{14}$$

$$S_{\rm b} = \frac{3}{2} \omega_p^2 \int \frac{d\Omega}{4\pi} \int d^4 X \int d^4 Y \operatorname{tr} \left[F_{\mu\lambda}(X) \langle X | \frac{v^{\mu} v_{\nu}}{(v \cdot \tilde{D})^2} | Y \rangle F^{\nu\lambda}(Y) \right], \quad (15)$$

where $\tilde{D} F_{\mu\nu} \equiv [D, F_{\mu\nu}]$, and the trace acts on color indices. Note that for undamped fields, the operator $v \cdot D$ never vanishes, and thus the inverse operators entering (14)–(15) are nonambiguous⁹. The action above coincides with the generating functional for HTL's derived in Refs.^{4,5} on the basis of gauge invariance. One sees from our analysis that S_{eff} may be given the physical interpretation of the classical action which describes long wavelength excitations of the hot quark-gluon plasma, at leading order in the coupling g, but only as long as Landau damping is inoperative. In cases where the boundary conditions matter, (e.g., for computing the field energy loss in the plasma), we have to recourse to the kinetic equations (5)–(7) and solve for the induced sources appropriate to the given physical conditions.

Acknowledgements

It is a pleasure to thank the organizers of the Banff workshop, especially Randy Kobes and Gabor Kunstatter, for their kind hospitality.

- R.D. Pisarski, Phys. Rev. Lett.63 (1989) 1129; E. Braaten and R.D. Pisarski, Nucl. Phys. B337 (1990) 569; ibid. B339 (1990) 310; Phys. Rev. Lett.64 (1990) 1338; Phys. Rev. D42 (1990) 2156; ibid. D46 (1992) 1829.
- V. Klimov, Sov. J. Nucl. Phys. 33 (1981) 934; Sov. Phys. JETP55 (1982) 199; H.A. Weldon, Phys. Rev. D26 (1982) 1394; ibid. 2789.
- 3. J. Frenkel and J.C. Taylor, Nucl. Phys. **B334** (1990) 199.
- 4. J.C. Taylor and S.M.H. Wong, Nucl. Phys. **B346** (1990) 115.
- E. Braaten and R. Pisarski, *Phys. Rev.* D45 (1992) 1827; J. Frenkel and J.C. Taylor, *Nucl. Phys.* B374 (1992) 156.
- 6. H.A. Weldon, Canadian J. Phys. 71 (1993) 300; R. Efraty and V.P. Nair, Phys. Rev. Lett.68 (1992) 2891; R. Jackiw and V.P. Nair, Columbia-MIT preprint CU-TP-594, April 1993; V.P. Nair, Columbia preprint CU-TP-601, June 1993; N.P. Chang, Contributed talk at the 3rd Thermal Fields Workshop, Banff, Aug 1993, these proceedings.
- 7. V.P. Silin, Sov. Phys. JETP11 (1960) 1136.
- 8. G. Baym, H. Monien, C.J. Pethick, and D.G. Ravenhall, *Phys. Rev. Lett.***64** (1990) 1867; H. Heiselberg and C.J. Pethick, *Phys. Rev.* **D47** (1993) R769.
- 9. J.P. Blaizot and E. Iancu, Nucl. Phys. **B390** (1993) 589; Phys. Rev. Lett.**70** (1993) 3376; Saclay preprint T93/064.