Perturbation theory in the nonperturbative QCD vacuum

Yu.A.Simonov Institute of Theoretical and Experimental Physics 117259, Moscow

Abstract

Perturbative expansion in the nonperturbative confining QCD background is formulated. The properly renormalized $\alpha_S(R)$ is shown to be finite at large distances, with the string tension playing the role of an infrared regulator. The infra–red renormalons are shown to be absent in the new perturbation series.

1 Introduction

The perturbation theory (PTH) expansion in QCD is an extremely useful tool, which has provided numerous links between theory and experiment [1].

The physical foundation of PTH in QCD is the celebrated asymptotic freedom phenomenon [2], ensuring the smallness of the physical coupling constant α_s at large momenta (small distances).

A formal extrapolation of the lowest order (one-loop) expressions for the renormalized α_s to large distances (small momenta) is known to be plagued by the ghost pole, which signals an uncontrolled growth of the strength of interaction at large distances. It is believed that PTH expansion breaks down even before the Landau ghost pole is reached, so that the latter has no physical meaning, similarly to the case of QED [3,4].

The impossibility of treating large distances in PTH of QCD strongly limits applications of QCD and from the formal point of view signifies the incompleteness of the theory.

The logarithmic growth of the renormalized α_s at large distances leads also to a basic difficulty, which is known as the infrared (IR) renormalon problem [5]. The IR renormalons appear as singularities on the positive real axis of the Borel parameter and destroy an attempt to sum up the PHT series.

Thus the consistency of the PHT approach is seriously in doubt, when one expands around the free unperturbed vacuum.

Alternatively it was understood during the last decades, that nonperturbative (NP) contributions are very important at large distances. In the framework of the QCD sum rules [6] the NP contributions haven been parametrized as local field vacuum averages using the operator product expansion (OPE) [7].

The phenomenologically found characteristics of the QCD vacuum clearly demonstrate that the NP fields in the vacuum are large (as compared to Λ_{QCD}). E.g. the gluonic condensate $\langle G^aG^a \rangle \approx (600MeV)^4 \gg \Lambda_{QCD}^4$ [6]; lattice calculations yield even larger values [8].

The NP effects coded as power corrections in OPE [6] crucially change the behavior of the OPE series (as compared to the PTH series) when one is leaving the asymptotic freedom region and approaching large distances. To quote the seminal paper [6]: "The basic idea behind all applications is that it is the power terms (not higher orders in the α_s series) that limit asymptotic freedom, if one tries to extend the short-distance approach to larger distances".

It is specific to the <u>local OPE</u>, that power terms explode at large distances, demonstrating the importance of the NP effects but also the difficulty of the method to penetrate the region beyond the confinement radius.

In this paper we propose to make a step forward in our strategy and instead of expansion in local condensates to consider from the beginning the PTH in the NP background of a general form. The framework for this exists for a long time [9-11]. Gluonic field is split into the background B_{μ} and perturbative gluon field a_{μ} and the new PTH is roughly an expansion in powers of g a_{μ} . The propagators of a_{μ} are exact Green's functions $G_{\mu\nu}$ in the background field B_{μ} .

A new element, crucial for the method, is that we are using a closed form

for $G_{\mu\nu}$ based on the so-called Feynman-Schwinger representation (FSR) [12]. This allows to display explicit dependence on B_{μ} , which enters through the parallel transporters P exp $ig \int_{y}^{x} B_{\mu} dz_{\mu}$. In the limit of large N_{c} any term of the new perturbative series contains B_{μ} through the Wilson loops W(C, B). Averaging over an ensemble of background fields $\{B_{\mu}\}$ then reduces the graph to a product of averaged Wilson loops if $N_{c} \to \infty$, and all the nonperturbative series is unambigiously defined by the expression for the averaged Wilson loop $\langle W(C, B) \rangle$. At large distances the latter obeys the area law [13], and one obtains therefore a one-parameter description of NP dynamics in the new perturbation series.

As a result one obtains a gauge-invariant PTH, even though the background gauge fixing term is used in the Lagrangian together with Faddeev-Popov ghosts [14]. It is not a purpose of this paper to formulate Feynman rules for diagrams of the new PTH and to study systematically the renormalization procedure in the case of the NP background. Instead we concentrate here on the lowest one-loop contribution to the charge renormalization. In this way we attempt to answer the question: how the asymptotic freedom (AF) behaviour of α_s , which is generated by the lowest gluon and ghost corrections in the background field, is maintained and how background parameters enter into the final expression for α_s .

To do that one first must define the charge renormalization in a gauge-invariant and NP nonlocal formulation. We propose 3 definitions and two of them study quantitatively.

We find that the NP background fields change the renormalized charge in a drastic way for small momenta (large distances).

The usual logarithmic growth of $\alpha_s(R)$ at large distances in the one-loop approximation disappears and $\alpha_s(R)$ saturates instead at rather moderate values. The mass scale m (or distance m^{-1}) which governs this change of asymptotics is large, $m \sim 1 Gev$, and physically is connected to the hybrid mass – or in the string terms – to the excitation mass of the transverse string vibration, $m^2 \sim 2\pi\sigma \sim 1 GeV^2$.

As a result the asymptotic freedom can be kept at any distances in the Euclidean space and expresses itself as a saturation of α_s at $\alpha_s(max) \leq 0.5$.

We always use Eucledian metrics and do not analytically continue perturbative expansions into the Minkowskian region, where physical thresholds appears. This will be a topic of separate publication.

This property drastically changes the problem of the summation of the

PTH series. We show that the IR renormalons do not appear in the background field, and the corresponding set of graphs has a finite sum.

Hence the only obstacle left is instantons (instanton - antiinstanton pairs) , which require an explicit treatment .

We stress that it not the purpose of our method to suggest an explicit way of splitting the fields into the background and perturbative ones. Instead we choose from the beginning the NP background fields, they are considered as given and this input is parametrized, e.g. by the full set of correlators $\langle F(x_1)F(x_2)...F(x_n) \rangle$.

The problem one has to solve in this case is how to find the perturbative series in this background, and we attempt to do it in the paper. Happily it appears that at large distances the only characteristics of the background which enters the answer is the string tension, and we take it to have the known phenomenological value.

The plan of the paper is as follows In section 2 we introduce background field Lagrangian, gauge-fixing and ghost terms following the usual prescriptions [9-11] and write explicitly the gluon propagator. In section 3 three definitions of the running coupling are compared, and in Section 4 one of it – defined through the energy of static quarks, is studied in detail. Another definition, yielding the running coupling $\alpha_s(p)$ in the momentum space, is discussed in Section 5.

Section 6 is devoted to the renormalons and the problem of summation of the perturbative series.

In the conclusion we summarize results of the paper and outline possible future developments.

Three appendices devoted to the explicit derivations of formulas used or discussed in the text. In Appendix 1 we consider the contribution of the term L_1 , which directly connects NP background and the perturbative field a_{μ} , to the propagator of A_{μ} .

Appendix 2 contains some detailes of derivation of the renormalization $\alpha_s(R)$ using the Wilson-loop approach.

Appendix 3 contains a detailed disscussion of the one-loop graph in presence of the confining background; most of the results used in the main text are based on the formulas explicitly derived there.

In Appendix 4 we treat the gluon spin part of the gluon propagator in the background field. Finnaly, Appendix 5 is devoted to the explicit calculation of the IR renormalon contribution.

2 Background Lagrangian and propagators

We split the gluonic field A_{μ} into the background B_{μ} and perturbative part a_{μ}

$$A_{\mu} = B_{\mu} + a_{\mu} \tag{1}$$

As we stressed in the Introduction we do not specify any principle of this division. For the purposes of the paper the exact properties of separation are immaterial, and we shall only assume that the NP fields B_{μ} ensure the area low of the Wilson loop $\langle W(B) \rangle$ with roughly the same string tension σ as known phenomenologically, $\sigma \approx 0.2 GeV$. To develope the PTH in the background field B_{μ} we use the well established background field formalism [9-11], specifically we extensively use method and notations of the paper [11], adjusted to the Euclidean space-time.

The generating functional can be written as

$$Z(J) = \int Da \ det(\frac{\delta G^a}{\delta w^b}) exp \int d^4x [L(a) - \frac{1}{2}(G^a)^2 + J^a_\mu a^a_\mu]$$
 (2)

where $L(a) = L_0 + L_1(a) + L_2(a) + L_{int}(a)$

$$L_{2}(a) = +\frac{1}{2}a_{\nu}(\hat{D}_{\lambda}^{2}\delta_{\mu\nu} - \hat{D}_{\mu}\hat{D}_{\nu} + ig\hat{F}_{\mu\nu})a_{\mu} =$$

$$= \frac{1}{2}a_{\nu}^{c}[D_{\lambda}^{ca}D_{\lambda}^{ad}\delta_{\mu\nu} - D_{\mu}^{ca}D_{\nu}^{ad} - g f^{cad}F_{\mu\nu}^{a}]a_{\mu}^{d} , \qquad (3)$$

$$\begin{split} D_{\lambda}^{ca} &= \partial_{\lambda} \cdot \delta_{ca} + g \ f^{cba} B_{\lambda}^{b} \equiv \hat{D}_{\lambda} \\ L_{0} &= -\frac{1}{4} (F_{\mu\nu}^{a}(B))^{2} \ ; \quad L_{1} = a_{\nu}^{c} D_{\mu}^{ca}(B) F_{\mu\nu}^{a} \\ L_{int} &= -\frac{1}{2} (D_{\mu}(B) a_{\nu} - D_{\nu}(B) a_{\mu})^{a} g \ f^{abc} a_{\mu}^{b} a_{\nu}^{c} - \frac{1}{4} g^{2} f^{abc} a_{\mu}^{b} a_{\nu}^{c} f^{aef} a_{\mu}^{e} a_{\nu}^{f} (4) \end{split}$$

The background gauge condition is

$$G^{a} = \partial_{\mu} a^{a}_{\mu} + g f^{abc} B^{b}_{\mu} a^{c}_{\mu} = (D_{\mu} a_{\mu})^{a}$$
 (5)

and the ghost vertex [14] is obtained from $\frac{\delta G^a}{\delta \omega^b} = (D_{\mu}(B)D_{\mu}(B+a))_{ab}$ to be

$$L_{ghost} = -\theta_a^+(D_\mu(B)D_\mu(B+a))_{ab}\theta_b \tag{6}$$

The linear part part of the Lagrangian L_1 disappears if B_{μ} satisfies classical equations of motion. We do not impose on B_{μ} this condition, and we show in the Appendix 1 that L_1 gives no contribution to the effect we are interested first of all – the modification of the AF logarithm.

We now can identify the propagator of a_{μ} from the quadratic terms in Lagrangian $L_2(a) - \frac{1}{2\xi}(G^a)^2$

$$G_{\nu\mu}^{ab} = [\hat{D}_{\lambda}^{2}\delta_{\mu\nu} - \hat{D}_{\mu}\hat{D}_{\nu} + ig\hat{F}_{\mu\nu} + \frac{1}{\xi}\hat{D}_{\nu}\hat{D}_{\mu}]_{ab}^{-1}$$
 (7)

It will be convenient sometimes to choose $\xi=1$ and end up with the well-known form of propagator in – what one would call – the background Feynman gauge

$$G_{\nu\mu}^{ab} = (\hat{D}_{\lambda}^2 \cdot \delta_{\mu\nu} - 2ig\hat{F}_{\mu\nu})^{-1}$$
 (8)

3 Definitions of the running coupling

In case of the free vacuum one can use different schemes of regularizations e.g. MS or $\overline{M}S$ schemes) and renormalize the coupling g^2 , fixing the value of the vertex at some Euclidean momentum. For, example one may require that the quark (or gluon) propagator has the form of the free propagator at some momentum p, $p^2 = -\mu^2$ [15].

In the case of NP vacuum this kind of definition is not possible anymore and one should instead introduce a new NP notion of the regularized coupling. We consider here three examples of such definition .

i) Definition of the running coupling through the interaction energy of heavy quarks (HQ)

The interaction energy of a heavy quark and antiquark E(R) at a distance R between them can be written as

$$E(R) = E_{NP}(R) - \frac{4\bar{\alpha}_s(R)}{3R}$$
(9)

where E_{NP} is the purely NP part of the energy, while $\bar{\alpha}_s(R)$ is a renormalized running coupling at distance R

$$\bar{\alpha}_s(R) = \frac{\bar{g}^2(R)}{4\pi} = \frac{g_0^2}{4\pi} (1 + \frac{g_0^2}{4\pi} f(R) + \dots$$
 (10)

and g_0 is the bare coupling, f(R) is to be calculated from the 1-loop diagrams in the background NP field.

In case of free vacuum $\bar{\alpha}_s^{(0)}(R)$ has been computed to one loop in [16] through the bare coupling

$$g^{2}(R) = g_{0}^{2} + g_{0}^{4} \frac{11}{48\pi^{2}} C_{2} \ln(\frac{R}{\delta})^{2}$$
(11)

where $\delta \sim 1/M$ is the UV cut-off.

Renormalizing via \overline{MS} scheme with the normalization mass μ one gets [17]

$$\alpha_{\bar{M}\bar{S}}(\mu,R) = \alpha_{\bar{M}\bar{S}} \left\{ 1 + \frac{\alpha_s}{\pi} \left[\frac{b_0}{2} (\ln \mu R + \gamma_E) + \frac{5}{12} b_0 - \frac{2}{3} C_A \right] \right\}$$
(12)

with $b_0 = \frac{11}{3}N_c - \frac{2}{3}N_f$, $\gamma_E = 0.5777...$

The main feature of (11) and (12) is the logarithmic grouth of $\alpha_s(R)$ at large distances, and this feature produces both the Landau ghost pole [3,4] and the IR renormalon problem [5]. In the next sections we shall demonstrate that the strong confining NP background eliminates the logarithmic growth of $\alpha_s(R)$ and is this way solves both the ghost pole and IR renormalon problem.

To calculate $\bar{\alpha}_s(R)$ in the NP background we shall use the gauge-invariant definition

$$E(R) = -\lim_{T \to \infty} \{ \frac{1}{T} ln < W_C(B+a) >_{B,a}^{reg} \}$$
 (13)

where contour C of the Wilson loop $\langle W_C \rangle$ is a rectangular $R \times T$ shown in Fig. 1.

The averaging over fields B_{μ} , a_{μ} and regularization is assumed in (13) and will be specified in the next section. Note that E(R) (13) contains both PTH and NP contributions as in (9). We again stress at this point, that in our approach there is no problem of separation of a given E(R) into E_{NP} and a PTH piece (which is of course not well defined). Instead we assume that an ensemble of fields $\{B_{\mu}\}$ is given to us, it is characterized by correlators

$$g^n < F_{\mu_1\nu_1}(B, x_1)...F_{\mu_n\nu_n}(B, x_2) >_B$$
 (14)

and in particular by the string tension σ , which can be expressed through the correlators. Our task is to construct the PTH in this given background and to express in particular $\bar{\alpha}_s(R)$ through the nonperturbative parameters. To perform renormalization in the background field formalism, one may use its gauge invariance [10]. One can visualize that all terms in (2) are gauge invariant with respect to gauge transformations "rotating" both B_{μ} and a_{μ} . It is convenient to split gauge transformations as follows [11]

$$B_{\mu} \to U^{+}(B_{\mu} + \frac{i}{g}\partial_{\mu})U$$
,
 $a_{\mu} \to U^{+}a_{\mu}U$, $B_{\mu} = B_{\mu}^{a} + t^{a}$. (15)

If we consider following [11] only amplitudes with no external a_{μ} lines (and we need only those), then renormalization of a_{μ} and θ lines is inessential (corresponding Z factors compensate each other), while g, B, ξ renormalize as

$$g_0 = Z_q g , B_0 = Z_B^{1/2} B, \xi_0 = Z' \xi$$
 (16)

Now because of the explicit gauge invariance, Z_g and Z_B are connected [11]

$$Z_a^{-1} = Z_B^{1/2} (17)$$

as a consequence, the quantities gB_{μ} and $gF_{\mu\nu}(B)$ are RG invariant, and therefore all characteristics of NP fields, like correlators (14) or string tention are RG invariant and should be considered on the same footing as the external momenta in the amplitudes.

ii) Definition through the B-field two-point function

This is equivalent to the method used in [11], where two 1-loop diagrams have been used of Fig. 2 and 3 with two external B lines. To make averaging over fields $\{B_{\mu}\}$ we multiply each diagram in the coordinate space with the parallel transporter in the adjoint representation,

$$\hat{\Phi}(x,y;B) = Pexp \ ig \ \int_{y}^{x} \hat{B}_{\mu} dz_{\mu}$$
 (18)

and consider both gluon and ghost propagating in the field B_{μ} with propagators $G_{\mu\nu}^{ab}(x,y,B)$ and $G^{ab}(xy;B)$ respectively. The product

$$H(x,y) \equiv \langle \hat{\Phi}^{ef}(x,y,B) \Gamma^{eac}_{\mu} G^{ab}_{\mu\nu}(x,y,B) G^{cd}(x,y,B) \Gamma^{fbd}_{\nu} \rangle_{B}$$
 (19)

depicted in Fig. 4, yields the one-loop correction to the renormalized charge, which in absence of field B_{μ} reduces to the second term on the r. h. s. of (11). Using results of the next section one can again prove that dependence like $\ln|x-y|$ disappears in (19) when $|x-y| \gg m^{-1}$, m > 1 GeV. This will be clear from discussion in the next two sections.

In a similar way one calculates two- and more loop corrections, using the same two-point diagrams as in [11].

iii) Definition through the finite box of size L.

This definision is similar to that of refs. [18]. One fills the box L^4 with the NP field B_{μ} and considers one-loop correction to g^2 . As in [18] one gets

$$\bar{g}^2(L) = g_0^2 + n_1(L)g_0^4 + \dots {20}$$

with $n_1 = h_0 - \frac{1}{2}h_1$,

$$h_0 = \frac{\frac{\partial}{\partial \eta} \ln \det \bar{G}}{\frac{\partial}{\partial \eta} \Gamma_0(B)} , h_1 = \frac{\frac{\partial}{\partial \eta} \ln \det G}{\frac{\partial}{\partial \eta} \Gamma_0(B)}$$
 (21)

Here η is a dimensionless parameter characterising strength of field B_{μ} , $\Gamma_0(B)$ is the effective action $(=\frac{1}{4}\int F_{\mu\nu}^2(B)d^4x)$.

For the abelian (nonconfining) field used in [18] one obtains the asymptotics [18]

$$n_1(L) \sim_{L \to \infty} \sum_{n=0}^{\infty} (r_n + s_n \ln L) / L^n$$
 (22)

with r_n, s_n some constants, $s_0 = \frac{11}{12\pi^2}$. Thus one recovers the usual free-vacuum renormalization with the infra-red logarithmic growth. If instead one uses the confining field B_{μ} , the asymptotics of $n_1(L)$ changes and the logarithm lnL disappears (the study of lndetG in case of confining background was done in [19] for the case of nonzero temperature T, a similar analysis for T = 0 will be published elsewhere).

In the next sections we shall study in detail the first and second definitions of the running coupling and shall find its one-loop renormalization. As will be clear both definitions yield the same qualitative phenomenon: the running coupling $\alpha_s(R)$ saturates at large R at some finite value, and the mass parameter m, which cuts off the logarithmic growth of $\alpha_s(R)$, is of the same order in both definitions.

4 Calculation of $\alpha_s(R)$ through the Wilson loop formalism.

In this section we shall follow the way outlined in the first definition of $\alpha_s(R)$, i.e. we shall use E(R) given by (13). To this end we need the PTH expansion of the Wilson loop $\langle W(B+a) \rangle$, i.e. expansion of the powers in gluon operator a_{μ} . We start with the Wilson contour C of an arbitrary configuration and divide the contour into N pieces $\Delta x_{\mu}(n)$, writing identically

$$W(B+a) \equiv \frac{1}{N_c} tr P \exp ig \int_C (B_\mu + a_\mu) dx_\mu =$$

$$= \frac{1}{N_c} tr P \prod_{n=1}^N [1 + ig(B_\mu(x(n)) + a_\mu(x(n))) \Delta x_\mu(n)] =$$

$$= \frac{1}{N_c} tr P \left\{ \prod_{n=1}^N (1 + igB_\mu \Delta x_\mu) + \prod_{n=1}^{k-1} (1 + igB_\mu \Delta x_\mu) iga_\mu(k) \Delta x_\mu(k) \times \prod_{n=k+1}^N (1 + igB_\mu \Delta x_\mu(n)) + \dots \right\}$$

One can represent the sum (23) as a sum of Wilson loops $W^{(n)}(B)$ with n insertions of the field $iga_{\mu}(k)$ on the contour C:

$$W(B+a) = W(B) + \sum_{n=1}^{\infty} (ig)^n W^{(n)}(B; X(1)...X(n)) dx_{\mu_1}(1)...dx_{\mu_n}(n)$$
 (24)

Now we must integrate over the fields Da_{μ} with the weight given in (2). In the lowest order $\langle W(B) \rangle_a = W(B)$ for the properly normalized weight, and in the order $O(g^2)$ one has a one-"gluon" exchange diagram shown in Fig. 5.

$$-g^{2} < W^{(2)} >_{a} dxdy = -g^{2} \int \Phi^{\alpha\beta}(x, y; B) G^{\delta\alpha;\beta\gamma}_{\mu\nu}(x, y; B) \Phi^{\gamma\delta}(y, x, B) dx_{\mu} dy_{\nu}(25)$$

where $\Phi^{\alpha\beta}$ are parallel transporters (18) with fundamental indices written explicitly, $\alpha, \beta = 1, ...N_c$

$$G^{\delta\alpha,\beta\gamma}_{\mu\nu} = t_a^{\delta\alpha} t_b^{\beta\gamma} G^{ab}_{\mu\nu} \quad , \tag{26}$$

and $G_{\mu\nu}^{ab}$ is the gluon propagator in the NP background field B_{μ} given in (7) and (8).

It is appropriate at this point to specify the notion of the gluon in the background and to distinguish it from the free gluon, interacting only perturbatively. We propose to call it the "hybrid-gluon", or h-gluon, since as we shall see, in most cases it will aquire a mass and behaves as a hybrid.

In the next order, $0(g^4)$, one obtains five diagrams shown in Figs. 6-10.

Next step is ti average over the ensemble of NP fields $\{B_{\mu}\}$. It is not necessary to specify the ensemble and prescribe the measure; we can use the cluster expansion to obtain for the simplest term [13].

$$< W(B) >_{B} = exp \sum_{n=1}^{\infty} \frac{(ig)^{n}}{n!} \int d\sigma(1)...d\sigma(n) < F(1)...F(n) >$$
 (27)

Exact gauge invariant form of the correlator $\langle F(1)...F(n) \rangle$ is discussed in [13]. The lowest order term in (27) e.g. has two independent structures the Kronecker one and the full derivative term [13]

$$g^{2} < F_{\mu\nu}(u)\Phi(u,v)F_{\rho\sigma}(v)\Phi(v,u) > = (\delta_{\mu\rho}\delta_{\nu\sigma} - \delta_{\mu\sigma}\delta_{\nu\rho})D(u-v) + \partial(D_{1})$$
 (28)

It is D and the Kronecker structures of higher correlators that bring about the area law of the Wilson loop [13]:

$$\langle W(B) \rangle_B \cong exp(-\sigma S_{min})$$
, S large (29)

where

$$\sigma = \frac{1}{2} \int \int_{-\infty}^{\infty} d^2 u D(|u|) + \int \langle FFFFF \rangle + \dots$$
 (30)

Thus any finite member of terms in the sum (27) or the whole sum, if it converges, yields the area law for the Wilson loop. In what follows we shall assume that the area law with $\sigma \cong \sigma_{phen}$ holds for all Wilson loops provided their size (both width and length) is large [13, 20], i.e. for the rectangular $R \times T$ in Fig. 1.

$$R, T \gg T_a$$
 (31)

where T_g is the gluonic correlation length [20], defined by the range of correlators D(u) etc.

We stress at this point that the assumption (29) is very natural from the point of view of the cluster expansion [13], and moreover it is supported by the lattice calculations [21].

With the help of (27) one easily obtains the first term in (9); using [22] one has

$$E_{NP}(R) = 2R \int_0^R d\lambda \int_0^\infty d\nu D(\lambda, \nu) + \int_0^R \lambda d\lambda \int_0^\infty d\nu [-2D(\lambda, \nu) + D_1(\lambda, \nu)] + \text{higher correlators.}(32)$$

We note that at small R, $R < T_g$, [22,23]

$$E_{NP}(R) \sim CR^2 + 0(R^4)$$
 (33)

where $0(R^4)$ comes from higher order correlators, while at large R [22]

$$E_{NP}(R) = \sigma R - C_0 \tag{34}$$

We refer the reader to ref. [24] for detailed discussion.

Now we turn to the h-gluon exchange diagrams of Fig. 5-10. First we must regularize them, following the method of [25]. It was shown there that all divergencies of smooth Wilson loops are regularized though introduction of some minimal distance δ on the contour C or by dimensional regularization and renormalized) by introduction of a common Z factor (heavy mass renormalization) and charge renormalization, generated by divergencies in diagrams of Fig. 6,7,9,10.

We shall follow the same procedure as in [25], but we are interested not only in the divergent parts of diagrams as in [25], but also in the large R dependence with and without background field B_{μ} .

We start with the simplest h-gluon exchange diagram, Fig.5, given by (25). The first step is to use for $G_{\mu\nu}$ the Feynman-Schwinger representation (FSR) [12]

$$G_{\mu\nu}^{ab}(x,y;B) = \int_0^\infty ds e^K(Dz)_{xy} \hat{\Phi}_F(x,y;B)_{\mu\nu}^{ab}$$
 (35)

where

$$K = \frac{1}{4} \int_0^S \dot{z}_{\mu}^2(\lambda) d(\lambda);$$

$$\Phi_F(x,y,B) = P_F P \exp ig \int_y^x \hat{B}_{\mu} dz_{\mu} \exp 2 ig \int_0^S \hat{F}_{\lambda\rho}(z(\lambda)) d\lambda \qquad (36)$$

and Φ_F differs from the usual parallel transporter (18) by the ordered (due to P_F) insertions of the gluon color magnetic moment interaction $2g\hat{F}_{\lambda\rho}$.

The use of FSR is the central point of the method since it enables one to write explicitly dependence on B_{μ} - it enters via Φ_F . As a next step we must average (25) over the ensemble $\{B_{\mu}\}$, i.e. to compute an average of an expression of the form.

$$\langle W^{(2)} \rangle_B \sim \langle \Phi t \hat{\Phi}_F t \Phi \rangle_B$$
 (37)

where $\hat{\Phi}_F$ is in the adjoint while Φ in the fundamental representation. To understand better the structure of (37) let us take the limit $N_c \to \infty$ and represent the adjoint line in $\hat{\Phi}_F$ by the double fundamental line, as suggested by 'tHooft [26]. We obtain two adjacent closed contours C_1 and C_2 as shown in Fig. 11. When both contours large, i.e. for $R, T \gg T_g$, and $N_C \to \infty$ one has

$$< W^{(2)} >_{B} \sim < W_{C_1}(B) >_{B} < W_{C_2} >_{B} + 0(\frac{1}{N_c^2}) \sim exp(-\sigma(S_1 + S_2))$$
 (38)

where S_1, S_2 are minimal areas inside contours C_1, C_2 .

Combining now (25), (35) and (38) we obtain

$$-g^{2} < W^{(2)} >_{B} = -g^{2} \int_{0}^{\infty} ds (Dz)_{xy} dx_{\mu} dy_{\nu} exp(-K - \sigma(S_{1} + S_{2})) \delta_{\mu\nu}$$
 (39)

(Note that we have omitted for simplicity the $\hat{F}_{\lambda\rho}$ term in (36), which brings about spin-orbit corrections, see Appendix 4 for discussion.)

Eq. (39) describes propagation of a scalar particle interacting with the contour via two strings, the world surface of which is given by S_1, S_2 .

We now show that for our choice of contour in Fig. 5,11, with the horizontal size T much larger than vertical size R, the resulting interaction tends to zero. Indeed for a given point $z(\lambda)$ on the line of propagator, see Fig. 5, 11 the interaction does not depend on the $z_4(\lambda)$, $z_1(\lambda)$ which are in the plane of the contour $R \times T$, and depends only on the transverse coordinates z_2 , z_3 . The interaction at the "time" λ can be written as

$$V \equiv \sigma(\sqrt{z_2^2 + z_3^2 + z_4^2} + \sqrt{z_2^2 + z_3^2 + (T - z_4)^2}) - \sigma T$$
 (40)

where we have normalized to the case with no h-gluon exchange. It is easy to see that for a generic case $T-z_4, z_4 \gg z_2, z_3$ (according to (13) we must finally take the limit $T \to \infty$) one has

$$V \approx \frac{\sigma}{2} \left(\frac{z_2^2 + z_3^2}{|z_4|} + \frac{z_2^2 + z_3^2}{|T - z_4|} \right) \to 0 \tag{41}$$

Thus we come back to the case of a free gluon exchange and obtain from (39)

$$-g^{2} \frac{\langle W^{(2)} \rangle_{B}}{\langle W \rangle_{B}} = -\frac{g^{2} C_{2}(f)}{4\pi^{2}} \int \int \frac{dx_{4} dy_{4}}{(x_{4} - y_{4})^{2} + R^{2}} = \frac{g^{2} C_{2}(f)}{4\pi R}$$
(42)

In a similar way we treat the diagrams with more gluon exchanges, like in Figs. 8, 12 and exponentiate the result (42), obtaining the lowest order term $\left(-\frac{4}{3}\frac{\bar{\alpha}_s(R)}{R}\right)$ in (9). One should mention that in multiple gluon exchanges like Fig. 12 there appear in general correlated gluon pairs, triplets etc. with average distance between gluons of the order of $(\sigma)^{-1/2}$. These correlated exchanges are not taken into account during exponentiation and should be accounted for separately, yielding a Yukawa-type correction to the perturbative part of E(R) starting with the order $O(g^4)$. We shall discuss these corrections in a separate publication, they do not change results of this paper concerning large distance logarithmic behaviour.

Now we come to the fourth order diagrams, needed to renormalize α_s shown in Figs. 6, 7, 9, 10.

First we discuss the structure of divergencies and logarithms in case of no background B_{μ} . Referring the reader to the Appendix 2 for details, we can e.g. write the resulting integral for the diagram of Fig. 10:

$$\frac{(f(R))}{R}_{Fig.10} = \frac{-3(1-\xi)}{16\pi^2} C_2 \int \frac{d^3x}{|\vec{x}||\bar{x}-R|^3} = \frac{-3(1-\xi)}{4\pi} \frac{C_2}{R} ln \frac{Re}{\delta}$$
(43)

In dimentional regularization one would have

$$ln\frac{R}{\delta} \to \frac{C}{\varepsilon} + ln\mu R + const$$
 (44)

For Figs. 6 and 7 one obtains (again $B_{\mu} = 0$)

$$\left(\frac{f(R)}{R}\right)_{Fig.6,7} = (13 - 3\xi)C_2 \sim \frac{d^4r}{|\bar{R} - \bar{r}|} \prod_{j=1}^{\infty} (r^2) \to \frac{13 - 3\xi}{12\pi} C_2 \ln\frac{R}{\delta}$$
(45)

where $\Pi(r^2)$ is the trasverse self-energy part of gluon and ghost in the coordinate space:

$$\prod(r^2) = \frac{1}{(\bar{r}^2 + r_4^2)^2} \tag{46}$$

Integrating over d^4r we obtain the same logarithm as in (43).

Some more care is needed for the graph of Fig. 9; leaving details to the Appendix 2 we mention only the final result

$$\left(\frac{f(R)}{R}\right)_{Fig.9} = \frac{(3-\xi)C_2}{2\pi R} ln\frac{R}{\delta} \tag{47}$$

Summing (43),(45) and (47) together we obtain

$$f(R) = \frac{b_0}{2\pi} ln \frac{R}{\delta} \tag{48}$$

in agreement with (11) and (12).

Consider now the case of nonzero B_{μ} , and let us average each of the diagram $W^{(4)}(B)$ in Fig.6,7,9,10 over $\{B_{\mu}\}$. We again use the limit $N_C \to \infty$ and replace each gluon line by a double fundamental line. Choosing the gauge fixing constant $\xi = 3$ we eliminate the logarithmic part (and divergence) of Fig. 9 and disregard it in what follows, since now we are only interested in the question: what happens with logarithms in (43),(45) and (47) when the NP confining field is included.

Following the same line of reasoning as we used for the diagram of Fig. 11 and imposing the confining field B_{μ} we obtain for diagrams of Fig. 6,7 and that of Fig. 10 three surfaces in Figs. 13 and 14 respectively, filled with confining film due to the area law. As in (39), we must consider again horizontal size T of diagrams in Figs.13,14 large and a result the surfaces S_1 and S_2 do not produce any force on the gluon line when $T \to \infty$. This is not true, however for S_3 , which provides a strong confining force between the fundamental lines which border S_3 .

As a result we must consider the behaviour of the self-energy part $\prod(x,y)$ in Fig. 13, defined by the FS representation:

$$\prod(x,y) \sim \int_0^\infty ds \int_0^\infty d\bar{s} Dz D\bar{z} exp(-K - \bar{K} - \sigma S_3)$$
 (49)

It is easy to show (for details see Appendix 3) that $\prod(x,y)$ behaves as $\frac{1}{(x-y)^4}$ at small $|x-y|, |x-y| \ll T_g$, and has an exponential fall-off at large |x-y|

$$\prod(x,y) \sim exp(-m_1|x-y|), |x-y| \to \infty$$
 (50)

Introducing this cut-off behaviour in (46) one obtains the following integral which replaces the original logarithmic expression in (45) (see details in Appendix 2).

$$\frac{1}{R} \ln \frac{R}{\delta} \to I(R) \equiv \int_{\delta}^{\infty} \frac{dr}{r} e^{-m_1 r} \left(\frac{1}{R} \theta(R - r) + \frac{1}{r} \theta(r - R)\right) \tag{51}$$

There are two regimes of behaviour of I(R): the region of the asympthotic freedom :

$$m_1 R \ll 1, I(R) \approx \frac{1}{R} ln \frac{Re}{\delta}$$
 (52)

and the infrared region

$$m_1 R \gg 1, I(R) \approx \frac{1}{R} \int_{\delta}^{R} \frac{dr}{r} e^{-m_1 r} \approx \frac{1}{R} ln \frac{1}{m_1 \delta}$$
 (53)

The same situation occurs for the diagram of Fig.14. Here the cut-off function $exp(-m|\vec{R}-\vec{x}|)$ appears under the integral in (43), producing the same integral I(R) as in (51).

As a final result of this section we formulate the new expression for f(R) taking the NP confining into account (modulo constant (nonlogarithmic) terms)

$$\frac{1}{R}f_{NP}(R) = \frac{b_0}{2\pi}I(R)$$
 (54)

with I(R) defined in (51).

5 Renormalization of α_s in the definition ii)

In the second nonperturbative definition of α_s through the two-point function H(x,y) (19), depicted in Fig. 4, we can again exploit the $N_c \to \infty$ limit. In this case each adjoint line, including $\hat{\Phi}(x,y,B)$, is represented as double fundamental line [26] and each closed fundamental contour $W_c(B)$, after averaging over fields $\{B_{\mu}\}$, acquires the area-law factor $exp(-\sigma S_{min}(C))$ - in agreement with our assumption (29). Thus each of the diagrams of Fig. 4 can be depicted as in Fig. 15, the difference between (a) and (b) is that for the gluon Green's function there are in addition gluon spin operators inserted on the gluon line, leading to the spin-dependent effects, which we disregard in the first approximation.

The diagrams of Fig. 4 can be written in the form similar to (49), which is discussed in detail in the Appendix 3

$$H_{\mu\nu}(x,y) = \frac{g^2}{(4\pi)^2} b_0 (\partial_\mu \partial_\nu - \partial^2 \delta_{\mu\nu}) \bar{\prod}(x,y)$$
 (55)

where $\bar{\prod}$ is defined as

$$\bar{\prod}(x,y) = \int_0^\infty ds \int_0^\infty d\bar{s} \ e^{-K-\bar{K}} (Dz)_{xy} (D\bar{z})_{xy} \ e^{-\bar{V}}$$
 (56)

with

$$\bar{V} = S_{min}^{(1)} + S_{min}^{(2)} + S_{min}^{(3)} \tag{57}$$

and $S_{min}^{(i)}$ correspond to the minimal areas inside the 3 contours depicted in Fig.15.

We have omitted in (57) for simplicity the gluon spin terms in the Green's function G in (19).

At small distances |x - y| one can follow analysis of Appendix 3 and obtain that

$$\bar{\prod}(x,y) \sim \frac{1}{(x-y)^{2d-4}}$$
 (58)

This leads to the familiar answer for the Fourier-transform of $H_{\mu\nu}$ [11]

$$H_{\mu\nu}(p) = \frac{g^2 b_0}{(4\pi)^2} (p_{\mu} p_{\nu} - p^2 \delta_{\mu\nu}) (N_{\varepsilon} - \ln \frac{p^2}{\mu^2} + const)$$
 (59)

with

$$b_0 = \frac{11}{3} N_c$$
, $N_{\varepsilon} = \frac{1}{\varepsilon} - C + \ln 4\pi$,

C is the Euler constant, C = 0.577.

At large |x-y| one must take into account the confining dynamics present in \bar{V} and find the corresponding mass. To this end one must look at the expressions (55), (56) more carefully.

To understand the structure of H(x, y), one can dissect the graph in Fig. 15 along the dash-dotted line and one obtains a system of two gluons 1 and 2 connected by a pair of strings, as shown in Fig. 16. One of the strings goes through the fixed point 3 which appears due to the dissection of the straight-line parallel transporter $\Phi(x, y; B)$ – see Fig. 15.

One can show that the mass of the system of two gluons with two strings in the configuration of Fig. 16 is larger than the corresponding mass without the additional condition of string passing through the point 3:

$$m(1,2;3) > m(1,2) \equiv m_2$$
 (60)

Thus we come to the conclusion that H describes the Green's function of a two-gluon glueball with a quantum nember $J^{PC} = 1^{-+}$ (and with an additional "heavy gluon" propagating along the straight line of $\hat{\Phi}(x, y, B)$, if we keep the point 3 in Fig. 16, and without it if we calculate a more light system according to the inequality (60)).

An estimate of m(1,2) with $J^{PC} = 1^{-+}$ has been done within the proper time Hamiltonian in [27]

$$m(1,2) \cong 2GeV \pm 0.15GeV \tag{61}$$

A similar estimate has been obtained in the lattice calculations, where usually two neighbouring members of the same multiplet L=1, S=1 are studied [28].

Using the arguments given in the Appendix 3, one can approximate $\bar{\Pi}(x,y)$ as

$$\bar{\prod}(x,y) \simeq C_0 \frac{exp(-m_2 \mid x - y \mid)}{(x-y)^{2D-4}}$$
 (62)

which after renormalization yields the Fourier-transform

$$\bar{\prod}(p) \approx C_0 \ln \frac{m_2^2 + p^2}{\mu^2} + const \tag{63}$$

A comparison of (59) and (63) shows that up to power corrections two expressions have the same μ dependence, and hence the same Z factors. Actually all the change from (59) to (63) can be described as introduction of a new (fifth) external momentum component $p_5^2 \equiv m_2^2$, so that the total momentum squared is in case of nonperturbative background equal to $p^2+m_2^2$. This fact does not change the Gell-Mann-Low equations which to one-loop order are [15]

$$\frac{dg}{d\ln\mu} = -b_0 \frac{g^3}{16\pi^3} \tag{64}$$

and solving (64) with boundary conditions as in (55) and (63) (neglecting power terms as before for large $(m^2 + p^2)x_0^2 \gg 1$) one obtains

$$\alpha_s(p) = \frac{\alpha_s(\mu)}{1 + \frac{b_0}{4\pi}\alpha_s(\mu) \ln \frac{m_2^2 + p^2}{\mu^2}}$$
 (65)

Introducing the Λ parameter (65) is rewritten as

$$\alpha_s(p) = \frac{4\pi}{b_0 \ln \frac{m_2^2 + p^2}{\Lambda_{NP}^2}}$$
 (66)

We have used notation Λ_{NP} to stress that this new Λ parameter may differ from the free vacuum Λ_{QCD} and needs a new determination from the experimental data. We stress here again that the appearance of m_2^2 together with p^2 is possible because of renormalization invariance of gB_{μ} [11], which in its turn yields invariance of σ and $m_2 \sim \sqrt{\sigma}$.

At large p one obtains in (66) the familiar AF logarith. This logarithm is still large for p=0 if one chooses a low normalization parameter $\Lambda_{NP}, \Lambda_{NP} \sim \Lambda_{QCD}$. Thus the AF phenomenon can be maintained down to very low p in the Euclidean region. This is similar to the situation which we have observed in the Wilson-loop renormalization scheme, eg. (53).

There, however, the mass m_1 corresponds to the one-Wilson-loop dynamics, while in our case actually a doubled film defines dynamics for m(1,2), with $\sigma_a \cong 2\sigma$, so that the mass $m_2 \cong \sqrt{2}m_1$. This fact demonstrates that the IR cutoff is scheme dependent. (Note that our definition of the renormalization scheme differs from the usual one, and the scheme dependence actually means the dependence on the physical process in question).

6 Renormalons and asymptotics of the perturbative series

In this section we study the consequences of the modified AF (66) for the properties of the PTH series. We shall concentrate for definiteness on the Euclidean correlator of e.m. currents $\Pi^{e.m.}(Q^2)$ as in [29, 30] and shall write

it as in [30] in a form exhibiting the contribution of the renormalon [5]

$$\prod^{e.m.}(Q^2) = \frac{1}{2\pi^2 b_0} \left(\frac{1}{\alpha_s(\mu^2)} - \frac{1}{\alpha_s(Q^2)}\right) + \frac{1}{2\pi^2 b_0} \ln\left(\frac{\alpha_s(Q^2)}{\alpha_s(\mu^2)}\right) + \sum \tilde{p}_n \alpha_s^n(Q^2) + \Delta \prod_{s=0}^{e.m.} (67)$$

where we have separated out of the PTH series the contribution of the set of diagrams shown in Fig. 17, denoted by $\Delta \Pi$. We shall first discuss the infrared (IR) renormalon, [5], i.e. the contribution from the graphs of Fig.17 from the domain of integration over k with $k \ll Q$. If the coupling constant α_s is normalized at Q^2 (we shall take it at $Q^2 + m^2$ and (66) is used for α_s , we have

$$\Delta \prod = \frac{\alpha_s(Q^2)}{8\pi^3} \sum_n \left(\frac{b_0 \alpha_s(Q^2)}{4\pi}\right)^n \int_0^{Q^2} \frac{k^2 dk^2}{Q^4} \ln^n \left(\frac{Q^2 + m^2}{k^2 + m^2}\right)$$
(68)

The analysis of the integral (68) made in the Appendix 5, yields

$$I = \int_0^{Q^2} \frac{k^2 dk^2}{Q^4} \ln^n \left(\frac{Q^2 + m^2}{k^2 + m^2} \right) = \begin{cases} \frac{n!}{2^{n+1}}, & n < 2n_0 \\ \eta \frac{n_0^{n+1}}{n+1}, & n > 2n_0, & \eta < 1 \end{cases}$$
 (69)

with the notation $n_0 = \ln \frac{Q^2 + m^2}{m^2}$.

As a result we obtain the following contribution of the graphs in Fig.17

$$\Delta \prod = \frac{4\pi}{8b_0 \pi^3} \sum_{n \gg 1}^{\infty} \left(\frac{\alpha_s(Q^2)b_0}{8\pi}\right)^n q_n \tag{70}$$

where

$$q_n = \begin{cases} (n-1)! & , & n < 2n_0 \\ \eta \frac{(2n_0)^n}{n} & , & n > 2n_0 \end{cases}$$
 (71)

We remind that the estimate (71) is an upper limit for q_n .

First we discuss a connection with the previous results [29,30] on the IR renormalons. In this case $m^2 \to 0$ (no background field) and $n_0 \to \infty$; q_n is defined as in (71) for any n and the Borel transform of $\Delta \prod$ is as in [29,30]

$$\Delta \prod = (16\pi^3)^{-1} \int_0^\infty \frac{exp(-\frac{t}{\alpha_s(Q^2)})dt}{1 - \frac{tb_0}{8\pi}}$$
 (72)

the Borel integral (72) exibits IR renormalon pole at $t = \frac{8\pi}{b_0}$ which makes the Borel summation of the PTH series prohibitive.

In our case, when n_0 is finite, the situation is drastically changed. Namely, having in mind that the estimate (71) is an upper limit of q_n both for $n > 2n_0$ and for $n < 2n_0$, we can take the sum in (70) which does not diverge and even does not need Borelization. Leaving details for the Appendix 5 we quote only the final result

$$\Delta \prod = \frac{\alpha_s}{8\pi^3} \sum_{n=1}^{\infty} \int_0^{n_0} \frac{\alpha_s(Q^2)tb_0}{4\pi} e^{-2t} dt \approx \frac{b_0 \alpha_s^2(Q^2)}{32\pi^4}$$
 (73)

There are no poles in the plane of the Borel variable, as can be easily seen from the absense of the factorial growth of the coefficient q_n at arbitrarily large n. We thus conclude that the convergence of PTH series is radically improved when the strong nonperturbative background is taken into account, and in particular the IR renormalons disappear.

We now turn to the ultraviolet (UV) renormalons [5]. Those come from the same series of graphs, Fig.17, but from the integration region with $k^2 \gg Q^2$. It is easy to understand that the UV renormalons do not change due to the modified AF logarithm (66), since in this kinematical region $k^2 \gg m^2$ and the change is immaterial. Hence we have as in [30] the estimate

$$\Delta \prod (UV) = -\frac{1}{6\pi^3} \sum_{n} \alpha_s^n (-\frac{b_0}{4\pi})^{n-1} (n-1)!$$
 (74)

which brings a pole at negative value of the Borel parameter, $t = -\frac{4\pi}{b_0}$, and hence the Borel sum of the PTH series is well defined.

7 Conclusion

We have introduced in this paper a new perturbation theory, based on the explicit separation of NP and perturbative fields, Eq.(1).

The former are defined by their vacuum correlators as in (27), which at large distances enter only as one constant - the string tension σ (30).

This is the input of the theory. The PTH expansion in powers of perturbative field a_{μ} is then defined unambiguously in terms of the NP input and coupling constant α_s .

The main emphasis of the paper is the drastic change in the behaviour of renormalized α_s at large distances as compared to the usual (free) PTH.

Two different NP definitions of the running coupling constant are used, Eqs.(9,13) and (19), and in both cases the one-loop contribution is shown to be bounded at large distances, in contrast to the logarithmic growth of the free one-loop result.

It is important that Gell-Mann-Low equations are not changed and the resulting behaviour of $\alpha_s(p)$ is characterized by only one new parameter, m^2 as in (65-66), which is RG invariant. The mass m depends on the process and in the typical situation of a gluon exchange in the e^+e^- annihilation, is of the order of one GeV. As a result the magnitude of $\alpha_s(p)$ everywhere in the Euclidean region of p^2 is limited and the Landau ghost pole is not inevitable. Rough estimates show that $\alpha_s(p)$ is less than unity in the Euclidean region; enabling one to use perturbation expansion at all distances.

As a consequence there appears a solution of the long standing problem of IR renormalons. It is shown in Section 6, that IR renormalons are absent when the running coupling has the form (66). Hence the Borel summation of the new perturbative series is not prohibited by the IR renormalons, and one encounters in the right Borel-parameter half-plane only the distant singularities due to instanton-antiinstanton $I\bar{I}$ pairs.

We do not consider here important questions, e.g. how inclusion of $I\bar{I}$ pairs into the NP vacuum fields influences the properties of the new perturbative series.

We also do not discuss the behaviour of $\alpha_s(p)$ in the Minkowskian region, $p^2 < 0$, which shall be presented in a subsequent paper.

During the earliest stage of this study the author intensely discussed with P. van Baal and G.'tHooft. Their useful remarks and suggestions are gratefully acknowledged.

The author is grateful for useful discussions to A.M.Badalyan, K. G.Boreskov, A.Yu.Dubin, B.L.Ioffe, A.B.Kaidalov, L.N.Lipatov, M.I.Polikarpov and K.A.Ter-Martirosyan.

APPENDIX 1

The contribution of the term L_1

Let us calculate the contribution of the term L_1 to the gluon propagator. If one denotes by $\langle \rangle_a$ the integral Da_μ with the weight L(a) as in (2), we obtain

$$\Delta_{\mu_1\mu_2}^B(x_1, x_2) \equiv \langle a_{\mu_1}(x_1) a_{\mu_2}(x_2) \rangle_a =$$

$$\int d^4 y_1 d^4 y_2 G_{\mu_1\nu_1}(x_1, y_1) D_{\rho} F_{\rho\nu_1}(y_1) D_{\lambda} F_{\lambda\nu_2}(y_2) G_{\nu_2\mu_2}(y_2, x_2) \tag{A1.1}$$

The gluon Green's function $G_{\mu\nu}$ is given in (8) and depends on the background field B_{μ} , as well as D_{μ} and $F_{\rho\lambda}$. Averaging over $\{B_{\mu}\}$ is done in (A.1.1) using cluster expansion. To get a simple estimate of Δ we replace $G_{\mu\nu}$ by the free Green's function $G_{\mu\nu}^{(0)}$ and take into account that

$$< D_{\rho} F_{\rho\nu_1}(y_1) D_{\lambda} F_{\lambda\nu_2}(y_2) >_B \Rightarrow \frac{\partial}{\partial y_{1\rho}} \frac{\partial}{\partial y_{2\lambda}} < F_{\rho\nu_1}(y_1) F_{\lambda\nu_2}(y_2) >$$
 (A1.2)

and for the latter we use the representation [13] in terms of two independent Lorentz structures, $D(y_1 - y_2)$ and $D_1(y_1 - y_2)$. The contribution of D is (that of D_1 is of similar character) in the momentum space

$$\Delta_{\mu_1 \mu_2}^B(k) \sim \frac{(k^2 \delta_{\mu_1 \mu_2} - k_{\mu_1} k_{\mu_2})}{k^4} D(k)$$
 (A1.3)

where

$$D(k) = \int d^4 y e^{iky} D(y) \tag{A1.4}$$

Insertion in (A.1.4) of the exponential fall-off for D(y) found in lattice calculations [31] yields

$$D(k) = \frac{\langle F^2(0) \rangle}{(N_c^2 - 1)} \cdot \frac{\pi^2 \mu}{(\mu^2 + k^2)^{5/2}} . \tag{A1.5}$$

With $\mu \approx 1 GeV$, $D(k) \approx 0.12$. Thus $\Delta^B(k)$ is a soft correction fast decreasing with k as k^{-5} to the perturbative gluon propagator.

APPENDIX 2

Calculation of the Wilson-loop charge renormalization

Here we discuss in more detail calculation of graphs, corresponding to Figs.6-10, which contribute to the charge renormalization in the "Wilson" renormalization scheme, Eq.(9).

We start with the graph of Fig.6. A typical term contributing to the amplitude can be written as

$$J^{(6)} = \int \Delta_{\mu_1 \mu}(y_1, x) \frac{\partial}{\partial x_{\nu}} \Delta_{\mu \lambda}(x - y) \Delta_{\nu \rho}(x - y) \frac{\vec{\partial}}{\partial y_{\rho}} \Delta_{\lambda \mu_2}(y, y_2) \times dy_{1\mu_1} dy_{2\mu_2} d^4 x d^4 y$$
(A2.1)

Let the upper and lower horizontal lines in Fig.6 correspond to $y_{14} \in [0, T], y_{1i} = 0$ and $y_{24} \in [0, T], y_{21} = R$, respectively. According to our discussion in Section 4, the first and the last gluon propagators in (A 2.1) are not affected by confinement when $T \to \infty$ and one obtains in (A 2.1)

$$J^{(6)} \sim dx_4 \frac{d^3x}{|\vec{x}|} \partial_{\nu} \prod (x, y) \vec{\partial}_{\nu} \frac{d^4y}{|\vec{R} - \vec{y}|}.$$
 (A2.2)

Applying Fourier transformation to both propagators in (A 2.2) yields

$$J^{(6)} \sim T \frac{d^4 r \prod(r)}{|\vec{R} - \vec{r}|}$$
 (A2.3)

In case of no background field $\Pi(r)$ is simply a product of two free gluon propagators and one gets

$$J^{(6)} \sim \frac{T d^3 \vec{r}}{|\vec{R} - \vec{r}| |\vec{r}|^3} \sim \frac{T}{R} \ln \frac{R}{\delta}$$
 (A2.4)

where δ^{-1} is the UV cut-off parameter.

We introduce the following notation for the one-loop renormalization of α_s in the scheme (i)

$$\alpha_s(R) = \alpha_s^{(0)}(1 + f(R)\alpha_s^{(0)} + \dots) \tag{A2.5}$$

In the case of no background the diverging contribution of graphs Fig.6 and 7 to f(R) is [25]

$$f(R) = \frac{13 - 3\xi}{12\pi} C_2 \ln \frac{R}{\delta}$$
 (A2.6)

This should be compared to the momentum space calculation [17] of no background case (Feynman gauge, $\xi = 1$)

$$\alpha_s(Q) = \alpha_s^{(0)} \left(1 + \frac{\alpha_s^{(0)}}{4\pi} \left[\left(-\frac{5}{3}C_2\right) \left(-\frac{2}{\varepsilon} + C - \ln 4\pi + \ln \frac{Q^2}{\mu^2}\right) + \frac{31}{9}C_2 \right]$$
(A2.7)

where C=0.577 in the Euler constant and in the \overline{MS} scheme one should omit the factor $(\frac{2}{\varepsilon}+C-\ln 4\pi)$ in (A.2.7). We now consider the case of nonzero background field, and take $\Pi(r)$ as estimated in Appendix 3, Eq. (A3.21)

$$\prod(r) \approx \frac{exp(-m_1\sqrt{\bar{r}^2 + r_4^2})}{(\bar{r}^2 + r_4^2)^2}$$
 (A2.8)

Insertion of $\Pi(r)$ (A2.8) into (A2.3) yields instead of (A2.4)

$$J^{(6)} = \frac{1}{4\pi} \int \frac{Td^3\vec{r}}{|\bar{R} - \bar{r}| |\bar{r}|^3} \int_0^\infty \frac{e^{-m_1 r ch x} dx}{(ch x)^3} = TI(R)$$
 (A2.9)

where

$$I(R) = \int_0^\infty \frac{dx}{(chx)^3} \int_\delta^\infty \frac{dr}{r} e^{-m_1 r chx} \left(\frac{1}{R} \Theta(R-r) + \frac{1}{r} \Theta(r-R)\right) \quad (A2.10)$$

It is clear that the integral over dx converses at $x \sim 1$ and therefore one can replace in the integral over r, $m_1 chx \to m^*, m^* > m_1$. It is interesting to study two limiting cases: 1) $m^*R \ll 1$ and 2) $m^*R \gg 1$. One has

$$I(R) \approx \frac{1}{R} \ln \frac{Re}{\delta} , \ m^*R \ll 1$$

$$I(R) \approx \frac{1}{R} \ln \frac{1}{m^*\delta} , \ m^*R \gg 1$$
(A2.11)

We turn now to the grphs of Fig. 8 and 9. Due to the nonabelian character of fields the graph of Fig. 9 is proportional to

$$tr(t^a t^b t^a t^b) = (\frac{N^2 - 1}{2N})^2 N - \frac{N}{4}(N^2 - 1)$$
 (A2.12)

The first term on the r.h.s. of (A2.12) can be considered as the "abelian extrapolation" of the graph of Fig. 8 to the crossed kinematics of Fig. 9, while the second term on the r.h.s. of (A2.12) represents the pure nonabelian contribution. In the case of no background one obtains therefore the following nonabelian contribution to the Wilson loop from the graph of Fig. 9

$$\frac{1}{N}W^{(2)} = -\frac{g^4(N^2 - 1)}{(4\pi^2)^2 8} \int \frac{dy_{1\mu}dy_{2\mu}dx_{1\nu}dx_{2\nu}}{(y_1 - y_2)^2 (x_1 - x_2)^2}$$
(A2.13)

which gives the following divergent contribution to f(R) (cf(A2.5))

$$f^{(9)}(R) = \frac{(3-\xi)C_2}{2\pi R} \ln \frac{R}{\delta}$$
 (A2.14)

This should be compared with the momentum space calculation of [17] (we keep only divergent (logarithmic) term)

$$\Delta \alpha_s(Q) = -\frac{(\alpha_s^{(0)})^2}{4\pi} \frac{N^2 - 1}{2N} N \cdot (3 - \xi) \ln \frac{Q^2}{\lambda^2}$$
 (A2.15)

To simplify our discussion we shall use now the gauge $\xi=3$ and therefore one can disregard the graph of Fig. 9. This is true also in case of nonzero background, since we are interested here only in the modification of the divergent term $\ln \frac{R}{\delta}$ due to the background.

We now turn to the graph of Fig. 10.

Its contribution to f(R) is

$$\frac{f^{(10)}(R)}{R} = -\frac{3(1-\xi)}{16\pi^2} C_2 \int \frac{d^3x}{|\vec{x}||\vec{x}-\vec{R}|^3}$$
 (A2.16)

where $|\vec{x}|^{-1}$ enters from the static gluon propagator $\Delta(y_1 - x)$ while the favctor $|\vec{x} - \vec{R}|^{-3}$ is from the triangle part $\Gamma(x, y_2, y_3)$ of Fig. 10. In case of the confining background one has to insert in the integral in (A2.16) the factor, $exp(-m_1 | \bar{x} - \bar{R}|)$, which yields again the function I(R) as in (A2.9). This factor appears in the asymptotics of $\Gamma(x, y_2, y_3)$ due to the confining area law with the surface S_3 , shown in Fig. 14. The asymptotics holds true when $|\bar{x} - \bar{R}|$ is large and the mass m_1 is the same as in (A2.8), since at large $|x-y_2|$, $|x-y_3|$ and fixed $|y_2-y_3|$ the function $\Gamma(x, y_2, y_3)$ behaves as the two-point function $\Pi(x, y)$ discussed in Appendix 3.

Combining all contributions from all graphs of Fig. 6-10 one obtains finally

$$\frac{1}{R}f^{(total)}(R) = \frac{b_0}{2\pi}I(R) , b_0 = \frac{11}{3}N$$
 (A2.17)

For small R, $m_1R \ll 1$, one has due to (A2.11) the usual renormalization factor

$$\alpha_s(R) = \alpha_s^{(0)} (1 + \alpha_s^{(0)} \frac{b_0}{2\pi} \ln \frac{R}{\delta} + ...)$$
 (A2.18)

while for large R, $m_1R \gg 1$, the function RI(R) tends to a constant and the unlimited growth of the renormalized charge $\alpha_s(R)$ is frozen.

APPENDIX 3

Properties of the self-energy function $\prod(x,y)$

In this appendix we study the properties of the self-energy part $\Pi(x, y)$, Fig. , both in the coordinate and in the momentum space. We consider for simplicity spinless particles in the loop, which is true for the ghost loop, inclusion of spin is discussed in Appendix 4.

The FSR for $\prod(x,y)$ is [12]

$$\prod(x,y) = \int_0^\infty ds \int_0^\infty d\bar{s} (Dz)_{xy} (D\bar{z})_{xy} e^{-K-\bar{K}-V}$$
 (A3.1)

where V reduces to σS_{min} for large loops [13]

$$K = \int_0^s \frac{\dot{z}_\mu^2(\lambda)d\lambda}{4} \quad , \quad \bar{K} = \int_0^{\bar{s}} \frac{\dot{z}_\mu^2(\bar{\lambda})d\bar{\lambda}}{4} \quad ; \tag{A3.2}$$

Introducing as in [24] "center-of-mass" and relative coordinates, R_{μ} and r_{μ}

$$R_{\mu} = \frac{\bar{s}}{s + \bar{s}} z_{\mu}(\lambda) + \frac{s}{s + \bar{s}} \bar{z}_{\mu}(\lambda) \quad ; \quad r_{\mu} = z_{\mu}(\lambda) - \bar{z}_{\mu}(\lambda) \tag{A3.3}$$

and new variables $\mu, \bar{\mu}$ instead of s, \bar{s}

$$\mu = \frac{T}{2s} , \quad \bar{\mu} = \frac{T}{2\bar{s}} , \quad \tilde{\mu} = \frac{\mu\bar{\mu}}{\mu + \bar{\mu}}$$
 (A3.4)

we obtain

$$\prod(x,y) = T^2 \int_0^\infty \frac{d\mu}{2\mu^2} \int_0^\infty \frac{d\bar{\mu}}{2\bar{\mu}^2} DR \cdot Dre^{\{-\int_0^T d\tau \frac{(\mu+\bar{\mu})}{2} \dot{R}^2 + \frac{\bar{\mu}}{2} \int_0^T \dot{r}^2 d\tau + V\}}$$
(A3.5)

where $T \equiv |x - y|$. The boundary conditions for trajectories $R_{\mu}(\tau), r_{\mu}(\tau)$ are:

$$R_{\mu}(0) = y_{\mu}; \quad R_{\mu}(T) = x_{\mu}; r_{\mu}(0) = r_{\mu}(T) = 0$$
 (A3.6)

as disscussed in [24], one can replace $V = \sigma S_{min}$ by the expression $V = \sigma |\vec{r}(\tau)|$, with the accuracy of $\sim 5\%$ for the mass eigenvalues. In this case one can go over to the Hamiltonian form in (A 3.5) using the relation

$$\int DR \ Dr \ e^{-\int Ld\tau} = \langle R = x, r = 0 | e^{-\int Hd\tau} | R = y; r = 0 \rangle$$
 (A3.7)

Since the motion in R_{μ} and r_4 coordinates in free one can easily calculate its contribution and the result for $\prod(x,y)$ is

$$\prod(x,y) = \frac{1}{4(2\pi)^{5/2}} \int \int_0^\infty \frac{d\mu_1 d\mu_2 e^{-\frac{\mu_1 + \mu_2}{2}T}}{\tilde{\mu}^{3/2} \sqrt{T}} G(0,0;T)$$
(A3.8)

where $G(r_f, r_{in}; T)$ is the Green's function of relative quark-antiquark (or gluon-gluon) motion,

$$G(0,0;T) = \sum_{n} C_n(\tilde{\mu}) e^{-\varepsilon_n(\tilde{\mu})T}, \qquad (A3.9)$$

 $C_n(\tilde{\mu}) = |\psi_n(0)|^2, \psi_n$ and $\varepsilon_n(\tilde{\mu})$ are eigenvalues of the Hamiltonian

$$H\psi_n = \varepsilon_n \psi_n; H = \frac{\vec{p}^2}{2\tilde{\mu}} + \sigma |\vec{r}| . \tag{A3.10}$$

For linear potential $C_n(\tilde{\mu}) = \frac{\tilde{\mu}\sigma}{2\pi}$ and does not depend on n, while $\varepsilon_n(\tilde{\mu})$ is

$$\varepsilon_n(\tilde{\mu}) = (2\tilde{\mu})^{-1/3} \sigma^{2/3} a(n) \tag{A3.11}$$

where a(n) are numbers, obtained from the solution of the dimensioless equation, with the asymptotics

$$a(n) = (\frac{3\pi}{2})^{2/3} (n + \frac{1}{2})^{2/3} , \quad n \to \infty$$
 (A3.12)

One may get a false impression that expressions (A 3.9- A 3.12) are nonrelativistic. In fact relativism is contained here in the $\tilde{\mu}$ variable which finally yields spectrum very close to that of the relativistic quark model [24] or 1+1 QCD.

Insertion of (A 3.9) into (A 3.8) yields

$$\prod(x,y) = A \sum_{n} \int \frac{d\mu_1 d\mu_2}{\sqrt{\tilde{\mu}T}} e^{-M_n(\mu_1,\mu_2)T}$$
(A3.13)

where

$$M_n(\mu_1, \mu_2) = \frac{\mu_1 + \mu_2}{2} + \varepsilon_n(\tilde{\mu}) ,$$
 (A3.14)

with

$$A \equiv \frac{\sigma}{4(2\pi)^{7/2}}$$

At large $T \equiv |x - y|$ the asymptotics of (A 3.13) is obtained performing the $d\mu_1, d\mu_2$ integrations via the steepest descent method [24], which yields

$$\prod(x,y) = \frac{A\pi\sqrt{3}}{T^{3/2}} \sum_{n} \sqrt{\bar{M}_{n}} e^{-\bar{M}_{n}T}$$
(A3.15)

where $\bar{M}_n = 4\sqrt{\sigma}(\frac{a(n)}{3})^{3/4}$ is obtained from the minimum of $M_n(\mu_1, \mu_2)$ in variables μ_1, μ_2 .

Therefore at large T the correlator \prod decays exponentially

$$\prod (x,y) \sim \frac{exp(-\bar{M}_0 T)}{T^{3/2}}$$
 (A3.16)

where \bar{M}_0 is the minimal mass value.

At small T one should use (A 3.13) and take into account, that in this situation the sum can be replaced by the integral

$$\sum_{n} e^{-\varepsilon_{n}(\tilde{\mu})T} \to \int dn e^{-\lambda(\tilde{\mu})T(n+\frac{1}{2})^{2/3}} =$$

$$= \frac{3}{2} \frac{\Gamma(\frac{3}{2})}{(\lambda T)^{3/2}} , \quad \lambda = (2\tilde{\mu})^{-1/3} \sigma^{2/3} (\frac{3\pi}{2})^{2/3} . \tag{A3.17}$$

Insertion of (A 3.17) into (A 3.13) yields

$$\prod(x,y) = \frac{1}{4(2\pi)^4 T^2} \int_0^\infty d\mu_1 \int_0^\infty d\mu_2 e^{-\frac{(\mu_1 + \mu_2)T}{2}} = \frac{1}{16\pi^4 T^4}$$
 (A3.18)

This is the exact answer for the free scalar gluon loop diagram. It is remarkable that the sum over bound states reproduces the correct small distance behaviour.

Finally we shall write the answer for the momentum space. Choosing $p = (\vec{0}, p_4)$ one has

$$\prod(p) = \bar{A} \sum_{n} \int_{-\pi/2}^{\pi/2} \cos^2 \theta d\theta \sum_{n} \frac{\sqrt{M_n}}{(M_n - ip_4 \sin \theta)^{D-3/2}}$$
(A3.19)

with
$$M_n = m\sqrt{n + \frac{1}{2}}$$
, $m = \frac{4\sqrt{\sigma}}{3^{3/4}}(\frac{3\pi}{2})^{1/2}$, $\bar{A} = \frac{\sqrt{3}\sigma}{(2\pi)^{3/2}}\Gamma(D - \frac{3}{2})$.

At large n one can replace the sum by the integral of the type $(D = 4 + \varepsilon)$.

$$\int_{m_0}^{\infty} \frac{dM}{(M-ia)^{D-3}} = \frac{1}{\varepsilon} - \ln(m_0 - ip_4 \sin \theta)$$
 (A3.20)

Integration over $d\theta$ of the logarithmic term in (A 3.20) yields finally the effective logarithm which appears in (63)

$$\int_{-\pi/2}^{\pi/2} \cos^2 \theta d\theta \cdot \ln(m_0 - ip_4 \sin \theta) = \frac{\pi}{4} \ln(m_0^2 + p_4^2 \sin^2 \bar{\theta})$$
 (A3.21)

On another hand each term with a given n contributes a simple pole in $\Pi(p)$, which can be separated out looking more closely at the integration region near $\theta \approx \pm \frac{\pi}{2}$. Indeed writing $\sin \theta = \cos \eta \approx 1 - \frac{\eta^2}{2}$ one can rewrite the integral in (A 3.19) as

$$\int_0^1 \frac{\eta^2 d\eta}{(\delta + M_n \frac{\eta^2}{2})^{5/2}} \approx \frac{1}{\delta} (\frac{2}{M_n})^{3/2} \cdot \frac{1}{3}$$
 (A3.22)

where $\delta = M_n - ip_4$. Finally the contribution of the poles is

$$\prod(p) = \sum_{n} \frac{\sqrt{3}\sigma}{2\pi(M_n^2 + p^2)}$$
 (A3.23)

The sum (A 3.23) is diverging for large n, calling for the regularization which we have explicitly done in (A 3.20).

The fact that $\Pi(p)$ has only simple poles equidistantly placed at $-p^2 = M_n^2 = m^2(n+\frac{1}{2})$ tells us that the difference $\Pi(p) - \Pi(p_0)$ is proportional to the ψ function, $\psi(z) = \frac{\Gamma'(z)}{\Gamma(z)}$.

Indeed, one can write

$$\Pi(p) - \Pi(p_0) \sim \psi(\frac{p^2 + \frac{1}{2}m^2}{m^2})$$
 (A3.24)

which asymptotically at large $\frac{p^2}{m^2} \gg 1$ behaves as

$$\psi(z) \sim \ln z - \frac{1}{2z} - \sum_{n=1}^{\infty} \frac{B_{2n}}{2nz^{2n}}$$
 (A3.25)

where B_n are Bernulli numbers. In this way we again obtain the logarithmic term similar to (A 3.21).

Finally we derive one more form of $\Pi(x,y)$ which also was used in the main text. One can rewrite (A 3.15) separating in \bar{M}_n the constant term, not depending on n:

$$\bar{M}_n = m_0 + M(n) \tag{A3.26}$$

such that at $n=n_{min}, M(n_{min})=0$. Having in mind that $d\bar{M}_n^2=m^2dn$, one can rewrite (A 3.15) as

$$\prod(x,y) = \frac{A\sqrt{3}\pi}{T^{3/2}}e^{-m_0T} \int dM^2 M^{1/2}e^{-MT} \cong \frac{A\sqrt{3}\pi\Gamma(\frac{5}{2})e^{-m_0T}}{T^4}$$
(A3.27)

This is exactly the form used in the text, e.g. in Eq.(62), and in the Appendix 2, Eq.(A. 2.8).

APPENDIX 4

Gluon spin term in the gluon propagator

The gluon propagator in the background field, Eq.(8) can be written in the form

$$G = (\hat{D}_{\lambda}^2 \cdot \hat{1} + ig\hat{\Sigma}F)^{-1}$$
(A4.1)

where $\hat{1}$ and $\hat{\Sigma}$ are unit and spin matrices of the gluon respectively

$$ig(\sum F)_{\mu\nu} = -2igF_{\mu\nu} \tag{A4.2}$$

In the situation, when G enters in the gauge-invariant combination with two parallel transporters as in (26), (37), one can write the Wilson loop average as in (37) using the cluster expantion. E.g. for the adjoint Wilson loop

$$\langle W_{adj} \rangle = \langle \hat{\Phi}_F(x, y, c_1) \hat{\Phi}_F(y, x, c_2) \rangle$$
 (A4.3)

where $\hat{\Phi}_F$ is defined in (36), and the paths C_1, C_2 form the closed loop $C = C_1 + C_2$ one obtains [32]

$$< W_{adj} > = exp \sum_{n=1}^{\infty} \frac{(ig)^n}{n!} \int \Delta(1)...\Delta(n) \ll F(1)...F(n) \gg$$
 (A4.4)

We have used the following compact notation

$$\Delta(k) = d\sigma(u_k) - \sum_{k=0}^{(1)} d\tau_k + \sum_{k=0}^{(2)} d\bar{\tau}_k$$
 (A4.5)

with the condition, that e.g. the combination

$$\sum_{k=0}^{(i)} d\tau_k \ll F(1)...F(k)...F(n) \gg = \sum_{k=0}^{(i)} d\tau_k \ll F(1)...F(z(\tau_k))...F(n) \gg (A4.6)$$

where $z(\tau_k)$ lies on the path C_i . At the same time the combination $d\sigma(u_k)$ is multiplied with $F(u_k)$ where u_k lies on the surface S_{min} inside the contour C. Therefore keeping in $\Delta(k)$ only the first term $d\sigma(u_k)$ one obtains the usual area law at large distances [13]:

$$\langle W_{adj} \rangle_{no \ spin} = exp(-\sigma_{adj}S_{min})$$
 (A4.7)

In contrast to that the terms with $\Sigma^{(1)}$ or $\Sigma^{(2)}$ generate perimeter-type contribution. Indeed the cumulants $\ll F(1)...F(n) \gg$ are effectively nonzero when all coordinates are within the interval of the size T_g , vacuum correlation length, $T_g \approx 0.2 fm$ [31].

length, $T_g \approx 0.2 fm$ [31]. The operator $\sum^{(i)} d\tau_k$ enforces, as we discussed above, one of the coordinates of the cumulant to lie on the contour and therefore all others are within the distance T_g from the contour, leading to the perimeter- type contribution

$$\langle W_{adj} \rangle_{spin} = exp(-\sigma_{adj}S_{min} - bL)$$
 (A4.8)

where b is some constant and L – the length of the contour C.

For large contours C, which are studied in this paper, one can therefore neglect in the first approximation the gluon spin contribution as compared to the area-law term.

APPENDIX 5

IR renormalons and background fields

The integral in (68) can be rewritten as

$$I \equiv \int_0^{Q^2} \frac{k^2 dk^2}{Q^4} \ln^n \frac{Q^2 + m^2}{k^2 + m^2} = \int_0^{n_0} t^n e^{-2t} dt - \frac{m^2}{Q^2} \int_0^{n_0} t^n e^{-t} dt$$
 (A5.1)

where $n_0 = \ln \frac{Q^2 + m^2}{m^2}$.

The sum in (68) can be now calculated easily (the interchange of the order of summing and integration is possible because the integral is in finite limits, and the sum is converging uniformly for $t \leq n_0$). One has

$$\sum_{n=1}^{\infty} (\alpha_s(Q^2) t \frac{b_0}{4\pi})^n = \frac{\alpha_s(Q^2) t \cdot b_0 / 4\pi}{1 - \alpha_s(Q^2) \frac{b_0}{4\pi} t}$$
(A5.2)

and the contribution of I in the sum over n is

$$\frac{b_0}{4\pi}\alpha_s(Q^2)\int_0^{n_0} \frac{e^{-2t}tdt}{1 - \alpha_s(Q^2)\frac{b_0}{4\pi}t} = \int_0^{n_0} \frac{tdte^{-2t}}{t_p - t} \sim \frac{1}{t_p}$$
(A5.3)

where

$$t_p = \frac{4\pi}{b_0 \alpha_s(Q^2)} = \ln \frac{Q^2 + m^2}{\Lambda^2} > n_0 ,$$
 (A5.4)

since $m > \Lambda$.

Insertion of (A 5.3) into (68) finally yields

$$\Delta \prod = \frac{\alpha_s(Q^2)}{8\pi^3} \cdot \frac{1}{t_n} \cong \frac{b_0 \alpha_s^2(Q^2)}{32\pi^4}$$
(A5.5)

This result is used in Eq. (73) of the main text.

References

- [1] L.V.Gribov, E.M.Levin and M.G.Ryskin, Phys. Rep. 100 (1983) 1.
- [2] D.J.Gross and F.Wilczek, Phys. Rev. Lett. 30, 1323 (1973)
 H.D.Politzer, Phys.Rev. Lett. 30, 1346 (1973).
- [3] L.D.Landau, A.A.Abrikosov and I.M.Khalatnikov, Dokl. AN SSSR, 1954, 95, 773.
- [4] V.B.Berestetsky, E.M.Lifshitz and L.P.Pitaevski, Quantum Electrodynamics, Moscow, Nauka, 1980.
- [5] G.'tHooft, in "The Whys of Subnuclear Physics", Ecice 1977, ed. A.Zichichi, Plenum, NY and London, p.94.
 G.Parisi, phys. Lett. B.76, 65 (1977)
 B.Lautrup, Phys. Lett. 69, 109 (1977).
- [6] M.A.Shifman, A.I.Vainshtein and V.I.Zakharov, Nucl. Phys. B147, 385 (1979).
- [7] K.Wilson, Phys. Rev. **179**, 2499 (1969).
- [8] M.Campostrini, A.Di Giacomo, Phys. Lett. B 225, 393 (1989).
- [9] B.S.De Witt, Phys Rev. 162, 1195 and 1239 (1967)
 J.Honerkamp, Nucl. Phys. B48, 269 (1972).
- [10] G.'tHooft, Nuc. Phys. 62, 444 (1973)
 G.'tHooft, Lecture given at Karpacz (1975) in: Acta Universitatis Wratislaviensis, 368, 345 (1976).
- [11] L.F.Abbot, Nuc. Phys. **185**, 189 (1981).
- [12] Yu. A.Simonov, Nucl. Phys. 307, 512 (1988), Yad. Fiz. 54, 192 (1991) and refs. therein.
- [13] H.G.Dosch, Phys. Lett. **B190**, 177 (1987)
 H.G.Dosch and Yu. A.Simonov, Phys. Lett. **205** (1988) 339.
- [14] L.D.Faddeev, V.N.Popov, Phys. Lett. **B25**, 29 (1967).

- [15] L.D.Faddeev and A.A.Slavnov, Introduction into quantum gauge theory, Moscow, Nauka, 1978.
 M.B.Voloshin and K.A.Ter-Martirosyan, Gauge theory of elementary particle, Moscow, Energoatomizdat, 1984.
- [16] T.D.Lee, Particle Physics and Introduction to Field Theory, Harwood, Chur, London, NY, p. 447.
- [17] S.N.Gupta, S.F.Redford and W.W.Repko, Phys. Rev. **D24** 2309 (1981), ibid. **D26**, 3305 (1982).
- [18] M.Lüscher et al. DESY 92-025 and 92-157.
- [19] Yu.A.Simonov, preprint ITEP 3-93.
- [20] V.Marquard and H.G.Dosch, Phys. Rev. **35** D, 2238 (1987)
- [21] M.Creutz, Quarks, Gluons and Lattices, Cambridge Univ. Press, Cambridge, 1983.
- [22] Yu.A.Simonov, Nucl. Phys. **B324**, 67 (1989)
- [23] I.I.Balitsky, Nucl. Phys. **B254**, 166 (1985).
- [24] Yu.A.Simonov, Yad. Fiz. **54**, 192 (1991).
- [25] V.S.Dotsenko and S.N.Vergeles, Nucl. Phys. **169**, 527 (1980).
- [26] G.'tHooft, Nuc. Phys. **B72**, 461 (1974).
- [27] Yu.A.Simonov, preprint TPI-MINN-90/19-T (unpublished) Phys. Lett **249** 514 (1990).
- [28] D.K.Sinclair, in Lattice '89 Conference, Nucl. Phys. B(Proc.
- [29] A.H.Mueller, Nucl. Phys. **B250**, 327 (1985).
- [30] V.I.Zakharov, MPI-Ph/92-33.
- [31] A.Di Giacomo and H.Panagopoulos, Phys.Lett. B285 (1992) 133; M.Campostrini, A.Di Giacomo and G.Mussardo, Z.Phys. C25 (1984) 173.
- [32] H.G.Dosch and Yu. A. Simonov, HD-THEP-92-23.

Figure captions

- Fig.1. The rectangular Wilson loop of the size $R \times T$ used to define the static quark energy E(R)
- Fig. 2. The gluon loop with external lines corresponding to the background field B_{μ} , used in [11] to calculate β function.
 - Fig. 3. The same as in Fig.2 but for the ghost loop.
- Fig. 4. The nonperturbative generalization of the diagrams of Fig.2 and 3 with the full account of the background in the propagators of the gluon (a) and the ghost (b).
- Fig. 5. One-gluon-exchange in the Wilson loop between the points x and y. Indices α, δ and β, γ are the color fundamental indices of the gluon at points x and y respectively.
- Fig. 6. The gluon-loop diagram for the gluon exchange inside Wilson loop.
 - Fig. 7. The same for the ghost-loop diagram.
- Fig. 8. Two-gluon exchange diagram from iteration of one-gluon-exchange interaction.
- Fig. 9. Two-gluon exchange crossed diagram, containing both iteration of one-gluon exchange (abelian part) and the nonabelian part.
 - Fig. 10. Gluon exchange with the three-gluon vertex in the Wilson loop.
- Fig.11. At large N_c the one-gluon-exchange diagram of Fig. 5 goes over into the product of two Wilson loops.
 - Fig. 12. Multiple iterative gluon exchanges.
- Fig.13. At large N_c the diagrams of Fig.6 and 7 contain three Wilson loops with areas S_1, S_2, S_3 .
- Fig.14. At large N_c the diagram of Fig.10 contains three Wilson loops with areas S_1, S_2, S_3 .
 - Fig.15. The diagrams of Fig.4 contain at large N_c three Wilson loops.
- Fig.16. Cutting the diagram of Fig.15 along the dash-dotted line reveals two gluons 1 and 2, connected by two fundamental strings. One of the strings passes necessarily through the point 3, lying on the parallel transporter $\Phi(B)$, shown in Fig. 4 and Fig. 15.
- Fig.17. Diagrams containing the renormalon contribution to the correlator of e.m. currents.

This figure "fig1-1.png" is available in "png" format from:

http://arxiv.org/ps/hep-ph/9311247v1

This figure "fig1-2.png" is available in "png" format from:

http://arxiv.org/ps/hep-ph/9311247v1

This figure "fig1-3.png" is available in "png" format from:

http://arxiv.org/ps/hep-ph/9311247v1