

Curvature Radiation by Ultrarelativistic Protons*

V. Berezhinsky ^{a,b}, A. Dolgov ^{c,d} and M. Kachelrieß ^{a,e}

^a INFN, Laboratori Nazionali del Gran Sasso, I-67010 Assergi (AQ), Italy

^b Physikalisches Institut der Universität Bonn, D-53115 Bonn, Germany

^c Institut for Experimental and Theoretical Physics, 117259 Moscow, Russia

^d Dept. Física Teórica, Universidad de Valencia, E-46100 Burjassot, Spain

^e Theoretische Physik I, Ruhr-Universität Bochum, D-44780 Bochum, Germany

We study pion curvature radiation by a proton, i.e. pion emission by a proton moving along a curved trajectory. We suggest an approximate semiclassical solution and the exact solution for which we assume that a proton moves in a fictitious magnetic field with the Larmor radius equal to the curvature radius of the real trajectory. As possible application we consider the pion radiation by ultrahigh energy protons moving along curved magnetic field lines. Such situation can occur in the magnetosphere of a young pulsar, in the magnetosphere of the accretion disk around a black hole, and in the vicinity of a superconducting cosmic string. The decay products of these pions, such as high energy photons or neutrinos, can give the observable consequences of the considered mechanism.

I. INTRODUCTION

According to the commonly used terminology, curvature radiation is electromagnetic radiation by a charged particle moving along a curved trajectory. Naturally, the implied mechanism can be generalized to the emission of any kind of particles. A simple example is pion curvature radiation by a proton. Protons can move along a curved trajectory in the gravitational field, e.g. in the vicinity of a black hole. Another example is the movement of a proton along a magnetic field line in the case of a strong field, e.g. in the magnetosphere of a pulsar or of an accretion disc. The transverse momentum of a proton in this case is suppressed due to synchrotron radiation.

This formulation of the problem is semiclassical by its nature: we define the interaction between particles in a quantum way, but introduce at the same time the classical concept of a trajectory, which does not exist in quantum field theory. We suggest here some solutions to the problem for the case of π^0 emission by a proton and indicate possible implications.

II. KINEMATICS AND DYNAMICAL PARAMETER OF THE PROBLEM

To make a quantitative insight into the problem let us consider a proton moving along a curved trajectory with local curvature radius R_C . We introduce the coherent interaction length, l_{int} , along the trajectory on which the pion is radiated. The four-momentum of the proton at the beginning and the end of l_{int} is denoted by (E_i, \vec{p}_i) and (E_f, \vec{p}_f) respectively, and that of a pion by (ϵ, \vec{k}) . We describe the effect of the curved trajectory in the interaction region by the longitudinal and transverse (relative to \vec{p}_i) momenta q_{\parallel} and q_{\perp} given by some external field and neglect the change of proton energy ($|q| \ll |p|$). Then we have

$$E_f = E_i - \epsilon, \quad \vec{p}_f = \vec{p}_i - \vec{k} + \vec{q}. \quad (1)$$

The interaction length is determined by q_{\parallel} , $l_{int} \approx 1/q_{\parallel}$. From the set of equations

$$q_{\perp} \approx p_i \theta, \quad q_{\parallel} \approx p_i(1 - \cos \theta) \approx \frac{1}{2} p_i \theta^2 \quad \text{and} \quad \theta \approx l_{int}/R_C \approx 1/(q_{\parallel} R_C), \quad (2)$$

we find q_{\parallel} and q_{\perp} as

$$q_{\perp} \approx m \left(\frac{p_i^2}{m^3 R_C} \right)^{1/3} \quad \text{and} \quad q_{\parallel} \approx \frac{1}{2} \frac{q_{\perp}^2}{p_i}, \quad (3)$$

*Accepted for publication in *Physics Letters B*

where m is the proton mass.

Naively, the natural parameter of the problem is $1/(mR_C)$. However, we shall choose as a parameter the quantity proportional to this value, namely the quantity in the brackets of Eq. (3):

$$\chi = E_i^2/(m^3 R_C) . \quad (4)$$

The motivation for this choice is given by the connection with the fictitious magnetic field method. Let us assume that the considered part of the trajectory is caused by a fictitious magnetic field B perpendicular to the trajectory plane. This means that the Larmor radius $R_L = E/(eB)$ equals the local curvature radius R_C . Then Eq. (4) gives

$$\chi = (E_i/m)(B_\perp/B_{cr}) , \quad (5)$$

where $B_{cr} = m^2/e$ and χ is the standard parameter [1,2]

$$\chi = e\sqrt{(F_{\mu\nu}p^\mu)^2}/m^3 \quad (6)$$

relevant for processes in strong electromagnetic fields. In this paper we shall confine ourselves to the quantum case $\chi \gg 1$ when the curvature radius is small, $R_C \ll E^2/m^3$.

Two remarks are in order.

The critical magnetic field in our problem is connected with the proton mass and is equal to $B_{cr} \approx 1.5 \cdot 10^{20}$ G.

One should distinguish between the parameter χ for the curvature radiation given by Eq. (4) and the parameter χ_{em} relevant for the real magnetic field in the considered system. The latter is given by Eq. (6) and results in Eq. (5) with B_\perp being the component of the *real* magnetic field B perpendicular to the proton momentum. The difference between these two parameters, χ given by Eq. (4) and χ_{em} given by Eq. (5), is very essential for a proton moving along a curved magnetic field line. The synchrotron radiation of the proton in the real magnetic field results in the loss of transverse momentum in the regime when $\chi_{em} \ll 1$:

$$\frac{1}{p_\perp} \frac{dp_\perp}{dt} = \frac{2}{3} \frac{e^2 m}{\gamma} \chi_{em}^2 , \quad (7)$$

where γ is the Lorentz factor of the proton. For $R_C < 10^8$ cm the lifetime given by (7) in the regime $\chi_{em} \ll 1$ is shorter than the characteristic time of flight and therefore a proton moves along the magnetic field line [3]. At the same time the curvature radiation parameter (4) can be larger (or much larger) than 1, providing thus pion production due to curvature radiation. In other words, this movement is characterized by $\chi_{em} \ll 1$ and $\chi > 1$ (or $\chi \gg 1$).

III. SEMICLASSICAL SOLUTION

Let us consider a proton moving along a curved trajectory. As it is described in the Introduction we single out the interaction region according to Eq. (3) and take the wavefunctions of the proton at the boundaries as plane waves ψ_i and ψ_f . Then the matrix element describing the interaction is

$$M = g\bar{\psi}_f(p_f)\gamma^5\psi_i(p_i)\phi(k) , \quad (8)$$

where p_i, p_f and k are determined by Eq. (1). The probability of the transition is calculated in the standard way as

$$d\Gamma = \frac{(2\pi)^4}{2E_i} \frac{1}{2} |M|^2 \frac{d^3k}{(2\pi)^3 2\epsilon} \frac{d^3p_f}{(2\pi)^3 2E_f} \delta(E_i - E_f - \epsilon) \delta(\vec{p}_i - \vec{p}_f - \vec{k} + \vec{q}) . \quad (9)$$

In the evaluation of the spinor traces we use the kinematical relations given by Eqs. (1) and (3). As a result we obtain

$$\Gamma = \frac{3g^2}{64\pi E_i} q_\perp^2 = \frac{3}{16} \alpha_s \frac{m^2}{E_i} \chi^{2/3} , \quad (10)$$

where $\alpha_s = g^2/4\pi$ and χ is given by Eq. (4) or in terms of the fictitious magnetic field by Eq. (5).

IV. FICTITIOUS MAGNETIC FIELD METHOD

In this approach we assume that the curved trajectory is caused by a fictitious magnetic field B which is chosen so that $R_L = R_C$. Therefore we now have to calculate pion radiation of a proton in a strong magnetic field. This process was already considered in Ref. [4] using the crossed field method [1] (see also Ref. [5]). Here, we shall calculate the probability of this process describing it as a transition of a proton from the Landau level N_i with polarization $\tau_i = \pm 1$ into the Landau level N_f due to the interaction $\mathcal{L}_{int} = g\psi_f\gamma^5\psi_i\phi$. The transition probability for a proton from the Landau level N_i to N_f is given by

$$\Gamma_{\pm}^{N_i \rightarrow N_f} = \frac{\alpha_s}{16^2} \sum_{\tau_f} \int_0^\pi d\theta \sin \theta \, k \, |\mathcal{M}|^2, \quad (11)$$

where θ is the angle between the pion momentum \vec{k} and the magnetic field. The matrix element squared $|\mathcal{M}|^2$ is given in terms of I -functions. For the very large Landau quantum numbers considered here, we can use an asymptotic expansion in terms of modified Bessel functions (for the notations used here and hereafter see Ref. [2]):

$$I_{N_f, N_i}(x) = \frac{(1 + \xi y)^{1/2}}{\pi \sqrt{3}} \sqrt{\mu} K_{1/3}(z) \quad (12)$$

where

$$x = \frac{k_\perp^2}{2eB}, \quad z = \frac{2}{3} (x_0^2 N_i N_f)^{1/4} \left(1 - \frac{x}{x_0}\right)^{3/2}, \quad x_0 = \left(\sqrt{N_i} - \sqrt{N_f}\right)^2 \quad (13)$$

$$\xi = \frac{3}{2} \frac{B}{B_{cr}} \frac{E_i}{m} = \frac{3}{2} \chi, \quad y = \frac{2E_i \epsilon \mu^{3/2}}{3eB(1 - \epsilon/E_i)} \quad (14)$$

$$\mu = 1 - \frac{2N_i eB}{E_i^2} \sin^2 \theta \quad \text{and} \quad \mu_0 = 1 - \frac{2N_i eB}{E_i^2}. \quad (15)$$

Furthermore, since pions are emitted mostly in the forward direction, i.e. the transition probability is strongly peaked at $\theta = \pi/2$, we can approximate all functions except functions of $1 - x/x_0$ by their values at $\theta = \pi/2$. Then we take the ultrarelativistic limit of the normalization constants of the spinors and obtain

$$\begin{aligned} |\mathcal{M}|^2 = & \frac{1 + \xi y}{3\pi^2} \left[\frac{1}{2} (1 - \tau_i \tau_f) D^2 \mu K_{1/3}^2(z) \right. \\ & - \frac{\xi y}{2} [(1 - \tau_i \tau_f) D^2 - \tau_i (1 - \tau_i \tau_f) D(E - 1)] \mu^{3/2} K_{1/3}(z) K_{2/3}(z) \\ & \left. + \frac{(\xi y)^2}{8} [(1 - \tau_i \tau_f)(1 - 2E + D^2 + E^2) - 2\tau_i (1 - \tau_i \tau_f) D(E - 1)] \mu^2 K_{2/3}^2(z) \right], \end{aligned} \quad (16)$$

with $D = m/(2E_f) - m/(2E_i)$ and $E = m/(4E_i E_f)$. Note that because of the pseudoscalar nature of the interaction only spin-flip transitions occur in the ultrarelativistic case.

Using now some identities for the Bessel functions and summing over the polarization states of the proton we obtain for the spin averaged emission rate

$$\Gamma^{N_i \rightarrow N_f} = \frac{\alpha_s}{4} k \frac{y \xi^2 \mu_0^{5/2} (1 + \xi y)}{3\sqrt{3}\pi} \left(\int_y^\infty dx K_{5/3}(x) + K_{2/3}(y) \right). \quad (17)$$

For the case $\xi \gg 1$ we can use for the functions $K_\mu(y)$ their asymptotic expression $K_\mu(y) = 2^{\mu-1} \Gamma(\mu) y^{-\mu}$ for $y \rightarrow 0$. Then, summing over all final Landau levels yields the total decay width Γ

$$\Gamma = \frac{2^{2/3} \Gamma^2(\frac{2}{3}) \Gamma(\frac{1}{3})}{18\sqrt{3}\pi} \alpha_s \frac{m^2}{E_i} \xi^{2/3} \approx \frac{1}{12.6} \alpha_s \frac{m^2}{E_i} \xi^{2/3}. \quad (18)$$

Our width (18) differs only by a numerical factor from the result of Ref. [4]. This factor, $9\Gamma(2/3)\Gamma(1/3)/(2\sqrt{3}\pi) \approx 3.0$, is however rather large.

On the other hand, the agreement between our semiclassical solution (10) and the Eq. (18) is surprisingly good. Both methods give not only the same functional dependence on E_i and χ , but differ only by a numerical factor 1.8. Because of the uncertainties in the kinematical relations used in the semiclassical calculation we estimate the uncertainty in the decay width by a factor 4 – 5. However, for two other processes in strong magnetic fields, $\nu \rightarrow \nu + e^+ + e^-$ and $\gamma \rightarrow e^+ + e^-$, we obtained excellent agreement between both methods.

V. ASTROPHYSICAL APPLICATIONS

The pion curvature radiation needs protons of extremely high energies. From condition $\chi \geq 1$ one obtains from (7)

$$E_i \geq 6.5 \cdot 10^9 R_6^{1/2} \text{ GeV},$$

where $R_6 = R_C/10^6$ cm is the curvature radius in units of 10^6 cm. In cosmic rays the particles are observed with energies up to $2 \cdot 10^{11}$ GeV. The particles with the highest energies are most probably protons. In the strong magnetic field inside a source protons of these energies move along the magnetic field lines [3]. Since the proton momentum is essentially parallel to the magnetic field the electromagnetic parameter, $\chi_{em} \ll 1$ and pion production in the strong magnetic field is absent. However, the curvature parameter (7) can be large if the curvature radius of the magnetic field line is sufficiently small and therefore the pion curvature radiation is efficient. Decays of pions result in the production of gamma and neutrino radiation.

An example can be given by young pulsar.

Independently from the concrete model of pulsar magnetospheres, the maximum potential ϕ_{max} on the surface of the pulsar is [6]:

$$\phi_{max} \geq 8.3 \cdot 10^{17} (\Omega/10^3 \text{ s}^{-1})^2 (B_s/10^3 \text{ G}) (R_s/10^6 \text{ cm})^3 \text{ V}, \quad (19)$$

where R_s is the star radius, Ω is the angular velocity and B_s is the surface magnetic field.

Therefore, pulsars with $\Omega \geq 3 \cdot 10^3 \text{ s}^{-1}$ can, in principle, accelerate protons to energies higher than $1 \cdot 10^{10}$ GeV. If this potential drop occurs nearby the surface of the pulsar (which is actually not the case of the modern models) the ultrahigh energy protons escape from the pulsar magnetosphere mostly along the first-open magnetic field line [6]. The curvature radius of this line is

$$R_C = 4/3 (R_S R_L)^{1/2}, \quad (20)$$

where $R_L = c/\Omega$ is the radius of the light cylinder. For $\Omega = 3 \cdot 10^3 \text{ s}^{-1}$ and $R_S = 1.2 \cdot 10^6$ cm, one obtains from Eq. (19) that the energy of accelerated protons could be higher than $1.3 \cdot 10^{10}$ GeV and the parameter (4) is $\chi \geq 1$.

Another example is an accretion disc, with a strong magnetic field, around a black hole with a mass $M \sim M_\odot$. The potential at the inner edge of the disc is [7]

$$\phi = \frac{1}{c} \int \vec{E} d\vec{l} = \frac{1}{c} \int v(r) B_\phi(r) dr = \frac{1}{\sqrt{6}} B_0 r_0 \ln \frac{R}{r_0}, \quad (21)$$

where r_0 , R and B_0 are the radii of the inner and outer edges of the disc and the magnetic field at the inner edge, respectively. Taking $E_i \sim e\phi$, $r_0 \sim R_C \sim 10^6$ cm and $B_0 \sim 10^{10}$ G we again obtain $\chi \geq 1$.

Finally, let us discuss superconducting cosmic strings. Cosmic strings are topological defects caused by symmetry breaking in the early universe. Under certain assumptions the string can become superconducting [8]. Due to the expansion of the universe the primordial magnetic field induces an electric current J in the string which grows up to the maximal value $J_{max} \approx eM/2\pi$, where M is the scale of the GUT symmetry breaking. Starting from this moment, the particles with GUT masses are evaporating from the string. Decays of these particles produce leptons and hadrons with maximum energies up to $M_{GUT} \approx 10^{16}$ GeV.

The string current produces a magnetic field with Biot-Savart geometry. The initially produced particles have momenta perpendicular to the magnetic field. Therefore, at a first stage the processes in the strong magnetic field dominate [9]. After losing energy, particles are captured in circular orbits [10] and the pion curvature radiation becomes one of the main energy loss mechanisms of high energy protons.

VI. CONCLUSIONS

We have studied the curvature radius of pion by a proton. The parameter of this problem is $\chi = E_i^2/(m^3 R_C)$, where R_C is the curvature radius of the proton trajectory and E_i is the proton energy.

For the case $\chi \gg 1$ we found the quasiclassical solution assuming that in the region of coherent interaction the trajectory gives the additional momentum \vec{q} to the proton. The parallel component of this momentum, \vec{q}_{\parallel} , determines the length of coherent interaction $l_{int} \sim 1/q_{\parallel}$. Then the perpendicular momentum, q_{\perp} , is found as $q_{\perp} \approx p_i \theta$, where p_i is the proton momentum and $\theta \approx l_{int}/R_C$. The wavefunctions of the proton at the boundaries of the interaction region, l_{int} , are taken as plane waves and the matrix element is calculated as $g\bar{u}(p_f)\gamma_5 u(p_i)\phi$ with p_f and p_i related as $\vec{p}_f = \vec{p}_i - \vec{k} + \vec{q}$, where \vec{k} is the pion momentum. The width of the proton decay, $p \rightarrow p + \pi^0$ is given by

$$\Gamma \approx 0.19 \alpha_s m^2 \chi^{2/3} / E_i. \quad (22)$$

We estimate the uncertainties of this method by a factor 4 – 5.

The other way of calculation is given by the method of fictitious magnetic field. Here we assume that the proton moves along the trajectory, due to a magnetic field, so that locally $R_L = R_C$, where R_L is the Larmor radius of the trajectory. We calculated the width of proton decay in this case as a probability of a transition between different Landau levels and found

$$\Gamma \approx 0.104 \alpha_s m^2 \chi^{2/3} / E_i \quad (23)$$

in reasonable agreement with the quasiclassical method.

However, the early calculations of Ref. [4] for the $p \rightarrow p + \pi^0$ decay in the strong magnetic field, made using a different approach, differ from our calculations by factor 3.

As applications we consider the astrophysical sources where high energy protons are moving along magnetic field lines. Since the proton momentum is essentially parallel to the magnetic field, the electromagnetic parameter $\chi_{em} \ll 1$, while the curvature radiation parameter can be $\chi > 1$ (in the case of sufficiently small R_C). These conditions are met in such sources as young pulsars, accretion discs around black holes, and superconductive cosmic strings.

ACKNOWLEDGMENTS

We are grateful to A. I. Nikishov for comments and for pointing out Ref. [4]. V.B. thanks the Institute of Physics of the Bonn University for hospitality and the Humboldt Foundation for the award of 1991. M.K. acknowledges a grant by Deutscher Akademischer Austauschdienst.

-
- [1] A. I. Nikishov and V. I. Ritus, JETP 46 (1964) 529 and 1191; V. I. Ritus and A. I. Nikishov, Proceedings of the P. N. Lebedev Institute (ed. N. G. Basov), v. 111, Quantum Electrodynamics in Strong Fields (1975, in Russian).
 - [2] A. A. Sokolov and I. M. Ternov, Synchrotron Radiation (Akademie-Verlag, Berlin 1968); Radiation from Relativistic Electrons (AIP, New York 1986).
 - [3] P. G. Sutherland, Fundamentals of Cosmic Physics 4 (1979) 95.
 - [4] G. F. Zharkov, Sov. J. Nucl. Phys. 1 (1965) 173.
 - [5] V. L. Ginzburg and G. F. Zharkov, JETP 47 (1964) 2279.
 - [6] V. S. Berezhinskii, S. V. Bulanov, V. A. Dogiel, V. L. Ginzburg (ed.) and V. S. Ptuskin, Astrophysics of Cosmic Rays (North-Holland, Amsterdam 1990).
 - [7] R. D. Blandford, Mon. Not. R. Soc. 176 (1977) 465; R. D. Blandford and R. L. Znajek, Mon. Not. R. Soc. 179 (1977) 433.
 - [8] E. Witten, Nucl. Phys. B 249 (1985) 557.
 - [9] V. Berezhinsky and H. R. Rubinstein, Nucl. Phys. B 323 (1989) 95.
 - [10] W. B. Perkins, Nucl. Phys. B 364 (1991) 451.