## Istituto Nazionale Fisica Nucleare

Sezione SANITÀ Istituto Superiore di Sanità Viale Regina Elena 299 I-00161 Roma, Italy

> INFN-ISS 97/2 April 1997

## Rare decays $B \to (K, K^*)(\ell^+ \ell^-, \nu \bar{\nu})$ in the quark model<sup>a</sup>

D. Melikhov<sup>(\*)</sup>, N. Nikitin<sup>(\*)</sup> and S. Simula<sup>(\*\*)</sup>

 (\*)Nuclear Physics Institute, Moscow State University Moscow, 119899, Russia
 (\*\*)Istituto Nazionale di Fisica Nucleare, Sezione Sanità Viale Regina Elena 299, I-00161 Roma, Italy

## Abstract

Long-distance effects in exclusive rare semileptonic transitions  $B \to (K, K^*)$  are analysed within a relativistic quark model. The meson transition form factors, describing the meson amplitudes of the effective weak Hamiltonian, are calculated within the dispersion formulation of the quark model as relativistic double spectral representations through the wave functions of the initial and final mesons. The dilepton spectra and lepton asymmetries are considered within the framework provided by the Standard Model. It is found that, while the non-resonant decay rates are very sensitive to quark model parameters, the model dependence of the predicted dilepton forward-backward and lepton polarization asymmetries is remarkably small, providing only an overall ~ 10% uncertainty.

PACS numbers: 13.20.He,12.39.Ki, 12.39.Pn

Keywords: decays of bottom mesons; relativistic quark models.

<sup>&</sup>lt;sup>a</sup>To appear in Physics Letters B.

The investigation of rare semileptonic decays of the B meson induced by the flavourchanging neutral current transitions  $b \to (s, d)$  provides an important test of the Standard Model (SM) and its possible extensions. Rare decays are forbidden at tree level and occur at the lowest order only through one-loop diagrams. This fact opens the possibility to probe at comparatively low energies the structure of the theory at large mass scales, thanks to the contributions of virtual particles in the loops. However, in order to reliably separate (perturbative) short-distance effects, the (non-perturbative) long-distance contributions, entering the amplitudes of exclusive rare decays, should be known with enough accuracy. The theoretical investigation of these contributions encounters the problem of describing the hadron structure, which yields the main uncertainty in the predictions of exclusive rare decays. In this letter exclusive rare semileptonic transitions  $B \to (K, K^*)$  are analysed within the framework of the SM and adopting a relativistic constituent quark (CQ) model. It is shown that the differential decay rates are very sensitive to quark model parameters, whereas the model dependence of the predicted dilepton forward-backward and lepton polarization asymmetries is quite small, providing only an overall ~ 10% uncertainty.

1. Operator basis. The effective weak Hamiltonian, which describes the  $b \to s\ell^+\ell^-$  transition, has the following form [1]

$$\mathcal{H}_{eff} = -\frac{4G_F}{\sqrt{2}} V_{tb} V_{ts}^* \sum_i C_i(\mu) \ O_i(\mu) \tag{1}$$

where  $G_F$  is the universal Fermi constant, the quantities  $C_i(\mu)$  are the Wilson coefficients, obtained after integrating out the heavy particles, and the  $O_i$ 's are the basis operators; within the SM, the operators providing the main contribution to rare decays are [2, 3]

$$O_{1} = \frac{1}{4} (\bar{s}_{\alpha} \gamma^{\mu} (1 - \gamma_{5}) b_{\alpha}) (\bar{c}_{\beta} \gamma_{\mu} (1 - \gamma_{5}) c_{\beta}),$$

$$O_{2} = \frac{1}{4} (\bar{s}_{\alpha} \gamma^{\mu} (1 - \gamma_{5}) b_{\beta}) (\bar{c}_{\beta} \gamma_{\mu} (1 - \gamma_{5}) c_{\alpha}),$$

$$O_{7} = \frac{e}{32\pi^{2}} \bar{s}_{\alpha} \sigma_{\mu\nu} [m_{b} (1 + \gamma_{5}) + m_{s} (1 - \gamma_{5})] b_{\alpha} F^{\mu\nu},$$

$$O_{9} = \frac{e^{2}}{32\pi^{2}} (\bar{s}_{\alpha} \gamma^{\mu} (1 - \gamma_{5}) b_{\alpha}) (\bar{\ell} \gamma_{\mu} \ell),$$

$$O_{10} = \frac{e^{2}}{32\pi^{2}} (\bar{s}_{\alpha} \gamma^{\mu} (1 - \gamma_{5}) b_{\alpha}) (\bar{\ell} \gamma_{\mu} \gamma_{5} \ell),$$
(2)

In Eq. (1) the renormalization scale  $\mu$  is usually chosen to be  $\mu \simeq m_b$  in order to avoid large logarithms in the matrix elements of the operators  $O_i$ . The Wilson coefficients  $C_i$  reflect the specific features of the theory at large mass scales; they are calculated at the scale  $\mu \simeq M_W$ and then evolved down to  $\mu = m_b$  by the renormalization group equations. The analytic expressions for  $C_i(\mu)$  in the SM can be found, e.g., in [2]. In what follows, the values of the Wilson coefficients at the scale  $\mu = m_b = 5 \text{ GeV}$  are [3]:  $C_1(m_b) = -0.235$ ,  $C_2(m_b) = 1.1$ ,  $C_7(m_b) = -0.333$ ,  $C_9(m_b) = 4.09$  and  $C_{10}(m_b) = -4.32$ . The four-quark operators  $O_1$  and  $O_2$  generate both short- and long-distance contributions to the effective weak Hamiltonian (1). Both contributions can be taken into account by replacing  $C_9(m_b)$  with an effective coefficient  $C_9^{eff}(m_b, q^2)$  given by [3]

$$C_{9}^{eff}(m_{b},q^{2}) = C_{9}(m_{b}) + \left[3C_{1}(m_{b}) + C_{2}(m_{b})\right] \cdot \left[h(\frac{m_{c}}{m_{b}},\frac{q^{2}}{m_{b}^{2}}) + \frac{3}{\alpha_{em}^{2}}\kappa \sum_{V_{i}=J/\psi,\psi',\dots}\frac{\pi\Gamma(V_{i}\to\ell\ell)M_{V_{i}}}{M_{V_{i}}^{2}-q^{2}-iM_{V_{i}}\Gamma_{V_{i}}}\right]$$
(3)

where  $q^2$  is the invariant mass squared of the lepton pair. The short-distance contributions are contained in the function  $h(m_c/m_b, q^2/m_b^2)$ , which describes the one-loop matrix element of the four-quark operators  $O_1$  and  $O_2$  (see, e.g., [2] for its explicit expression). The long-distance contribution, related to the formation of intermediate  $c\bar{c}$  bound states, is usually estimated by combining the factorization hypothesis and the Vector Meson Dominance (VMD) assumption [3, 4]; phenomenological analyses [4] suggest that in order to reproduce correctly the branching ratio BR $(B \to J/\psi X \to \ell^+ \ell^- X) = BR(B \to J/\psi X) \cdot BR(J/\psi \to \ell^+ \ell^-)$  the fudge factor  $\kappa$ , which appears in Eq. (3) to correct phenomenologically for inadequacies of the factorization + VMD framework, should satisfy the approximate relation:  $\kappa [3C_1(m_b) + C_2(m_b)] \approx 1$ . To sum up, the effective weak Hamiltonian has the following structure (cf. [2, 3, 5])

$$\mathcal{H}_{eff}(b \to s\ell^{+}\ell^{-}) = \frac{G_{F}}{\sqrt{2}} \frac{\alpha_{em}}{2\pi} V_{ts}^{*} V_{tb} \left[ -2i \frac{m_{b}}{q^{2}} C_{7}(m_{b}) (\bar{s}\sigma_{\mu\nu}q^{\nu}(1+\gamma_{5})b) (\bar{\ell}\gamma^{\mu}\ell) + C_{9}^{eff}(m_{b}) (\bar{s}\gamma_{\mu}(1-\gamma_{5})b) (\bar{\ell}\gamma^{\mu}\ell) + C_{10}(m_{b}) (\bar{s}\gamma_{\mu}(1-\gamma_{5})b) (\bar{\ell}\gamma^{\mu}\gamma_{5}\ell) \right],$$
  
$$\mathcal{H}_{eff}(b \to s\nu\bar{\nu}) = \frac{G_{F}}{\sqrt{2}} \frac{\alpha_{em}}{2\pi\sin^{2}\theta_{W}} V_{tb} V_{ts}^{*} X(x_{t}) (\bar{s}\gamma_{\mu}(1-\gamma_{5})b) (\bar{\nu}\gamma^{\mu}(1-\gamma_{5})\nu)$$
(4)

where  $x_t = (m_t/M_W)^2$  and  $X(x_t)$  is given in [5]. At  $m_t = 176 \ GeV$  one has  $X(x_t) = 2.02$ .

2. Meson form factors. The long-distance contribution to  $B \to (K, K^*)$  decays is contained in the meson matrix elements of the bilinear quark currents appearing in  $\mathcal{H}_{eff}$ , i.e. in the relativistic invariant transition form factors of the vector, axial-vector and tensor currents<sup>b</sup>. The amplitudes of meson decays are induced by the quark transition  $b \to s$  through the vector  $V_{\mu} = \bar{s}\gamma_{\mu}b$ , axial-vector  $A_{\mu} = \bar{s}\gamma_{\mu}\gamma^5 b$  and tensor  $T_{\mu\nu} = \bar{s}\sigma_{\mu\nu}b$  currents, with the following covariant structure [6]

$$\langle P(M_{2}, p_{2})|V_{\mu}(0)|P(M_{1}, p_{1}) \rangle = f_{+}(q^{2}) P_{\mu} + f_{-}(q^{2}) q_{\mu}, \langle V(M_{2}, p_{2}, e)|V_{\mu}(0)|P(M_{1}, p_{1}) \rangle = 2g(q^{2}) \epsilon_{\mu\nu\alpha\beta} e^{*\nu} p_{1}^{\alpha} p_{2}^{\beta}, \langle V(M_{2}, p_{2}, e)|A_{\mu}(0)|P(M_{1}, p_{1}) \rangle = ie^{*\alpha} [f(q^{2}) g_{\mu\alpha} + a_{+}(q^{2}) p_{1\alpha} P_{\mu} + a_{-}(q^{2}) p_{1\alpha} q_{\mu}], \langle P(M_{2}, p_{2})|T_{\mu\nu}(0)|P(M_{1}, p_{1}) \rangle = -2i s(q^{2}) (p_{1\mu}p_{2\nu} - p_{1\nu}p_{2\mu}), \langle V(M_{2}, p_{2}, e)|T_{\mu\nu}(0)|P(M_{1}, p_{1}) \rangle = ie^{*\alpha} [g_{+}(q^{2}) \epsilon_{\mu\nu\alpha\beta} P^{\beta} + g_{-}(q^{2}) \epsilon_{\mu\nu\alpha\beta} q^{\beta} + h(q^{2}) p_{1\alpha} \epsilon_{\mu\nu\beta\gamma} p_{1}^{\beta} p_{2}^{\gamma}],$$

$$(5)$$

<sup>&</sup>lt;sup>b</sup>In rare semileptonic decays there is another long-distance effect, known as the weak annihilation, which is caused by the Cabibbo-suppressed part of the four-fermion operators not included in the operator basis (1). However, the impact of this process in  $B \to (K, K^*)$  transitions is negligible [3].

where  $q = p_1 - p_2$  and  $P = p_1 + p_2$ .

The relativistic invariant form factors, appearing in Eq. (5), contain information on the non-perturbative aspects of the decay processes, so that they should be calculated within a nonperturbative approach for any particular initial and final mesons. To this end various theoretical methods have been adopted, like: the light-cone quark model (LCQM) [7], the constituent quark picture [8], the heavy-quark symmetry (HQS) relations [9], the three-point sum rules (3pSR)[10] and the light-cone sum rules (LCSR) [11]. The main outcome of existing analyses is the remarkable model-dependence of the predicted form factors, illustrated in Table 1 in terms of another set of frequently used form factors, namely:  $F_1(q^2) = f_+(q^2), F_0(q^2) = f_+(q^2) + f_+(q^2)$  $q^{2}f_{-}(q^{2})/(Pq), F_{T}(q^{2}) = -(M_{1}+M_{2})s(q^{2}), V(q^{2}) = (M_{1}+M_{2})g(q^{2}), A_{1}(q^{2}) = f(q^{2})/(M_{1}+M_{2}), A_{2}(q^{2}) = f(q^{2})/(M_{1}+M_{2}$  $A_2(q^2) = -(M_1 + M_2)a_+(q^2), A_0(q^2) = [q^2a_-(q^2) + f(q^2) + (Pq)a_+(q^2)]/2M_2, T_1(q^2) = -g_+(q^2), A_0(q^2) = [q^2a_-(q^2) + f(q^2) + (Pq)a_+(q^2)]/2M_2, T_1(q^2) = -g_+(q^2), A_0(q^2) = [q^2a_-(q^2) + f(q^2) + (Pq)a_+(q^2)]/2M_2, T_1(q^2) = -g_+(q^2), A_0(q^2) = [q^2a_-(q^2) + f(q^2) + (Pq)a_+(q^2)]/2M_2, T_1(q^2) = -g_+(q^2), A_0(q^2) = [q^2a_-(q^2) + f(q^2) + (Pq)a_+(q^2)]/2M_2, T_1(q^2) = -g_+(q^2), A_0(q^2) = [q^2a_-(q^2) + f(q^2) + (Pq)a_+(q^2)]/2M_2, T_1(q^2) = -g_+(q^2), A_0(q^2) = -g_+(q^2), A_0(q^2)$  $T_2(q^2) = -g_+(q^2) - q^2 g_-(q^2)/(Pq)$  and  $T_3(q^2) = (M_1 + M_2)^2 [g_-(q^2)/(Pq) - h(q^2)/2]$ . This fact causes quite uncertain predictions for branching ratios and, in particular, for dilepton spectra and asymmetries within the SM [3, 7, 9, 10, 11, 12]. It is clear that such an uncertainty may become an obstacle for extracting information on the Wilson coefficients (particularly, their signs) and for analysing possible deviations from SM predictions, like those expected in SUSYmodels [3, 13].

We have investigated the relevant meson form factors within a relativistic CQ model adopting a dispersion formulation, which has proved to be successful in describing semileptonic decays of heavy mesons [14]. The dispersion formulation of the quark model has several contact points with the LCQM (see, e.g., [15]). However, the LCQM has the problem of a direct application at time-like values of  $q^2$  because of the contribution arising from the so-called nonpartonic diagram, which cannot be killed at  $q^2 > 0$  by an appropriate choice of the reference frame. The dispersion formulation overcomes this difficulty. Indeed, the LCQM form factors at  $q^2 < 0$  are re-written as double spectral representations in the invariant masses of the initial and final  $q\bar{q}$  pairs, and, then, an analytical continuation is performed to reach the time-like region  $q^2 > 0$ .

Let us consider the transition from the initial meson  $q(m_2)\bar{q}(m_3)$  with mass  $M_1$  to the final meson  $q(m_1)\bar{q}(m_3)$  with mass  $M_2$ , induced by the quark transition  $m_2 \to m_1$  through the current  $\bar{q}(m_1)J_{\mu(\nu)}q(m_2)$ . For the transition  $B_u \to (K, K^*)$  one has  $m_2 = m_b$ ,  $m_1 = m_s$  and  $m_3 = m_u$ . The CQ structure of the initial and final mesons is described by the vertices  $\Gamma_1$ and  $\Gamma_2$ , respectively. The initial *B*-meson vertex has the spinorial structure  $\Gamma_1 = i\gamma_5 G_1/\sqrt{N_c}$ , where  $N_c$  is the number of colours; the final meson vertex has the structure  $\Gamma_2 = i\gamma_5 G_2/\sqrt{N_c}$ for a pseudoscalar state and  $\Gamma_{2\mu} = [A\gamma_{\mu} + B(k_1 - k_3)_{\mu}] G_2/\sqrt{N_c}$ , with A = -1 and  $B = 1/(\sqrt{s_2} + m_1 + m_3)$  for an *S*-wave vector meson. At  $q^2 < 0$  the form factors are given by the following spectral representation [14]

$$f_i(q^2) = \int_{(m_1+m_3)^2}^{\infty} \frac{ds_2 \ G_2(s_2)}{\pi(s_2 - M_2^2)} \int_{s_1^-(s_2, q^2)}^{s_1^+(s_2, q^2)} \frac{ds_1 \ G_1(s_1)}{\pi(s_1 - M_1^2)} \frac{\tilde{f}_i(s_1, s_2, q^2)}{16\lambda^{1/2}(s_1, s_2, q^2)}$$
(6)

where  $s_1^{\pm}(s_2, q^2) \equiv [s_2(m_1^2 + m_2^2 - q^2) + q^2(m_1^2 + m_3^2) - (m_1^2 - m_2^2)(m_1^2 - m_3^2) \pm \lambda^{1/2}(s_2, m_3^2, m_1^2) \\ \lambda^{1/2}(q^2, m_1^2, m_2^2)]/2m_1^2 \text{ and } \lambda(s_1, s_2, s_3) \equiv (s_1 + s_2 - s_3)^2 - 4s_1s_2 \text{ is the triangle function. Equation}$ 

(6) corresponds only to the contribution of the two-particle singularities in the Feynman graphs. For pseudoscalar and vector mesons with mass M, built up of CQ's with masses  $m_q$  and  $m_{\bar{q}}$ , the function G(s) can be written as

$$G(s) = \frac{\pi}{\sqrt{2}} \frac{\sqrt{s^2 - (m_q^2 - m_{\bar{q}}^2)^2}}{\sqrt{s - (m_q - m_{\bar{q}})^2}} \frac{s - M^2}{s^{3/4}} w(k^2)$$
(7)

where  $k = \lambda^{1/2}(s, m_q^2, m_{\bar{q}}^2)/2\sqrt{s}$  and  $w(k^2)$  is the ground-state S-wave radial wave function, normalized as  $\int_0^\infty dk k^2 |w(k^2)|^2 = 1$ . The double spectral densities  $\tilde{f}_i(s_1, s_2, q^2)$ , appearing in Eq. (6), are explicitly given by

$$\begin{split} \tilde{f}_{+} + \tilde{f}_{-} &= 4[m_1m_2\alpha_1 - m_2m_3\alpha_1 + m_1m_3(1 - \alpha_1) - m_3^2(1 - \alpha_1) + \alpha_2s_2], \\ \tilde{f}_{+} - \tilde{f}_{-} &= 4[m_1m_2\alpha_2 - m_1m_3\alpha_2 + m_2m_3(1 - \alpha_2) - m_3^2(1 - \alpha_2) + \alpha_1s_1], \\ \tilde{g} &= -2A \left[m_1\alpha_2 + m_2\alpha_1 + m_3(1 - \alpha_1 - \alpha_2)\right] - 4B\beta, \\ \tilde{a}_{+} + \tilde{a}_{-} &= -4A[2m_2\alpha_{11} + 2m_3(\alpha_1 - \alpha_{11})] + 4B[C_1\alpha_1 + C_3\alpha_{11}], \\ \tilde{a}_{+} - \tilde{a}_{-} &= -4A[-m_1\alpha_2 - m_2(\alpha_1 - 2\alpha_{12}) - m_3(1 - \alpha_1 - \alpha_2 + 2\alpha_{12})] + 4B[C_2\alpha_1 + C_3\alpha_{12}], \\ \tilde{f} &= \tilde{f}_D + (M_1^2 - s_1 + M_2^2 - s_2)\tilde{g}, \\ \tilde{s} &= 2[m_1\alpha_2 + m_2\alpha_1 + m_3(1 - \alpha_1 - \alpha_2)], \\ \tilde{g}_{+} + \tilde{g}_{-} &= 4A[m_3(m_1 - m_3) + \alpha_1(m_1 - m_3)(m_2 - m_3) + \alpha_2s_2 + 2\beta] + 8B(m_1 + m_3)\beta, \\ \tilde{g}_{+} - \tilde{g}_{-} &= 4A[m_3(m_2 - m_3) + \alpha_2(m_1 - m_3)(m_2 - m_3) + \alpha_1s_1] + 8B(m_2 - m_3)\beta, \\ \tilde{h} &= -8A\alpha_{12} - 8B[-m_3\alpha_1 + (m_3 - m_2)\alpha_{11} + (m_3 + m_1)\alpha_{12}], \end{split}$$

where  $\tilde{f}_D = -4A[m_1m_2m_3 + m_2(s_2 - m_1^2 - m_3^2)/2 + m_1(s_1 - m_2^2 - m_3^2)/2 - m_3(s_3 - m_1^2 - m_2^2)/2 + 2\beta(m_2 - m_3)] + 4BC_3\beta$ ,  $\alpha_1 = [(s_1 + s_2 - s_3)(s_2 - m_1^2 + m_3^2) - 2s_2(s_1 - m_2^2 + m_3^2)]/\lambda(s_1, s_2, s_3)$ ,  $\alpha_2 = [(s_1 + s_2 - s_3)(s_1 - m_2^2 + m_3^2) - 2s_1(s_2 - m_1^2 + m_3^2)]/\lambda(s_1, s_2, s_3)$ ,  $\beta = [2m_3^2 - \alpha_1(s_1 - m_2^2 + m_3^2) - \alpha_2(s_2 - m_1^2 + m_3^2)]/4$ ,  $\alpha_{11} = \alpha_1^2 + 4\beta(s_2)/\lambda(s_1, s_2, s_3)$ ,  $\alpha_{12} = \alpha_1\alpha_2 - 2\beta(s_1 + s_2 - s_3)/\lambda(s_1, s_2, s_3)$ ,  $C_1 = s_2 - (m_1 + m_3)^2$ ,  $C_2 = s_1 - (m_2 - m_3)^2$  and  $C_3 = s_3 - (m_1 + m_2)^2 - C_1 - C_2$ .

At  $q^2 < 0$  the representation (6) with the spectral densities (8) for all the form factors but  $f(q^2)$  coincide with the corresponding LCQM expressions (see, e.g., [15]). This is due to the fact that within the dispersion approach all the form factors but  $f(q^2)$  are given by a double dispersion representations without subtractions, whereas in order to construct the form factor  $f(q^2)$  from its double spectral density a subtraction procedure is necessary. We fix this procedure by requiring that, in case of meson transitions induced by a heavy-to-heavy quark transition, all the form factors, including  $f(q^2)$ , satisfy the leading-order Isgur-Wise (*IW*) relations [6] as well as the subleading  $O(1/m_Q)$  relations of the  $1/m_Q$  expansion [16], which are model-independent consequences of QCD. This is fulfilled by the form factors given by Eq. (6) with the spectral densities (8), provided that the functions  $G_i(s_i)$  are localized near the  $q\bar{q}$  threshold with a width of the order  $\Lambda_{QCD}$  [17]. Moreover, for meson decays induced by a heavy-to-light quark transition the dispersion formulation satisfies the leading-order relations between the form factors of the vector and tensor currents given in [6]. The analytical continuation to the time-like region  $q^2 > 0$  generates two contributions: the first one is the normal contribution, which is just the expression (6) taken at  $q^2 > 0$ , and the second one is an additional anomalous contribution, described explicitly in [14]. The normal contribution dominates the form factors at small  $q^2$  and vanishes when  $q^2 = (m_2 - m_1)^2$ , while the anomalous contribution is negligible at small  $q^2$  and steeply rises as  $q^2 \rightarrow (m_2 - m_1)^2$ .

3. Results. We ran calculations adopting two different CQ models, which will be referred to as Set 1 and Set 2. In the former the simple Gaussian ansätz of the ISGW2 model [18] is used for  $w(k^2)$  in Eq. (7), whereas in the latter model the radial function  $w(k^2)$  is the variational solution of the effective  $q\bar{q}$  semi-relativistic Hamiltonian of Godfrey and Isgur (GI) [19]. The values of the CQ masses as well as the average of the internal momentum squared,  $\langle k^2 \rangle$ , are reported in Table 2. The main difference between the radial functions  $w(k^2)$  in the two CQ models relies in their behaviour at high values of the internal momentum k (see Table 2 and cf. [20, 21]): while the Gaussian ansätz yields a soft wave function, which takes into account mainly the effects of the confinement size, the GI wave functions exhibit high-momentum components generated by the effective one-gluon-exchange of the GI potential. The impact of these components on the predicted universal IW function and on the form factor  $f_+(q^2)$  for heavy-to-light transitions has been analyzed in [21] and found not negligible.

The results of our calculations of the form factors have been fitted in terms of a simple  $q^2$  behaviour of the form  $f_i(q^2) = f_i(0)/[1 - \sigma_1 q^2 + \sigma_2 q^4]$ ; the resulting values of the parameters  $f_i(0)$ ,  $\sigma_1$  and  $\sigma_2$  are reported in Table 3. It can clearly be seen that the form factors are very sensitive to the choice of the quark model. In particular, the form factors obtained with the GI wave function (Set 2) are systematically larger than those corresponding to the Gaussian-like ansätz (Set 1). This feature is related both to the larger content of high-momentum components and to the lower values of the CQ masses characterizing the model of Ref. [19] with respect to the ISGW2 model (see Table 2).

The behaviour of the form factors of interest is known in case of heavy parent and daughter quarks inducing the heavy-to-heavy meson transition  $M_1 \rightarrow M_2$ . The leading-order  $1/m_Q$  relations read as [6]

$$f_{+}(\omega) = -g_{+}(\omega) = \frac{M_{1} + M_{2}}{2\sqrt{M_{1}M_{2}}} \xi(\omega), \qquad f_{-}(\omega) = -g_{-}(\omega) = -\frac{M_{1} - M_{2}}{2\sqrt{M_{1}M_{2}}} \xi(\omega),$$

$$s(\omega) = g(\omega) = a_{-}(\omega) = -a_{+}(\omega) = \frac{1}{2\sqrt{M_{1}M_{2}}} \xi(\omega),$$

$$f(\omega) = \sqrt{M_{1}M_{2}} (1 + \omega) \xi(\omega), \qquad h(\omega) = \frac{1}{\sqrt{M_{1}M_{2}}} O(1/m_{Q}), \qquad (9)$$

where  $\omega = (M_1^2 + M_2^2 - q^2)/2M_1M_2$  and  $\xi(\omega)$  is the *IW* function. One should not expect these relations to work with high accuracy for the  $B \to (K, K^*)$  transitions, because the daughter *s*quark cannot be considered heavy enough. As a matter of fact, using the *IW* function calculated in [21], we have checked that large violations of Eq. (9) occur, especially far from the zero-recoil point. However, Eq. (9) implies some (approximate) *HQS* relations among the form factors, namely:  $F_1(q^2) \simeq V(q^2) \simeq A_0(q^2) \simeq A_2(q^2) \simeq -F_T(q^2) \simeq T_1(q^2) \simeq T_3(q^2)$  and  $F_0(q^2) \simeq$  $A_1(q^2) \simeq T_2(q^2)$ , which turn out to be fulfilled within ~ 20% for both *Set* 1 and *Set* 2 in the whole kinematical range.

The averages of our predictions at  $q^2 = 0$  obtained within Set 1 and Set 2 is reported in Table 1. It can be seen that our form factors are consistent with the LCSR results. At the same time, a striking disagreement with some of the 3pSR results has been found, especially in case of the form factors  $F_T$  and  $T_3$ . Indeed, the form factor  $T_3$  of Ref. [10] has an opposite sign and is large in absolute value compared with  $T_1$ , as well as the form factor  $|F_T|$  is small compared with  $F_1$ , at variance with the approximate HQS relations. We want to mention that in Ref. [10] both  $F_T$  and  $T_3$  are claimed to be constructed in terms of some of the other form factors applying equations of motion. However, it can be checked that the expression given in [10] for  $T_3$  does not satisfy the HQS requirements in the heavy-quark limit.

The differential decay rates and asymmetries have been calculated using the expressions given in Refs. [10, 12, 13]. The predictions for the dilepton distribution in  $B \to (K, K^*) \mu^+ \mu^$ decays are reported in Fig. 1, where the non-resonant contributions are shown separately. The total decay rates turn out to be at least one order of magnitude larger than the non-resonant decay rates. However, the resonant contributions are strongly peaked in narrow regions around their masses, so that outside these regions the resonance influence is almost negligible. This fact allows to reliably separate the resonant contributions from the non-resonant one, which contains the relevant information on the Wilson coefficients. In Table 4 our predictions for the non-resonant decay rates and branching ratios are listed. The dependence on the chosen quark model is strong, yielding a large uncertainty in the predictions; such a drawback may be limited by testing the same CQ model in several semileptonic decays, including in particular the shape of differential decay rates and lepton spectra, which are expected to be sensitive to the form of the CQ model wave functions. Finally, note that the transitions  $B \to K^* \mu^+ \mu^-$  and  $B \to K^* e^+ e^-$  have different rates, because the amplitude  $B \to K^* \ell^+ \ell^-$  has a kinematical pole at  $q^2 = 0$ , which makes the corresponding decay rate very sensitive to the lower boundary of the phase space volume  $(q^2 = 4m_\ell^2)$ , while the amplitude  $B \to K\ell^+\ell^-$  is regular at  $q^2 = 0$  and, therefore, insensitive to the mass of the light lepton.

Our results for total decay rates and branching ratios are summarized in Table 5 and compared with the predictions of other approaches. It can be seen that our results are consistent with those of Ref. [3], which are based on the application of the HQS relations for the form factors, whereas they differ from the predictions of Ref. [10]. On one hand, the present level of model dependence does not provide the opportunity to extract precise values of  $V_{ts}$  and to study possible effects beyond the SM. On the other hand, the situation with lepton asymmetries looks much more optimistic. Our results for the dilepton forward-backward and lepton polarization asymmetries are presented in Figs. 2 and 3. The asymmetries calculated within different quark models turn out to differ only by a very small amount (~ 10%) in the whole kinematical range. This is due to the following facts: i) the asymmetries depend on ratios among form factors; ii) our form factors obey approximate HQS relations, which means that their ratios are only slightly model-dependent. We point out that our dilepton asymmetry is approximately a factor of 2 lower than the result of Ref. [10]<sup>c</sup> and this disagreement can be traced back again to the large difference in the form factor  $T_3$  calculated in [10] and in the present work (see Table 1).

<sup>&</sup>lt;sup>c</sup>We have taken the resonance phase in accordance with Ref. [4], whereas an opposite sign is used in [10].

In conclusion, we have analysed rare semileptonic transitions  $B \to (K, K^*)$  within a relativistic constituent quark model, formulated in a dispersion form. The differential decay rates are found to be strongly sensitive to the choice of the particular quark model, obtaining an overall ~ 50% uncertainty in the predictions. However, at the same time, the asymmetries of lepton distributions result to be almost insensitive to quark model parameters, so that they can be predicted within the framework of the Standard Model with little uncertainty (~ 10%). Our predictions for lepton asymmetries in rare semileptonic decays may provide a reliable starting point for investigating possible violations of Standard Model predictions, like those expected in SUSY models.

Acknowldgments. We are grateful to B. Stech and K.A. Ter-Martirosyan for useful discussions and to L. Smirnova for her interest in this work, which was supported by the *RFBR* under grant 96-02-18121a.

## References

- [1] B. Grinstein, M.B. Wise and M.J. Savage: Nucl. Phys. **B319** (1989) 271.
- [2] A. Buras and M. Münz: Phys. Rev. **D52** (1995) 186.
- [3] A. Ali, T. Mannel and T. Morozumi: Phys. Lett. B273 (1991) 505. A. Ali: Acta Phys. Pol. B27 (1996) 3529; Nucl. Instrum. Meth. A384 (1996) 8.
- [4] C.S. Lim, T. Morozumi and A.T. Sanda: Phys. Lett. B218 (1989) 343. P.J. O'Donnell and H.K.K. Tung: Phys. Rev. D43 (1991) R2067.
- [5] T. Inami and C. S. Lim: Prog. Theor. Phys. 65 (1981) 287. G. Buchalla and A.J. Buras: Nucl. Phys. B400 (1993) 225.
- [6] N. Isgur and M.B. Wise: Phys. Lett. B232 (1989) 113; Phys. Lett. B237 (1990) 527; Phys. Rev. D42 (1990) 2388.
- [7] W. Jaus and D. Wyler: Phys. Rev. **D41** (1990) 3405.
- [8] B. Stech: Phys. Lett. **B354** (1995) 447; Z. Phys. **C75** (1997) 245.
- [9] G. Burdman: Phys. Rev. **D52** (1995) 6400. W. Roberts: Phys. Rev. **D54** (1996) 863.
- [10] P. Colangelo *et al.*: Phys. Rev. **D53** (1996) 3672; Phys. Lett. **B395** (1997) 339.
- [11] T.M. Aliev *et al.*: Phys. Lett. **B400** (1997) 194 and hep-ph 9612480.
- [12] C.Q. Geng and C.P. Kao: Phys. Rev. **D54** (1996) 5636.
- [13] C. Greub, A. Ioannissian and D. Wyler: Phys. Lett. **B346** (1995) 349.

- [14] D. Melikhov: Phys. Rev. D53 (1996) 2460; Phys. Lett. B380 (1996) 363; Phys. Lett. B394 (1997) 385.
- [15] W. Jaus: Phys. Rev. **D41** (1990) 3394; Phys. Rev. **D53** (1996) 1349.
- [16] M. Luke: Phys. Lett. B252 (1990) 447. M. Neubert and V. Rieckert: Nucl. Phys. B382 (1992) 97.
- [17] D. Melikhov and N. Nikitin: hep-ph/9609503 (unpublished).
- [18] D. Scora and N. Isgur: Phys. Rev. **D52** (1995) 2783.
- [19] S. Godfrey and N. Isgur: Phys. Rev. **D32** (1985) 189.
- [20] F. Cardarelli *et al.*: Phys. Lett. B332 (1994) 1; Phys. Lett. B349 (1995) 393; Phys. Lett. B359 (1995) 1; Few-Body Syst. Suppl. 9 (1995) 267; Phys. Rev. D53 (1996) 6682.
- [21] S. Simula: Phys. Lett. **B373** (1996) 193. I. Grach *et al.*: Phys. Lett. **B385** (1996) 317.
- [22] R. Ammar *et al.*: Phys. Rev. Lett. **71** (1993) 674 and CLEO CONF 96-05 (1996).
- [23] T. Skwarnicki: Proc. of the 17<sup>th</sup> Int. Symp. on Lepton-Photon Interactions, Beijing (China), August 1995, eds. Z. Zhi-Peng and C. He-Sheng (World Scientific), p. 238.
- [24] C. Anway-Wiese et al.: FERMILAB CONF-95-201-E (1995). T. Speer et al.: FERMILAB CONF-96-320-E (1996).

**Table 1.** Transition form factors for the semileptonic decays  $B \to (K, K^*)$  evaluated at  $q^2 = 0$  within various theoretical methods. The last row represents the average of the results obtained in the present work adopting the two CQ models, Set 1 (ISGW2 wave function [18]) and Set 2 (GI wave function [19]), described in the text.

Ref.	$F_1(0)$	$F_T(0)$	V(0)	$A_1(0)$	$A_2(0)$	$T_1(0)$	$T_3(0)$
LCQM [7]	0.30	-0.30	0.35	0.26	0.24	0.31	0.32
3pSR [10]	$0.25\pm0.03$	$-0.14\pm0.03$	$0.47\pm0.03$	$0.37\pm0.03$	$0.40\pm0.03$	$0.38\pm0.06$	-1.96
LCSR [11]	0.29	-0.31	0.45	0.36	0.40	0.36	0.39
This work	$0.43\pm0.07$	$-0.40\pm0.06$	$0.40\pm0.10$	$0.34\pm0.08$	$0.27\pm0.05$	$0.37\pm0.09$	$0.35\pm0.09$

**Table 2.** Values of the CQ masses and of the average internal momentum squared  $\langle k^2 \rangle$ , in  $(GeV/c)^2$ , for the two CQ models adopted in this work.

Ref.	$m_u$	$m_s$	$m_c$	$m_b$	$< k^2 >_K$	$< k^2 >_{K^*}$	$< k^2 >_{B_u}$
$Set \ 1 \ [18]$	0.33	0.55	1.82	5.2	0.29	0.16	0.28
$Set \ 2 \ [20]$	0.22	0.42	1.65	5.0	0.77	0.34	0.60

**Table 3.** Parameters of the fit  $f_i(q^2) = f_i(0)/[1 - \sigma_1 q^2 + \sigma_2 q^4]$  to the  $B \to (K, K^*)$  transition form factors calculated within the two CQ models adopted.

Decays	$B \to K$			$B \to K^*$						
	$f_{+}(0)$	$f_{-}(0)$	s(0)	g(0)	f(0)	$a_{+}(0)$	$a_{-}(0)$	h(0)	$g_{+}(0)$	$g_{-}(0)$
Ref.	$\sigma_1$	$\sigma_1$	$\sigma_1$	$\sigma_1$	$\sigma_1$	$\sigma_1$	$\sigma_1$	$\sigma_1$	$\sigma_1$	$\sigma_1$
	$\sigma_2$	$\sigma_2$	$\sigma_2$	$\sigma_2$	$\sigma_2$	$\sigma_2$	$\sigma_2$	$\sigma_2$	$\sigma_2$	$\sigma_2$
Set 1	0.36	-0.30	0.06	0.048	1.60	-0.036	0.041	0.0037	-0.28	0.24
	0.048	0.050	0.049	0.057	0.0288	0.053	0.055	0.075	0.058	0.059
	0.00063	0.00061	0.00064	0.00085	0.00028	0.00082	0.00088	0.0016	0.0009	0.00096
Set 2	0.50	-0.42	0.080	0.083	2.62	-0.052	0.067	0.0085	-0.47	0.43
	0.035	0.036	0.035	0.042	0.011	0.030	0.040	0.044	0.042	0.043
	0.00017	0.00018	0.00017	0.00036	0.00015	-0.00025	0.00016	0.00001	0.00036	0.00037

**Table 4.** Non-resonant decay rates and branching ratios of the decays  $B \to (K, K^*)$   $(\ell^+ \ell^-, \nu \bar{\nu})$ . Decay rates are given in units  $|V_{ts}^* V_{tb}|^2 \times 10^8 \ s^{-1}$ , while branching ratios are in units  $|V_{ts}^* V_{tb}|^2 \times 10^{-4}$ .

Ref.		$B \to K \ell^+ \ell^-$	$B \to K \sum \nu_i \bar{\nu}_i$	$B \to K^* e^+ e^-$	$B \to K^* \mu^+ \mu^-$	$B \to K^* \sum \nu_i \bar{\nu}_i$
$Set \ 1$	Γ	2.0	25	5.2	4.1	51
	$\mathcal{BR}$	2.9	37	7.6	6.0	75
$Set \ 2$	Γ	3.3	39	11.5	8.5	108
	$\mathcal{BR}$	4.8	57	17.0	12.5	160

**Table 5.** Non-resonant decay rates and branching ratios of rare radiative and semileptonic decays of *B*-meson. Theoretical predictions are given in units  $|V_{ts}/0.033|^2$ . The results of Ref. [10] have been recalculated replacing the value  $|V_{ts}/0.04|^2$  with  $|V_{ts}/0.033|^2$ . The uncertainties in  $V_{ts}$  are not included in the error bars.

Ref.	This work	[3]	[10]	Exp.	
Decay	$\Gamma(s^{-1})$	$\Gamma(s^{-1})$	$\Gamma(s^{-1})$	$\Gamma(s^{-1})$	
mode	$\mathcal{BR}$	$\mathcal{BR}$	$\mathcal{BR}$	$\mathcal{BR}$	
$B \to K^* \gamma$	$(2.9 \pm 1.3) \times 10^7$				
	$(3.9 \pm 1.7) \times 10^{-5}$	$(4.9 \pm 2.0) \times 10^{-5}$		$(4.2 \pm 1.0) \times 10^{-5} [22]$	
$B \to K \ell^+ \ell^-$	$(2.9 \pm 0.7) \times 10^5$				
	$(4.2 \pm 0.9) \times 10^{-7}$	$(4.0 \pm 1.5) \times 10^{-7}$	$2 \times 10^{-7}$	$< 0.9 \times 10^{-5} [23]$	
$B \to K \sum \nu_i \bar{\nu}_i$	$(3.5 \pm 0.8) \times 10^6$				
	$(5.2 \pm 1.1) \times 10^{-6}$	$(3.2 \pm 1.6) \times 10^{-6}$	$(1.6 \pm 0.4) \times 10^{-6}$		
$B \to K^* e^+ e^-$	$(9.2 \pm 3.5) \times 10^5$				
	$(1.4 \pm 0.5) \times 10^{-6}$	$(2.3 \pm 0.9) \times 10^{-6}$	$0.7 \times 10^{-6}$	$< 1.6 \times 10^{-5} [23]$	
$B \to K^* \mu^+ \mu^-$	$(6.9 \pm 2.4) \times 10^5$				
	$(1.0 \pm 0.4) \times 10^{-6}$	$(1.5 \pm 0.6) \times 10^{-6}$	$0.7 \times 10^{-6}$	$< 2.5 \times 10^{-5} [24]$	
$B \to \overline{K^*} \sum \nu_i \bar{\nu}_i$	$(8.7 \pm 3.1) \times 10^6$				
	$(1.3 \pm 0.5) \times 10^{-5}$	$(1.1 \pm 0.55) \times 10^{-5}$	$(3.5 \pm 0.5) \times 10^{-6}$		



Figure 1. Differential decay rate for the transitions  $B \to K\ell^+\ell^-$  (a) and  $B \to K^*\mu^+\mu^-$  (b) versus the invariant mass squared of the lepton pair,  $q^2$ , divided by the *B*-meson mass squared,  $M_B^2$ . Dashed and solid lines correspond to our calculations obtained using the two CQ models Set 1 (ISGW2 wave function [18]) and Set 2 (GI wave function [19]), respectively. The thin lines represent the non-resonant contribution only, whereas the thick lines are the sum of non-resonant and resonant contributions.



Figure 2. The dilepton forward-backward asymmetry  $(A_{FB})$  for the decay  $B \to K^* \mu^+ \mu^-$  versus  $q^2/M_B^2$ . Dashed and solid lines are as in Fig. 1. The non-resonant contribution (a) and the sum of non-resonant and resonant contributions (b) are separately shown.



Figure 3. The same as in Fig. 2, but for the lepton longitudinal polarization asymmetry  $(P_L)$  for the decays  $B \to K \ell^+ \ell^-$  (a, b) and  $B \to K^* \mu^+ \mu^-$  (c, d), respectively.