Derivative expansions of Euclidean thermal effective actions

T.S.Evans*

Theoretical Physics, Blackett Laboratory, Imperial College, Prince Consort Road, London SW7 2BZ, U.K. (22nd February 1998, revised 16th July 1998)

I compute the derivative expansion of an effective action at finite temperature using the imaginary time approach. I show that it is a well behaved expansion giving a unique seriers contrary to previous results. This disparity is shown to originate in the choice of thermal Green functions used in the calculations.

PACS: 11.10.Wx, 11.10.Lm. Preprint Imperial/TP/97-98/21, hep-ph/9808382

Effective actions provide a powerful tool for analysing complex systems. They enable the complicated effects of different sectors of the theory to be incorporated in a compact way into a description of the one sector of interest. For example, in thermal field theory the effective action encodes the effects that the quantum and thermal fluctuations of several fields have on the specific field under consideration. However, as effective actions of quantum fields are often very complicated, it is common to consider the low energy/momentum behaviour of one field by expanding the effective action in terms of the derivatives of that field. This is a well trodden path in zero temperature quantum field theory [1–3].

It is therefore surprising that similar calculations of derivative expansions of thermal effective actions give strange results [4–11]. It has been shown in the literature that series is not unique as it depends on the order in which one performs the time and space derivative expansions. In particular the lowest order term, which is usually identified as the effective potential or free energy [1–3], varies significantly. The importance of this fact has been stressed in the condensed matter literature, for instance [4,11]. In relativistic particle physics, such derivative expansions are also frequently used but at finite temperature it is common to address only the spatial derivatives [12]. However, many practical problems are dynamical ones, such as tunnelling rates, or the slow roll of fields in inflation, so temporal derivatives are required.

In this letter I show how to calculate a well behaved derivative expansion of an effective action using the Euclidean approach to thermal field theory. The key idea is that such expansions describe field configurations which vary *slowly* in time and space and hence are not thermal equilibrium configurations. By explicit calculation, I will show that the retarded thermal Green functions used in previous analyses are *not* relevant to this problem. These Green functions describe how the system responds to a *sudden* impulse as linear response theory shows. Important physical processes, especially Landau damping, do contribute to my unique derivative expansion whereas they caused the problems in the literature where retarded Green functions were incorrectly used.

Recently, the problem was clearly demonstrated by Das and Hott [10] in a simple relativistic model, putting earlier comments [4–9] into context. Following [10] I use a model with two scalar fields

$$\mathcal{L}[\phi,\eta] = \frac{1}{2}\eta\Delta^{-1}\eta - \frac{1}{2}g\phi\eta^2 + \mathcal{L}_0[\phi]$$

where Δ is the η field propagator and $\mathcal{L}_0[\phi]$ is the free Lagrangian for ϕ with arbitrary dispersion relation and source. Treating ϕ as the background field, whose effective action I wish to calculate, and integrating out the η field fluctuations gives a generating functional of the form

$$Z = \int D\phi \, \exp\{i \int d^4x \mathcal{L}_0[\phi] + iS'_{\text{eff}}[\phi]\}$$
$$S'_{\text{eff}}[\phi] = \frac{i}{2} \operatorname{Tr} \{\ln\left[1 - g\Delta(x, x')\phi(x')\right]\}$$

Four-vectors are always Minkowskii, e.g. $k^{\mu} = (E, \vec{k}), k^2 = E^2 - \vec{k}^2$, with $E \in \mathcal{R}e$ for physical values.

The only non-trivial contribution to the effective action comes from the non-local logarithmic term S'_{eff} which I expand as a series in $g\phi$, $S'_{\text{eff}} = \sum_{p=1}^{\infty} S^{(p)}_{\text{eff}}$. The non-locality is first clearly visible in the $O(g^2\phi^2)$ term, $S^{(2)}_{\text{eff}}$,

$$S_{\text{eff}}^{(2)} = \frac{-ig^2}{4} \int d^4x \, \int d^4x' \, \phi(x)\Delta(x,x')\phi(x')\Delta(x',x) \tag{1}$$

so we will focus on this term c.f. [10].

^{*}email: T.Evans@ic.ac.uk, WWW: http://euclid.tp.ph.ic.ac.uk/~time

The derivative expansion is an attempt to represent such non-local terms as a series of local terms. Thus $\phi(x')$ of (1) is expressed in terms of $\phi(x)$ through a Taylor series, which we may write as

$$\phi(x') = \exp\{\lambda[(x'-x).\frac{\partial}{\partial x''}]\}\phi(x'')\Big|_{x''=x,\lambda=1}$$
(2)

In (2) the formal counting parameter λ allows us to generate the terms of the derivative expansion as a λ power series. This gives

$$S_{\text{eff}}^{(2)} = -\frac{1}{2} \int d^4 x \, \phi(x) B(-i\lambda \partial_\mu) \phi(x) \Big|_{\lambda=1}$$

$$-iB(p) = \frac{(-ig)^2}{2} \int \frac{d^4 k}{(2\pi)^4} \, i\Delta(k) i\Delta(k+p)$$
(3)

Thus, all the information needed about the contribution of the η field fluctuations to the derivative expansion of $S_{\text{eff}}^{(2)}$ comes from a small four-momentum expansion of the bubble diagram B(p). The formal parameter $p^{\mu} = -i\lambda\partial^{\mu}$ is not in general small so such a derivative expansion of the bubble diagram B can only have any meaning if $\phi(p)$ dies off sufficiently fast as p^{μ} moves away from zero, i.e. the expansion is valid only for *small* deviations from space-time independent ϕ field configurations. At zero temperature with real Minkowskii or Euclidean times and energies, this works perfectly [1–3].

Now let us turn to finite temperature and use the Euclidean time approach [13]. At first sight the arguments which led from (1) to (3) work equally well in thermal field theory provided one changes the time integrations to run from τ_i to $\tau_i - i\beta$, uses scalar fields which are periodic over this range, and alters the propagators appropriately.

However, the precise meaning of B is not clear in the thermal context because the same diagram represents many different Green functions with distinct values [14,15]. Existing analyses of the effective action in this language [4–7,10] (but not [9]) assume without proof that retarded functions should be used. This is the most natural for Euclidean thermal field theory [14,15] and the effective potential at finite temperature is most easily linked to an expansion in terms of retarded diagrams [7]. For later comparison, let us see what this gives for $S_{\text{eff}}^{(2)}$. In (1) we set $B = B_{\text{ret}}$, the retarded bubble diagram, which can be calculated with any thermal field formalism to give

$$B_{\rm ret}(p_{\mu} = (E, \vec{p}); \beta) = -\frac{g^2}{2} \int \frac{d^3 \vec{k}}{(2\pi)^3} \sum_{s_0, s_1 = \pm 1} \frac{s_0 s_1}{4\omega\Omega} \frac{1}{E + s_0 \omega + s_1\Omega} \frac{(e^{\beta(s_0 \omega + s_1\Omega)} - 1)}{(e^{\beta s_0 \omega} - 1)(e^{\beta s_1\Omega} - 1)}$$
(4)

where $\omega = \omega(\vec{k})$ and $\Omega = \omega(\vec{k} + \vec{p})$ and we have assumed $\Delta(E, \vec{k}) = (E^2 - \omega^2)^{-1}$.

The problem with this thermal retarded diagram is that it does not have a unique value in the $p^{\mu} \to 0$ limit [4–11,16]. In (4), it comes from a divergence in the denominator when $E + s_0\omega + s_1\Omega \to 0$ as $p^{\mu} \to 0$ for the terms $s_0 = -s_1$ when $\beta < \infty$. A more familiar form for the retarded bubble is obtained if we now assume a relativistic dispersion relation and then perform the angular integration, but this still has the same problems in the limit of $p^{\mu} \to 0$. This strange behaviour of the retarded bubble diagram at zero momentum is *not* due to any technical calculational problems [7,16]. From a physical point of view the origin of this branch point is well known and arises from the physical processes known as Landau damping [13]. These processes have particles from the heat bath involved in the *in* or *out* states and thus are only present at non-zero temperature.

Physically, using B_{ret} implies that the behaviour of fields described by an effective action is *very* sensitive to the way I approach the zero momentum point. It implies that *almost constant* fields, which vary only on scales much, much bigger than any physical scale in the problem, have a significantly different behaviour when they vary faster in space than time as compared to the opposite case.

It is therefore worth performing the calculation more carefully, focusing on the temporal aspect as this changes most radically at finite temperature. Starting from (1) I find

$$S_{\text{eff}}^{(2)} = \frac{-ig^2}{4} \int_{\tau_i}^{\tau_i - i\beta} d\tau \int \frac{d^3 \vec{p}}{(2\pi)^3} \phi(\tau, \vec{p}).$$
$$\int_{\tau_i}^{\tau_i - i\beta} d\tau' \int \frac{d^3 \vec{k}}{(2\pi)^3} \Delta(\tau, \tau'; \vec{k}) \phi(\tau', -\vec{p}) \Delta(\tau', \tau; \vec{k} + \lambda \vec{p}) \bigg|_{\lambda = 1}$$
(5)

The propagators are for the η field in thermal equilibrium. For illustrative purposes we again work with a relativistic field for which the most convenient representation here is

$$\Delta(\tau,\tau';\vec{k}) = \frac{i}{\beta} \sum_{n \in \mathcal{Z}} \sum_{s=\pm 1} e^{-iE(\tau-\tau')} \frac{s}{2\omega} \frac{1}{E - s\omega(\vec{k})} \bigg|_{E=2\pi i n/\beta}$$

The key point comes in the need to expand the second field in (5) as in (2)

$$\phi(\tau', -\vec{p}) = \exp\{\lambda(\tau' - \tau)\frac{\partial}{\partial\tau''}\}\phi(\tau'', -\vec{p})\Big|_{\tau''=\tau, \lambda=1}$$
(6)

Performing the τ' integration in $S_{\text{eff}}^{(2)}$ (5) gives

$$I = \int_{\tau_i}^{\tau_i - i\beta} d\tau' \ e^{i(E'' + E - E')\tau'} e^{-i(E'' + E - E')\tau} = \frac{(e^{(E'' + E - E')\beta} - 1)}{i(E'' + E - E')} e^{i(E'' + E - E')(\tau_i - \tau)}$$
(7)

where $E'' \equiv -i\lambda(\partial/\partial \tau'')$ coming from (6). When we use the result I in (5), E and E' come from the two propagators and so are both identical to Matsubara frequencies $2\pi i n/\beta$, $n \in \mathbb{Z}$. If and only if E'' is also limited to Matsubara frequencies does I then reduce to the usual energy conserving delta function $I = -i\beta\delta_{n'',n'-n}$. From (6) it is clear that taking $E'' \equiv -i\lambda(\partial/\partial \tau'')$ identically equal to one of the Matsubara frequencies is equivalent to saying that the background field ϕ must be periodic in imaginary time.

However, I intend to truncate the derivative expansion of the ϕ field. This is a disaster in the compact time coordinate because the truncated Taylor series of a periodic function is not necessarily periodic! A perturbation of interest to a periodic background field $\phi(\tau'')$ may be small and periodic itself, but keeping only a finite numbers of terms in the Taylor series (6) leaves us with a non periodic perturbation. Thus I am forced to consider energies of the background ϕ field, E'' in (7), which are merely in the neighbourhood of zero energy and are not a pure Matsubara frequency, so that $\exp\{\beta E''\} \neq 1$. The derivative expansion is good for studying *small, but arbitrary* deviations from an constant background. This includes non-periodic, i.e. non-equilibrium, perturbations where for a short time the heat bath will remain in equilibrium.

To put the point more forcefully, existing analyses assume $\exp\{\beta E''\} = 1$ and are therefore only correct if they impose equilibrium rigorously. Consequently they are correct only for static configurations and expanding such results about zero energy is invalid. One exception is [9] where it is also argued that the limitation of E''to Matsubara frequencies makes no sense. However, in [9] no rigorous algorithm for the choice of the analytic continuation in energy away from discrete E'' is provided c.f. [14,15], and Weldon [16] has shown the use of "Feynman parameterisation" in [9] is incorrect.

With non-periodic perturbations allowed, one can show that the calculation is no longer expected to be independent of the initial time τ_i , which is the time at which the temperature and density matrix were defined [17,19]. Taking great care over factors such as $\exp{\{\beta E\}}$ and over the analytic properties of integrands at large energies when energy sums over E, E' are performed, contour integration techniques [13] give

$$S_{\text{eff}}^{(2)} = -\frac{1}{2} \int_{\tau_{i}}^{\tau_{i} - i\beta} d\tau \int \frac{d^{3}\vec{p}}{(2\pi)^{3}} \phi(\tau, \vec{p}) \mathcal{B}(-i\lambda \frac{\partial}{\partial \tau''}, \lambda \vec{p}; m; \tau - \tau_{i}) \phi(\tau'', -\vec{p}) \Big|_{\tau'' = \tau, \lambda = 1}$$
(8)
$$\mathcal{B}(E'', \vec{p}; m; \tau - \tau_{i}) = \frac{g^{2}}{2} \int \frac{d^{3}\vec{k}}{(2\pi)^{3}} \sum_{s_{0}, s_{1} = \pm 1} \frac{s_{0}s_{1}}{4\omega\Omega} \frac{1}{(E'' + s_{0}\omega + s_{1}\Omega)} \frac{1}{(e^{\beta s_{0}\omega} - 1)(e^{\beta s_{1}\Omega} - 1)} \cdot \left\{ (e^{\beta(E'' + s_{0}\omega + s_{1}\Omega)} - 1) + (e^{-i(\tau - \tau_{i})(E'' + s_{0}\omega + s_{1}\Omega)} - 1) \cdot (e^{\beta E''} - 1)e^{\beta(s_{0}\omega_{0} + s_{1}\Omega)} \right\}$$

Note that the earlier problems with the derivative expansion in (4) stemmed from the $E'' + s_0 \omega + s_1 \Omega$ denominator, which in the limit of static and homogeneous fields $(E'' \equiv -i\lambda(\partial/\partial\tau'') = 0, \vec{p} = 0)$ is divergent. The same denominator is present in (8) but now the numerators are zero in this limit. Thus the space and time derivative expansion of this expression is well defined. It also means that there are no branch points at $E''^2 = \vec{p}^2$ or $E''^2 = 4m^2 + \vec{p}^2$ though the latter can never be seen in a small E'', \vec{p} expansion.

Equation (8) shows that $\mathcal{B} = B_{\text{ret}}$ of (4) only if I set $\exp\{\beta E''\} = 1$, i.e. limit myself to exactly periodic ϕ field configurations. This again emphasises the fact that the use of the retarded bubble diagram in the derivative expansion for effective actions is valid only for static configurations.

In order to make the spatial derivative expansion, I need the form of the dispersion relation of the η field. Taking this to be the usual relativistic form, $\omega(\vec{k})^2 = |\vec{k}|^2 + m^2$, it is straight forward to generate the expansion. Up to first order I have

$$S_{\text{eff}}^{(2)'} = \frac{g^2}{32\pi^2} \int_{\tau_i}^{\tau_i - i\beta} d\tau \int d^3 \vec{x} \ \phi(\tau, \vec{x}) \left[\gamma_{0,0} - i\beta\gamma_{1,0}(\tau - \tau_i)\frac{\partial}{\partial\tau} + O(\frac{\partial^2}{\partial\tau^2}, \vec{\nabla}^2) \right] \phi(\tau, \vec{x}) \tag{9}$$

$$\gamma_{0,0} = \int_0^\infty dk \ \frac{-k^2}{\omega^3} \left[\frac{e^{\beta\omega} + 1}{e^{\beta\omega} - 1} + 2\beta\omega \frac{e^{\beta\omega}}{(e^{\beta\omega} - 1)^2} \right] = \frac{k^2}{\omega} \frac{d}{d\omega} \left(\frac{1}{\omega} \coth(\frac{1}{2}\beta\omega) \right)$$

$$\gamma_{1,0}(\tau) = \frac{1}{2}\gamma_{0,0} + \int_0^\infty dk \ \left[\frac{2k^2}{\omega^2} \frac{e^{\beta\omega}}{(e^{\beta\omega} - 1)^2} (-i\tau) + \frac{k^2}{2\omega^3} \frac{(e^{-2i\tau\omega} - 1)(1 + e^{2i(\tau - i\beta)\omega})}{(e^{\beta\omega} - 1)^2} \right]$$

This first order expansion illustrates several features of the general result (8). Galliean invariance tells us that there can be no terms odd in $\vec{\nabla}$, and no such terms are present in (8) or (9). Terms with no time derivatives correspond to pure static perturbations in the background fields. These are trivially periodic and hence [17,19] must be independent of $\tau - \tau_i$. $\gamma_{0,0}$ and the explicit form (8) confirm this. Interestingly, there is a term linear in the time derivative. This unfamiliar term is a Lorentz scalar at finite temperature because of the existence of an extra four-vector, u^{μ} , the four velocity of the observer with respect to the heat bath. I have implicitly chosen to measure quantities with respect to the heat bath rest frame so that $u^{\mu} = (1, \vec{0})^{\mu}$ and the relevant Lorentz scalars are $\partial_{\mu}\partial^{\mu}$ and $\partial/\partial\tau \equiv \partial_{\mu}.u^{\mu}$.

Most importantly, the lowest order term is the contribution to the effective potential and $\gamma_{0,0}$ is indeed found to be precisely that obtained in standard calculations of the effective potential at finite temperature, be they real- or imaginary-time calculations. Only now with this analysis and its well behaved derivative expansion can we justify the use such a free energy term as a potential in an evolution equation for quantum fields in a heat bath, something which has often been assumed at in the past.

Contrary to previous suggestions, this result shows that Landau damping does not destroy the derivative expansion. Indeed, Landau damping terms are present in the above calculation as they are the $s_1 = -s_2$ terms with $\omega - \Omega$ factors in (8).

Further confirmation that this approach to derivative expansions of thermal effective actions is correct comes from a real-time calculation [17] and from one using a general path-ordered formalism [18,19] calculation, both of which give the same result as this Euclidean calculation. Likewise, the generalization to other types of field is straight forward [19].

Finally, this analysis of thermal effective actions is not in conflict with linear response [13], where retarded Green functions are relevant. In linear response one considers perturbations which are space-time delta functions. This is the exact opposite of the nearly constant field configurations described by the derivative expansion of approximation to thermal effective actions. Thus we should not be surprised when we find Green functions like \mathcal{B} of (8) and not retarded Green functions control the dynamics of slowly varying fields.

I would like to thank I.Aitchison, M.Asprouli, V.Galan Gonzalez, R.Rivers and D.Steer for useful discussions. This work was supported by the Royal Society, and by the European Commission under their Human Capital and Mobility programme, contract number CHRX-CT94-0423.

- [1] S.Coleman, Aspects of Symmetry (Cambridge University Press, Cambridge, 1985, ISBN 0 521 31827 0).
- [2] C.M.Fraser, Z.Phys.C 28 101 (1985).
- [3] A.Dobado, A.Gómez-Nicola, A.L.Maroto, J.R.Peláez, Effective Lagrangians for the Standard Model (Springer-Verlag, Berlin, 1997, ISBN 3-540-62570-4).
- [4] E.Abrahams and T.Tsuneto, Phys.Rev. 152 416 (1966).
- [5] Y.Fujimoto, in Quantum Field Theory, edited by F.Mancini (Elsevier Science Publishers, Amsterdam, 1986), p.435.
- [6] T.S. Evans, Z.Phys.C **41** (1988) 333.
- [7] T.S.Evans, Can.J.Phys. **71** 241 (1993).
- [8] P.Elmfors, K.Enqvist, and I.Vilja, Phys.Rev.Lett. 71 480 (1993) (hep-ph/9303269).
- [9] P.S.Gribosky and B.R.Holstein, Z.Phys.C 47 214 (1990).
- [10] A.Das and M.Hott, Phys.Rev.D 50 6655 (1994).
- [11] I.J.R.Aitchison and D.J.Lee, Phys.Rev.B 56 8303 (1997).
- [12] I.Moss, D.Toms and A.Wight, Phys.Rev.D 46 1671 (1992).
- [13] M.LeBellac, Thermal Field Theory (Cambridge University Press, Cambridge, 1996, ISBN 0 521 46040 9).
- [14] G.Baym and N.D.Mermin, J.Math.Phys. 2 232 (1961).
- [15] T.S.Evans, Nucl. Phys. **B374** 340 (1992).
- [16] H.A.Weldon, Phys.Rev.D 47 (1993) 594.
- [17] M.Asprouli and V. Galan Gonzalez, Derivative expansions of real-time thermal effective actions, Imperial College, Imperial/TP/97-98/026, Report No. hep-ph/9802319.
- [18] T.S.Evans, The Unique Derivative Expansion for Thermal Effective Actions, talk given at the 5th International Workshop on Thermal Field Theory, Regensburg, Germany, 10-14 August, 1998. Original overheads available at http://euclid.tp.ph.ic.ac.uk/~time/papers/reg/regtalk.ps, Imperial/TP/97-98/69, Report No. hep-ph/98mmnnn.
- [19] T.S.Evans, in preparation.