

Neutrino Magnetic Moment and Solar Neutrino Experiments

1

Ana M. Mourão^{a *} and Anna Rossi^{b †}

^aCENTRA - Centro Multidisciplinar de Astrofísica and Dep. of Physics, I.S.T.
Avenida Rovisco Pais, 1, 1096 Lisboa Codex - Portugal

^bCFIF - Centro de Física das Interações Fundamentais - I.S.T.
Avenida Rovisco Pais, 1, 1096 Lisboa Codex - Portugal

We have studied the effect of a non-vanishing neutrino magnetic moment (μ_ν) on the ν_x ($x=e, \mu, \tau$) elastic scattering off electrons for the Super-Kamiokande detector. The bounds on the μ_ν we have obtained are comparable to that extracted from laboratory experiments. Furthermore, we outline the potential of the Borexino experiment which may be sensitive to neutrino magnetic moments $\lesssim 10^{-10} \mu_B$. In our analysis we have considered both cases of Majorana and Dirac neutrinos.

1. Introduction

The solar neutrino problem (SNP) is nowadays regarded as a direct evidence for physics beyond the standard electroweak model. This is due to the fact that the observed deficit of electron neutrinos in all solar neutrino experiments can only be explained assuming that non-zero neutrino masses and/or neutrino magnetic moments might lead to flavour, spin or spin-flavour neutrino oscillations of the solar left-handed neutrinos ν_{eL} [1]-[6]. At the same time a lot of work has been done also to understand the implications of the uncertainties in helioseismology [7] and nuclear physics [8] for the SNP.

As was already pointed out in the literature [9]-[13] a non-vanishing neutrino magnetic moment, μ_ν , can also affect the neutrino elastic scattering off electrons through which solar ν 's are detected in Super-Kamiokande (SK). Therefore the expected signal in such a detector may depend also on the electromagnetic properties of the neutrinos.

In previous works [12,13] bounds have been obtained on μ_ν from Kamiokande and Super-Kamiokande data, taking into account the restrictions on the ν_e survival probability imposed by Homestake, Gallex and SAGE experiments.

In this work we present an updated analysis of the effect of a non-vanishing magnetic moment on the scattering off electrons for the Super-

Kamiokande detector, similarly to the study carried on in [13]. We have considered both the cases of Dirac and Majorana neutrinos.

For definiteness we examine the two neutrino system $\nu_e - \nu_x$ ($x=\mu, \tau$) with a non-zero mass difference δm^2 . In the solar interior the spin-flavour resonant conversion [5] $\nu_{eL} \rightarrow \nu_{xR}$, with probability P_R , occurs at higher matter density with respect to the usual MSW resonant conversion $\nu_{eL} \rightarrow \nu_{xL}$ [6], characterised by the probability P_L ¹. As a result of both conversions we have three different neutrino ‘flavours’ reaching the Earth:

$$\begin{aligned}\Phi_{\nu_{xR}} &= P_R \Phi_{SSM}, \\ \Phi_{\nu_{xL}} &= P_L(1 - P_R) \Phi_{SSM} \equiv P_{xL} \Phi_{SSM}, \\ \Phi_{\nu_{eL}} &= (1 - P_L)(1 - P_R) \Phi_{SSM} \equiv P_{eL} \Phi_{SSM},\end{aligned}\tag{1}$$

here Φ_{SSM} is the standard solar model (SSM) prediction for a certain component of the neutrino flux [14]. In the above relations² there are only two independent quantities, *e.g.* P_{eL} and P_R . Hence $P_L = 1 - \frac{P_{eL}}{1 - P_R}$ where $P_R \leq 1 - P_{eL}$. On the basis of this picture we have calculated the expected signal in SK experiment.

For the sake of simplicity, we assume that all the probabilities are energy independent. In particular the present SK data on the ν_{eL} energy spectrum do not yet exclude that P_{eL} is energy

¹We remind that the $\nu_{xL} \rightarrow \nu_{eR}$ spin-flavour resonance would occur at a much higher matter density, for given δm^2 . We assume therefore that it does not take place in the solar interior.

²The picture envisaged from the eq. (1) could appear, for example, when the MSW and spin-flavour resonances lie far away from each other to be treated separately. This may be the case in the central region of the sun.

*This work was in part supported by JNICT projects PRAXIS/PCEX/P/FIS/4/96 and ESO/P/PRO/1127/96, Fundação Oriente and GTAE-Lisbon.

[†]Supported by a grant from GTAE - Lisbon.

independent for $E_\nu \geq 6.5$ MeV [15].

2. Signal in the Super-Kamiokande detector

The total signal in the Super-Kamiokande experiment, for the case of Dirac neutrinos, can be written as

$$\begin{aligned} R_{SK}^{\text{total}} &= R_{SK}^{\text{w+em}} \\ &= P_{eL} [\langle \sigma_{\nu_{eL}}^{\text{w}} \rangle + \langle \sigma_{\nu_{eL}}^{\text{em}} \rangle] \\ &\quad + P_{xL} [\langle \sigma_{\nu_{xL}}^{\text{w}} \rangle + \langle \sigma_{\nu_{xL}}^{\text{em}} \rangle] \\ &\quad + P_R \langle \sigma_{\nu_{xR}}^{\text{em}} \rangle \end{aligned} \quad (2)$$

where the averaged total $\nu - e$ cross sections are

$$\langle \sigma_\alpha^i \rangle = \int dE_\nu \Phi_{\text{SSM}}^{\text{8B}}(E_\nu) \sigma_\alpha^i(E_\nu), \quad (3)$$

where $\alpha = \nu_{eL}, \nu_{xL,R}$ and $i=\text{em}$ for electromagnetic and $i=\text{w}$ for weak cross sections. $\Phi_{\text{SSM}}^{\text{8B}}(E_\nu)$ is the ^8B solar neutrino flux from BP98 [14]. In the calculation of the cross sections $\sigma_\alpha^i(E_\nu)$ we have taken into account the energy resolution of the detector [16]. The electromagnetic cross section can be taken e.g. from Kerimov *et al* [17]. Taking into account that $\langle \sigma_\alpha^{\text{em}} \rangle \propto \mu_\nu^2$ it is easy to see that for $P_{eL} = 1, P_R = P_{xL} = 0$, and $\mu_{\nu_e} = 0$ we obtain the SSM expectation for the signal.

In our analysis we use the most recent SK data for the ^8B solar neutrino flux, $\Phi_{SK}^{\text{exp}} = (2.44 \pm 0.05) \times 10^6 \text{ cm}^2\text{s}^{-1}$ [18], normalised to the SSM prediction - $\Phi_{\text{BP98}}^{\text{th}} = (5.15 \pm 0.98) \times 10^6 \text{ cm}^2\text{s}^{-1}$ [14], namely

$$Z_K = \frac{\Phi_{SK}^{\text{exp}}}{\Phi_{\text{BP98}}^{\text{th}}} = 0.47 \pm 0.09. \quad (4)$$

As opposite to the Super-Kamiokande experiment, the Homestake detector is only sensitive to the ν_{eL} component of the solar neutrino flux and the total rate is mainly due to ^8B neutrinos. Considering the experimental data [19] $Z_{\text{Cl}} = 0.28 \pm 0.03$, we can assume that for the higher energy spectrum of ^8B neutrinos, $E_\nu \geq 6$ MeV, the neutrino survival probability is $P_{eL} \sim 30\%$. Therefore this implies a total depletion of the intermediate energy ^7Be neutrinos as the present understanding of the SNP points to [1]-[3].

We can now find the values of the neutrino magnetic moment compatible with solar neutrino experiments, i.e needed to obtain the signal observed at the Superkamiokande.

3. Limits on neutrino magnetic moments

We have studied the impact of non-vanishing μ_ν in the SK signal taking into account the contribution of the several neutrino ‘flavours’ as shown in (1). In Fig. 1 we show the contour-plots for $Z_K = 0.47$ in the parameter space (P_R, μ_ν) , where $\mu_\nu \equiv \mu_{\nu_e} = \mu_{\nu_x}$. We repeat our analysis with other values of Z_K , just to understand implications of uncertainties in the SSM used in the evaluation of μ_ν . Notice that the larger the value of P_R the larger the value of μ_ν needed to saturate Z_K in order to compensate for the loss of the ν_{xL} component (recall that P_{eL} is fixed at 0.3). The present experimental Z_K implies the bound $\mu_\nu \lesssim (2 \div 5) \times 10^{-10} \mu_B$, almost independently of P_R .

We have considered also the case with $\mu_\nu \equiv \mu_{\nu_x}$ and $\mu_{\nu_e} = 0$. In this case we obtain a similar plot as that in Fig.1 and the limit $\mu_\nu \lesssim 5 \times 10^{-10} \mu_B$.

For the sake of completeness we have also studied the case of Majorana neutrinos, for which the *antineutrino* state $\nu_{xR} = \nu_{xL}^c \equiv \tilde{\nu}_x$, is ‘active’, having both electromagnetic and standard weak interactions. Hence in the eq. (2) one more term is to be added, *i.e.* $P_R \langle \sigma_{\tilde{\nu}_x}^{\text{w}} \rangle$. We remember that for Dirac neutrinos both diagonal or transition μ_ν can be generated [20], while for Majorana neutrinos only transition (off-diagonal) magnetic moments are allowed [21].

In Fig. 2 we show our result in this scenario: the upper bound for Majorana neutrinos comes out to be similar, $\mu_\nu \lesssim (1.5 \div 4.0) \times 10^{-10} \mu_B$ for the extreme case $P_R = 0.7$. Note that for $P_R < 0.7$, smaller values of μ_ν are in principle tolerated as the $\tilde{\nu}_x$ ’s contribute to the weak cross section (compare with the Dirac case shown in Fig. 1). Our upper limit was obtained assuming vanishing vacuum mixing angle (and then $P_L = 0$), thereby satisfying the experimental constraints on the $\tilde{\nu}_e$ which could emerge from the subsequent vacuum oscillation $\tilde{\nu}_x \rightarrow \tilde{\nu}_e$ [22].

We conclude that the bounds we have obtained




Figure 1. The contour plots of the expected signal versus the SSM prediction Z_K in Super-Kamiokande in the (P_R, μ_ν) parameter space in the case of Dirac neutrinos. The dotted, solid, dashed and dot-dashed curves correspond to $Z_K = 0.55, 0.47, 0.4$ and $Z_K = 0.37$, respectively. μ_ν is given in units of Bohr magneton.

are slightly more restrictive than those from accelerator experiments, namely in the case of μ_{ν_μ} it is $\mu_{\nu_\mu} < 7.4 \times 10^{-10} \mu_B$ [25]. However our bounds are still not comparable to those from reactor experiments, - $\mu_{\nu_e} < 1.8 \times 10^{-10} \mu_B$. Needless to say that our results are more stringent in the case of μ_{ν_τ} for which $\mu_{\nu_\tau} < 5.4 \times 10^{-7} \mu_B$ [24].

Finally, we have discussed the potential of the future Borexino experiment [23] which will detect ^7Be neutrinos through $\nu-e$ elastic scattering. For $\mu_\nu = 10^{-10} \mu_B$, the cross section σ^{em} can be comparable to $\sigma_{\nu_e}^w$ for $E_\nu \leq 1$ MeV. Therefore we can expect a substantial signal in Borexino even in the case of complete conversion of the initial ^7Be - ν_e 's into ν_{xR} or $\tilde{\nu}_x$. In Fig. 3 we have plotted the energy distribution of the events for Borexino in the case of complete conversion $\nu_e \rightarrow \tilde{\nu}_x$ (dotted line). We note that in this distribution we have taken into account the contributions from all solar neutrino components. However the ^7Be neutrinos contribute to more than 90% of the signal. Other important contribution is given by the *pep* flux




Figure 2. The same as in Fig.1 but for the Majorana case.

of solar neutrinos. For comparison we also shown the SSM distribution (solid line). We can expect $\sim 50\%$ of the SSM prediction and a specific distortion of the spectrum. This is in contrast with the case of pure MSW conversion ($\mu_\nu = 0$) $\nu_e \rightarrow \nu_x$ that would imply a $(20 \div 25)\%$ reduction in the signal.

4. Conclusions

In this contribution we have updated the analysis on the effect of a non-vanishing neutrino magnetic moment on the $\nu-e$ cross section in Super-Kamiokande experiment. The limits we achieved $-\mu_\nu \lesssim (2 \div 5) \times 10^{-10} \mu_B$ - remain comparable to that extracted from the previous Kamiokande data as the electromagnetic cross section is smaller than the weak one for the energy range involved, $E \geq 6$ MeV. Therefore experiments with a much lower energy threshold - such as Borexino or Hellaz - could exhibit a much better sensitivity to a non-zero μ_ν and consequently provide a better testing of the spin-flavour resonant conversion itself as a solution to the SNP[23,26].




Figure 3. The energy distribution of the events in Borexino experiment. The dotted curve represents the expected signal in the case of complete $\nu_e \rightarrow \bar{\nu}_x$ (Majorana case) with $\mu_\nu = 10^{-10} \mu_B$. The solid line corresponds to the SSM distribution.

Acknowledgments

We are grateful to E. Akhmedov, L. Bento, D. Gough and Y. Suzuki for very useful discussions. One of us (A.M.) would like to thank the organisers for the invitation to present this work and for the very stimulating $\nu 98$ Conference.

REFERENCES

1. V. Castellani, S. Degl'Innocenti and G. Fiorentini, *Astron. Astrophys.* 271 (1993) 601.
2. K. M. Heeger and R. G. H. Robertson, *Phys. Rev. Lett.* 77 (1996) 3720.
3. J. N. Bahcall, P. I. Krastev and A. Yu. Smirnov, *Phys. Rev.* D58 (1998) to appear.
4. A. Cisneros, *Astrophys. Space Sci.* 10 (1971) 87; M. B. Voloshin, M. I. Vysotsky, L. B. Okun, *Sov. J. Nucl. Phys.* 44 (1986) 440.
5. C. S. Lim and W. J. Marciano, *Phys. Rev.* D37 (1988) 1368; E. Kh. Akhmedov, *Sov. J. Nucl. Phys.* 48 (1988) 382; E. Kh. Akhmedov, *hep-ph/9705451*; H. Minakata and H. Nunokawa *Phys. Rev. Lett.* 63 (1989) 121.
6. L. Wolfenstein, *Phys. Rev.* D17 (1978) 2369; D20 (1979) 2364; S. P. Mikheyev, A. Smirnov, *Sov. J. Nucl. Phys.* 42 (1985) 913.
7. D. Gough, in these Proceedings.
8. W. C. Haxton, in these Proceedings; E. G. Adelberger et al., *astro-ph/9805121*.
9. R. Barbieri and G. Fiorentini, *Nucl. Phys.* B304 (1988) 909; A. Suzuki et al., *Phys. Rev.* D43 (1991) 3557.
10. A. B. Balantekin, P. J. Hatchell and F. Loreti, *Phys. Rev.* D41 (1990) 3583.
11. K.S. Babu, N. Mohapatra and I. Z. Rothstein, *Phys. Rev.* D44 (1991) 2265.
12. A. M. Mourão, J. Pulido, J. Ralston, *Phys. Lett.* B285 (1992) 364; Erratum-ibid. 288 (1992) 421.
13. J. Pulido and A. M. Mourão, *Phys. Rev.* D57 (1998) 1794.
14. J. N. Bahcall, S. Basu and M. H. Pinsonneault, *Phys. Lett.* B433 (1998) 1.
15. Y. Fukuda et al.- Super-Kamiokande Coll., *Phys. Rev. Lett.* 81 (1998) 1158.
16. J. N. Bahcall, *Neutrino Astrophysics*, CUP, 1993.
17. B. Kerimov, M. Ya. Safin, H. Nazih, *Izvestiya Akademii Nauk SSSR, Ser. Fiz.* 52 (1988) 126; P. Vogel, J. Engel, *Phys. Rev.* D39(1989) 3378.
18. Y. Suzuki, in these Proceedings.
19. B. Cleveland et al., *Nucl. Phys.* B38 (1995) 47.
20. K. Fujikawa and R. Shrock, *Phys. Rev. Lett.* 45 (1980) 963.
21. J. Schechter and J. W. F. Valle, *Phys. Rev.* D24 (1981) 1883.
22. R. Barbieri et al., *Phys. Lett.* B259 (1991) 119; M. Aglietta et al., *JETP Lett.* 63 (1996) 791; A. A. Bykov et al., *hep-ph/9808342*.
23. C. Arpesella et al., (Borexino Collaboration), Proposal of BOREXINO (1991); Z. G. Berezhiani and A. Rossi *Phys. Rev.* D51 (1995) 5229.
24. Review of Particle Properties, C. Caso et al., *EPJ* C3 (1998) 1.
25. D. A. Krakauer et al. *Phys. Lett.* B 252 (1990) 177.
26. S. Pastor et al., *hep-ph/9803378*.