

An Alternative to See-Saw

J. I. Silva-Marcos[‡]

NIKHEF, Kruislaan 409, 1098 SJ Amsterdam, The Netherlands

We give a new mechanism for generating very small, and almost degenerate, neutrino masses, without resorting to the see-saw mechanism or unnatural small Yukawa couplings. It requires the existence of at least 4 families with an almost democratic structure for the Yukawa couplings. It is also proven that this structure can account for large lepton mixings of the three light leptons.

16.60Pq, 12.15Ff

As is well known, the famous see-saw mechanism [1] is particularly attractive for explaining the smallness of the three neutrino masses. However, another mechanism might be responsible for very small neutrino masses. Within the framework of 4 families [2], an almost democratic structure for the Yukawa couplings is sufficient to explain the smallness of the masses of the three known neutrinos. One can see this by considering the democratic limit for the neutrino mass matrix,

$$m_\nu = \frac{\mu_\nu}{4} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix} \quad (1)$$

Obviously, the eigenvalue spectrum of this mass matrix consists of three neutrinos with zero mass and one massive heavy neutrino, $(0, 0, 0, \mu_\nu)$. The democratic structure for the neutrino mass matrix is very appealing. On the one hand, all Yukawa couplings can be as large as any of the other fermion Yukawa couplings, while, on the other hand, all, but one, eigenvalues of the mass matrix are zero, or practically zero in the case of an almost democratic structure. Furthermore, democratic mass matrices are related to exact S_n family symmetries. The existence of three very light neutrinos, leads us to consider, at least, 4 families and an almost democratic structure. Thus, we can explain very small neutrino masses without resorting to the see-saw mechanism or unnatural small Yukawa couplings.

The universal strength for Yukawa couplings (USY) [3] is a natural framework to study small deviations from the democratic limit [4]. It was proven [5], that for three generations, USY can account for almost degenerate neutrinos [6] and for the strong hierarchy in the charged lepton sector. In this paper we shall further generalize these results to 4 generations of Dirac neutrinos. With regard to lepton mixing, we shall consider here only cases where the mixing (mainly) occurs for the light leptons. Thus the mixings of the fourth heavy neutrino with the three light charged leptons and the mixings of the fourth heavy charged lepton with the three light neutrinos will be very small.

As a first step, we analyze two preliminary limit situations, one for the neutrino and the other for the charged leptons, which represent small deviations from the exact democratic limit in Eq.(1), and which serve as simplifications of a more realistic scenario given further on.

For the neutrinos, in this first step, the three light neutrinos acquire a mass but maintain their degeneracy. We achieve this by allowing a very small phase α for the diagonal elements of the neutrino mass matrix, such that the democratic limit of Eq.(1) becomes,

$$m_\nu = \frac{\mu_\nu}{4} \begin{bmatrix} e^{i\alpha} & 1 & 1 & 1 \\ 1 & e^{i\alpha} & 1 & 1 \\ 1 & 1 & e^{i\alpha} & 1 \\ 1 & 1 & 1 & e^{i\alpha} \end{bmatrix} \quad (2)$$

As can be readily verified, the mass spectrum of this neutrino mass matrix consists of three light neutrinos with equal mass, and one heavy neutrino:

$$m_{\nu_i} = \frac{1}{2} \mu_\nu \left| \sin\left(\frac{\alpha}{2}\right) \right| \quad , \quad i = 1, 2, 3 \quad ; \quad m_{\nu_4} = \mu_\nu \sqrt{1 - \frac{3}{4} \sin^2\left(\frac{\alpha}{2}\right)} \quad (3)$$

It is clear that one can choose the mass of the light neutrinos as small as one wants by taking the phase α to be very small. Thus the breaking of the S_4 symmetries will be very small. The diagonalization of this neutrino mass matrix is done with:

$$F_\nu = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{6}} & \frac{-1}{\sqrt{12}} & \frac{1}{2} \\ 0 & \frac{2}{\sqrt{6}} & \frac{-1}{\sqrt{12}} & \frac{1}{2} \\ 0 & 0 & \frac{3}{\sqrt{12}} & \frac{1}{2} \\ \frac{-1}{\sqrt{2}} & \frac{-1}{\sqrt{6}} & \frac{-1}{\sqrt{12}} & \frac{1}{2} \end{bmatrix} \quad (4)$$

One must realize that, at this stage, that there is much freedom in the choice of the first three column vectors of the unitary matrix of Eq.(4), because of the degeneracy of the three light eigenvalues. Therefore, these three vectors are (for now) given, up to modulo 3×3 unitary rotations, which, of course, will be fixed when the light neutrino degeneracy is lifted.

With regard to the charged leptons, we consider in this first step the limit when the electron and muon are massless while the tau acquires a mass (as well as the heavy fourth heavy charged lepton). One can accomplish this with just one phase r ,

$$m_\ell = \frac{\mu_\ell}{4} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & e^{ir} \end{bmatrix} \quad (5)$$

and one gets the following masses for the for the tau and the heavy charged lepton,

$$m_\tau = \frac{3}{4\sqrt{2}} \mu_\ell \frac{|\sin(\frac{r}{2})|}{\sqrt{1+(1-\frac{9}{16}\sin^2(\frac{r}{2}))^{1/2}}} \quad ; \quad m_{\ell_4} = \frac{1}{\sqrt{2}} \mu_\ell \sqrt{1+(1-\frac{9}{16}\sin^2(\frac{r}{2}))^{1/2}} \quad (6)$$

From Eq.(6) one finds, in first order in r , that $m_\tau/m_{\ell_4} = 3|r|/16$, which for a value of $m_{\ell_4} = 5 \text{ TeV}$ would fix $r = 1.9 \times 10^{-3}$. The diagonalization of m_ℓ in Eq.(5) requires a matrix

$$F_\ell = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{6}} & \frac{-1}{\sqrt{12}} & \frac{1}{2} \\ \frac{-1}{\sqrt{2}} & \frac{-1}{\sqrt{6}} & \frac{-1}{\sqrt{12}} & \frac{1}{2} \\ 0 & \frac{2}{\sqrt{6}} & \frac{-1}{\sqrt{12}} & \frac{1}{2} \\ 0 & 0 & \frac{3}{\sqrt{12}} & \frac{1}{2} \end{bmatrix} \quad (7)$$

but again, here one must bear in mind that the first two charged lepton masses are still zero, and that therefore, the first two column vectors of F_ℓ are, in principal, given up to modulo unitary 2×2 rotations, which shall be fixed when the electron and the muon acquire a mass. The unitary matrix F_ℓ of Eq.(7) is not the full matrix which diagonalizes the charged lepton mass matrix of Eq.(5) because of mixing between the tau and the heavy charged lepton. However, in first order, it is a very good approximation. It can be readily verified that the mixing between the tau and the charged lepton is of the order of $o(m_\tau/m_{\ell_4})$. The lepton mixing matrix is given (in first order) by,

$$V_{\text{mixing}}^{\text{lepton}} = F_\ell^\dagger \cdot F_\nu = \begin{bmatrix} \frac{1}{2} & \frac{-\sqrt{3}}{2} & 0 & 0 \\ \frac{-\sqrt{3}}{6} & \frac{-1}{6} & \frac{4\sqrt{2}}{6} & 0 \\ \frac{-\sqrt{6}}{3} & \frac{-\sqrt{2}}{3} & \frac{-1}{3} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad (8)$$

One sees that relevant mixing between the leptons only occurs for the light species.

The limit cases which we have considered so far for the neutrino as well as the charged lepton mass matrices and diagonalizations are crucial. In essence, they contain the first order approximations of a realistic example which we present now, and where the neutrino degeneracy is lifted and the electron and muon acquire mass. As an example of a realistic ansatz we propose:

$$m_\nu = \frac{\mu_\nu}{4} \begin{bmatrix} e^{i\alpha} & 1 & 1 & 1 \\ 1 & e^{i\alpha(1+\varepsilon)} & 1 & 1 \\ 1 & 1 & e^{i\alpha(1+\delta)} & 1 \\ 1 & 1 & 1 & e^{i\alpha} \end{bmatrix} \quad ; \quad m_\ell = \frac{\mu_\ell}{4} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & e^{ip} & 1 & 1 \\ 1 & 1 & e^{iq} & 1 \\ 1 & 1 & 1 & e^{ir} \end{bmatrix} \quad (9)$$

The slight deviation from the limit case of the neutrino mass matrix in Eq.(2), induced by $\delta, \varepsilon \neq 0$, will generate the neutrino mass differences Δm_{32}^2 and Δm_{21}^2 . One finds that $\delta = o(\Delta m_{32}^2/m_{\nu_3}^2)$ and $\varepsilon = o(\Delta m_{21}^2/m_{\nu_3}^2)$. This mass

matrix has one exact light neutrino mass eigenvalue as given in Eq.(3). For the charged lepton mass matrix, $q \neq 0$ and $p \neq 0$ will generate masses respectively for the muon and electron. One has $q = o(m_\mu/m_{\ell_4})$ and $p = o(m_e/m_{\ell_4})$ [7]. The realistic ansatz of Eq.(9) and its choice of phases, will fix the explained arbitrary unitary rotations associated with the degeneracy of the three light neutrinos and the zero masses of the electron and muon, because of the explicit lifting of that degeneracy and the non-zero masses. However, this choice of phases is such that these rotations are equal to the identity. Therefore, in first order, we shall have precisely the same diagonalization matrices as in Eqs.(4, 7) and a lepton mixing matrix as in Eq.(8). To complete our ansatz, we give an exact numerical example with input:

$$\begin{aligned} \alpha &= 2 \times 10^{-8} & \delta &= 2 \times 10^{-3} & \varepsilon &= 3 \times 10^{-6} \\ r &= 1.9 \times 10^{-3} & q &= 1.3 \times 10^{-4} & p &= 8.2 \times 10^{-7} \end{aligned} \quad (10)$$

which corresponds with the following output:

$$\begin{aligned} \Delta m_{21}^2 &= 9 \times 10^{-6} \text{ eV}^2 & \Delta m_{32}^2 &= 6.7 \times 10^{-3} \text{ eV}^2 & m_{\nu_3} &= 1.5 \text{ eV} \\ m_e &= 511 \text{ KeV} & m_\mu &= 105.7 \text{ MeV} & m_\tau &= 1777 \text{ MeV} \end{aligned} \quad (11)$$

In addition, there will be two heavy leptons associated with (the large values of) μ_ν and μ_ℓ . One obtains $\mu_\nu = m_{\nu_4} = 300 \text{ GeV}$ and $\mu_\ell = m_{\ell_4} = 5 \text{ TeV}$. The mass of the heavy neutrino is far above the LEP bound $M_Z/2$. The lepton mixing matrix will also have contributions of higher orders,

$$\left| V_{\text{mixing}}^{\text{lepton}} \right| = \begin{bmatrix} 0.501 & 0.865 & 1.1 \times 10^{-3} & 1.4 \times 10^{-7} \\ 0.305 & 0.176 & 0.936 & 1.7 \times 10^{-5} \\ 0.810 & 0.469 & 0.353 & 2.0 \times 10^{-4} \\ 1.7 \times 10^{-4} & 9.6 \times 10^{-5} & 5.4 \times 10^{-5} & 1. \end{bmatrix} \quad (12)$$

To summarize, we have given a new mechanism for generating very small, and almost degenerate, neutrino masses, without resorting to the see-saw mechanism or unnatural small Yukawa couplings. This new mechanism requires the existence of at least 4 families [8]. Within the framework of the universal strength for Yukawa couplings one only has very small phases, which may be associated with almost exact S_4 symmetries. It was also proven that this structure for the Yukawa couplings can account for large lepton mixing of the three light leptons.

This work was partially supported by the Fundação para a Ciência e Tecnologia of the Portuguese Ministry of Science and Technology.

[†] On leave from the Centro de Física das Interações Fundamentais of the Instituto Superior Técnico of Lisbon. E-mail address: juca@nikhef.nl

- [1] T. Yanagida, in *Proc. of the Workshop on Unified Theory and Baryon Number in the Universe*, KEK report 79-18, 1979; M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, North Holland, 1979.
- [2] Recently P.Hung constructed a model with 4 families, having also one massive fourth neutrino, but, where at tree level, all light neutrinos are massless; P. Hung, hep-ph/9810419.
- [3] G.C. Branco, J.I. Silva-Marcos and M.N. Rebelo, Phys. Lett. B 237 (1990) 446; G. C. Branco and J. I. Silva-Marcos, Phys. Lett. B 359 (1995) 166; G.C. Branco, D. Emmanuel-Costa and J.I. Silva-Marcos, Phys. Rev. D 56 (1997) 107;
- [4] For other deviations from the democratic limit see J.I. Silva-Marcos, hep-ph/9807549.
- [5] G.C. Branco, M.N. Rebelo and J.I. Silva-Marcos, Phys. Lett. B 428 (1998) 136.
- [6] If one assumes that the neutrinos constitute the hot dark matter of the universe, having a mass a few eV, then one is lead to highly degenerate neutrinos. A systematic analysis of Majorana neutrino degeneracy was proposed by G.C. Branco, M.N. Rebelo and J.I. Silva-Marcos, hep-ph/9810328.
- [7] In a forthcoming paper we shall specify the mass ratio dependence of the phases.
- [8] The generalization of the mechanism here proposed, for more families, is straightforward.