

Positive Order H^2 Mass Corrections and Ack-Dine Baryogenesis

John McDonald¹

Department of Physics and Astronomy, University of Glasgow, Glasgow G12 8QQ, SCOTLAND

February 5, 2020

Abstract

It is usually assumed that the order H^2 corrections to the SUSY-breaking mass squared terms in the early Universe must be negative in order to allow the Ack-Dine mechanism to work. We reconsider this assumption in the context of D-term inflation models for the case where the mass squared term has a correction cH^2 with $c > 0$. We show that, in general, the baryon asymmetry is likely to be too small if $c > 9/16$. However, for c as large as 0.5 the observed baryon asymmetry can be readily generated; in particular, for $d = 6$ directions the observed asymmetry can be produced for a wide range of reheating temperatures, in contrast with the case of negative H^2 corrections which require a reheating temperature around 1 GeV. Thus positive H^2 corrections do not rule out the Ack-Dine mechanism and can even greatly broaden its applicability to inflation models.

¹mcDonald@physics.gla.ac.uk

1 Introduction

Ack-Dine (AD) baryogenesis [1] is a natural possibility for generating the baryon asymmetry in the context of the MSSM. It is based on inducing a baryon asymmetry in a coherently oscillating squark condensate. In the modern view of the AD mechanism [2, 3], the initial value of the AD field and its subsequent evolution depends crucially on the order H^2 corrections to the SUSY breaking mass squared terms that are expected once the energy density in the early Universe and non-minimal Kahler couplings are taken into account [2, 4]. The natural assumption to make is that such corrections must have a negative sign, so that the minimum of the AD potential is non-zero at early times when the expansion rate is large, turning positive (and so allowing the AD scalar to coherently oscillate) only once the gravity-mediated SUSY breaking terms come to dominate. Positive order H^2 mass corrections are commonly thought to result in an AD field with an amplitude too small to account for the B asymmetry. However, it is not obvious that this is generally true and it is this issue that we wish to explore here. The case for negative order H^2 corrections was originally put forward in the context of F-term inflation models, under the assumption that the sign of the H^2 corrections during and after inflation are the same [3]. In this case, if the H^2 terms were positive, the AD scalar would be exponentially damped nearly to zero during inflation. It was later assumed [5] that the same condition had to hold for D-term inflation models [6]. However, these have no order H^2 corrections to scalar masses along D-flat directions during inflation. We will show that in D-term inflation models with positive order H^2 corrections (and also in F-term inflation models which have a change in sign of the H^2 correction from negative to positive after inflation) it is possible to have successful AD baryogenesis, with a much wider applicability to inflation models than in the case of negative H^2 corrections.

In the AD mechanism, a baryon asymmetry is induced when the A-term in the scalar potential produces a phase shift between the real and imaginary parts of the AD field [7]. In D-term inflation models, the A-term receives no order H correction either during or after inflation [5, 6]. Thus the initial phase of the AD scalar is random and so

is typically of the order of one relative to the real direction. The baryon asymmetry will be fixed once $H < m$, where $m \approx 100 \text{ GeV}$ is the gravity-mediated SUSY breaking scalar mass term, at which point the B violating terms become suppressed relative to the B conserving terms in the potential. Thus in order to calculate the baryon asymmetry we need to calculate the amplitude and the phase shift of the AD field at $H = m$.

2 Amplitude at $H = m$.

The scalar potential of the AD field along a dimension d at direction has the general form [3],

$$V = (m^2 + cH^2)j j^2 + \frac{A}{dM^{d-3}} + \text{h.c.} + \frac{2j j^{2(d-1)}}{M^{2(d-3)}}; \quad (1)$$

where $M = M_{\text{Pl}} = \frac{P}{8}$ and typically $1 = (d-1)!$ for Planck-scale non-renormalizable terms. We will focus on R-parity conserving models, for which d is even [3], and consider the cases $d = 4$ and $d = 6$. The evolution of the amplitude depends on whether the $j j^2$ term or the non-renormalizable term is dominant, with the value of the AD scalar below which the potential is j^2 dominated, c_c , being given by

$$c_c = \frac{P}{2} - \frac{c}{2} \frac{1}{2^{(d-2)}} H^2 M^{2(d-3) - \frac{1}{2^{(d-2)}}}; \quad (2)$$

where $j j^2 = \frac{P}{2}$.

Evolution for $c < c_c$: When $c < c_c$ the scalar field evolves according to

$$+ 3H \dot{\phi} = -cH^2 \phi; \quad (3)$$

After inflation (but prior to reheating), the Universe is matter dominated by inflation oscillations. For a matter dominated Universe Eq. (3) has the solution

$$\phi/a = \frac{1}{2} \frac{4}{2} \frac{3}{2} \frac{9}{4} 4c^5; \quad (4)$$

where a is the scale factor. There are then two distinct cases. If $c > 9/16 = 0.56$, the square root is imaginary and we have a damped oscillating solution of the form

$$= \frac{a_0}{a} e^{-\frac{3}{4} \log \frac{a_0}{a}} \cos \left(\frac{1}{2} \sqrt{\frac{9}{4c} - \frac{9}{4}} \log \frac{a_0}{a} \right); \quad (5)$$

Note that the amplitude falls off less rapidly than the corresponding amplitude for a coherently oscillating scalar with a constant mass term, for which $\propto a^{-3/2}$. If, on the other hand, $c < 9/16$, then the field is damped but not oscillating, with a magnitude given by

$$= \frac{a_0}{a} \quad (6)$$

Evolution for $c > c_c$: For $c > c_c$, the field is effectively oscillating in a $a^{2(d-1)}$ potential. The energy density evolves as $\propto a^{6(d-1)/d}$ and so the amplitude of the oscillation evolves as

$$\propto a^{\frac{3}{d}} \quad (7)$$

Thus for a field which starts oscillating at $\phi_0 > \phi_c$, the amplitude at $m < H < H_c$ is given by (from now we use ϕ)

$$= \frac{H}{H_c}^{\frac{2}{3}} \frac{H_c}{H_0}^{\frac{2}{d}} \phi_0; \quad (8)$$

where H_c is the value at which $\phi = \phi_c$,

$$H_c^{4/d} = \frac{2^{1/d-2}}{c} \frac{1}{H_0}^{2(d-2)} \frac{1}{M}^{d(d-3)} \left(\frac{\phi_0}{2}\right)^{d(d-2)} \quad (9)$$

and H_0 is the value at the end of inflation. For the case $\phi_0 < \phi_c$, $H_c \neq H_0$ in Eq. (8).

For the $d = 4$ case H_c is not fixed by Eq. (9), because both ϕ and ϕ_c for a given value of H are proportional to $H^{1/2}$. Thus if the value of ϕ at the end of inflation, ϕ_0 , is less than ϕ_c then it will remain so until $H = m$. ϕ_c at H_0 is given by

$$\phi_c = 7 \cdot 10^{15} \frac{c}{2}^{1/4} \frac{H_0}{10^{13} \text{ GeV}}^{1/2} \text{ GeV}; \quad (10)$$

where we are considering the typical value at the end of inflation to be $H_0 = 10^{13} \text{ GeV}$, in accordance with the observed cosmic microwave background perturbations [9, 10]. For the $d = 6$ case, H_c and ϕ_c are given by

$$H_c = 1.7 \cdot 10^{27} \frac{c}{2}^{3/2} \frac{H_0}{10^{13} \text{ GeV}}^4 \frac{10^{16} \text{ GeV}}{\phi_0}^{1/12} \text{ GeV} \quad (11)$$

and

$$\phi_c = 5.5 \cdot 10^{17} \frac{c}{2}^{1/2} \frac{H_0}{10^{13} \text{ GeV}} \frac{10^{17} \text{ GeV}}{\phi_0}^{1/3} \text{ GeV}; \quad (12)$$

We first consider the case with the largest degree of damping, $c > 9=16 \approx 0.56$. In this case the amplitude of the oscillations evolves as $\propto a^{-3/4}$. We will see that $\rho_{\phi} < \rho_c$ must be satisfied in order to have acceptable energy density perturbations in both the $d=4$ and $d=6$ cases. In this case the field at H_{min} is generally given by

$$(H_{\text{min}})^3 \approx 10^{10} \frac{10^{13} \text{ GeV}}{H_0} \frac{1}{10^{16} \text{ GeV}} \quad (13)$$

Thus the amplitude of the AD field at H_{min} is not very small so long as the initial value of the AD field at the end of inflation is large enough. In fact, the amplitude of Eq. (13) is typically of the order of the amplitude expected in the $d=4$ case with negative H^2 corrections, $(M_{\text{pl}})^{1/2} \approx 10^{10} \text{ GeV}$. Therefore so long as the phase shift is sufficiently large we should be able to generate the observed asymmetry.

3 Phase Shift

In general, the phase shift will be the product of an initial CP violating phase and a "kinematical factor" coming from the time evolution of the real and imaginary components of the AD field. Assuming an initial phase of order one, the phase shift will be given by

$$\Delta \phi = \int_{t_{\text{min}}}^{t_{\text{max}}} \Delta m \, dt; \quad (14)$$

where Δm is the difference in the effective mass terms for the scalars along the real and imaginary directions of the AD field ($\phi = (\phi_R + i \phi_I) e^{i\theta}$). We will consider the phase shift for $\rho_{\phi} < \rho_c$; it is straightforward to show that the contribution to the phase shift from $\rho_{\phi} > \rho_c$ is generally negligible compared with this.

From the equations of motion for ϕ_R and ϕ_I , there is a shift in the effective mass squared term, $m_{\text{eff}}^2 \rightarrow m_{\text{eff}}^2 + m^2$ due to the A-term contribution, where $m_{\text{eff}}^2 = \partial^2 V / \partial \phi^2$ and where

$$m^2 = \frac{A}{2^{d-2} M^{d-3}} \frac{H_0^2}{H_0} \frac{H_0^{2/3(d-2)}}{H_0} \quad (15)$$

For small m^2 we therefore have

$$\Delta \phi \approx \Delta m^2 = 2m_{\text{eff}}^2 = K_d H_0^{2/3(d-2)-1}; \quad (16)$$

where

$$K_d = \frac{A}{2^{d-2} c^{1/2} M_{\text{pl}}^{d-3}} \frac{1}{H_0^{\frac{2}{3}(d-2)}} : \quad (17)$$

Thus the phase shift is given by

$$= \frac{2K_d}{3^{\frac{2}{3}(d-2)-2}} (H_0^{\frac{2}{3}(d-2)-2} - H^{\frac{2}{3}(d-2)-2}) : \quad (18)$$

In general this will be dominated by the contribution at H_{min} . The B asymmetry density at H_{min} is then $n_B = m^2 S_{\text{in}}$. This is related to the present baryon-to-photon ratio η_B by

$$\eta_B = \frac{n_B}{2} \frac{H_{\text{pl}}^2 M_{\text{pl}}^2}{T_R} : \quad (19)$$

We next calculate η_B for the $d = 4$ and $d = 6$ cases.

$d = 4$: In this case the phase shift is given by

$$\approx \frac{1}{2} \frac{10^{13} \text{ GeV}}{H_0} \frac{1}{10^{16} \text{ GeV}} 10^{11} : \quad (20)$$

where we have taken $A_{\text{min}} = 100 \text{ GeV}$. Thus typically the phase is of order one.

η_B is then given by

$$\frac{\eta_B}{10^{-10}} \approx 40 \frac{10^{13} \text{ GeV}}{H_0} \frac{1}{10^{16} \text{ GeV}} \frac{T_R}{10^9 \text{ GeV}} 10^{11} : \quad (21)$$

For the case $c > 9=16$ (corresponding to $\alpha = 3=4$) and with $H_0 = 10^{16} \text{ GeV}$ (the upper limit from scale-invariant density perturbations, as discussed below) this is marginally able to account for the observed B asymmetry ($\eta_B \approx (3-8) \times 10^{-10}$ [1]) if $T_R > 10^8 \text{ GeV}$, close to the upper bound from thermal gravitino production, $T_R < 10^9 \text{ GeV}$. As discussed in detail for the $d = 6$ case below, the asymmetry can be enhanced (corresponding to lower values of T_R being compatible with the observed B asymmetry) if c is even slightly less than $9=16$; for example, $c = 0.5$ gives a 5×10^3 enhancement of the B asymmetry. However, in the context of D-term inflation models, we will see that the $d = 4$ condensate, as in the case of negative order H^2 corrections [5], is unlikely to survive thermalization by the first stage of reheating after inflation.

$d = 6$: In this case the phase shift is given by

$$10^{-16} \frac{1}{2} \frac{10^{13} \text{ GeV}}{H_0} \frac{1}{10^{16} \text{ GeV}} 10^{11} : \quad (22)$$

Thus typically the phase shift is no larger than around 10^{-10} for $c > 9=16$. (This is much smaller than in the case of negative H^2 corrections, for which δ_B is then given by

$$\frac{\delta_B}{10^{-10}} \approx 3 \cdot 10^{16} \frac{1}{2 \cdot 8 \cdot 3} \frac{10^{13} \text{ GeV}}{H_0} \frac{1}{10^{16} \text{ GeV}} \frac{T_R}{10^9 \text{ GeV}} 10^{11 \cdot 3 \cdot 4} : \quad (23)$$

From this we see that even with $\phi_0 = 10^{17} \text{ GeV}$, which we will show is the upper limit from the scale-invariance of the density perturbation spectrum, the highly suppressed phase shift in the case $c > 9=16$ means that we cannot produce a value of δ_B much above 10^{-20} . Thus there is no possibility of accounting for the observed B asymmetry in this case. However, once $c < 9=16$ there is a rapid enhancement of the B asymmetry. This can be seen from the "enhancement factor" $(10^{11})^{3 \cdot 4}$ in Eq. (23). This gives the observed asymmetry for $T_R = 10^8 \text{ GeV}$ and $\phi_0 = 10^{17} \text{ GeV}$ if $c = 0.5$. The corresponding values for other ϕ_0 and T_R are given in Table 1. From this we see that, with values of c between 0.2 and 0.5, the observed asymmetry can be accommodated by a wide range of reheating temperatures.

Table 1. $d=6$ B asymmetry enhancement factor.

c	Enhancement factor	$T_R (\phi_0 = 10^{17} \text{ GeV})$	$T_R (\phi_0 = 10^{16} \text{ GeV})$
0.51	$3 \cdot 10^{10}$	10^9 GeV	10^{15} GeV
0.50	$1 \cdot 10^{11}$	10^8 GeV	10^{14} GeV
0.45	$5 \cdot 10^{14}$	10^4 GeV	10^{10} GeV
0.40	$5 \cdot 10^{17}$	10 GeV	10^7 GeV
0.35	$2 \cdot 10^{20}$	0.1 GeV	10^5 GeV
0.30	$4 \cdot 10^{22}$	10^3 GeV	10^3 GeV
0.20	$3 \cdot 10^{26}$	10^{-7} GeV	0.1 GeV

Therefore, although values of c close to or larger than 1 are indeed ruled out, for values as large as 0.5 the observed asymmetry can be generated along the $d = 6$ direction. Such a value of c seems entirely plausible and consistent with the expectation that c is of the order of one. Indeed, the observed asymmetry can be accommodated

by a much wider range of reheating temperatures than in the case of negative H^2 corrections, which requires T_R to be of the order of 1 GeV .

4 Density Perturbation Constraints

The magnitude of the B asymmetry depends on the initial value of the AD field. This cannot be too large during inflation, otherwise the energy density of the AD field will dominate the spectral index of the energy density perturbations, resulting in an unacceptable deviation from scale-invariance [12]. This has already been discussed for the $d = 6$ case in the context of D-term inflation, where it was shown that the requirement that the spectral index is dominated by the inflation field implies that [12]

$$\kappa_{\text{max}} < 0.48 \frac{g}{\tilde{g}}^{1=4} (M_{\text{Pl}})^{1=2} < 10^{17} \text{ GeV} ; \quad (24)$$

where g and \tilde{g} are the Fayet-Iliopoulos gauge coupling and superpotential coupling of the D-term inflation model and $M_{\text{Pl}} = 1.22 \times 10^{19} \text{ GeV}$ [6, 10]. For the $d = 4$ case the analogous requirement is that

$$\kappa_{\text{max}} < 0.23 \frac{g}{\tilde{g}}^{1=2} < 10^{16} \text{ GeV} ; \quad (25)$$

where in general \tilde{g} cannot be very small compared with g , otherwise there would be efficient parametric resonant decay of the inflation and violation of the thermal gravitino production bound [13]. However, there is a generic problem for the $d = 4$ direction in the context of D-term inflation models [5]. D-term inflation models typically have a first stage of reheating immediately after the end of inflation from the decay of the field of the inflation sector, with $T_R \sim g^{1=2} M_{\text{Pl}} \sim 10^{15} \text{ GeV}$. In order to avoid thermalization by this radiation density, the AD field at the end of inflation must be greater than $H_{\text{inf}}^{3=8} M_{\text{Pl}}^{5=8} \sim 2 \times 10^{16} \text{ GeV}$ [5]. Thus the upper bound from scale-invariance implies that the $d = 4$ directions with positive H^2 corrections are thermalized, as in the case of negative H^2 corrections. The $d = 6$ direction can, however, avoid thermalization and still be consistent with scale-invariance.

5 F-term Inflation

So far we have been considering D-term inflation. However, the above discussion will also apply to F-term inflation models if some conditions are met. The main differences between D-term and F-term inflation are that the AD scalar will receive order H^2 corrections to its mass both during and after inflation and that the A-term will also receive order H corrections. The effect of the order H corrections to the A-term is that the initial CP violating phase will be set by the explicit CP violating phase appearing in the A-term (which can be much smaller than one) rather than by a random initial condition. The effect of the positive order H^2 mass correction during inflation is to exponentially drive the AD scalar to zero, so ruling out AD baryogenesis [3]. However, it is still possible that the order H^2 term could become positive after inflation, if there was a sign change of the H^2 correction at the end of inflation. This is possible, for example, if the energy density after inflation was made up of two fields with comparable energies. In this case the H^2 term would be the sum of contributions from each field, and the sum of the contributions could change as the energy is transferred away from the vacuum energy at the end of inflation. Thus it is conceivable that the H^2 correction could be negative during inflation but positive after. The D-term discussion would then apply to this case also, since the value of the fields at the end of inflation from minimizing the potential during inflation simply corresponds to c .

6 Conclusions

In D-term inflation models, the A- $\tilde{\chi}$ -Dine mechanism with a positive cH^2 correction to the SUSY breaking mass squared term can generate the observed B asymmetry if $c < 0.5$. The $d = 4$ direction is ruled out by the near scale-invariance of the density perturbations together with the thermalization of the AD field during the first stage of reheating. The $d = 6$ direction, on the other hand, can successfully generate the observed baryon asymmetry, and for a much wider range of reheating temperatures than in the case of negative H^2 corrections. Similar conclusions apply to the case of

F-term inflation models with a positive H^2 correction after inflation. Therefore the AD mechanism can have a much wider range of applicability to particle physics models than previously thought. In particular, since the reheating temperature in $d = 4$ and $d = 6$ models with a CP violating phase 1 is constrained to be very high or very low (10^8 GeV and 1 GeV respectively), in many inflation models the AD mechanism with positive H^2 corrections will be the natural option.

Acknowledgements

This work has been supported by the PPARC.

References

- [1] I A A eck and M D ine, Nucl. Phys. B 249 (1985) 361.
- [2] M D ine, L Randall and S Thomas, Phys. Rev. Lett. 75 (1995) 398.
- [3] M D ine, L Randall and S Thomas, Nucl. Phys. B 458 (1996) 291.
- [4] M D ine, W Fischler and D Nemeschansky, Phys. Lett. B 136 (1984) 169;
G D Coughlan, R Holman, P Ramond and G G Ross, Phys. Lett. B 140 (1984)
44; E D Copeland, A Liddle, D Lyth, E Stewart and D Wands, Phys. Rev. D 49
(1994) 6410; E D Stewart, Phys. Rev. D 51 (1995) 6847.
- [5] C Kolda and J March-Russell, hep-ph/9802358.
- [6] E Halvø, Phys. Lett. B 387 (1996) 43; P Binetruy and G Dvali, Phys. Lett. B 388
(1996) 241.
- [7] J McDonald, hep-ph/9901453.
- [8] M S Turner, Phys. Rev. D 28 (1983) 1243.
- [9] G F Smoot et al., Ap. J. 396 (1992) L1; A Stebbins, astro-ph/9705178.
- [10] D Lyth and A Riotto, Phys. Lett. B 412 (1997) 28.

- [1] S Sarkar, Rep. Prog. Phys. 59 (1996) 1493.
- [2] K Enqvist and JM Donald, hep-ph/9811412.
- [3] K Enqvist and JM Donald, Phys. Rev. Lett. 81 (1998) 3071.