Non-leptonic Two-meson Decays of *B*-Mesons in the Covariant Oscillator Quark Model with Factorization Ansatz

Rukmani MOHANTA, Anjan K. GIRI, Mohinder P. KHANNA Muneyuki ISHIDA,* Shin ISHIDA** and Masuho ODA***

Department of Physics, Panjab University, Chandigarh - 160014, India *Department of Physics, Tokyo Institute of Technology Tokyo 152-8551, Japan

**Atomic Energy Research Institute, College of Science and Technology Nihon University, Tokyo 101-0062, Japan

*** Faculty of Engineering, Kokushikan University, Tokyo 154-8515, Japan

(Received November 20, 1998)

Exclusive nonleptonic bottom meson decays are studied in the covariant osillator quark model using the factorization assumption. The main feature of this model is that it can simultaniously be applied to both heavy \rightarrow heavy and heavy \rightarrow light transitions, satisfying the constraints of the heavy quark effective theory (HQET) in the appropriate limit. The results obtained are in overall agreement with the present experimental data for various *B* decays.

§1. Introduction

The description of exclusive nonleptonic decays of B mesons represents an important and complicated theoretical problem. These decays are nonperturbative in nature and cannot be calculated reliably from the QCD Lagrangian. In contrast to exclusive semileptonic decays, where the weak current matrix elements between meson states are involved, nonleptonic decays require the evaluation of hadronic matrix elements of four local quark operators. To simplify the analysis it is usually assumed that the matrix element of the current-current weak interaction factorizes into the product of two single current matrix elements. Thus the problem reduces to the calculation of the meson form factors, which are contained in the hadronic matrix elements of weak currents as in the case of semileptonic decays, and of the meson decay constants, describing the leptonic decays. This makes the factorization hypothesis¹⁾ a very appealing assumption. Although it is very difficult to prove the factorization hypothesis theoretically within our present understanding of QCD nonperturbative effects, this hypothesis is expected to be valid to a rather good approximation in the case of transitions with large energy release, such as heavy B decays, since the final mesons carrying large momenta escape from the region of interaction, thereby minimizing the effects of a final state interaction. Several tests have been made to prove its validity phenomenologically,²⁾ and, it has been shown to work well for the description of B meson decays into a D or D^* and a light meson. Once the factorization assumption is made, nonleptonic decays are related to the corresponding semileptonic decays.

2 R. Mohanta, A. K. Giri, M. P. Khanna, M. Y. Ishida, S. Ishida and M. Oda

In this paper we wish to calculate the branching ratios of the exclusive nonleptonic two-meson decays of \bar{B}^0 , B^- and \bar{B}_s mesons in the framework of the covariant oscillator quark model (COQM)³⁾ on the basis of factorization. One of the most important motivations of the covariant oscillator quark model (COQM) is to describe covariantly the centre of mass motion of hadrons, retaining the considerable sucesses of the non-relativistic quark model with regard to the static properties of hadrons. A key element of the COQM for achieving this is its direct treatment of the squared masses of hadrons, in contrast to the mass itself as done in conventional approaches. This makes the covariant treatment simple. The COQM has been applied to various problems⁴⁾ with satisfactory results. Recently, Ishida et al have studied the semileptonic weak decays of heavy hadrons using this model⁵⁾ and derived the same relations of weak form factors as in HQET.⁶⁾ Furthermore, the predicted spectra for $B \to (D^*, D) l \nu$ were shown to fit experimental data quite well.⁷⁾ Keeping this success in mind, we extend application of the COQM to the nonleptonic decays of \bar{B}^0 , B^- and \bar{B}_s mesons.

This paper is organized as follows. In §2 we present the expressions for nonleptonic decay amplitudes in the factorization approximation. In §3 we present a brief description of the covariant oscillator quark model. Using this model we have evaluated the form factors and obtained the decay rates for various nonleptonic decay processes. The decay modes $B \rightarrow D^*V$ are considered in §4. Section 5 contains our results and discussion.

§2. General formalism

Neglecting the penguin contribution, the effective Hamiltonian describing the decays under consideration is given by

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{cb} V_{q_i q_j} \left[C_1(m_b) O_1 + C_2(m_b) O_2 \right]$$
(2.1)

with

$$O_1 = (\bar{q}_i q_j)^{\mu} (\bar{c}b)_{\mu}$$
 and $O_2 = (\bar{q}_i b)^{\mu} (\bar{c}q_j)_{\mu}$, (2.2)

where C_1 and C_2 are the Wilson coefficients, and the quark current $(\bar{q}_i q_j)_{\mu}$ denotes the usual (V-A) current. q_i and q_j are two types of quark flavors that are hadronized to the P or V mesons.

The factorization approach to two-body nonleptonic decays $B \rightarrow DM$ implies that the decay amplitude can be expressed by the product of one particle matrix elements:

$$\langle DM | \mathcal{H}_{\text{eff}} | B \rangle = \frac{G_F}{\sqrt{2}} V_{cb} V_{q_i q_j} \left[a_1 \langle D | (\bar{c}b)_\mu | B \rangle \langle M | (\bar{q}_i q_j)^\mu | 0 \rangle \right.$$
$$\left. + a_2 \langle M | (\bar{q}_i b)_\mu | B \rangle \langle D | (\bar{c}q_j)^\mu | 0 \rangle \right] .$$
(2.3)

Here $a_1 = C_1 + C_2/N_c$ and $a_2 = C_2 + C_1/N_c$, where N_c represents the number of colors.

It should be noted that in writing Eq. (2.3) we have discarded the contribution of color octet currents which emerges after the Fierz rearrangement of color singlet operators. Clearly these currents violate factorization since they cannot allow transition to the vacuum states. In the factorization approximation one can distinguish^{*}) three classes of B meson decays : the 'class I' transitions such as $\bar{B}^0 \to M_1^+ M_2^-$, where only the term a_1 contributes (both mesons produced by charged currents); 'class II' transitions, such as $\bar{B}^0 \to M_1^0 M_2^0$, where only the term a_2 contributes (both mesons are produced by neutral currents); and 'class III' transitions, such as $B^- \to M_1^0 M_2^-$, where both terms contribute coherently.

In order to evaluate the transition amplitudes we use the following matrix elements:

$$\langle P(p) | (\bar{q}_i q_j)^{\mu} | 0 \rangle = -i f_P \ p^{\mu}, \langle V(p, \epsilon) | (\bar{q}_i q_j)^{\mu} | 0 \rangle = M_V \ f_V \ \epsilon^{\mu}, \langle a_1(p, \epsilon) | (\bar{q}_i q_j)^{\mu} | 0 \rangle = M_{a_1} \ f_{a_1} \ \epsilon^{\mu},$$
(2.4)

where P, V and a_1 represent the pseudoscalar, vector and the axial vector mesons, respectively. To evaluate the hadronic form factors we use the COQM. These are presented in the next section.

§3. Model Framework, hadronic form factors and decay width of $B \rightarrow PP$, $B \rightarrow PV$ and $B \rightarrow VP$

The general treatment of COQM may be called the "boosted *LS*-coupling scheme," and the wavefunctions being tensors in $\tilde{U}(4) \times O(3, 1)$ -space, reduce to those in $SU(2)_{\rm spin} \times O(3)_{\rm orbit}$ -space in the nonrelativistic quark model in the hadron rest frame. The spinor and space-time portion of the wave functions separately satisfy the respective covariant equations, the Bargmann-Wigner (BW) equation for the former and the covariant oscillator equation for the latter. The form of the meson wave function has been determined completely through the analysis of mass spectra.

In COQM, the meson states are described by bi-local fields $\Phi_A^B(x_{1\mu}, x_{2\mu})$, where $x_{1\mu}(x_{2\mu})$ is the space time coordinate of the constituent quark (antiquark), $A = (a, \alpha)$ $(B = (b, \beta))$ describing its flavor and covariant spinor. Here we write only the positive frequency part of the relevant ground state fields:

$$\Phi_A^B(x_{1\mu}, x_{2\mu}) = e^{iP \cdot X} U(P)_A^B f_{ab}(x_{\mu}; P) , \qquad (3.1)$$

where U and f are the covariant spinor and internal space-time wave functions respectively, satisfying the Bargmann-Wigner and oscillator wave equations. The quantity $x_{\mu}(X_{\mu})$ is the relative (CM) coordinate, $x_{\mu} \equiv x_{1\mu} - x_{2\mu}(X_{\mu} \equiv m_1 x_{1\mu} + m_2 x_{2\mu})/(m_1 + m_2)$, where the m_i represent the quark masses). The function U is

^{*)} The contributions due to quark annihilation processes, which are expected to be small, ¹⁾ are neglected in the calculation of this paper.

4 R. Mohanta, A. K. Giri, M. P. Khanna, M. Y. Ishida, S. Ishida and M. Oda

given by

$$U(P) = \frac{1}{2\sqrt{2}} \left[(-\gamma_5 P_s(v) + i\gamma_\mu V_\mu(v))(1 + iv \cdot \gamma) \right], \qquad (3.2)$$

where $P_s(V_s)$ represents the pseudoscalar (vector) meson field, and $v_{\mu} \equiv P_{\mu}/M$ $[P_{\mu}(M)$ is the four momentum (mass) of the meson]. The function U, being represented by the direct product of quark and antiquark Dirac spinors with the meson velocity, is reduced to the non-relativistic Pauli-spin function in the meson rest frame. The function f is given^{*)} by

$$f(x_{\mu}; P) = \frac{\beta}{\pi} \exp\left(-\frac{\beta}{2}\left(x_{\mu}^2 + 2\frac{(x \cdot P)^2}{M^2}\right)\right) . \tag{3.3}$$

The value of the parameter β is determined from the mass spectra⁸⁾ as $\beta_{\pi/\rho/a_1} = 0.14$, $\beta_{K/K^*} = 0.142$, $\beta_{D/D^*} = 0.148$, $\beta_{D_s} = 0.154$, $\beta_B = 0.151$ and $\beta_{B_s} = 0.160$ (in units of GeV²).

The effective action for weak interactions of mesons with W-bosons is given by

$$S_W = \int d^4 x_1 d^4 x_2 \langle \bar{\Phi}_{F,P'}(x_1, x_2) i \gamma_\mu (1 + \gamma_5) \Phi_{I,P}(x_1, x_2) \rangle W_{\mu,q}(x_1) , \qquad (3.4)$$

where we have denoted the interacting (spectator) quarks as 1 (2). The CKM matrix elements and the coupling constant are omitted. This equation is obtained from consideration of Lorentz covariance, assuming a quark current with the standard V - A form. In Eq. (3.4), $\Phi_{I,P}$ ($\bar{\Phi}_{F,P'}$) denotes the initial (final) meson with definite four momentum $P_{\mu}(P'_{\mu})$, and q_{μ} is the momentum of W boson. The function $\bar{\Phi}$ is defined by $\bar{\Phi} = -\gamma_4 \Phi^{\dagger} \gamma_4$, and $\langle \rangle$ represents the trace of Dirac spinor indices. Our relevant effective current $J_{\mu}(X)_{P',P}$ is obtained by identifying the above equation with

$$S_W = \int d^4 X J_\mu(X)_{P',P} W_\mu(X)_q .$$
 (3.5)

Then $J_{\mu}(X=0)_{P',P} \equiv J_{\mu}$ is explicitly given as ⁹⁾

$$J_{\mu} = I^{qb}(w)\sqrt{MM'} \times \left[\bar{P}_{s}(v')P_{s}(v)(v+v')_{\mu} + \bar{V}_{\lambda}(v')P_{s}(v)(\epsilon_{\mu\lambda\alpha\beta}v'_{\alpha}v_{\beta} - \delta_{\lambda\mu}(w+1) - v_{\lambda}v'_{\mu}\right], \quad (3.6)$$

where M(M') denotes the physical masses of the initial (final) mesons. It should be noted that in the pure confining limit, the masses of the ground state mesons are equal to the simple sums of their constituents, which are much different from the physical masses in the case of light quark pseudoscalar mesons, such as π and K. Therefore we do not consider the transitions $B \to \pi$ and $B \to K$ in our analysis as the reliability of the results is less for these transitions. The quantity $I^{qb}(w)$, which is

^{*)} In this paper we employ the pure-confining approximation, neglecting the effect of the onegluon-exchange potential, which is expected to be good for the heavy/light-quark meson system.

the overlapping of the initial and final wave functions, represents the universal form factor. It describes the confined effects of quarks and is given by

$$I^{qb}(w) = \frac{4\beta\beta'}{\beta + \beta'} \frac{1}{\sqrt{C(w)}} \exp(-G(w)); \qquad C(w) = (\beta - \beta')^2 + 4\beta\beta' w^2, \quad (3.7)$$

and

$$G(w) = \frac{m_n^2}{2C(w)} \left[\left(\beta + \beta'\right) \left\{ \left(\frac{M}{M_s}\right)^2 + \left(\frac{M'}{M_s'}\right)^2 - 2\frac{MM'}{M_sM_s'}w \right\} + 2\left\{ \beta'\left(\frac{M}{M_s}\right)^2 + \beta\left(\frac{M'}{M_s'}\right)^2 \right\} (w^2 - 1) \right], \qquad (3.8)$$

where $M_s(M'_s)$ represents the sum of the constituent quark masses of the initial (final) meson, and m_n is the spectator quark mass.

The form factor function $I^{cb}(w)$ for $B \to D(D^*)$ decays corresponds to the Isgur-Wise function $\xi(w)$ in HQET.⁶⁾ At the zero recoil point w = 1, the value of $I^{cb}(w)$ is given by

$$I^{cb}(w=1) = \frac{4\beta\beta'}{(\beta+\beta')^2} .$$
 (3.9)

In the heavy quark symmetry limit $\beta = \beta'$, so Eq. (3.9) correctly reproduces⁷ the normalization condition of HQET, i.e., $\xi(w = 1) = 1$. However, HQET, as it is, predicts nothing about the Isgur-Wise function except for the zero recoil point, while in COQM the form factor functions can be derived at any kinematical point of interest due to the fact that the center of mass motion of the meson there is treated covariantly, as was mentioned in §1. In addition, the COQM form factor $I^{qb}(w)$ is also applicable for the heavy-to-light transition processes, while HQET does not provide anything for this sector.

After obtaining the effective current in the COQM, the decay widths for various $B \to D$ and $B \to D^*$ decay modes can be obtained with Eqs. (2·3), (2·4) and (3·6). These are as follows:

$$\Gamma(B(v) \to D(v_1)P(v_2)) = \frac{G_F^2}{16\pi M_B^2} |V_{cb}V_{q_iq_j}|^2 |\mathbf{p}| \\
\times \left[a_1 f_P \sqrt{M_B M_D} (I_q^{cb}(w_1))(1+w_1) (M_B - M_D) \right. \\
+ a_2 f_D \sqrt{M_B M_P} (I_q^{qb}(w_2))(1+w_2) (M_B - M_P) \right]^2, \quad (3.10)$$

$$\Gamma(B(v) \to D^*(v_1)P(v_2)) = \frac{G_F^2}{16\pi M_B^2} |V_{cb}V_{q_iq_j}|^2 |\mathbf{p}|^3$$

6 R. Mohanta, A. K. Giri, M. P. Khanna, M. Y. Ishida, S. Ishida and M. Oda

$$\times \left[a_1 f_P \sqrt{\frac{M_B}{M_{D^*}}} (I_q^{cb}(w_1)) (M_B + M_{D^*}) + a_2 f_{D^*} \sqrt{\frac{M_B}{M_P}} (I_q^{qb}(w_2)) (M_B + M_P)\right]^2, \qquad (3.11)$$

$$\Gamma(B(v) \to D(v_1)V(v_2)) = \frac{G_F^2}{16\pi M_B^2} |V_{cb}V_{q_iq_j}|^2 |\mathbf{p}|^3 \\
\times \left[a_1 f_V \sqrt{\frac{M_B}{M_D}} (I_q^{cb}(w_1)) (M_B + M_D) \\
+ a_2 f_D \sqrt{\frac{M_B}{M_V}} (I_q^{qb}(w_2)) (M_B + M_V) \right]^2, \quad (3.12)$$

where we have taken $w_1 = v \cdot v_1$ and $w_2 = v \cdot v_2$. Here $|\mathbf{p}|$ is the c.m. momentum of the emitted particles in the rest frame of the *B* meson. As stated earlier, only the a_1 term contributes to class I decays, and only the a_2 term contributes to class II decays, while both a_1 and a_2 terms contribute coherently to class III decays.

The COQM is also applicable to heavy to light transitions, as well as heavy to heavy transitions, such as $B \to \rho$ and $B \to K^*$. In this case the above formulas are changed by replacing $(D^*(v_1), I_q^{cb}(w_1), V_{cb})$ in $B \to D^*$ transition with $(\rho(v_1), I_q^{nb}(w_1), V_{ub})$ and $(K^*(v_1), I_q^{sb}(w_1), V_{ub})$ in $B \to \rho$ and $B \to K^*$ transitions, respectively.

§4. Decay rate, polarization and angular correlation in the decays $B \rightarrow VV$

The helicity amplitude for the decay process $B(p) \to D^*(k_1, \epsilon_1)V(k_2, \epsilon_2)$ can be expressed by three invariant amplitudes, a, b and c. It is given following Ref. 10) as

$$H_{\lambda} = \epsilon_{1\mu}^{(\lambda)*} \epsilon_{2\nu}^{(\lambda)*} \left[a g^{\mu\nu} + \frac{b}{M_{D^*} M_V} p^{\mu} p^{\nu} + \frac{ic}{M_{D^*} M_V} \epsilon^{\mu\nu\alpha\beta} k_{1\alpha} p_{\beta} \right] . \tag{4.1}$$

The coefficients a, b and c describe the S-, P- and D- wave contributions to the two final vector particles. In the present framework these are given as

$$a = \frac{G_F}{\sqrt{2}} V_{cb} V_{q_i q_j} [a_1 f_V M_V \sqrt{M_B M_{D^*}} I^{cb}(w_1)(1+w_1) + a_2 f_{D^*} M_{D^*} \sqrt{M_B M_V} I^{qb}(w_2)(1+w_2)],$$

$$b = -\frac{G_F}{\sqrt{2}} V_{cb} V_{q_i q_j} \left[a_1 f_V M_V^2 \sqrt{\frac{M_{D^*}}{M_B}} I^{cb}(w_1) + a_2 f_{D^*} M_{D^*}^2 \sqrt{\frac{M_V}{M_B}} I^{qb}(w_2) \right],$$

$$c = -\frac{G_F}{\sqrt{2}} V_{cb} V_{q_i q_j} \left[a_1 f_V M_V^2 \sqrt{\frac{M_{D^*}}{M_B}} I^{cb}(w_1) + a_2 f_{D^*} M_{D^*}^2 \sqrt{\frac{M_V}{M_B}} I^{qb}(w_2) \right]. (4.2)$$

The helicity amplitudes are given as

$$H_{\pm 1} = a \pm \sqrt{x^2 - 1} c$$
 and $H_0 = -ax - b(x^2 - 1)$, (4.3)

where x is defined by

$$x \equiv \frac{k_1 \cdot k_2}{M_{D^*} M_V} = \frac{M_B^2 - M_{D^*}^2 - M_V^2}{2M_{D^*} M_V}$$
(4.4)

and obeys

$$x^{2} = 1 + \frac{M_{B}^{2} |\mathbf{p}|^{2}}{M_{D^{*}}^{2} M_{V}^{2}} .$$
(4.5)

The corresponding decay rate can be obtained as

$$\Gamma(B \to D^*V) = \frac{|\mathbf{p}|}{8\pi M_B^2} \left[2|a|^2 + |xa + (x^2 - 1)b|^2 + 2(x^2 - 1)|c|^2 \right].$$
(4.6)

The decay distribution is parametrized by the coefficients

$$\frac{\Gamma_T}{\Gamma} = \frac{|H_{+1}|^2 + |H_{-1}|^2}{|H_0|^2 + |H_{+1}|^2 + |H_{-1}|^2}, \qquad \qquad \frac{\Gamma_L}{\Gamma} = \frac{|H_0|^2}{|H_0|^2 + |H_{+1}|^2 + |H_{-1}|^2},
\alpha_1 = \frac{\operatorname{Re}(H_{+1}H_0^* + H_{-1}H_0^*)}{|H_0|^2 + |H_{+1}|^2 + |H_{-1}|^2}, \qquad \qquad \beta_1 = \frac{\operatorname{Im}(H_{+1}H_0^* - H_{-1}H_0^*)}{|H_0|^2 + |H_{+1}|^2 + |H_{-1}|^2},
\alpha_2 = \frac{\operatorname{Re}(H_{+1}H_{-1}^*)}{|H_0|^2 + |H_{+1}|^2 + |H_{-1}|^2}, \qquad \qquad \beta_2 = \frac{\operatorname{Im}(H_{+1}H_{-1}^*)}{|H_0|^2 + |H_{+1}|^2 + |H_{-1}|^2}. \quad (4.7)$$

In general, the dominant terms in the angular correlations are Γ_T/Γ , Γ_L/Γ , α_1 and α_2 . The terms β_1 and β_2 are small since they are nonvanishing only if the helicity amplitudes H_{+1} , H_{-1} and H_0 or the invariant amplitudes a, b and c, respectively, have different phases.

In the case of heavy to light transitions, $B \to \rho$ and $B \to K^*$, the corresponding formulas are obtained by the procedure explained as the end of the last section.

§5. Results and conclusion

In order to make the numerical estimate, we use the following values of various quantities. The quark masses (in GeV) are taken as $m_u = m_d = 0.4$, $m_s = 0.55$, $m_c = 1.7$ and $m_b = 5$.⁸⁾ The particle masses and lifetimes are taken from Ref. 15). The relevant CKM parameter values used are $V_{cb} = 0.0395$, $V_{cs} = 1.04$, $V_{cd} = 0.224$, $V_{ud} = 0.974$, $V_{us} = 0.2196$ and $V_{ub} = 0.08 \times V_{cb} = 0.00316$. The decay constants are taken as $f_{\pi} = 130.7$, $f_K = 159.8$; $f_{K^*} = 214$;¹¹⁾ $f_{\rho} = 210$;¹²⁾ $f_D = 220$, $f_{D^*} = 230$, $f_{D_s} = 240$, $f_{D_s^*} = 260^{13)}$ and $f_{a_1} = 205^{14)}$ (in MeV). The decay constant $f_{J/\psi}$ is determined from the value of $\Gamma(J/\psi \to e^+e^-)$:¹⁵⁾

$$f_{J/\psi} = \sqrt{\frac{9}{4}} \left(\frac{3}{4\pi\alpha^2}\right) \Gamma(J/\psi \to e^+ e^-) M_{J/\psi} = 404.5 \text{ MeV}.$$
 (5.1)

Table I. Branching ratios of nonleptonic \bar{B}^0 decays in the COQM. Note that the values do not include the possible contribution from the penguin diagram, which is generally expected to be of order $10^{-6} \sim 10^{-7}$. It is shown that the penguin diagram does not contribute to $\rho^+ D_s^-, \rho^+ D_s^{*-}, \bar{K}^{*0} \bar{D}^0$ and $\bar{K}^{*0} \bar{D}^{*0}$. The contribution to $\rho^+ \pi^-, \rho^+ \rho^-$ and $\rho^+ a_1^-$ is about 3×10^{-7} .¹⁾

\bar{B}^0 modes	This work	Expt. [15]
Class I		
$D^{+}\pi^{-}$	3.00×10^{-3}	$(3.0 \pm 0.4) \times 10^{-3}$
D^+K^-	2.28×10^{-4}	-
D^+D^-	4.345×10^{-4}	-
D^+D^s	10.9×10^{-3}	$(8.0 \pm 3.0) \times 10^{-3}$
$D^+ \rho^-$	7.406×10^{-3}	$(7.9 \pm 1.4) \times 10^{-3}$
$D^{+}K^{*-}$	3.84×10^{-4}	-
$D^{+}D^{*-}$	3.29×10^{-4}	$< 1.2 \times 10^{-3}$
$D^{+}D_{s}^{*-}$	0.86×10^{-2}	$(1.0 \pm 0.5) \times 10^{-2}$
$D^{+}a_{1}^{-}$	6.51×10^{-3}	$(6.0 \pm 3.3) \times 10^{-3}$
$D^{*+}\pi^{-}$	3.07×10^{-3}	$(2.76 \pm 0.21) \times 10^{-3}$
$D^{*+}K^{-}$	2.28×10^{-4}	-
$D^{*+}D^{-}$	3.14×10^{-4}	$< 1.8 \times 10^{-3}$
$D^{*+}D_{s}^{-}$	7.615×10^{-3}	$(9.6 \pm 3.4) \times 10^{-3}$
$D^{*+}\rho^{-}$	8.91×10^{-3}	$(6.7 \pm 3.3) \times 10^{-3}$
$D^{*+}K^{*-}$	4.89×10^{-4}	-
$D^{*+}D^{*-}$	8.74×10^{-4}	$< 2.2 \times 10^{-3}$
$D^{*+}D_{s}^{*-}$	2.49×10^{-2}	$(2.0 \pm 0.7) \times 10^{-2}$
$D^{*+}a_{1}^{-}$	0.99×10^{-2}	$(1.30 \pm 0.27) \times 10^{-2}$
$\rho^+ D_s^-$	2.17×10^{-5}	$< 7 \times 10^{-4}$
$\rho^+ D_s^{*-}$	4.63×10^{-5}	$< 8 \times 10^{-4}$
$ ho^+\pi^-$	6.53×10^{-6}	$< 8.8 \times 10^{-5}$
$ ho^+ ho^-$	1.83×10^{-5}	$< 2.2 \times 10^{-3}$
$ ho^+a_1^-$	1.94×10^{-5}	$< 3.4 \times 10^{-3}$
Class II		
$D^0 ho^0$	0.649×10^{-4}	$< 3.9 \times 10^{-4}$
$D^{*0} ho^0$	1.22×10^{-4}	$< 5.6 \times 10^{-4}$
$ar{K}^{*0}J/\psi$	1.504×10^{-3}	$(1.35 \pm 0.18) \times 10^{-3}$
$ar{K}^{*0}ar{D}^0$	1.16×10^{-6}	_
$ar{K}^{*0}ar{D}^{*0}$	2.25×10^{-6}	_

The parameters a_1 and a_2 appearing in these decays, which have recently been determined from the CLEO data, ¹⁶⁾ are $a_1 = 1.02$ and $a_2 = 0.23$.¹⁷⁾ Using these values we obtain the branching ratios for \bar{B}^0 , B^- and \bar{B}_s mesons, which are tabulated in Tables I, II and III, respectively. The overall agreement between the predicted and experimental data is quite remarkable. Here, it may be worthwhile to note that the relevant processes are relativistic and the form factor functions $I(\omega)$ play significant roles: For example, the velocity of the final D^+ in the process $\bar{B}^0 \to D^+\pi^-(\bar{B}^0 \to$ $D^+D_s^{*-})$ is $v_D = 0.78c(0.70c)$, and the value of I is I = 0.53(0.65). The polarization and the angular distribution parameters for $B \to D^*V$ decay modes are presented in Table IV. The polarization fraction (Γ_L/Γ) for $\bar{B}^0 \to D^{*+}\rho^-$ (0.883) agrees well with experimental value $0.93 \pm 0.05 \pm 0.05$.¹⁵⁾ For the decay mode $\bar{B}^0 \to \bar{K}^{*0}J/\psi$, Γ_L/Γ (0.428) also agrees with the recent CLEO data $0.52 \pm 0.07 \pm 0.04$.¹⁸⁾ Table II. Branching ratios of nonleptonic B^- decays in the COQM. The numbers for $D^0 a_1^-$, $D^{*0} a_1^-, D^0 \pi^-, D^{*0} \pi^-, D^0 K^-, D^{*0} K^-$, $\rho^0 \pi^-$ and $\rho^0 a_1^-$ in class III do not contain the contributions from the color-suppressed a_2 -term in Eq. (2.3). Note also that the values do not include the possible contribution from the penguin diagram, which are generally expected to be of order $10^{-6} \sim 10^{-7}$. It is shown that the penguin diagram does not contribute to $\rho^0 D_s^-, \rho^0 D_s^{*-}, \rho^- J/\psi, K^{*-} \bar{D}^0$ and $K^{*-} \bar{D}^{*0}$. The contribution to $\rho^0 \pi^-, \rho^0 \rho^-$ and $\rho^0 a_1^-$ is about 3×10^{-7} .¹

B^- modes	This work	Expt. [15]
Class I		
D^0D^-	4.53×10^{-3}	-
$D^0 D_s^-$	1.16×10^{-2}	$(1.3 \pm 0.4) \times 10^{-2}$
$D^{0}D^{*-}$	3.48×10^{-4}	-
$D^{0}D_{s}^{*-}$	9.07×10^{-3}	$(9.0 \pm 4.0) \times 10^{-3}$
$D^{*0}D^{-}$	3.34×10^{-4}	-
$D^{*0}D_{s}^{-}$	0.81×10^{-2}	$(1.2 \pm 0.5) \times 10^{-2}$
$D^{*0}D^{*-}$	9.23×10^{-4}	-
$D^{*0}D_{s}^{*-}$	2.625×10^{-2}	$(2.7 \pm 1.0) \times 10^{-2}$
$ ho^0 D_s^-$	1.15×10^{-5}	$< 4 \times 10^{-4}$
$ ho^0 D_s^{*-}$	2.45×10^{-5}	$< 5 \times 10^{-4}$
Class II		
$K^{*-}J/\psi$	1.587×10^{-3}	$(1.47 \pm 0.27) \times 10^{-3}$
$ ho^- J/\psi$	5.79×10^{-5}	$< 7.7 \times 10^{-4}$
$K^{*-}\bar{D}^0$	1.23×10^{-6}	_
$K^{*-}\bar{D}^{*0}$	2.38×10^{-6}	—
Class III		
$D^0 \rho^-$	1.004×10^{-2}	$(1.34 \pm 0.18) \times 10^{-2}$
$D^{*0}\rho^-$	1.264×10^{-2}	$(1.55 \pm 0.31) \times 10^{-2}$
$D^{0}a_{1}^{-}$	6.89×10^{-3}	$(5.0 \pm 4.0) \times 10^{-3}$
$D^{*0}a_{1}^{-}$	1.04×10^{-2}	$(1.9 \pm 0.5) \times 10^{-2}$
$D^0\pi^-$	3.18×10^{-3}	$(5.3 \pm 0.5) \times 10^{-3}$
$D^{*0}\pi^-$	3.25×10^{-3}	$(4.6 \pm 0.4) \times 10^{-3}$
D^0K^-	2.41×10^{-4}	_
$D^{0}K^{*-}$	5.34×10^{-4}	_
$D^{*0}K^{-}$	2.42×10^{-4}	-
$D^{*0}K^{*-}$	7.13×10^{-4}	_
$ ho^0\pi^-$	3.45×10^{-6}	$< 4.3 \times 10^{-5}$
$\rho^0 a_1^-$	1.03×10^{-5}	$< 6.2 \times 10^{-4}$
$ ho^0 ho^-$	1.45×10^{-5}	$< 1.0 \times 10^{-3}$

In this paper we have calculated the branching ratios of the exclusive nonleptonic two-meson decay of B mesons using the covariant oscillator quark model based on the factorization approximation. The applied form factors are consistent with the predictions of heavy quark symmetry. The overall agreement of our predictions for two meson nonleptonic decays of B mesons with the existing experimental data suggests that the factorization approximation works well and the estimation of confinement effects by the COQM is valid.

$\bar{B}_s^0 \mod$	This work	Expt. $[15]$
$D_s^+\pi^-$	0.248×10^{-2}	< 13%
$D_s^+K^-$	$1.89 imes 10^{-4}$	-
$D_s^+ D^-$	3.73×10^{-4}	-
$D_s^+ D_s^-$	9.57×10^{-3}	-
$D_s^+ \rho^-$	6.15×10^{-3}	-
$D_{s}^{+}K^{*-}$	3.196×10^{-4}	-
$D_{s}^{+}D^{*-}$	2.86×10^{-4}	-
$D_{s}^{+}D_{s}^{*-}$	7.48×10^{-3}	-
$D_{s}^{+}a_{1}^{-}$	4.86×10^{-3}	-
$D_{s}^{*+}\pi^{-}$	2.55×10^{-3}	-
$D_s^{*+}K^-$	1.9×10^{-4}	-
$D_{s}^{*+}D^{-}$	2.735×10^{-4}	-
$D_{s}^{*+}D_{s}^{-}$	6.68×10^{-3}	-
$D_s^{*+}\rho^-$	7.49×10^{-3}	-
$D_{s}^{*+}K^{*-}$	4.1×10^{-4}	-
$D_{s}^{*+}D^{*-}$	$6.85 imes 10^{-4}$	-
$D_{s}^{*+}D_{s}^{*-}$	2.216×10^{-2}	-
$D_{s}^{*+}a_{1}^{-}$	8.39×10^{-3}	-

Table III. Branching ratios of nonleptonic \bar{B}_s^0 decays in the COQM model.

Table IV. Polarization and angular correlation parameter of $B \to D^* V$ decays.

Decay modes	Γ_L/Γ	α_1	α_2
$\bar{B}^0 \to D^{*+} \rho^-$	0.883	-0.416	0.04
$\bar{B}^0 \to D^{*+} D^{*-}$	0.538	-0.662	0.176
$\bar{B}^0 \to D^{*+} D_s^{*-}$	0.515	-0.665	0.187
$\bar{B}^0 \to \bar{K}^{*0} J/\psi$	0.428	-0.605	0.142
$B^- \to D^{*0} \rho^-$	0.855	-0.446	0.044
$B^- \to D^{*0} D^{*-}$	0.538	-0.661	0.176
$B^- \to D^{*0} D_s^{*-}$	0.515	-0.665	0.187
$B^- \to K^{*-} J/\psi$	0.428	-0.605	0.141

Acknowledgements

R. M. would like to thank CSIR, the Govt. of India, for a fellowship. A. K. G. and M. P. K. acknowledge the financial support from DST, the Govt. of India. The authors are grateful to Dr. K. Yamada for informing them of the recent values of β obtained from the analyses of mass spectra.

References

- 1) M. Bauer, B. Stech and M Wirbel, Z. Phys. C34(1987), 103.
- 2) J. L. Rosner, Phys. Rev. D42 (1990), 3732.
 - M. Tanimato, K. Goda and K. Senba, Phys. Rev. D42 (1990), 3741.
 - A. Acker, S. Pakvasa, S. F. Tuan and S. P. Rosen, Phys. Rev. D43 (1991), 3083.
 - T. Mannel, W. Roberts and Z. Ryzak, Phys. Lett. **B259** (1991), 359.
 - M. J. Dugan and B. Grinstein, Phys. Lett. B255 (1991), 583.
 - A. Deandrea, N. Di Bartolomeo, R. Gatto and G. Nardulli, Phys. Lett. B318 (1993), 549.
 - D. Bortoletto and S. Stone, Phys. Rev. Lett. 65 (1990), 2951.
 - M. Neubert, V. Rieckert, B. Stech and Q. P. Xu, in *Heavy Flavors*, ed. A. J. Buras and M. Linder (World Scientific, Singapore, 1992).

- A. N. Kamal, Q. P. Xu and A. Czarnecki, Phys. Rev. D49 (1994), 1330.
- 3) H. Yukawa, Phys. Rev. 77 (1950), 219; 91 (1953), 415, 416.
 S. Ishida, Prog. Theor. Phys. 45 (1971), 1570, 1905.
 R. P. Feynmann, M. Kislinger and F. Ravndal, Phys. Rev. D3 (1971), 2706.
 S. Ishida, M. Y. Ishida and M. Oda, Prog. Theor. Phys. 93 (1995), 939.
 The COQM has a long history of development since the bilocal theory by H. Yukawa. On the preceding works see, the references given in the above last paper.
- S. Ishida et al., Prog. Theor. Phys. **71** (1984), 806.
 S. Ishida, K. Yamada and M. Oda, Phys. Rev. **D28** (1983), 2918.
 S. Ishida et al., Phys. Rev. **D20** (1979), 2906.
 - S. Ishida and K. Yamada, Phys. Rev. **D35** (1987), 265.
 - S. Ishida, A. Morikawa and M. Oda, Prog. Theor. Phys. **99** (1998), 257.
- 5) M. Oda, M. Y. Ishida and S. Ishida, Nihon Univ. Preprint NUP-A-94-7.
- N. Isgur and M. B. Wise, Phys. Lett. B232 (1990), 113; B237 (1990), 527; Nucl. Phys. B348 (1991), 276.
 - A. F. Falk, H. Georgi, B. Grinstein and M. B. Wise, Nucl. Phys. B343 (1991), 1.
 H. Georgi, Phys. Lett. B240 (1990), 447; Nucl. Phys. B348 (1991), 293.
 B. Grinstein, Nucl. Phys. 339 (1990), 253.
 M. Neubert, Phys. Lett. B264 (1991), 455; Phys. Rep. 245 (1994), 259.
- 7) M. Y. Ishida, S. Ishida and M. Oda, Prog. Theor. Phys. 98 (1997), 159.
- 8) S. Ishida and K. Yamada, Phys. Rev D35 (1987), 265.
 S. Ishida and M. Oda, Prog. Theor. Phys. 89 (1993), 1033.
 S. Ishida, M. Y. Ishida and K. Yamada, Prog. Theor. Phys. 91 (1994), 775.
- 9) S. Ishida, M. Y. Ishida and M. Oda, Prog. Theor. Phys. 93 (1995), 781.
- (1) S. Isinda, M. T. Isinda and M. Oda, 1163. Theor. Phys. 55 (1999), 101.
 (10) G. Valencia, Phys. D39 (1989), 3339.
 G. Kramer and W. F. Palmer, Phys. Rev. D45 (1992), 193; Phys. Lett. B279 (1992), 181; Phys. Rev. D46 (1992), 3197.
 E. Golowich and S. Pakvasa, Phys. Rev. D51 (1995), 1215.
- 11) A. Ali and C. Greub, Phys. Rev **D57** (1998), 2996.
- 12) T. Mannel and W. Roberts, Z. Phys. C59 (1993), 179.
- 13) M. Neubert et al., in Ref. 2).
- 14) C. Reader and N. Isgur, Phys. Rev. D47 (1993), 1007.
- 15) C. Caso et al., The Euro. Phys. J. C3 (1998), 1.
- 16) CLEO Collaboration, M. S. Alam et al, Phys. Rev. D50 (1994), 43.
- H. Y. Cheng and B. Tseng, Phys. Rev. D51 (1995), 6259.
 A. J. Buras, Nucl. Phys. B434 (1997), 606.
 M. Ciuchini et al., Nucl. Phys. B415 (1994), 403.
- 18) CLEO Collaboration, C. P. Jessop et al., Phys. Rev. Lett. 79 (1997), 4533.