

# M assive chiral random m atrix ensem bles at = 1 & 4 : Q C D Dirac operator spectra

Taro Nagao

Department of Physics, Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan

Shinsuke M . N ishigaki

Department of Physics, Faculty of Science, Tokyo Institute of Technology, Oh-okayama, Meguro, Tokyo 152-8551, Japan  
(March 1, 2000)

The zero momentum sectors in effective theories of QCD coupled to pseudoreal (two colors) and real (adjoint) quarks have alternative descriptions in terms of chiral orthogonal and symplectic ensembles of random matrices. Using this correspondence, we compute correlation functions of Dirac operator eigenvalues within a sector with an arbitrary topological charge in a presence of finite quark masses of the order of the smallest Dirac eigenvalue. These novel correlation functions, expressed in terms of Pfaans, interpolate between known results for the chiral and quenched limits as quark masses vary.

PACS number(s): 05.40.-a, 12.38.Aw, 12.38.Lg

## I. INTRODUCTION

Random matrix theory of disordered Hamiltonians [1,2] relies upon an ansatz that in the ergodic regime where the system size  $L$  is much larger than the elastic mean free path but much smaller than the localization length, details of the Hamiltonian are lost except for its time-reversal and internal symmetries. This ansatz was materialized by Efetov [3] who derived Wigner-Dyson statistics out of Anderson tight-binding model by retaining only the zero-momentum mode of his spectral nonlinear model (NL M).

By taking the flavor symmetry among quarks into consideration as an additional internal symmetry, Verbaarschot and collaborators [4-6] have reinterpreted this wisdom in the context of quantum chromodynamics (QCD). In this context, the spectral NL M is a supersymmetric extension [7,8] of the conventional NL M over the coset manifold associated with the chiral symmetry breaking [9,10]:

$Z$

$$Z^{(2)}(\beta; M) = \frac{dU}{Z_{SU(N_f)}} \exp(\text{Re} \operatorname{tr} e^{i\beta N_f M} U^Y); \quad (1a)$$

$$Z^{(1)}(\beta; M) = \frac{dU}{Z_{SU(2N_f)=Sp(2N_f)}} \exp(\text{Re} \operatorname{tr} e^{i\beta N_f M} U J U^T = 2); \quad (1b)$$

$$Z^{(4)}(\beta; M) = \frac{dU}{Z_{SU(N_f)=SO(N_f)}} \exp(\text{Re} \operatorname{tr} e^{i\beta (N_c N_f) M} U U^T); \quad (1c)$$

after retaining only the zero mode. Here the superscripts (2, 1, 4) of Dyson indices refer to the anti-unitary symmetry of Euclidean Dirac operators (which are considered as stochastic Hamiltonians) in three classes of QCD [11]:

= 2 :  $N_c = 3$ ;  $N_f$  fundamental Dirac fermions;

= 1 :  $N_c = 2$ ;  $N_f$  fundamental Dirac fermions; (2)

= 4 :  $N_c = 2$ ;  $N_f$  adjoint Majorana fermions:

The rescaled quark masses  $M$  are defined as

$$\begin{aligned} M &= \text{diag}(\lambda_1, \dots, \lambda_{N_f}) & (\lambda = 2, 1, 4); \\ M &= \text{diag}(\lambda_1, \dots, \lambda_{N_f}) J & (\lambda = 1); \\ i & L^4 m_i; J = \mathbb{1}_{N_f} & J; J = \begin{matrix} 0 & 1 \\ 1 & 0 \end{matrix}; \end{aligned} \quad (3)$$

with a limit

$$L \rightarrow 1; m_i \rightarrow 0; \beta_i \text{ fixed} \quad (4)$$

being assumed,  $\beta$  stands for the quark condensate in the chiral limit, and  $\beta_i$  stands for the vacuum angle. By the same token as the spectral NL M of the tight-binding model was derived from a conventional random matrix ensembles [12], these NL Ms have an alternative derivation from chiral random matrix ensembles (RM Es) [4,13]:

$$\begin{aligned} Z^{(\lambda)}(\beta; M) &= \int_0^{\pi} \frac{d\theta}{2} e^{i\theta} Z^{(\lambda)}(\theta; M) \\ &= \frac{1}{dW} \det \frac{W^Y}{\det \frac{m_i}{W^Y m_i}}; \quad (5) \end{aligned}$$

under a limit

$$N \rightarrow 1; m_i \rightarrow 0; \beta_i = (0)m_i \text{ fixed}; \quad (6)$$

Here the integrals are over complex, real, and quaternion real ( $N +$ )  $N$  matrices  $W$  for  $\lambda = 2, 1, 4$ , respectively, and (0) stands for the large- $N$  spectral density of the random matrix  $D = \begin{pmatrix} 0 & W \\ W^Y & 0 \end{pmatrix}$ :

$$(0) = \lim_{N \rightarrow 1} \text{htr}((iD)^{-1}); \quad (7)$$

at the origin. It is understood for  $\lambda = 4$  that two-fold degenerated eigenvalues in the determinant are only

counted once, and the topological charge is substituted by  $N_c$ . These RM Es are motivated by the microscopic theories (Euclidean QCD) on a lattice, with a crude simplification of replacing matrix elements of the antiHermitian Dirac operator  $D = (\partial + iA)$  by random numbers  $D$  generated according to the weight  $e^{-\text{tr} V(W^T W)}$ . Under this correspondence, the microscopic limit (6) is equivalent to Leutwyler-Smilga limit (4), since the size  $N$  of the matrix  $W$  is interpreted as the number of sites  $L^4$  of the lattice on which QCD is discretized, and the Dirac spectral density at zero virtuality

(0) is related to the quark condensate by Banks-Casher relation  $= \langle 0 | L^4 | 0 \rangle = 1$  [14]. Using the RME representation, various correlation functions of microscopically rescaled Dirac eigenvalues

$$N_1 ! ; ! 0 ; = \langle 0 | : \text{qed} : \rangle \quad (8)$$

have been computed for all three values of in the massless case  $= 0$  [15{23]. On the other hand, in a presence of finite 's, Dirac eigenvalues comparable to 's are expected to obey statistics that interpolate the chiral ( $= 0$ ) and quenched limits ( $= 1$  or  $N_f = 0$ ). These novel spectral correlation functions have so far been analytically computed solely for the chiral unitary ( $= 2$ ) ensemble [24{29]. Therefore we aim to treat the remaining cases, chiral orthogonal ( $= 1$ ) and symplectic ( $= 4$ ) ensembles with finite mass parameters, and compute Dirac eigenvalue correlation functions for these ensembles. We anticipate that advances in numerical simulations of lattice QCD with dynamical quarks [30,31] will confirm our analytical results, Eqs.(48) and (53) below, in a foreseeable future.

This Article is organized as follows. In Sect. 2 we compute the correlation functions for the chiral orthogonal ensemble, by utilizing the quaternion determinant method developed in Ref. [32]. In Sect. 3 we exhibit an explicit form of the correlation functions for the chiral symplectic ensemble, which was obtained by the Authors [33] as a corollary to the computation of the partition function. In Appendix A we collect definitions related to a quaternion determinant. In Appendices B and C we present alternative expressions of the quaternion kernels for the orthogonal ensemble and the symplectic ensembles with quadruply degenerated masses, respectively. It enables us to identify our results with those in a paper by Akemann and Kanzeper [34], which appeared after the Letter by the present Authors [33] and computed a few, rather limited cases of the correlation functions than ours.

## II. ORTHOGONAL ENSEMBLE

We start by expressing the partition function (5) of the RME in terms of eigenvalues  $x_i = \sqrt{\lambda_i}$  of the Wishart matrix  $W^T W$  (up to an overall constant):

$$Z^{(0)}(\text{fm } g) = \prod_{i=1}^N m_i \int_0^\infty \frac{1}{N!} dx_i P(\text{fxg}; \text{fm } g); \quad (9)$$

$$P(\text{fxg}; \text{fm } g) = \prod_{j=1}^N e^{-V(x_j)} x_j^{-\frac{j+1}{2}-1} \prod_{i=1}^N (x_j + m_i^2)^{-\frac{1}{2}} \prod_{j>k} x_k x_j : \quad (10)$$

The indices and are suppressed for simplicity. Since the partition function (5) is even under  $\gamma_5$ , we have set non-negative integer, without loss of generality. The p-level correlation function of the Wishart matrix  $W^T W$  is defined as

$$\langle x_1 ; \dots ; x_p ; \text{fm } g \rangle = \frac{\langle p(x_1 ; \dots ; x_p ; \text{fm } g) \rangle}{\langle 0(\text{fm } g) \rangle}; \quad (11)$$

$$\langle p(x_1 ; \dots ; x_p ; \text{fm } g) \rangle = \frac{1}{(N-p)!} \int_0^\infty \prod_{j=p+1}^N dx_j P(\text{fxg}; \text{fm } g); \quad (12)$$

Then the p-level correlation function of the block off-diagonal Hermitian matrix  $iD$ ,

$$\langle \gamma_1 ; \dots ; \gamma_p ; \text{fm } g \rangle = h \prod_{k=1}^p \text{tr} (\gamma_k iD) i; \quad (13)$$

is expressed in terms of multiplied by the Jacobian of the transformation  $\gamma_k x = z^2$ :

$$\langle \gamma_1 ; \dots ; \gamma_p ; \text{fm } g \rangle = 2^p \prod_{j=1}^p \int_{-\infty}^{\infty} dz_j \langle \gamma_1^2 ; \dots ; \gamma_p^2 ; \text{fm } g \rangle; \quad (14)$$

As the universality of correlation functions of the unitary ensemble in the microscopic limit (8) [35{37,26] is known to inherit to those of orthogonal and symplectic ensembles [38,39], it succeeds to concentrate on Laguerre ensembles,  $V(x) = x$ . This leads to Wigner's semicircle law

$$\langle \gamma \rangle = \frac{2^p}{2N} \overline{\int_{-2\sqrt{N}}^{2\sqrt{N}} z^p dz}; \quad (15)$$

Now we concentrate on  $= 1$ , and define new variables  $z_j$  as

$$z_j = \frac{x_j^2 - 1}{x_j} \quad (0); \quad j = 1, \dots, N; \quad (16)$$

Then the multiple integral (12) is expressed as

$$\begin{aligned} \langle p(z_1 ; \dots ; z_{p+1}) \rangle &= \prod_{j=1}^p \int_{-\infty}^{\infty} \frac{1}{w(z_j)} \prod_{j>k} \frac{dz_j}{z_j} \int_{-\infty}^{\infty} \frac{dz_{p+1}}{z_{p+1}} \prod_{j=1}^p \frac{w(z_j)}{z_j} \prod_{j>k} \frac{dz_j}{z_j} \\ &\quad \frac{1}{(N-p)!} \int_0^\infty \prod_{j=p+1}^N dz_j \end{aligned} \quad (17)$$

where

$$w(z) = \frac{1}{2} j^2 e^{2z} : \quad (18)$$

Eq.(17) resembles an  $(+p)$ -level correlation function of the conventional ( $n$  massless) Laguerre ensemble with  $+N$  levels. However, conventionally the levels  $z_1, \dots, z_{+p}$  are all positive, while in the present case some of them ( $z_1, \dots, z_p$ ) are negative. We carefully incorporate this fact into the following evaluation.

Let us denote the integrand in Eq.(17) as

$$p(z_1, \dots, z_p) = \prod_{j=1}^p \frac{q_j}{w(z_j)} \prod_{j>k} j z_j - z_k \quad (19)$$

with  $+N$ . It can be readily seen that

$$p(z_1, \dots, z_p) = \prod_{j=1}^p \frac{q_j}{w(z_j)} (z_j - z_k) \quad (20)$$

$$\begin{aligned} & \leq P f[\operatorname{sgn}(z_k - z_j)]_{j,k=1, \dots, p} \quad (\text{: even}) \\ & : P f[\operatorname{sgn}(z_k - z_j)]_{j,k=1, \dots, p} \quad [g_j]_{j=1, \dots, p} \quad (\text{: odd}); \\ & \quad [g_k]_{k=1, \dots, p} \end{aligned}$$

with  $g_j = g_k = 1$ . The  $P f$  ans in the above can be represented as quaternion determinants [40{42,22,23,32,43]. In doing so, we need to introduce monic skew-orthogonal polynomials  $R_n(z) = z^n + \dots$ , which satisfy the skew-orthogonality relation:

$$hR_{2n}; R_{2m+1}i_R = hR_{2m+1}; R_{2n}i_R = r_{n-m}; \text{others} = 0; \quad (21)$$

where

$$hf;gi_R = \int_0^{z_1} dz \frac{p}{w(z)} g(z) \int_0^{z_2} dz^0 \frac{p}{w(z^0)} f(z^0) \quad (f \neq g); \quad (22)$$

and (integrated) wave functions,

$$r_n(z) = \frac{p}{z} \frac{1}{w(z)} R_n(z); \quad (23a)$$

$$r_n(z) = \int_0^{z^0} dz^0 \operatorname{sgn}(z - z^0) \frac{p}{w(z^0)} R_n(z^0); \quad (23b)$$

Note that for negative  $z$ ,  $r_n(z)$  is a constant:

$$r_n(z < 0) = r_n(0) \quad s_n; \quad (24)$$

Now we present the following theorems:

Theorem 1

For even  $p$ , we can express  $p(z_1, \dots, z_p)$  as

$$p(z_1, \dots, z_p) = \prod_{j=0}^{\frac{p}{2}-1} r_j T \det[f(z_j; z_k)]_{j,k=1, \dots, p}; \quad (25)$$

The quaternion elements  $f(z; z^0)$  are represented as

$$f(z; z^0) = \frac{S(z; z^0)}{D(z; z^0)} \frac{I(z; z^0)}{S(z^0; z)} : \quad (26)$$

The functions  $S(z; z^0)$ ,  $D(z; z^0)$  and  $I(z; z^0)$  are given by

$$S(z; z^0) = \prod_{n=0}^{\frac{p}{2}-1} \frac{z_{2n}(z) z_{2n+1}(z^0) z_{2n+1}(z) z_{2n}(z^0)}{r_n}; \quad (27a)$$

$$D(z; z^0) = \prod_{n=0}^{\frac{p}{2}-1} \frac{z_{2n}(z) z_{2n+1}(z^0) z_{2n+1}(z) z_{2n}(z^0)}{r_n}; \quad (27b)$$

$$I(z; z^0) = \prod_{n=0}^{\frac{p}{2}-1} \frac{z_{2n}(z) z_{2n+1}(z^0) z_{2n+1}(z) z_{2n}(z^0)}{r_n}; \quad (27c)$$

Theorem 2

For odd  $p$ , we can express  $p(z_1, \dots, z_p)$  as

$$\begin{aligned} p(z_1, \dots, z_p) = & \prod_{j=0}^{\frac{p-1}{2}} r_j s_{j+1} T \det[f^{\text{odd}}(z_j; z_k)]_{j,k=1, \dots, p} : \end{aligned} \quad (28)$$

The quaternion elements are represented as

$$f^{\text{odd}}(z; z^0) = \frac{S^{\text{odd}}(z; z^0)}{D^{\text{odd}}(z; z^0)} \frac{I^{\text{odd}}(z; z^0)}{S^{\text{odd}}(z^0; z)} \quad (29)$$

and  $s_n$  is defined in Eq.(24). The functions  $S^{\text{odd}}$ ,  $D^{\text{odd}}$ , and  $I^{\text{odd}}$  are given in terms of  $S$ ,  $D$ , and  $I$  in Theorem 1 according to

$$S^{\text{odd}}(z; z^0) = S(z; z^0) + \frac{r_1(z^0)}{s_1}; \quad (30a)$$

$$D^{\text{odd}}(z; z^0) = D(z; z^0); \quad (30b)$$

$$I^{\text{odd}}(z; z^0) = I(z; z^0) + \frac{r_1(z) r_1(z^0)}{s_1}; \quad (30c)$$

Here  $\bar{s}_n$  stands for a substitution

$$R_n(z) \mapsto R_n(z) - \frac{s_n}{s_1} R_{n-1}(z) \quad (31)$$

for  $n = 0, \dots, p-2$ , associated with a change in the upper limit of the sum

$$= 2 \quad 1 \quad \bar{s}_1 \quad [= 2] \quad 1: \quad (32)$$

Theorem 3

Let the quaternion elements  $q_{jk}$  of a selfdual  $n \times n$  matrix  $Q_n$  depend on  $n$  real or complex variables  $z_1, \dots, z_n$  as

$$q_{jk} = f(z_j; z_k); \quad (33)$$

We assume that  $f(z; z^0)$  satisfies the following conditions.

$$f(z; z)d(z) = c; \quad (34)$$

$$f(z; z^0)f(z^0; z^0)d(z^0) = f(z; z^0) + f(z; z^0)f(z; z^0);$$

Here  $d(z)$  is a suitable measure,  $c$  is a constant scalar, and  $f(z; z^0)$  is a constant quaternion. Then we have

Z

$$T \det Q_n d(z_n) = (c - n + 1) T \det Q_{n-1}; \quad (35)$$

where  $Q_{n-1}$  is the  $(n-1) \times (n-1)$  matrix obtained by removing the row and the column which contain  $z_n$ . It is straightforward to show that the quaternion element  $f(z; z^0)$  and  $f^{odd}(z; z^0)$  in Theorem 1 and Theorem 2 both satisfy the conditions imposed on  $f(z; z^0)$  in Theorem 3 with  $d(z) = w(z)dz$ . This means that we can write

$$p(z_1; \dots; z_{+p}) = \frac{Q_{[(+N)=2]}}{w(z_j)} \frac{\prod_{j=0}^{j=p} r_j}{\prod_{j>k} j z_j z_k} \quad (36)$$

$$s_{+N-1} T \det [f(z_j; z_k)]_{j,k=1,\dots,+p} \quad (+N : \text{even}) \\ s_{+N} T \det [f^{odd}(z_j; z_k)]_{j,k=1,\dots,+p} \quad (+N : \text{odd})$$

Since the final result in the asymptotic limit  $N \rightarrow 1$  should be insensitive to the parity of  $N$ , we consider only even  $+N$  henceforth. Then the  $p$ -level correlation function (11) is finally written as

$$(x_1; \dots; x_p; m_1; \dots; m_p) = \frac{T \det [f(z_j; z_k)]_{j,k=1,\dots,+p}}{T \det [f(z_j; z_k)]_{j,k=1,\dots,+p}}; \quad (37)$$

Now we proceed to take the asymptotic limit of the correlation function, by making use of explicit forms for the skew-orthogonal polynomials and their norms associated with the weight (18) obtained by Nagao and Wadati [44]:

$$S_{++}(z; z^0) = \frac{1}{8N} S(z; z^0) - \frac{1}{4} \int_0^Z dt t^2 \int_0^Z ds s J_{-1}(ts) \frac{J(t^0)}{t^0} - J(ts) J_{-1}(t^0) + \frac{J(t^0)}{4t^0};$$

$$D_{++}(z; z^0) = \frac{1}{(8N)^2} D(z; z^0) - \frac{1}{16} \int_0^Z dt t^2 \int_0^Z ds s J_{-1}(ts) \frac{J(t^0)}{t^0} - J(t^0) J_{-1}(t^0);$$

$$I_{++}(z; z^0) = I(z; z^0) - \int_0^Z dt t^2 \int_0^Z du u J_{-1}(tu) J(tv) - J(tu) v J_{-1}(tv) - J(u) du \operatorname{sgn}(z^0);$$

$$(b) z < 0; z^0 > 0$$

We define microscopic variables and as

$$z = \frac{2}{8N}; \quad z^0 = \frac{2}{8N}; \quad (43)$$

$$R_{2n}(z) = \frac{(2n)!}{2^{2n+1}} \frac{d}{dz} L_{2n+1}^{(-1)}(2z);$$

$$R_{2n+1}(z) = \frac{(2n+1)!}{2^{2n+1}} L_{2n+1}^{(-1)}(2z);$$

$$\frac{(2n)!}{2^{2n+2}} (2n+1) \frac{d}{dz} L_{2n}^{(-1)}(2z);$$

$$r_n = 2^{-4n} (2n)!(2n+1)!;$$

expressed in terms of the Laguerre polynomials

$$L_n^{(a)}(z) = \frac{z^a e^z}{n!} \frac{d^n}{dz^n} (e^{-z} z^{n+a}); \quad (39)$$

We need to evaluate the local asymptotics of the quaternion function  $f(z; z^0)$ , whose constituent Laguerre polynomials tend to

$$L_n^{(a)}(z) \rightarrow \begin{cases} (n=z)^{a=2} J_a(2^p \frac{p}{n} \bar{z}) & (z > 0) \\ (n=\bar{z})^{a=2} I_a(2^p \frac{p}{n} \bar{z}) & (z < 0) \end{cases}; \quad (40)$$

as  $n \rightarrow 1$ ,  $z \rightarrow 0$ , with  $nz$  fixed. Three cases should be considered separately:

$$(a) z; z^0 > 0; \quad (b) z < 0; z^0 > 0; \quad (c) z; z^0 < 0$$

(the case  $z > 0, z^0 < 0$  is unnecessary because of the selfduality  $f(z; z^0) = \hat{f}(z^0; z)$ ).

$$(a) z; z^0 > 0$$

We define microscopic variables and as

$$z = \frac{2}{8N}; \quad z^0 = \frac{2}{8N}; \quad (41)$$

according to Eq.(8) with  $\langle 0 \rangle = 2^p \frac{p}{2N}$ . If both arguments are positive, the asymptotic limit is known to be [22]

$$S_{++}(z; z^0) = \frac{1}{8N} S(z; z^0) - \frac{1}{4} \int_0^Z dt t^2 \int_0^Z ds s J_{-1}(ts) \frac{J(t^0)}{t^0} - J(ts) J_{-1}(t^0) + \frac{J(t^0)}{4t^0}; \quad (42a)$$

$$D_{++}(z; z^0) = \frac{1}{(8N)^2} D(z; z^0) - \frac{1}{16} \int_0^Z dt t^2 \int_0^Z ds s J_{-1}(ts) \frac{J(t^0)}{t^0} - J(t^0) J_{-1}(t^0); \quad (42b)$$

$$I_{++}(z; z^0) = I(z; z^0) - \int_0^Z dt t^2 \int_0^Z du u J_{-1}(tu) J(tv) - J(tu) v J_{-1}(tv) - J(u) du \operatorname{sgn}(z^0); \quad (42c)$$

according to Eqs.(6) and (8). When one of the arguments is negative, the identity (24) should be taken into consideration. We find

$$S_+(z; z^0) = S_+(z) - \frac{1}{8N} S(z; z^0) - \frac{J(z)}{4}; \quad (44a)$$

$$S_+(z; z^0) = \frac{1}{8N} S(z^0; z) - \frac{1}{4} \int_0^{Z_1} dt t^2 \int_0^s ds J_{-1}(ts) \frac{I(t)}{4} - J(ts) I_{-1}(t) + \frac{I(z)}{4}; \quad (44b)$$

$$D_+(z; z^0) = \frac{1}{(8N)^2} D(z; z^0) - \frac{1}{16} \int_0^{Z_1} dt t^2 I_{-1}(t) \frac{J(t)}{4} - \frac{I(t)}{4} J_{-1}(t); \quad (44c)$$

$$I_+(z; z^0) = I(z) - I(z; z^0) - \int_0^s J(u) du + 1; \quad (44d)$$

(c)  $z; z^0 < 0$

We define microscopic variables and as

$$z = \frac{z^2}{8N}; \quad z^0 = \frac{z^{02}}{8N}; \quad (45)$$

We can readily derive

$$S(z; z^0) = S(z^0) - \frac{1}{8N} S(z; z^0) - \frac{I(z^0)}{4}; \quad (46a)$$

$$D(z; z^0) = \frac{1}{(8N)^2} D(z; z^0) - \frac{1}{16} \int_0^{Z_1} dt t^2 I_{-1}(t) \frac{I(z^0)}{4} - \frac{I(t)}{4} I_{-1}(t^0); \quad (46b)$$

$$I(z; z^0) = I(z; z^0) - \text{sgn}(z^0); \quad (46c)$$

In Eqs.(44a), (44d), and (46a), we have introduced symbols with one sign subscript (e.g.  $S_+$ ) in order to indicate that they depend only on the second arguments of those with two sign subscripts (e.g.  $S_+$ ).

The scaled correlation function  $s$  is defined as

$$s(z_1; \dots; z_p; z_{p+1}; \dots; z_N) = \frac{1}{(8N)^p} (Pf \frac{1}{8N}; \dots; Pf \frac{p}{8N}; Pf \frac{1}{8N}; \dots; Pf \frac{p}{8N}); \quad (47)$$

By making use of Eqs.(14), (37), and Dyson's equality (A 6), we finally obtain

$$\begin{aligned} s(z_1; \dots; z_p; z_{p+1}; \dots; z_N) &= 2^p \sum_{k=1}^2 \frac{Y^p}{j_k j} \frac{\begin{matrix} I \\ S \\ D \\ S^T \\ I^T \\ S^T \\ S^T \end{matrix}}{\begin{matrix} S \\ D \\ S^T \\ D^T \\ I^T \\ D^T \\ S^T \end{matrix}} \frac{\begin{matrix} I_+ \\ S_+ \\ S^T_+ \\ D^T_+ \\ S^T_+ \\ S^T_+ \end{matrix}}{\begin{matrix} S_+ \\ D_+ \\ I_+ \\ S_{++} \\ I_+ \\ D_{++} \end{matrix}} \frac{\begin{matrix} S_+ \\ D_+ \\ S_{++} \\ S_{++} \end{matrix}}{\begin{matrix} S_{++} \\ D_{++} \end{matrix}} \\ &= 2^p \sum_{k=1}^2 \frac{Y^p}{j_k j} \frac{\begin{matrix} D \\ S^T \\ I^T \\ S^T \\ D^T \\ S^T \end{matrix}}{\begin{matrix} Pf \\ Pf \\ Pf \\ Pf \\ Pf \\ Pf \end{matrix}} \frac{\begin{matrix} S \\ D \\ S^T \\ I_+ \\ S^T \\ D^T \end{matrix}}{\begin{matrix} D \\ S^T \\ I^T \\ S^T \\ D^T \\ S^T \end{matrix}} \frac{\begin{matrix} D_+ \\ S_{++} \\ I_+ \\ S_{++} \\ D_{++} \end{matrix}}{\begin{matrix} S_{++} \\ D_{++} \end{matrix}} \quad (\text{ : even}) \end{aligned} \quad (48a)$$

$$\begin{aligned} s(z_1; \dots; z_p; z_{p+1}; \dots; z_N) &= 2^p \sum_{k=1}^2 \frac{Y^p}{j_k j} \frac{\begin{matrix} 0 \\ S \\ D \\ S^T \\ I^T \\ S^T \end{matrix}}{\begin{matrix} Pf \\ Pf \\ Pf \\ Pf \\ Pf \\ Pf \end{matrix}} \frac{\begin{matrix} I_+ \\ S_+ \\ S^T_+ \\ I_+ \\ S^T_+ \\ D^T_+ \end{matrix}}{\begin{matrix} S \\ D \\ S^T \\ I_+ \\ S^T \\ D^T \end{matrix}} \frac{\begin{matrix} S_+ \\ D_+ \\ I_+ \\ S_{++} \\ S^T_+ \\ D_{++} \end{matrix}}{\begin{matrix} S_{++} \\ D_{++} \end{matrix}} \quad (\text{ : odd}): \end{aligned} \quad (48b)$$

The elements of the matrices  $S_0$ ,  $D_0$ ,  $I_0$  and the row vectors  $S$ ,  $I$  ( $; z^0 = +; z$ ) in the above are defined as

$$(S_{++})_{k\ell} = S_{++}(k; \cdot); (S_{+})_{i\ell} = S_{+}(i; \cdot); (S_{-})_{ij} = S_{-}(i; j); (S_{+})_{\ell} = S_{+}(\cdot); (S_{-})_j = S_{-}(j); \text{etc.} \quad (49)$$

where the subscripts take their values in  $i, j = 1, \dots, p$ , and  $k, \ell = 1, \dots, p$ . In the last two lines we have exploited a Pfa an identity that holds for antisymmetric matrices  $A, B$ , and a row vector  $v$ :

$$\begin{array}{c} 2 \\ | \\ \begin{array}{c} 6 \\ 4 \\ \hline v^T \end{array} \end{array} \begin{array}{c} A \\ | \\ \begin{array}{c} v \\ : \\ 7 \\ 5 \\ \hline vB \end{array} \end{array} \begin{array}{c} 3 \\ | \\ \begin{array}{c} 1 \\ 7 \\ 5 \\ \hline 1 \end{array} \end{array} = Pf[A]Pf[B] \quad (\text{rank}(A), \text{rank}(B) : \text{even}) \quad (50a)$$

$$\begin{array}{c} 2 \\ | \\ \begin{array}{c} 6 \\ 4 \\ \hline v^T \end{array} \end{array} \begin{array}{c} A \\ | \\ \begin{array}{c} v \\ : \\ 7 \\ 5 \\ \hline 1 \end{array} \end{array} \begin{array}{c} 3 \\ | \\ \begin{array}{c} 1 \\ 7 \\ 5 \\ \hline 1 \end{array} \end{array} = Pf \begin{array}{c} 6 \\ 4 \\ \hline v^T \end{array} \begin{array}{c} A \\ | \\ \begin{array}{c} 0 \\ v^T \\ B \\ \hline 1 \end{array} \end{array} \quad (\text{rank}(A), \text{rank}(B) : \text{odd}): \quad (50b)$$

In the quenched limit  $\lim_{\epsilon \rightarrow 0}$  when the ratio of two Pfans is replaced by a minor  $Pf \frac{I_{++}}{S_{++}} \frac{S_{++}}{D_{++}}$ , the correlation function tends to that of Laguerre orthogonal ensemble computed by Nagao and Forrester, Eqs.(2.21), (2.18), (2.19), and (3.20) in Ref. [22], with  $\epsilon = 2a + 1$ . By the same token, it satisfies a sequence

$$s(f(g; 1, \dots, p)) = \frac{1!}{1} s(f(g; 1, \dots, 1)) = \frac{1!}{1} \frac{1!}{1} s(f(g; 1, \dots, 2)) = \frac{2!}{2} \dots ; \quad (51)$$

as each of the masses are decoupled by sending to infinity. To illustrate this decoupling, we exhibit in Fig.1 a plot of the spectral density  $s(\cdot)$  ( $\epsilon = 0; p = 1; \epsilon = 1$ ) that interpolates between known results for the chiral and quenched limits.

### III. SYMPLECTIC ENSEMBLE

Although correlation functions of the massive chiral symplectic ensemble have previously been computed by the authors [33], we nevertheless present their explicit expressions for the sake of completeness. We concentrate on the case with an even  $N_f$  ( $= 2$ ) number of flavors and pairwise degenerated mass parameters, corresponding to adjoint Dirac fermions in the QCD context. The scaled p-level correlation functions, defined in Eq.(47), is expressed by construction as a ratio of partition functions with  $2$  and  $2 + 4p$  flavors [45,46],

$$s(1, \dots, p; f(g)) = C^{(p)} \frac{\prod_{k=1}^p \frac{Y^p}{(2k-2)^4} \prod_{k=1}^p \frac{Y^p}{j_k j^3} \prod_{i=1}^p \frac{Y}{(k+i)^2}}{Pf[\prod_{k=1}^p I_{++}]} \frac{Z^{(4)}(1, 1, \dots, 1; i_1, \dots, i_p, \dots, i_p)}{Z^{(4)}(1, 1, \dots, 1)} \frac{z_1^4 z_2^4 \dots z_p^4}{\prod_{k=1}^p \{z_k\}} \quad (52)$$

Here  $C^{(p)}$  stands for a constant to be fixed below. Using an explicit form of the partition function, Eq.(31) of Ref. [33], and taking convenient limits in  $i_k$ 's, we obtain

$$s(1, \dots, p; f(g)) = (-2)^p \frac{\prod_{k=1}^p \frac{Y^p}{j_k j} Pf[\prod_{k=1}^p I_{++}]}{Pf[\prod_{k=1}^p I_{++}]} \quad (\text{: even}) \quad (53a)$$

$$= (-2)^p \frac{\prod_{k=1}^p \frac{Y^p}{j_k j} Pf[\prod_{k=1}^p I_{++}]}{Pf[\prod_{k=1}^p Q^T]} \quad (\text{: odd}): \quad (53b)$$

The elements of the matrices  $S^0, D^0, I^0$  and the row vectors  $Q^0, P^0$  ( $Q^0 = +; P^0 = -$ ) in the above are defined as

$$\begin{aligned}
(I_{-})_{ij} &= I_{-}(i; j) \quad i, j \quad dt^0 \quad du \quad I_2(2t_i)I_2(2tu_j) \quad I_2(2tu_i)I_2(2t_j); \\
(I_{+})_{ij} &= I_{+}(i; j) \quad i, j \quad dt^0 \quad du \quad I_2(2t_i)J_2(2tu_j) \quad I_2(2tu_i)J_2(2t_j); \\
(S_{+})_{ij} &= S_{+}(i; j) \quad i, j \quad dt^2 \quad du \quad I_2(2t_i)uJ_2(2tu_j) \quad I_2(2tu_i)J_2(2t_j); \\
(I_{++})_{kj} &= I_{++}(k; j) \quad k, j \quad dt^0 \quad du \quad J_2(2t_k)J_2(2tu_j) \quad J_2(2tu_k)J_2(2t_j); \\
(S_{++})_{kj} &= S_{++}(k; j) \quad k, j \quad dt^2 \quad du \quad J_2(2t_k)uJ_2(2tu_j) \quad J_2(2tu_k)J_2(2t_j); \\
(D_{++})_{kj} &= D_{++}(k; j) \quad k, j \quad dt^3 \quad du \quad uJ_2(2t_k)J_2(2tu_j) \quad J_2(2tu_k)J_2(2t_j); \\
(Q_{-})_j &= \int_0^{\infty} dt I_2(2t_j); \quad (Q_{+})_j = \int_0^{\infty} dt J_2(2t_j); \quad (P_{+})_j = \int_0^{\infty} dt uJ_2(2t_j);
\end{aligned} \tag{54}$$

where the subscripts take their values in  $i, j = 1, \dots, p$ , and  $k, j' = 1, \dots, p$ . The overall constant is determined as the above by requiring that in the quenched limit  $\lim_{N \rightarrow \infty}$  the ratio of two Pfaffians is replaced by a minor Pfaffian  $\frac{I_{++}}{S_{++}} \frac{S_{++}}{D_{++}}$ , the correlation function tends to that of Laguerre symplectic ensemble computed by Nagao and Forrester, Eqs.(2.27), (2.25), and (4.7–9) in Ref. [22] (whose notations are related to ours via an unfolding change of variables)

$$\begin{aligned}
I_{++}(\cdot; \cdot) &= I_4 \frac{2}{8N}; \frac{0^2}{8N}; \\
S_{++}(\cdot; \cdot) &= \frac{1}{8N} S_4 \frac{2}{8N}; \frac{0^2}{8N}; \\
D_{++}(\cdot; \cdot) &= \frac{1}{(8N)^2} D_4 \frac{2}{8N}; \frac{0^2}{8N};
\end{aligned} \tag{55}$$

and  $\cdot = 2a - 1=2$ . It is easy to confirm that the correlation functions satisfies the decoupling sequence (51) as each of the masses are sent to infinity. To illustrate this decoupling, we exhibit in Fig. 2 a plot of the spectral density  $s(\cdot; \cdot)$  ( $a = 0; p = 1; j = 1$ ) that interpolates between known results for the chiral and quenched limits.

#### ACKNOWLEDGMENTS

This work was supported in part (SMN) by JSPS Research Fellowships for Young Scientists, and by Grant-in-Aid No. 411044 from the Ministry of Education, Science, and Culture, Japan.

#### APPENDIX A : QUATERNION DETERMINANT

A quaternion is defined as a linear combination of four basic units  $f_1; e_1; e_2; e_3$ :

$$q = q_0 + q_1 e_1 + q_2 e_2 + q_3 e_3; \tag{A 1}$$

Here the coefficients  $q_0; q_1; q_2$  and  $q_3$  are real or complex numbers. The first part  $q_0$  is called the scalar part of  $q$ . The quaternion basic units satisfy the multiplication laws

$$\begin{aligned}
1 \cdot 1 &= 1; 1 \cdot j \neq e_j \quad 1 = e_j; j = 1, 2, 3; \\
e_1^2 &= e_2^2 = e_3^2 = e_1 e_2 e_3 = 1;
\end{aligned} \tag{A 2}$$

The multiplication is associative and in general not commutative. The dual  $\hat{q}$  of a quaternion  $q$  is defined as

$$\hat{q} = q_0 - q_1 e_1 - q_2 e_2 - q_3 e_3; \tag{A 3}$$

For a selfdual  $N \times N$  matrix  $Q$  with quaternion elements  $q_{jk}$  has a dual matrix  $\hat{Q} = [q_{kj}]$ . The quaternion units can be represented as  $2 \times 2$  matrices

$$\begin{aligned}
1 &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}; \quad e_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}; \\
e_2 &= \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}; \quad e_3 = \begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix};
\end{aligned} \tag{A 4}$$

We define a quaternion determinant  $T \det Q$  of a selfdual  $Q$  (i.e.,  $Q = \hat{Q}$ ) as

$$T \det Q = \sum_{P} \frac{X}{(-1)^{N-1}} \frac{Y^1}{(-1)^{N-1}} (q_{ab} q_{bc} \dots q_{da}); \tag{A 5}$$

where  $P$  denotes any permutation of the indices  $(1, 2, \dots, N)$  consisting of 1 exclusive cycles of the form  $(a ! b ! c ! \dots ! d ! a)$  and  $(^N 1)$  is the parity of  $P$ . The subscript 0 means that the scalar part of the product is taken over each cycle. Note that a quaternion determinant of a selfdual quaternion matrix is always a scalar. The quaternion determinant can as well be represented by the  $2N \times 2N$  representation  $C(Q)$  [40]:

$$T \det Q = P f[J C(Q)]; \quad J = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}; \tag{A 6}$$

APPENDIX B : QUATERNION KERNEL FOR  
THE ORTHOGONAL ENSEMBLE

Recently Akemann and Kan zieper [34] derived the asymptotic correlation function in a special case  $p = \infty = 1$ . Though their result is clearly in agreement with that special case of ours, there is a difference in the appearance because they adopted alternative asymptotic formulas. Their formulas are based on an asymptotic relation for  $z; z^0 > 0$  and  $\epsilon \neq 1$  (see below):

$$S(z; z^0) = 2N \int_0^{\infty} dt t J_1(t) J_1(t^0) + \frac{J_1(0)}{z} \int_0^{\infty} ds s J_2(s) - 1 ; \quad (\text{B } 1)$$

$$\begin{aligned} t \int_0^{\infty} ds s J_1(ts) &= J_1(t) + (-1) \int_0^{\infty} ds J_1(ts); \\ t \int_0^{\infty} ds J_1(ts) &= J_1(t) + J_1(0) + (-1) \int_0^{\infty} \frac{ds}{s} J_1(ts); \end{aligned} \quad (\text{B } 3)$$

Substitution of Eq.(B 3) into (42a) yields

$$S(z; z^0) = 2N \left[ \int_0^{\infty} dt t J_1(t) J_1(t^0) + \int_0^{\infty} dt t (J_1(t) - J_1(0)) J_1(t^0) + \frac{J_1(0)}{z} \right] + \frac{1}{z} \int_0^{\infty} dt t \int_0^{\infty} ds s J_1(ts) J_1(t^0) - (-1) \int_0^{\infty} dt t \int_0^{\infty} \frac{ds}{s} J_1(ts) J_1(t^0) ; \quad (\text{B } 4)$$

By partial integrations, we find

$$\int_0^{\infty} dt t J_1(t) J_1(t^0) = \int_0^{\infty} dt t J_1(t) J_1(t^0) - J_1(0) \frac{J_1(0)}{z} \quad (\text{B } 5)$$

and

$$\frac{1}{z} \int_0^{\infty} dt t \int_0^{\infty} ds s J_1(ts) J_1(t^0) = \int_0^{\infty} dt t \int_0^{\infty} \frac{ds}{s} J_1(ts) J_1(t^0) - \int_0^{\infty} \frac{ds}{s} J_1(s) J_1(0) \frac{J_1(0)}{z} ; \quad (\text{B } 6)$$

We substitute (B 5) and (B 6) into (B 4) and obtain

$$S(z; z^0) = 2N \left[ \int_0^{\infty} dt t J_1(t) J_1(t^0) - J_1(0) \int_0^{\infty} dt t J_1(t^0) - J_1(0) + (-1) \int_0^{\infty} dt t \int_0^{\infty} \frac{ds}{s} J_1(s) J_1(0) - 1 \frac{J_1(0)}{z} \right] ; \quad (\text{B } 7)$$

Again by a partial integration, we find

$$J_1(0) + (-1) \int_0^{\infty} \frac{ds}{s} J_1(s) = J_1(0) + \int_0^{\infty} ds J_2(s) \quad (\text{B } 8)$$

and thus arrive at the desired result (B 1) provided that  $\epsilon$  is not equal to 1. Since  $J_1(0) = 1$ , the case  $\epsilon = 1$  is exceptional.

Similarly, for  $z < 0; z^0 > 0$  and  $\epsilon \neq 1$ , an alternative expression

$$S(z^0; z) = 2N \int_0^{\infty} dt t J_1(t) I_1(t) - \frac{I_1(0)}{z} \int_0^{\infty} ds s J_2(s) - 1 ; \quad (\text{B } 9)$$

where we have adopted the microscopic variables (41). This asymptotic relation was derived by Forrester, Nagao, and Honner [43] in a study of parametric random matrix ensembles. Note that the first integral in the above is equal to the Bessel kernel,

$$\frac{J(\epsilon) J_{-1}(0) - J_{-1}(\epsilon) J(0)}{2\pi^2} ; \quad (\text{B } 2)$$

From the derivation the equivalence of (42a) and (B 1) was well established. However, it is worth directly proving it here for an unambiguous identification.

Using the Bessel function identities, we can readily see that

$$\begin{aligned} t \int_0^{\infty} ds s J_1(ts) &= J_1(t) + (-1) \int_0^{\infty} ds J_1(ts); \\ t \int_0^{\infty} ds J_1(ts) &= J_1(t) + J_1(0) + (-1) \int_0^{\infty} \frac{ds}{s} J_1(ts); \end{aligned} \quad (\text{B } 3)$$

in terms of the microscopic variables (43) is available.

## APPENDIX C : QUATERNION KERNEL FOR THE SYMPLECTIC ENSEMBLE

In previous [33] and this works, the authors evaluated Dirac eigenvalue correlation functions for the symplectic ensemble with doubly degenerate masses. If masses are quadruply degenerate, the evaluation of the correlation functions is easier because the conventional integration method for the massless Laguerre ensemble works without any modification. In that case, the multiple integral we need to evaluate is

$$\begin{aligned} & p(x_1; \dots; x_p; fm g) \\ &= \frac{1}{(N-p)!} \int_0^{z_1} \int_0^{z_1} \dots \int_{j=p+1}^N dz_j \int_{j=1}^N x_j^{2p+1} e^{4x_j} \quad (C1) \\ & \quad \times \prod_{j=1}^N \prod_{i=1}^4 (x_j + m_i^2)^4 \prod_{j>k} j x_j - x_k \end{aligned}$$

where we set  $N_f = 4$  and

$$fm g = \left[ \frac{z_1}{m_1}, \dots, \frac{z_1}{m_1}, \dots, \frac{z_N}{m_1}, \dots, \frac{z_N}{m_1} \right]$$

The conventional massless' theory [42,22,23,32] tells us that the correlation functions are written in terms of quaternion determinants:

$$\begin{aligned} (x_1; \dots; x_p; fm g) &= \frac{p(z_1; \dots; z_{+p})}{0(z_1; \dots; z)} \\ &= \frac{T \det [f(z_j; z_k)]_{j,k=1,\dots,+p}}{T \det [f(z_j; z_k)]_{j,k=1,\dots,p}}; \quad (C2) \end{aligned}$$

where we have adopted the notation (16).

The quaternion function  $f(z; z^0)$  is represented as

$$f(z; z^0) = \begin{cases} S(z; z^0) & I(z; z^0) \\ D(z; z^0) & S(z^0; z) \end{cases} : \quad (C3)$$

We evaluate the asymptotic limit of the quaternion function  $f(z; z^0)$  in each of the three cases

$$(a) z; z^0 > 0; \quad (b) z < 0; z^0 > 0; \quad (c) z; z^0 < 0;$$

as in Sect.2. in terms of the microscopic variables  $; ; ;$  defined by Eqs.(41), (43), (45), respectively.

$$(a) z; z^0 > 0$$

In the case  $z; z^0 > 0$ , Nagao and Forrester [22] derived the asymptotic limit (55), that is

$$\begin{aligned} & \frac{1}{8N} S(z; z^0) = S_{++}(; ;^0); \\ & \frac{1}{(8N)^2} D(z; z^0) = D_{++}(; ;^0); \\ & I(z; z^0) = I_{++}(; ;^0); \end{aligned} \quad (C4)$$

where  $S_{++}$ ,  $D_{++}$ , and  $I_{++}$  are defined in Eq.(54).

$$(b) z < 0; z^0 > 0$$

Using the asymptotic formula for the Bessel function (40), we can similarly treat negative argument cases to obtain

$$\begin{aligned} & \frac{1}{8N} S(z; z^0) = S_{+}(; ); \\ & \frac{1}{8N} S(z^0; z) = \int_0^{z_1} \int_0^{z_1} dt t^2 \int_0^{2tu} du J_2(2tu) I_{2+1}(2t) - J_2(2t) u I_{2+1}(2tu); \\ & \frac{1}{(8N)^2} D(z; z^0) = \int_0^{z_1} \int_0^{z_1} dt t^3 \int_0^{2tu} du u I_{2+1}(2tu) J_{2+1}(2t) - I_{2+1}(2t) J_{2+1}(2tu) \\ & I(z; z^0) = I_{+}(; ); \end{aligned} \quad (C5)$$

where  $S_{+}$  and  $I_{+}$  are defined in Eq.(54).

$$(c) z; z^0 < 0$$

$$\begin{aligned} & \frac{1}{8N} S(z; z^0) = \int_0^{z_1} \int_0^{z_1} dt t^2 \int_0^{2tu} du I_2(2tu) I_{2+1}(2t^0) - I_2(2t) u I_{2+1}(2tu^0); \\ & \frac{1}{(8N)^2} D(z; z^0) = \int_0^{z_1} \int_0^{z_1} dt t^3 \int_0^{2tu} du u I_{2+1}(2tu) I_{2+1}(2t^0) - I_{2+1}(2t) I_{2+1}(2tu^0); \\ & I(z; z^0) = I_{-}(;^0); \end{aligned} \quad (C6)$$

where  $I_{++}$  is defined in Eq.(54).

We can use Dyson's equality (A 6) to see that the above quaternion determinant expression is identical to the limit of quadruple mass degeneracy of the general formula (53a) employing Pfaans. In this case, yet another equivalent asymptotic formula was recently presented by Akemann and Kanazieper [34]. Now we shall directly demonstrate the equivalence. We should firstly note that, under the change of the quaternion elements

$$S(z; z^0) = S(z; z^0) - S(z; z^0) + \frac{1}{W(z^0)} \frac{dW(z^0)}{dz^0} I(z; z^0);$$

$$\int_0^{Z_1} dt t \int_0^{Z_1} du J_2(2tu) J_2(2t^0) = \int_0^{Z_1} dt t \int_0^{Z_1} du \frac{du}{u} J_{2+1}(2tu) J_{2+1}(2t^0) + \frac{1}{2} J_2(2^0) \int_0^{Z_1} du \frac{du}{u} J_{2+1}(2u);$$

(C 9)

by a partial integration. The asymptotic formulas (C 4), together with the identity (C 9), yield

$$\begin{aligned} S(z; z^0) &= 2N \int_0^{Z_1} dt t \int_0^{Z_1} du J_{2+1}(2tu) J_2(2t^0) + (2+1) \int_0^{Z_1} du \frac{du}{u} J_{2+1}(2tu) \\ &\quad + 2 \int_0^{Z_1} dt t \int_0^{Z_1} du u J_{2+1}(2tu) J_2(2t^0) + (2+1) \int_0^{Z_1} du J_2(2tu) J_2(2t^0) \\ &\quad - (2+1) \int_0^{Z_1} du \frac{du}{u} J_{2+1}(2u); \end{aligned} \quad (C 10)$$

Partial integrations give rise to the Bessel function equalities

$$\begin{aligned} (2+1) \int_0^{Z_1} du \frac{du}{u} J_{2+1}(2tu) &= J_{2+1}(2t^0) + 2t \int_0^{Z_1} du J_2(2tu); \\ (2+1) \int_0^{Z_1} du J_2(2tu) J_2(2t^0) &= J_2(2t^0) + 2t^0 \int_0^{Z_1} du u J_{2+1}(2tu); \end{aligned} \quad (C 11)$$

Substituting (C 11) into (C 10) and again using the formula (B 6), we obtain

$$S(z; z^0) = 2N \int_0^{Z_1} dt t \int_0^{Z_1} du J_{2+1}(2tu) J_{2+1}(2t^0) + \frac{J_2(2^0)}{2} J_{2+1}(2^0) (2+1) \int_0^{Z_1} du \frac{du}{u} J_{2+1}(2u); \quad (C 12)$$

By a partial integration, we can rewrite it as

$$S(z; z^0) = 2N \int_0^{Z_1} dt t \int_0^{Z_1} du J_{2+1}(2tu) J_{2+1}(2t^0) - \frac{J_2(2^0)}{2} \int_0^{Z_2} ds J_{2+2}(s); \quad (C 13)$$

This is the asymptotic formula derived by Forrester, Nagao, and Honner [43] in a study of parametric random matrices and then applied by Akemann and Kanazieper [34] to the massive Dirac operator problem. Thus we established the equivalence in the case of quadruply degeneracy masses. We can straightforwardly extend it to the formulas with  $z$  and/or  $z^0$  negative.

$$\begin{aligned} I(z; z^0) &= I(z; z^0) - \int_z^{Z_1} S(z; z^0) dz^0; \\ D(z; z^0) &= D(z; z^0) - \frac{\partial}{\partial z} S(z; z^0); \end{aligned} \quad (C 7)$$

where

$$W(z) = j \cdot j^{+1=2} e^{2z}; \quad (C 8)$$

the quaternion determinant is unchanged. This transformation was introduced in Ref. [44].

For  $z; z^0 > 0$ , we find an identity

$$\int_0^{Z_1} dt t \int_0^{Z_1} du J_{2+1}(2tu) J_2(2t^0) + \frac{1}{2} J_2(2^0) \int_0^{Z_1} du \frac{du}{u} J_{2+1}(2u);$$

(C 9)

e-mail addresses:

nagao@sphinx.phys.sci.osaka-u.ac.jp,  
nishigak@th.phys.titech.ac.jp.

- [1] M. L. Mehta, Random Matrices, 2nd Ed., Academic Press (San Diego, 1991).
- [2] T. Guhr, A. M. Müller-Groeling, and H. A. Weidenmüller, Phys. Rep. 299, 189 (1998).
- [3] K. Efetov, Supersymmetry in Disorder and Chaos, Cambridge Univ. Press (Cambridge, 1997).
- [4] E. V. Shuryak and J. J. M. Verbaarschot, Nucl. Phys.

- A 560, 306 (1993).
- [5] J.J.M . Verbaarschot and I. Zahed, Phys. Rev. Lett. 70, 3852 (1993).
- [6] J.J.M . Verbaarschot, e-print hep-ph/9902394.
- [7] P.H . Damgaard, J.O . Stoen, D . Toublan, and J.J.M . Verbaarschot, Nucl. Phys. B 547, 305 (1999).
- [8] D . Toublan and J.J.M . Verbaarschot, Nucl. Phys. B 560, 259 (1999).
- [9] H . Leutwyler and A . Smilga, Phys. Rev. D 46, 5607 (1992).
- [10] A . Smilga and J.J.M . Verbaarschot, Phys. Rev. D 51, 829 (1995).
- [11] J.J.M . Verbaarschot, Phys. Rev. Lett. 72, 2531 (1994).
- [12] J.J.M . Verbaarschot, H . A . Wiedemann, and M . R . Zimmerman, Phys. Rep. 129, 367 (1985).
- [13] M . A . Halasz and J.J.M . Verbaarschot, Phys. Rev. D 52, 2563 (1995).
- [14] T . Banks and A . Casher, Nucl. Phys. B 169, 103 (1980).
- [15] T . Nagao and K . Slevin, J. Math. Phys. 34, 2075 (1993).
- [16] P . J . Forrester, Nucl. Phys. B 402, 709 (1993).
- [17] A . V . Andreev, B . D . Simons, and N . Taniguchi, Nucl. Phys. B 432, 487 (1994).
- [18] K . Slevin and T . Nagao, Phys. Rev. Lett. 70, 635 (1993).
- [19] P . J . Forrester and T . D . Hughes, J. Math. Phys. 35, 6736 (1994).
- [20] C . A . Tracy and H . W . Idom , Commun. Math. Phys. 161, 289 (1994).
- [21] J.J.M . Verbaarschot, Nucl. Phys. B 426, 559 (1994).
- [22] T . Nagao and P . J . Forrester, Nucl. Phys. B 435, 401 (1995).
- [23] T . Nagao and P . J . Forrester, Nucl. Phys. B 509, 561 (1998).
- [24] T . G . Uhr and T . W . Ettig, Nucl. Phys. B 506, 589 (1997).
- [25] A . D . Jackson, M . K . Sener, and J.J.M . Verbaarschot, Phys. Lett. B 387, 355, (1996).
- [26] P . H . Damgaard and S . M . Nishigaki, Nucl. Phys. B 518, 495 (1998).
- [27] T . W . Ilke, T . G . Uhr, and T . W . Ettig, Phys. Rev. D 57, 6486 (1998).
- [28] J . Jurkiewicz, M . Nowak, and I. Zahed, Nucl. Phys. B 513, 759 (1998).
- [29] S . M . Nishigaki, P . H . Damgaard, and T . W . Ettig, Phys. Rev. D 58, 087704 (1998).
- [30] M . E . Berbenni-Bitsch, S . M . Eyer, and T . W . Ettig, Phys. Rev. Lett. 80, 1146 (1998).
- [31] F . Farchioni, P . de Forcrand, I . Hipp, C . B . Lang, and K . Splittor, e-print hep-lat/9912004.
- [32] T . Nagao and P . J . Forrester, Nucl. Phys. B 530, 742 (1998); B 563, 547 (1999).
- [33] T . Nagao and S . M . Nishigaki, e-print hep-th/0001137.
- [34] G . A . kemann and E . K . Anzinger, e-print hep-th/0001188.
- [35] S . Nishigaki, Phys. Lett. B 387, 139 (1996).
- [36] G . A . kemann, P . H . Damgaard, U . M . Agnieszka, and S . Nishigaki, Nucl. Phys. B 487, 721 (1997).
- [37] E . K . Anzinger and V . Freilikher, Philos. Mag. B 77, 1161 (1998).
- [38] H . W . Idom , J. Stat. Phys. 94, 347 (1999).
- [39] M . K . Sener and J.J.M . Verbaarschot, Phys. Rev. Lett. 81, 248 (1998).
- [40] F . Dyson, Commun. Math. Phys. 19, 235 (1970)
- [41] M . L . Mehta, Matrix Theory, Les Editions de Physique (Paris, 1989).
- [42] G . M . Ahoux and M . L . Mehta, J. Phys. I (France) 1 (1991) 1093.
- [43] P . J . Forrester, T . Nagao, and G . Honner, Nucl. Phys. B 533, 601 (1999).
- [44] T . Nagao and M . W . Adati, J. Phys. Soc. Japan 60, 3298 (1991); 61, 78 (1992); 61, 1910 (1992).
- [45] P . H . Damgaard, Phys. Lett. B 424, 322 (1998).
- [46] G . A . kemann and P . H . Damgaard, Nucl. Phys. B 528, 411 (1998); Phys. Lett. B 432, 390 (1998).




FIG. 1. The scaled spectral density  $\rho_s(\zeta; \mu)$  for the chiral orthogonal ensemble with one ( $= 1$ ) flavor.




FIG. 2. The scaled spectral density  $\rho_s(\zeta; \mu, \mu)$  for the chiral symplectic ensemble with two degenerate ( $= 1$ ) flavors.