

Addendum-erratum to:
“Nonasymptotic critical behavior from field theory at $d = 3$. II. The ordered-phase case”
Phys. Rev. B35, 3585 (1987).

C. Bagnuls*

Service de Physique de l'Etat Condensé, C. E. Saclay, F91191 Gif-sur-Yvette Cedex, France

C. Bervillier†

Service de Physique Théorique, C. E. Saclay, F91191 Gif-sur-Yvette Cedex, France

D. I. Meiron‡

Applied and Computational Mathematics, MC 08-31, Caltech, Pasadena, CA 91125, USA

B. G. Nickel§

Department of Physics, University of Guelph, Guelph, Ontario, Canada N1G2W1

(January 11, 2002)

This note is intended to emphasize the existence of estimated Feynman integrals in three dimensions for the free energy of the $O(1)$ scalar theory up to five loops which may be useful for other work. We also correct some errors in the original paper.

05.70.Jk, 11.10.Hi

Addendum In tables I and II we present the corrected table I of the original paper¹. They display the values of the Feynman integrals of the $O(1)$ scalar theory contributing to the free energy up to five loops and their symmetry factors. The values differ from those Feynman integrals previously calculated in Ref. 2 due to the necessity of introducing a “soft” mass parameter instead of the usual renormalized (at zero-momentum) mass (see the text of the original paper¹ and Ref. 3 for details). Consequently, many of the estimates of Feynman integrals presented in table I of Ref. 1 have been extracted from Ref. 2 by accounting for a harmless 3-d renormalization in order to get a soft-mass parameter (characterizing a minimal subtraction scheme similar to that introduced in Ref. 4). Since the five-loop contributions to the free energy involve φ^3 vertices mixed to φ^4 vertices, Feynman integrals which are different or cannot be obtained from those considered in Ref. 2 have been estimated in three dimensions for the occasion⁵. These are:

- the four-loop integrals with Heap’s numbers 13–17 (column h in the following tables I and II, see table IV of Ref. 6). They involve only φ^3 vertices. They have been estimated and successfully compared to similar calculations extracted from Ref. 7.
- the five-loop integrals with Heap’s numbers 80–102 (see table V of Ref. 6). They involve φ^3 vertices mixed with a single φ^4 vertex. They have been calculated exclusively for this work.
- the five-loop integrals with Heap’s numbers 103–118 (see table V of Ref. 6). They involve only φ^3 vertices. They have been calculated for this work and successfully compared to similar calculations extracted from Ref. 8.

Errata Tables I and II (corrected table I of Ref. 1) account for two corrections compared to the original paper¹:

- for $b = 5$, $h = 15$, $l = 0$, $m = 1$, in the column “Value at $d = 3$ ”, one reads 6.24797746 instead of 4.21825152.
- Eq. (A13), which gives the contributions $b = 5$, $h = 49$, $l = 2$, $m = 1$ and $b = 5$, $h = 50$, $l = 2$, $m = 2$ in table I, should read:

$$4I_1^{(5,2)}(1) + 2I_2^{(5,2)}(1) = -0.1202442510 - 1.558293723 \ln(r'_0/g_0^2) \quad (\text{A13})$$

*e-mail: bagnuls@spec.saclay.cea.fr

†e-mail: bervil@spht.saclay.cea.fr

‡e-mail: dim@its.caltech.edu

§e-mail: bgn@physics.uoguelph.ca

As consequences, in table II of Ref. 1, the values of F_{500} and F_{510} should read:

$$\begin{aligned} F_{500} &= -0.45163891229 \times 10^{-7} \\ F_{510} &= 0.80687809188 \times 10^{-7} \end{aligned}$$

in agreement with table 1 of Ref. 9. In addition, as indicated in the note [12] of Ref. 10, the five loop coefficients of $X(v)$ and $F(v)$ given in table III of Ref. 1 should be modified but also that of $\tilde{F}(v)$, so that it is preferable to redone table III (see the present table III).

Moreover, the following equations were not correctly written, they should read:

$$\Gamma_{0,-}^{(0,0)}(r_0, g_0, M_0, \epsilon) = \frac{1}{2} r_0 M_0^2 + g_0 \frac{M_0^4}{24} + \sum_{b=1}^5 \Gamma_b(r_0, g_0, M_0, \epsilon) \quad (3.1)$$

$$H_{blk} = \left(2 - \frac{b}{2} - l\right) F_{blk} + (l+1) F_{b,l+1,k} - \frac{1}{2} (k+1) F_{b,l,k+1} \quad (3.20)$$

$$Q_{blk} = 2 \left(2 - \frac{b}{2} - l\right) H_{b,l-1,k} + (1+2l) H_{blk} - (k+1) H_{b,l-1,k+1} \quad (3.22)$$

To get the value of A^+/A^- given in table VI of Ref. 1 using Eq. (4.9), the values of $\tilde{F}(v^*)$ given in table V must be divided by v^* .

In table VII of Ref. 1, the values of Y_3 should read:

$$\begin{aligned} &-2.07825 \times 10^{-3} \\ &3.45588 \times 10^{-2} \end{aligned}$$

All these corrections have been accounted for in Ref. 12 where an updated calculation of the complete critical-to-classical crossover is presented. In addition, we have explicitly verified (see fig. 1) that the errors have had no important consequence on the final results as it could be clearly deduced from a comparison of our estimates of universal amplitude-combinations¹ with those of Guida and Zinn-Justin¹¹ who used the corrected series.

Acknowledgements We thank Prof. H. Kleinert for having incited us to publish, at last, the corrected table I of Feynman graphs.

¹ C. Bagnuls, C. Bervillier, D. I. Meiron and B. G. Nickel, Phys. Rev. **B35**, 3585 (1987).

² B. G. Nickel, D. I. Meiron and G. A. Baker, Jr, “*Compilation of 2-pt and 4-pt graphs for continuous spin models*”, Guelph University preprint, unpublished (1977). May be obtained via the web site http://www.physik.fu-berlin.de/~kleinert/kleiner_reb8/programs/programs.html.

³ C. Bagnuls and C. Bervillier, Phys. Rev. **B32**, 7209 (1985) and in “*Fluctuating Paths and Fields*”, p. 401, Ed. by W. Janke, A. Pelster, H.-J. Schmidt, M. Bachmann (World Scientific, Singapore, 2001).

⁴ S. Weinberg, Phys. Rev. **D8**, 3497 (1973).

⁵ In Ref. 2 Feynman graphs with two and four external legs of a pure φ^4 theory (hence, with regard to similar contributions to the free energy, involving two and four φ^3 vertices respectively) have been exclusively considered. Consequently, when the number of φ^3 vertices exceeds four, new integrals have to be estimated.

⁶ B. R. Heap, J. Math. Phys. **7**, 1582 (1966).

⁷ J. Reeve, A. J. Guttmann and B. Keck, Phys. Rev. **B26**, 3923 (1982).

⁸ J. Reeve, J. Phys. **A15**, L521 (1982).

⁹ F. J. Halfkann and V. Dohm, Z. Phys. **B89**, 79 (1992).

¹⁰ A. Pelissetto, P. Rossi and E. Vicari, Phys. Rev. **E58**, 7146 (1998).

¹¹ R. Guida and J. Zinn-Justin, Nucl. Phys. **B489**, 626 (1997); J. Phys. **A31**, 8103 (1998).

¹² C. Bagnuls and C. Bervillier, “*Classical-to-critical crossovers from field theory*”, hep-th/0112209, submitted to Phys. Rev. E (2001).




FIG. 1. Illustrations on the effective exponent $\gamma_{eff}(t)$ of the (small) effect of the corrections of the errors mentioned in the text. The two uncertainty bounds “max” and “min” corresponding to the resummation criteria of Ref. 1 are displayed. These curves have been obtained using the results of Ref. 12.

TABLE I. Corrected table I. See the caption in the original paper Ref. 1.

b	h	l	m	P_m^{-1}	Value at $d = 3$	b	h	l	m	P_m^{-1}	Value at $d = 3$
1	—	0	1	2	$-4/3$						
2	1	1	1	12	$-2 \ln(y)$	5	49	2	1	48	
						50		2	24		See Eq. (A.13)
3	1	0	1	48	-22.79417368	51		3	24		-0.007497124
					$+16 \ln(y)$						$-0.086950305 \ln(y)$
	2	1	1	8	4.107471254	52		4	32		0.36927278680
	3	2	1	16	0.5194312413	53		5	64		0.36927278680
	4		2	24	0.17390061070	54		6	8		0.22602937610
						55		7	8		0.26394187370
4	3	0	1	48	$-19.73920880 \ln(y)$	56		8	8		$0.97910169300 \times 10^{-1}$
	5	1	1	24	-0.2964527240	57		9	32		0.29097562780
					$-(4/3) \ln(y)$	58		10	8		0.26394187370
	6		2	16	2.0657193571	59		11	16		0.22602937610
	7		3	8	1.7234905497	60		12	16		0.22294544960
	8		4	8	1.2405960978	61		13	4		0.16404726530
						62		14	16		0.19621789690
	9	2	1	16	0.43051311360	63		15	8		$0.94895087300 \times 10^{-1}$
	10		2	4	0.31160313040	64		16	8		0.18956812860
	11		3	4	0.12578653970	65		17	8		$0.93486460600 \times 10^{-1}$
	12		4	8	$0.79516908900 \times 10^{-1}$	66		18	4		$0.94895087300 \times 10^{-1}$
						67		19	16		$0.97910169300 \times 10^{-1}$
	13	3	1	48	0.18361624610	68		20	8		0.18956812860
	14		2	16	$0.80721242900 \times 10^{-1}$	69		21	4		$0.66591760000 \times 10^{-1}$
	15		3	8	$0.37859728700 \times 10^{-1}$	70		22	8		$0.53216192900 \times 10^{-1}$
	16		4	12	$0.14620245800 \times 10^{-1}$	71		23	16		$0.49947540000 \times 10^{-1}$
	17		5	72	$0.12244670000 \times 10^{-1}$	72		24	4		0.17156245110
						73		25	4		$0.72746695800 \times 10^{-1}$
5	15	0	1	144	6.24797746	74		26	2		$0.59161184100 \times 10^{-1}$
					$+4.602913152 \ln(y)$	75		27	4		$0.56505190000 \times 10^{-1}$
					$+4 \ln^2(y)$	76		28	8		$0.89556123600 \times 10^{-1}$
	16		2	128	22.90931839	77		29	12		$0.32950940000 \times 10^{-1}$
	17		3	32	16.60229522	78		30	2		$0.34817630000 \times 10^{-1}$
						79		31	16		$0.34355390400 \times 10^{-1}$
	31	1	1	48	-0.142393552	80	3	1	32		0.16930103539
					$-1.038862482 \ln(y)$	81		2	16		$0.70750283580 \times 10^{-1}$
	32		2	12	-0.122911141	82		3	16		$0.33556467810 \times 10^{-1}$
					$-0.767152192 \ln(y)$	83		4	16		$0.67245000060 \times 10^{-1}$
	33		3	32	1.373092004	84		5	8		$0.70750283580 \times 10^{-1}$
	34		4	16	2.449689513	85		6	8		0.11737660585
	35		5	8	1.072299357	86		7	4		$0.30521594800 \times 10^{-1}$
	36		6	16	0.629580783	87		8	8		$0.33556467810 \times 10^{-1}$
	37		7	8	0.785125191	88		9	4		$0.50133149830 \times 10^{-1}$
	38		8	4	0.729050922	89		10	8		$0.29350397320 \times 10^{-1}$
	39		9	8	0.853563223	90		11	4		$0.30521594800 \times 10^{-1}$
	40		10	4	0.489725124	91		12	4		$0.23674907200 \times 10^{-1}$
	41		11	12	0.31815728						

TABLE II. Corrected table I continued.

b	h	l	m	P_m^{-1}	Value at $d = 3$	b	h	l	m	P_m^{-1}	Value at $d = 3$
5	92	2	13	4	$0.12127180970 \times 10^{-1}$	5	105	4	3	96	$0.27143831355 \times 10^{-1}$
	93		14	2	$0.11619507150 \times 10^{-1}$		106		4	16	$0.16765892110 \times 10^{-1}$
	94		15	8	$0.99110117700 \times 10^{-2}$		107		5	8	$0.11447149611 \times 10^{-1}$
	95		16	8	$0.38967085760 \times 10^{-1}$		108		6	16	$0.13776551025 \times 10^{-1}$
	96		17	4	$0.17420445300 \times 10^{-1}$		109		7	32	$0.10033568785 \times 10^{-1}$
	97		18	4	$0.16062044550 \times 10^{-1}$		110		8	8	$0.65847074272 \times 10^{-2}$
	98		19	4	$0.18506772840 \times 10^{-1}$		111		9	4	$0.41535270157 \times 10^{-2}$
	99		20	2	$0.68328380000 \times 10^{-2}$		112		10	8	$0.47864827139 \times 10^{-2}$
	100		21	8	$0.79486181900 \times 10^{-2}$		113		11	16	$0.36515474403 \times 10^{-2}$
	101		22	4	$0.57948451700 \times 10^{-2}$		114		12	16	$0.31588757411 \times 10^{-2}$
	102		23	4	$0.54193873300 \times 10^{-2}$		115		13	4	$0.16563666713 \times 10^{-2}$
							116		14	12	$0.13970600340 \times 10^{-2}$
	103	4	1	128	$0.99256755397 \times 10^{-1}$		117		15	48	$0.12151455340 \times 10^{-2}$
	104		2	16	$0.34440788678 \times 10^{-1}$		118		16	16	$0.11715681490 \times 10^{-2}$

TABLE III. Modified table III of Ref. 1.

X	S	\tilde{F}	F
1.5	1.0	4.5	-0.25
0.0	0.0	0.0	0.0
0.0	-0.16666666	0.33333333	$-7.1299755 \times 10^{-3}$
2.9928535×10^{-2}	2.9018571×10^{-2}	-0.17775128	3.8703438×10^{-3}
$-2.3069166 \times 10^{-2}$	1.6869642×10^{-3}	6.8988036×10^{-2}	$-3.2942831 \times 10^{-3}$
2.0211486×10^{-2}	4.3368472×10^{-2}	-0.12771112	4.1253272×10^{-3}