Jack superpolynomials, superpartition ordering and determinantal formulas

Patrick Desrosiers*

Département de Physique,

Université Laval,

Québec, Canada, G1K 7P4.

Luc Lapointe[†]

Department of Mathematics and Statistics,

McGill University,

Montréal, Québec H3A 2K6, Canada.

Pierre Mathieu[‡]
Département de Physique,
Université Laval,
Québec, Canada, G1K 7P4.

April 2001

Abstract

We call superpartitions the indices of the eigenfunctions of the supersymmetric extension of the trigonometric Calogero-Moser-Sutherland model. We obtain an ordering on superpartitions from the explicit action of the model's Hamiltonian on monomial superfunctions. This allows to define Jack superpolynomials as the unique eigenfunctions of the model that decompose triangularly, with respect to this ordering, on the basis of monomial superfunctions. This further leads to a simple and explicit determinantal expression for the Jack superpolynomials.

^{*}pdesrosi@phy.ulaval.ca

[†]lapointe@math.mcgill.ca

[‡]pmathieu@phy.ulaval.ca

1 INTRODUCTION 2

1 Introduction

Orthogonal polynomials that are eigenfunctions of Hamiltonians of physical interest certainly deserve consideration. Among this class of orthogonal polynomials are the eigenfunctions of integrable quantum many-body systems of the Calogero-Moser-Sutherland (CMS) type [1, 2, 3, 4], such as the Jack polynomials [5, 6], the eigenfunctions of the trigonometric CMS model.

In [7], we launched the study of a generalized version of Jack polynomials that are eigenfunctions of the supersymmetric extension of the trigonometric CMS model [8] (stCMS model). In addition to the bosonic variables x_i ($i = 1, \dots, N$, where N is the number of particles), this model contains fermionic degrees of freedom described by the variables θ_i . Therefore, the generalized version of Jack polynomials incorporates Grassmannian (i.e., fermionic) variables.

The usual Jack polynomials are not uniquely defined as being eigenfunctions of the trigonometric CMS model. An extra ingredient is required to characterize them: they need to decompose triangularly on the basis of monomial functions. Such a triangular decomposition entails an ordering. In this case, it is the standard dominance ordering [6, 9].

A similar situation holds for the superanalogues: in order to characterize precisely the Jack superpolynomials, it is necessary to require a triangular decomposition in terms of some sort of monomial superfunctions. The first step in this case thus amounts to finding a fermionic extension of the monomial functions. Fortunately, there is a rather natural way of defining the symmetric monomial superfunctions once the general symmetry properties of the stCMS eigenfunctions are understood. These monomial superfunctions are indexed by superpartitions (see [7] and sect. 2), which are, roughly, ordered pairs of partitions (with one of the partitions required to have distinct parts). The second step then reduces to the formulation of a proper ordering among superpartitions allowing to characterize the triangular decomposition of the Jack superpolynomials. Relying on the usual dominance ordering on standard partitions, it proved possible to find such an ordering [7].

Although this dominance ordering is technically sufficient to calculate the Jack superpolynomials, it does not provide the most precise characterization of the Jack superpolynomials. Indeed, it does not rule out monomials that do not actually appear in the decomposition of a given Jack superpolynomial.

The aim of the present paper is to improve this situation by introducing a weaker ordering (i.e., in the sense that fewer elements are comparable) on superpartitions. The basic clue to the discovery of the ordering lies in the following observation. If a triangular decomposition exists for the Jack superpolynomials in terms of the symmetric monomial superfunctions, the action of the Hamiltonian itself on the basis of symmetric monomial superfunctions is likely to be triangular (and the converse is also true). The key point then is that there are usually fewer (and certainly never more) terms in the expansion of the action of the Hamiltonian on a given symmetric supermonomial (in terms of symmetric superpolynomials) than there are in the expansion of this superpolynomial (again in terms of symmetric supermonomials). Therefore, it is simpler to unravel the ordering at work by considering the action of the Hamiltonian on supermonomials. This is how the ordering presented here was found.

The analysis of the action of the Hamiltonian on the supermonomial basis had an important offshoot: all the expansion coefficients could be calculated explicitly. This provided a rather nontrivial generalization of the results of [10], which are recovered here as a special case.

¹In the terminology of this paper, the terms that can appear in the action of the Hamiltonian on the supermonomial basis are obtained from a single rearrangement. In the Jack superpolynomial decomposition, there are in addition terms associated to superpartitions generated from multiple rearrangements.

As already stated, the triangular nature of the action of the Hamiltonian on the supermonomial basis implies a triangular decomposition of the Jack superpolynomials. This, combined with the fact that the superpartitions entering in the decomposition of a given Jack superpolynomial on the monomial superfunction basis are associated to distinct eigenvalues, readily implies that the Jack superpolynomials can be written as determinants. The entries of the determinants being the known eigenvalues and the known expansion coefficients of the action of the Hamiltonian on the supermonomial basis, one thus ends up with a remarkably simple and rather explicit expression for the Jack superpolynomials that generalizes the one obtained in [11] in the non-supersymmetric case.

The article is organized as follows. In section 2, we briefly review the stCMS model as well as some results of [7] that are required for the present article, namely superpartitions, the symmetric supermonomials and the Jack superpolynomials. In section 3 we present our main result: the explicit action of the Hamiltonian on the supermonomial basis. The resulting monomials are characterized by partitions that can be obtained from the original one via the action of a two-particle lowering operator $R_{ij}^{(\ell)}$ (the labels i and j being those of the particles on which the operator acts - with 'intensity' specified by ℓ). The new ordering is also formulated directly in terms of this lowering operator. Section 4 is devoted to computing explicitly the action of the stCMS Hamiltonian on supermonomials and thereby proving the result stated in section 3. In the last section, we reformulate the definition of the Jack superpolynomials in terms of this new ordering and present the determinantal formula.

We would like to stress that all our results provide genuine extensions of non-supersymmetric ones, the latter being associated to the zero-fermion sector. In this sector, a superpartition reduces to a partition and a monomial superfunction reduces to an ordinary monomial symmetric function. In this regard, the detailed proof of the action of the stCMS Hamiltonian on the supermonomial basis includes an explicit proof of the main result of [10].

2 The supersymmetric CMS model and Jack superpolynomials

2.1 The Hamiltonian

The Hamiltonian of the stCMS model reads [8]:

$$\mathcal{H} = -\frac{1}{2} \sum_{i=1}^{N} \partial_{x_i}^2 + \left(\frac{\pi}{L}\right)^2 \sum_{i < j} \frac{\beta(\beta - 1 + \theta_{ij}\theta_{ij}^{\dagger})}{\sin^2(\pi x_{ij}/L)} - \left(\frac{\pi\beta}{L}\right)^2 \frac{N(N^2 - 1)}{6},\tag{1}$$

where $x_{ij} = x_i - x_j$, $\theta_{ij} = \theta_i - \theta_j$ and $\theta_{ij}^{\dagger} = \partial_{\theta_i} - \partial_{\theta_j}$. The θ_i 's, with $i = 1, \dots, N$, are the anticommuting variables that, added to the standard variables x_i , make the model supersymmetric. That is to say, for $i, j \in \{1, \dots, N\}$, we have

$$[x_i, x_j] = 0, \qquad \{\theta_i, \theta_j\} = \theta_i \theta_j + \theta_j \theta_i = 0, \qquad [x_i, \theta_j] = 0. \tag{2}$$

The supersymmetric invariance is enforced in the very construction of this Hamiltonian, i.e., in defining it as the anticommutator of two supersymmetric charges Q and Q^{\dagger} :

$$\mathcal{H} = \frac{1}{2} \{ Q, Q^{\dagger} \} \quad \text{with} \quad Q = \sum_{j} \partial_{\theta_{j}} (\partial_{x_{j}} - i\Phi_{j}(x)), \quad Q^{\dagger} = \sum_{j} \theta_{j} (\partial_{x_{j}} + i\Phi_{j}(x)) \quad (3)$$

where

$$\Phi_j(x) = \frac{\pi\beta}{L} \sum_{k \neq j} \cot\left(\pi \frac{x_{ij}}{L}\right) . \tag{4}$$

The supersymmetric charges are fixed by the requirement that they be nilpotent and that, in the absence of fermionic variables (that is, by setting $\theta_i = \partial_{\theta_i} = 0$), the Hamiltonian reduce to the tCMS Hamiltonian.

An observation that proves to be crucial in [7] is that the term encapsulating the dependency upon the fermionic variables is a fermionic-exchange operator [8]:

$$\kappa_{ij} \equiv 1 - \theta_{ij}\theta_{ij}^{\dagger} = 1 - (\theta_i - \theta_j)(\partial_{\theta_i} - \partial_{\theta_j}). \tag{5}$$

This means that its action on any monomial function $f(\theta_i, \theta_i)$ is such that

$$\kappa_{ij} f(\theta_i, \theta_j) = f(\theta_j, \theta_i) \kappa_{ij}. \tag{6}$$

This allowed the powerful exchange-operator formalism [12] to be used to study the properties of the model in [7]. Note that when we refer to exchange-operators $(\kappa_{ij}, K_{ij}, K_{ij})$ in this article) we understand operators that satisfy relations of the type:

$$\kappa_{ij} = \kappa_{ji} \quad , \quad \kappa_{ij}\kappa_{jk} = \kappa_{ik}\kappa_{ij} = \kappa_{jk}\kappa_{ki} \quad , \quad \kappa_{ij}^2 = 1.$$
(7)

It proves convenient to remove the contribution of the ground-state wave function,

$$\psi_0(x) = \Delta^{\beta}(x) \equiv \prod_{j < k} \sin^{\beta} \left(\frac{\pi x_{jk}}{L} \right) , \tag{8}$$

from the Hamiltonian. The transformed Hamiltonian (which is still supersymmetric) becomes then

$$\bar{\mathcal{H}} \equiv \frac{1}{2} \left(\frac{L}{\pi} \right)^2 \Delta^{-\beta} \mathcal{H} \Delta^{\beta} \,. \tag{9}$$

When written in terms of the new bosonic variables $z_j = e^{2\pi i x_j/L}$, it finally reads

$$\bar{\mathcal{H}} = \sum_{i} (z_i \partial_i)^2 + \beta \sum_{i < j} \frac{z^{ij}}{z_{ij}} (z_i \partial_i - z_j \partial_j) - 2\beta \sum_{i < j} \frac{z_i z_j}{z_{ij}^2} (1 - \kappa_{ij}). \tag{10}$$

where $\partial_i = \partial_{z_i}$ and $z^{ij} = z_i + z_j$. The Jack superpolynomials will be eigenfunctions of this Hamiltonian.

2.2 Superpartitions and the monomial symmetric superfunctions

As explained in [7], since \mathcal{H} leaves invariant the space of polynomials of a given degree in z and a given degree in θ , we can look for eigenfunctions of the form:

$$\mathcal{A}^{(m)}(z,\theta;\beta) = \sum_{1 \le i_1 < i_2 < \dots < i_m \le N} \theta_{i_1} \cdots \theta_{i_m} A^{(i_1 \dots i_m)}(z;\beta), \qquad m = 0, 1, 2, 3, \dots,$$
 (11)

where $A^{(i_1...i_m)}$ is a homogeneous polynomial in z. Due to the presence of m fermionic variables in its expansion, $A^{(m)}$ is said to belong to the m-fermion sector.

The solutions $\mathcal{A}^{(m)}$, being symmetric superpolynomials, must be invariant under the combined action of κ_{ij} and K_{ij} , the exchange operator acting on the z_i variables:

$$K_{ij}f(z_i, z_j) = f(z_j, z_i,)K_{ij}$$
 (12)

In other words, $\mathcal{A}^{(m)}$ must be invariant under the action of \mathcal{K}_{ij} , where

$$\mathcal{K}_{ij} \equiv \kappa_{ij} K_{ij}. \tag{13}$$

Given that the θ products are antisymmetric, the functions $A^{(i_1...i_m)}$ must satisfy

$$K_{ij}A^{(i_1\dots i_m)}(z;\beta) = -A^{(i_1\dots i_m)}(z;\beta) \quad \forall \quad i \text{ and } j \in \{i_1\dots i_m\},$$

$$K_{ij}A^{(i_1\dots i_m)}(z;\beta) = A^{(i_1\dots i_m)}(z;\beta) \quad \forall \quad i \text{ and } j \notin \{i_1\dots i_m\}.$$

$$(14)$$

i.e., $A^{(i_1...i_m)}$ must be completely antisymmetric in the variables $\{z_{i_1}, \ldots, z_{i_m}\}$, and totally symmetric in the remaining variables $z/\{z_{i_1}, \ldots, z_{i_m}\}$.

Superpartitions (to be denoted by capital Greek letters) provide the proper labelling of symmetric superpolynomials [7]. A superpartition in the m-fermion sector is a sequence of integers that generates two standard partitions separated by a semicolon:

$$\Lambda = (\Lambda_1, \dots, \Lambda_m; \Lambda_{m+1}, \dots, \Lambda_N) = (\lambda^a; \lambda^s), \tag{15}$$

the first one being associated to an antisymmetric function, meaning that its parts are all distinct:

$$\lambda^a = (\Lambda_1, \dots, \Lambda_m), \qquad \Lambda_i > \Lambda_{i+1} \ge 0, \qquad i = 1, \dots m - 1, \tag{16}$$

and the second one, to a symmetric function:

$$\lambda^{s} = (\Lambda_{m+1}, \dots, \Lambda_{N}), \qquad \Lambda_{i} \ge \Lambda_{i+1} \ge 0, \quad i = m+1, \dots, N-1, \tag{17}$$

with equal parts then being allowed. In the zero-fermion sector, the semicolon is omitted and Λ reduces to λ^s . We often write the degree of a superpartition as $n = |\Lambda| = \sum_{i=1}^{N} \Lambda_i$. Note finally that to any superpartition Λ , we associate a unique standard partition Λ^* obtained by rearranging the parts of the superpartition in decreasing order. For instance, the rearrangement of $\Lambda = (4, 2, 1; 5, 3, 3, 1)$ is the partition $\Lambda^* = (5, 4, 3, 3, 2, 1, 1)$.

The monomial symmetric superpolynomials have been introduced in [7] as a natural basis of the ring of symmetric superfunctions (we will also refer to this basis as the supermonomial basis):

$$m_{\Lambda}(z,\theta) = m_{(\Lambda_1,\dots,\Lambda_m;\Lambda_{m+1},\dots,\Lambda_N)}(z,\theta) = \sum_{\sigma \in S_N} {'} \theta^{\sigma(1,\dots,m)} z^{\sigma(\Lambda)}, \tag{18}$$

where the prime indicates that the summation is restricted to distinct terms, and where

$$z^{\sigma(\Lambda)} = z_1^{\Lambda_{\sigma(1)}} \cdots z_m^{\Lambda_{\sigma(m)}} z_{m+1}^{\Lambda_{\sigma(m+1)}} \cdots z_N^{\Lambda_{\sigma(N)}} \quad \text{and} \quad \theta^{\sigma(1,\dots,m)} = \theta_{\sigma(1)} \cdots \theta_{\sigma(m)} \,. \tag{19}$$

Given a reduced decomposition $\sigma = \sigma_{i_1} \cdots \sigma_{i_n}$ of an element σ of the symmetric group S_N , let \mathcal{K}_{σ} stand for $\mathcal{K}_{i_1,i_1+1} \cdots \mathcal{K}_{i_n,i_n+1}$. In this notation, we can rewrite the monomial superfunction m_{Λ} as

$$m_{\Lambda} = \frac{1}{f_{\Lambda}} \sum_{\sigma \in S_N} \mathcal{K}_{\sigma} \left(\theta_1 \cdots \theta_m z^{\Lambda} \right) , \qquad (20)$$

where the normalization constant f_{Λ} is

$$f_{\Lambda} = f_{\lambda^s} = n_{\lambda^s}(0)! \, n_{\lambda^s}(1)! \, n_{\lambda^s}(2)! \cdots, \tag{21}$$

with $n_{\lambda^s}(i)$ the number of i's in λ^s , the symmetric part of $\Lambda = (\lambda^a; \lambda^s)$.

2.3 Jack superpolynomials

The Jack superpolynomials can now be defined.

Definition 1. The Jack superpolynomials are the unique functions satisfying the following two properties [7]:

(i):
$$\bar{\mathcal{H}} \mathcal{J}_{\Lambda}(z,\theta;\beta) = \varepsilon_{\Lambda} \mathcal{J}_{\Lambda}(z,\theta;\beta)$$
,
(ii): $\mathcal{J}_{\Lambda}(z,\theta;\beta) = m_{\Lambda}(z,\theta) + \sum_{\Omega: \Omega^{*} < \Lambda^{*}} c_{\Lambda,\Omega}(\beta) m_{\Omega}(z,\theta)$. (22)

In the definition, the ordering on partitions is the usual dominance ordering,

$$\lambda \ge \omega \quad \text{iff} \quad \lambda_1 + \lambda_2 + \dots + \lambda_i \ge \omega_1 + \omega_2 + \dots + \omega_i \,, \quad \forall i \,,$$
 (23)

and the eigenvalue ε_{Λ} , which turns out to be independent of the fermionic sector, is given by:

$$\varepsilon_{\Lambda} = \varepsilon_{\Lambda^*} = \sum_{j} \left[\Lambda_j^{*2} + \beta (N + 1 - 2j) \Lambda_j^* \right]. \tag{24}$$

where $\Lambda_i^* \equiv (\Lambda^*)_i$. The uniqueness property, rather obvious from a computational point of view, is established in sect. 5. Many explicit examples of Jack superpolynomials can be found in [7].

As stressed in [7], the mere existence and uniqueness of functions satisfying the two conditions of Definition 1 is remarkable. Later in this article, we will show that this definition does in fact provide a genuine characterization of the Jack superpolynomials. However, it was also observed that the dominance ordering of the partitions Λ^* is not precise enough in the sense that some monomials m_{Ω} with $\Omega^* < \Lambda^*$ do not appear in the sum. A partial ordering among superpartitions that fixes this problem will be presented in the following section.

3 Action on the monomial basis and ordering of superpartitions

In order to motivate the dominance ordering presented at the end of this section, we first display the explicit action of $\bar{\mathcal{H}}$ on the monomial superpolynomials. This, in turn, will lead to a simple expression for the Jack superpolynomials in the form of determinants (cf. sect. 5).

As in the previous section, let $\Lambda = (\lambda^a; \lambda^s) = (\Lambda_1, \dots, \Lambda_m; \Lambda_{m+1}, \dots, \Lambda_N)$ be a superpartition in the *m*-fermion sector.

In order to characterize the monomials that are generated by the action of $\bar{\mathcal{H}}$ on m_{Λ} , we introduce the operator² $R_{ij}^{(\ell)}$ whose action, for i < j and $\ell \ge 0$, is given by:

$$R_{ij}^{(\ell)}(\Lambda_1, \dots, \Lambda_i, \dots, \Lambda_j, \dots, \Lambda_N) = \begin{cases} (\Lambda_1, \dots, \Lambda_i - \ell, \dots, \Lambda_j + \ell, \dots, \Lambda_N) & \text{if } \Lambda_i > \Lambda_j, \\ (\Lambda_1, \dots, \Lambda_i + \ell, \dots, \Lambda_j - \ell, \dots, \Lambda_N) & \text{if } \Lambda_j > \Lambda_i. \end{cases}$$
(25)

²The non-supersymmetric version of this operator was introduced in [9] in the case $\ell = 1$ (more precisely, it is the raising version of the operator that is used), and in [11] in the general case.

This action of $R_{ij}^{(\ell)}$ is non-zero only in the following cases:

$$\begin{split} & \mathrm{I}: \qquad i,j \in \{1,\ldots,m\} \quad \text{and} \quad \lfloor \frac{\Lambda_i - \Lambda_j - 1}{2} \rfloor \geq \ell \,, \\ & \mathrm{II}: \qquad i \in \{1,\ldots,m\} \,, j \in \{m+1,\ldots,N\} \quad \text{and} \quad |\Lambda_i - \Lambda_j| - 1 \geq \ell \,, \\ & \mathrm{III}: \qquad i,j \in \{m+1,\ldots,N\} \quad \text{and} \quad \lfloor \frac{\Lambda_i - \Lambda_j}{2} \rfloor \geq \ell \,. \end{split} \tag{26}$$

where $\lfloor x \rfloor$ stands for the largest integer smaller or equal to x. Otherwise, it is understood that $R_{ij}^{(\ell)}$ annihilates the superpartition Λ , i.e., $R_{ij}^{(\ell)} \Lambda = \emptyset$. In the following, we will say that a pair (i,j) is of type I if $i,j \in \{1,\ldots,m\}$, of type II if $i \in \{1,\ldots,m\}$ and $j \in \{m+1,\ldots,N\}$, and of type III if $i,j \in \{m+1,\ldots,N\}$.

In the *m*-fermion sector, given a sequence $\gamma = (\gamma_1, \dots, \gamma_m; \gamma_{m+1}, \dots, \gamma_N)$, we will denote by $\overline{\gamma}$ the superpartition whose antisymmetric part is the rearrangement of $(\gamma_1, \dots, \gamma_m)$ and whose symmetric part is the rearrangement of $(\gamma_{m+1}, \dots, \gamma_N)$. For example, we have

$$\overline{(1,3,2;2,3,1,2)} = (3,2,1;3,2,2,1).$$
(27)

Also, σ_{γ} will stand for the element of S_N that sends γ to $\overline{\gamma}$, that is $\sigma_{\gamma}\gamma = \overline{\gamma}$. Note that we can always choose σ_{γ} such that $\sigma_{\gamma} = \sigma_{\gamma}^a \sigma_{\gamma}^s$, with σ_{γ}^a and σ_{γ}^s permutations of $\{1, \ldots, m\}$ and $\{m+1, \ldots, N\}$ respectively.

We now state the explicit action of the stCMS Hamiltonian on monomials. The proof will be presented in the following section.

Theorem 2. $\bar{\mathcal{H}}$ acts on $m_{\Lambda}(z,\theta)$ as follows:

$$\bar{\mathcal{H}} \, m_{\Lambda}(z,\theta) = \varepsilon_{\Lambda} m_{\Lambda}(z,\theta) + \sum_{\Omega \neq \Lambda} v_{\Lambda\Omega}(\beta) \, m_{\Omega}(z,\theta) \tag{28}$$

where ε_{Λ} is given in (24) and $v_{\Lambda\Omega}(\beta)$ is non-zero only if $\Omega = (\omega^a; \omega^s) = \overline{R_{ij}^{(\ell)} \Lambda}$, for a given $R_{ij}^{(\ell)}$ with $\ell > 0$. In this case, the coefficient $v_{\Lambda\Omega}(\beta)$ reads:

$$v_{\Lambda\Omega}(\beta) = \begin{cases} 2\beta \operatorname{sgn}(\sigma_{R_{ij}^{(\ell)}\Lambda}^{a}) (\Lambda_i - \Lambda_j - \delta\ell) n(\Lambda_i - \ell, \Lambda_j + \ell) & \text{if } \Lambda_i > \Lambda_j \\ 2\beta \operatorname{sgn}(\sigma_{R_{ij}^{(\ell)}\Lambda}^{a}) (\Lambda_j - \Lambda_i - \delta\ell) n(\Lambda_i + \ell, \Lambda_j - \ell) & \text{if } \Lambda_j > \Lambda_i \end{cases}$$
(29)

where $\operatorname{sgn}(\sigma_{R_{ij}^{(\ell)}\Lambda}^a)$ stands for the sign of the permutation $\sigma_{R_{ij}^{(\ell)}\Lambda}^a$. The parameter δ is equal to 2,1 or 0 if the pair (i,j) is of type I, II or III in (26) respectively. Finally n(a,b) is a symmetry factor given by:

$$n(a,b) = \begin{cases} 1 & i, j \text{ of type I} \\ n_{\omega^s}(b) & i, j \text{ of type II} \\ n_{\omega^s}(a)n_{\omega^s}(b) & i, j \text{ of type III and } a \neq b \\ \frac{1}{2}n_{\omega^s}(a)\left(n_{\omega^s}(a) - 1\right) & i, j \text{ of type III and } a = b \end{cases}$$
 (30)

Since in (29), $1 \le \ell \le \max\{\Lambda_i, \Lambda_j\}$, the symmetry factor n(a, b) of the non-diagonal coefficients is never invoked with arguments a or b equal to zero. This has the important consequence that the coefficients $v_{\Lambda\Omega}(\beta)$ in (28) do not depend on the number N of variables, as long as N is large enough so that $m_{\Omega}(z,\theta) \ne 0$. Indeed, with $\Omega = (\omega^a; \omega^s)$, the number of variables only influences the number of parts equal to zero in the partition ω^s , which as we said does not influence the value of the symmetry factor n(a,b).

Remark 3. If $\Omega = \overline{R_{ij}^{(\ell)}\Lambda} = \overline{R_{i'j'}^{(\ell')}\Lambda}$, we must have $\ell = \ell'$, $\Lambda_i = \Lambda_{i'}$ and $\Lambda_j = \Lambda_{j'}$. The choice of i, j or i', j' is thus irrelevant in the computation of $v_{\Lambda\Omega}(\beta)$.

We now look at an example. For $N \geq 5$, we have

$$\bar{\mathcal{H}} m_{(2,1;4,2)} = \varepsilon_{(2,1;4,2)} m_{(2,1;4,2)} + 2\beta m_{(3,1;3,2)} - 4\beta m_{(3,2;2,2)} + 4\beta m_{(2,1;3,3)} + 8\beta m_{(2,1;3,2,1)} + 24\beta m_{(2,1;2,2,2)} + 4\beta m_{(2,1;4,1,1)}.$$
(31)

The coefficient -4β in front of $m_{(3,2;2,2)}$ is explained first by identifying the proper rearrangement that links (3,2;2,2) to (2,1;4,2), namely

$$(3,2;2,2) = \overline{R_{2,3}^{(2)}(2,1;4,2)} = \overline{(2,3;2,2)}$$
(32)

(which is a type-II case) and then by evaluating the corresponding coefficient $v_{\Lambda,\Omega}$ from Theorem 2:

$$2\beta \operatorname{sgn}(\sigma_{(2,3;2,2)}^a)(4-1-2)n(3,2) = -4\beta, \tag{33}$$

since $n(3,2) = n_{(2,2)}(2) = 2$ (there are two 2's in (2,2)) and the sign of the permutation that sends (2,3) to (3,2) is -1.

Knowing the action of $\bar{\mathcal{H}}$ on the monomial basis allows us to define a partial ordering on superpartitions.

Definition 4. We say that $\Lambda \geq^s \Omega$ iff $\Omega = \overline{R_{i_k,j_k}^{(\ell_k)} \dots \overline{R_{i_1,j_1}^{(\ell_1)} \Lambda}}$, for a given sequence of operators $R_{i_1,j_1}^{(\ell_1)},\dots,R_{i_k,j_k}^{(\ell_k)}$ (the equality occurs in the case of the null sequence).

Proposition 5. The relation \geq^s provides a partial ordering on superpartitions.

Proof: To show that $>^s$ is a genuine ordering, we have to verify the following three properties:

- (1) reflexivity: $\Lambda \geq^s \Lambda$;
- (2) transitivity: if $\Gamma \geq^s \Omega$ and $\Omega \geq^s \Lambda$, then $\Gamma \geq^s \Lambda$;
- (3) antisymmetry: if $\Lambda \geq^s \Omega$ and $\Omega \geq^s \Lambda$, then $\Lambda = \Omega$.

Reflexivity follows from the consideration of a null sequence of lowering operators. The transitivity property is also immediate since two sequences can be glued together to form a longer sequence. In order to prove antisymmetry, we first make the following two simple observations from (25):

$$(i): \Lambda \geq^s \Omega \Longrightarrow \Lambda^* \geq \Omega^*, \quad (ii): \Lambda \geq^s \Omega \quad \text{and} \quad \Lambda^* = \Omega^* \Longrightarrow \Lambda = \Omega,$$
 (34)

where as usual Λ^* and Ω^* are the partitions associated to the superpartitions Λ and Ω respectively. Now, let $\Lambda \geq^s \Omega$ and $\Omega \geq^s \Lambda$. Then, from the first observation, $\Lambda^* \geq \Omega^*$ and $\Omega^* \geq \Lambda^*$, i.e., $\Lambda^* = \Omega^*$. Therefore, we have $\Lambda \geq^s \Omega$ and $\Lambda^* = \Omega^*$, which lead to $\Lambda = \Omega$ from the second observation.

Using this new partial ordering on superpartitions, we now have that the action (28) of $\bar{\mathcal{H}}$ on $m_{\Lambda}(z,\theta)$ is triangular.

Corollary 6.

$$\bar{\mathcal{H}} \, m_{\Lambda}(z,\theta) = \varepsilon_{\Lambda} m_{\Lambda}(z,\theta) + \sum_{\Omega < s_{\Lambda}} v_{\Lambda\Omega}(\beta) \, m_{\Omega}(z,\theta) \,. \tag{35}$$

Notice that not all Ω 's such that $\Omega <^s \Lambda$ appear in this expansion with a non-zero coefficient. Only those that can be obtained from Λ by the application of a *single* lowering operator have non-zero coefficients.

³The dominance ordering on partitions is a special case of this ordering on superpartitions. We draw attention to the technical importance of formulating an ordering using the lowering operator $R_{ij}^{(\ell)}$, especially in the case of the usual dominance ordering (as opposed to its formulation in terms of inequalities).

4 The action of $\overline{\mathcal{H}}$ on the monomial basis: a detailed proof

We will first focus on the non-diagonal coefficients in the action of \mathcal{H} on $m_{\Lambda}(z,\theta)$, that is, on the coefficients $v_{\Lambda,\Omega}$ for $\Lambda \neq \Omega$, and wait until the end of the section before computing the diagonal coefficient ε_{Λ} . We will show that the coefficients $v_{\Lambda,\Omega}$ are indeed given by the expression displayed in Theorem 2 for the special case N=2, for the pairs (i,j) of types I, II and III. And, in a second step, we will prove the theorem in general. The reason why the two-particle case plays such a central role is that the operator $R_{ij}^{(\ell)}$ acts only on two parts of a partition, that is, on N=2 particles.

4.1 Non-diagonal coefficients: the case N=2

We will use the following notation:

$$B_{ij} = \frac{z^{ij}}{z_{ij}} (z_i \partial_i - z_j \partial_j), \qquad F_{ij} = -2 \frac{z_i z_j}{z_{ij}^2} (1 - \kappa_{ij}),$$
 (36)

(B for boson and F for fermion). The part of the Hamiltonian acting non-diagonally on the monomial basis is $\beta(B+F)$, where

$$B = \sum_{1 \le i < j \le N} B_{ij}, \qquad F = \sum_{1 \le i < j \le N} F_{ij}.$$
 (37)

In order to simplify the notation, we will set $\Lambda_1 = r$, $\Lambda_2 = s$.

Case III:

In this case, we must have m=0. This gives $\Lambda=(r,s)=(r,s)$, and

$$m_{(r,s)} = \frac{1}{f_{(r,s)}} (1 + \mathcal{K}_{12}) z_1^r z_2^s = \frac{1}{f_{(r,s)}} (z_1^r z_2^s + z_1^s z_2^r).$$
 (38)

Clearly, the action of F_{12} vanishes since there are no θ terms. The action of B_{12} is simply

$$B_{12}m_{(r,s)} = \frac{1}{f_{(r,s)}} \frac{z^{12}}{z_{12}} (r-s)(z_1 z_2)^s (z_1^{r-s} - z_2^{r-s})$$

$$= \frac{z^{12}}{z_{12}} (r-s)(z_1 z_2)^s z_{12} (z_1^{r-s-1} + z_1^{r-s-2} z_2 + \dots + z_1 z_2^{r-s-2} + z_2^{r-s-1})$$

$$= (r-s)(z_1 z_2)^s (z_1^{r-s} + 2[z_1^{r-s-1} z_2 + \dots + z_1 z_2^{r-s-1}] + z_2^{r-s})$$
(39)

where we have set $f_{(r,s)} = 1$, since the action vanishes in the case r = s. Therefore, by recombining to construct monomial symmetric functions, we get

$$\beta B_{12} m_{(r,s)} = \beta B m_{(r,s)} = \beta (r-s) m_{(r,s)} + \sum_{\ell=1}^{\lfloor (r-s)/2 \rfloor} 2\beta (r-s) m_{(r-\ell,s+\ell)}. \tag{40}$$

If we let $\Omega = (r - \ell, s + \ell)$, we have $\Omega = R_{12}^{(\ell)} \Lambda = \overline{R_{12}^{(\ell)} \Lambda}$. We thus recover the right value $2\beta(\Lambda_1 - \Lambda_2) = 2\beta(r - s)$ for the coefficient $v_{\Lambda,\Omega}$, for $1 \le \ell \le \lfloor (\Lambda_1 - \Lambda_2)/2 \rfloor$ (see (29) with $\delta = 0$).

Case II:

We now have m=1. Consider first the action of B+F on $m_{(r;s)}$, supposing that r>s. Given

$$m_{(r;s)} = \frac{1}{f_{(r;s)}} (1 + \mathcal{K}_{12}) \,\theta_1 z_1^r z_2^s = \theta_1 z_1^r z_2^s + \theta_2 z_1^s z_2^r \tag{41}$$

(since $f_{(r;s)} = 1$), we get

$$B_{12}m_{(r;s)} = \frac{z^{12}}{z_{12}}(r-s)(z_1z_2)^s(\theta_1z_1^{r-s} - \theta_2z_2^{r-s})$$
(42)

and

$$F_{12}m_{(r;s)} = -2\frac{z_1 z_2}{z_{12}^2} (\theta_1 - \theta_2)(z_1^r z_2^s - z_1^s z_2^r)$$

$$= -\frac{2}{z_{12}} (\theta_1 - \theta_2)(z_1 z_2)^s (z_1^{r-s} z_2 + \dots + z_1 z_2^{r-s}).$$
(43)

In the sum $B_{12} + F_{12}$, we now concentrate on the term θ_1 :

$$(B_{12} + F_{12})m_{(r;s)}\big|_{\theta_1} = \frac{\theta_1}{z_{12}}(z_1 z_2)^s \{ (r-s)z^{12}z_1^{r-s} - 2(z_1^{r-s}z_2 + \dots + z_1 z_2^{r-s}) \}.$$
 (44)

To proceed further, we use the following identity proved in the appendix.

Identity 7.

$$\{(r-s)z^{12}z_1^{r-s} - 2(z_1^{r-s}z_2 + \dots + z_1z_2^{r-s})\} = z_{12}\{(r-s)z_1^{r-s} + \sum_{\ell=1}^{r-s-1} 2(r-s-\ell)z_1^{r-s-\ell}z_2^{\ell}\}.$$
(45)

Taking into account the θ_2 terms (which are recovered by symmetry from the θ_1 terms under $z_1 \leftrightarrow z_2$), we have:

$$\beta(B_{12} + F_{12})m_{(r;s)} = \beta(B + F)m_{(r;s)} = \beta(r - s)m_{(r;s)} + \sum_{\ell=1}^{r-s-1} 2\beta(r - s - \ell)m_{(r-\ell;s+\ell)}. \tag{46}$$

We thus get the right coefficient, namely $2\beta(\Lambda_1 - \Lambda_2 - \delta\ell)$, with $\delta = 1$. Moreover, the upper limit on ℓ shows that $\Lambda_1 - \Lambda_2 > \ell$, as it should. The derivation in the case r < s is identical.

Case I:

Since we now have m=2, we consider the action of B+F on $m_{(r,s;0)}$. Given that, for r>s,

$$m_{(r,s;0)} = \frac{1}{f_{(r,s;0)}} (1 + \mathcal{K}_{12}) \,\theta_1 \theta_2 z_1^r z_2^s = \theta_1 \theta_2 (z_1^r z_2^s - z_1^s z_2^r) \,, \tag{47}$$

a direct computation yields

$$B_{12}m_{(r,s;0)} = \theta_1 \theta_2 \frac{z^{12}}{z_{12}} (r-s)(z_1 z_2)^s (z_1^{r-s} + z_2^{r-s}), \qquad (48)$$

and since $(1 - \kappa_{12})\theta_1\theta_2 = 2\theta_1\theta_2$,

$$F_{12}m_{(r,s;0)} = -4\theta_1\theta_2 \frac{z_1 z_2}{z_{12}^2} (z_1^r z_2^s - z_1^s z_2^r)$$

$$= -4\theta_1\theta_2 \frac{z_1 z_2}{z_{12}^2} (z_1 z_2)^s z_{12} (z_1^{r-s-1} + z_1^{r-s-2} z_2 + \dots + z_2^{r-s-1})$$

$$= -4\theta_1\theta_2 \frac{1}{z_{12}} (z_1 z_2)^s (z_1^{r-s} z_2 + z_1^{r-s-1} z_2^2 + \dots + z_1 z_2^{r-s}). \tag{49}$$

Therefore, we obtain

$$(B_{12} + F_{12})m_{(r,s;0)} = \frac{\theta_1\theta_2}{z_{12}}(z_1z_2)^s \{ (r-s)z^{12}(z_1^{r-s} + z_2^{r-s}) - 4(z_1^{r-s}z_2 + \dots + z_1z_2^{r-s}) \}.$$
 (50)

To proceed, we need the following identity also proved in the appendix.

Identity 8.

$$\{(r-s)z^{12}(z_1^{r-s}+z_2^{r-s})-4(z_1^{r-s}z_2+\cdots+z_1z_2^{r-s})\}$$

$$=z_{12}\{(r-s)(z_1^{r-s}-z_2^{r-s})+\sum_{\ell=1}^{\lfloor (r-s-1)/2\rfloor}2(r-s-2\ell)(z_1^{r-s-\ell}z_2^{\ell}-z_1^{\ell}z_2^{r-s-\ell})\}. (51)$$

Using the identity, we get

$$\beta(B_{12} + F_{12})m_{(r,s;0)} = \beta(B+F)m_{(r,s;0)} = \beta(r-s)m_{(r,s;0)} + \sum_{\ell=1}^{\lfloor (r-s-1)/2\rfloor} 2\beta(r-s-2\ell)m_{(r-\ell,s+\ell;0)}.$$
(52)

From this expression, the value of the coefficient $v_{\Lambda,\Omega}$ is given immediately. It is indeed $2\beta(\Lambda_1 - \Lambda_2 - \delta\ell)$, with $\delta = 2$. The upper limit in the summation implies that $\lfloor (\Lambda_1 - \Lambda_2 - 1)/2 \rfloor \geq \ell$, as it should.

4.2 Non-diagonal coefficients: the case N > 2

Let Λ_{ij} denote the restriction of a superpartition to its i^{th} and j^{th} entries, that is,

$$\Lambda_{ij} = \begin{cases}
(\Lambda_i, \Lambda_j; 0) & \text{if } i, j \text{ is of type I} \\
(\Lambda_i; \Lambda_j) & \text{if } i, j \text{ is of type II} \\
(\Lambda_i, \Lambda_j) & \text{if } i, j \text{ is of type III}
\end{cases}$$
(53)

Also, in accordance with (26), we define S_{ij}^{Λ} , to be

$$S_{ij}^{\Lambda} = \begin{cases} \lfloor \frac{\Lambda_i - \Lambda_j - 1}{2} \rfloor & \text{if } i, j \text{ of type I} \\ |\Lambda_i - \Lambda_j| - 1 & \text{if } i, j \text{ of type II} \\ |\frac{\Lambda_i - \Lambda_j}{2}| & \text{if } i, j \text{ of type III} \end{cases}$$
(54)

For an arbitrary number of particules, we have thus

$$\beta(B+F) m_{\Lambda} = \beta \sum_{1 \leq i < j \leq N} (B_{ij} + F_{ij}) \frac{1}{f_{\Lambda}} \sum_{\sigma \in S_{N}} \mathcal{K}_{\sigma} \left(\theta_{1} \cdots \theta_{m} z^{\Lambda} \right)$$

$$= \frac{\beta}{f_{\Lambda}} \sum_{\sigma \in S_{N}} \mathcal{K}_{\sigma} \left(\sum_{1 \leq i < j \leq N} (B_{ij} + F_{ij}) \theta_{1} \cdots \theta_{m} z^{\Lambda} \right)$$

$$= \frac{\beta}{f_{\Lambda}} \sum_{1 \leq i < j \leq N} \sum_{\sigma \in S_{N}} \mathcal{K}_{\sigma} \left((B_{ij} + F_{ij}) \theta_{1} \cdots \theta_{m} z^{\Lambda} \right)$$

$$= \frac{1}{2} \frac{\beta}{f_{\Lambda}} \sum_{1 \leq i < j \leq N} \sum_{\sigma \in S_{N}} \mathcal{K}_{\sigma} \left((B_{ij} + F_{ij}) (1 + \mathcal{K}_{ij}) \theta_{1} \cdots \theta_{m} z^{\Lambda} \right)$$

$$= \frac{1}{2} \frac{1}{f_{\Lambda}} \sum_{1 \leq i < j \leq N} \sum_{\ell = 0}^{S_{ij}^{\Lambda}} \mathcal{V}_{\Lambda_{ij}} \left(R_{ij}^{(\ell)} \Lambda \right)_{ij} \sum_{\sigma \in S_{N}} \mathcal{K}_{\sigma} \left(\frac{1}{f \left(R_{ij}^{(\ell)} \Lambda \right)_{ij}} (1 + \mathcal{K}_{ij}) \theta_{1} \cdots \theta_{m} z^{R_{ij}^{(\ell)}} \Lambda \right)$$

$$= \frac{1}{f_{\Lambda}} \sum_{1 \leq i < j \leq N} \sum_{\ell = 0}^{S_{ij}^{\Lambda}} \frac{\mathcal{V}_{\Lambda_{ij}} \left(R_{ij}^{(\ell)} \Lambda \right)_{ij}}{f \left(R_{ij}^{(\ell)} \Lambda \right)_{ij}} \sum_{\sigma \in S_{N}} \mathcal{K}_{\sigma} \left(\theta_{1} \cdots \theta_{m} z^{R_{ij}^{(\ell)}} \Lambda \right)$$

$$= \frac{1}{f_{\Lambda}} \sum_{1 \leq i < j \leq N} \sum_{\ell = 0}^{S_{ij}^{\Lambda}} \frac{\mathcal{V}_{\Lambda_{ij}} \left(R_{ij}^{(\ell)} \Lambda \right)_{ij}}{f \left(R_{ij}^{(\ell)} \Lambda \right)_{ij}} \sum_{\sigma \in S_{N}} \sup \left(\sigma_{R_{ij}^{(\ell)}}^{\alpha} \Lambda \right) \mathcal{K}_{\sigma} \mathcal{K}_{\sigma_{R_{ij}^{\ell}}^{\alpha}} \mathcal{K}_{\sigma_{R_{ij}^{\ell}}^{\beta}} \left(\theta_{1} \cdots \theta_{m} z^{R_{ij}^{\ell}} \Lambda \right)$$

$$= \frac{1}{f_{\Lambda}} \sum_{1 \leq i < j \leq N} \sum_{\ell = 0}^{S_{ij}^{\Lambda}} \sup \left(\sigma_{R_{ij}^{\ell}}^{\alpha} \Lambda \right) \frac{\mathcal{V}_{\Lambda_{ij}} \left(R_{ij}^{(\ell)} \Lambda \right)_{ij}}{f \left(R_{ij}^{(\ell)} \Lambda \right)_{ij}} \sum_{\sigma' \in S_{N}} \mathcal{K}_{\sigma'} \left(\theta_{1} \cdots \theta_{m} z^{R_{ij}^{\ell}} \Lambda \right)$$

$$= \frac{1}{f_{\Lambda}} \sum_{1 \leq i < j \leq N} \sum_{\ell = 0}^{S_{ij}^{\Lambda}} \sup \left(\sigma_{R_{ij}^{\ell}}^{\alpha} \Lambda \right) \frac{\mathcal{V}_{\Lambda_{ij}} \left(R_{ij}^{\ell} \Lambda \right)_{ij}}{f \left(R_{ij}^{\ell} \Lambda \right)_{ij}} \int_{\sigma' \in S_{N}} \mathcal{K}_{\sigma'} \left(\theta_{1} \cdots \theta_{m} z^{R_{ij}^{\ell}} \Lambda \right)$$

$$= \frac{1}{f_{\Lambda}} \sum_{1 \leq i < j \leq N} \sum_{\ell = 0}^{S_{ij}^{\Lambda}} \sup \left(\sigma_{R_{ij}^{\ell}}^{\alpha} \Lambda \right) \frac{\mathcal{V}_{\Lambda_{ij}} \left(R_{ij}^{\ell} \Lambda \right)_{ij}}{f \left(R_{ij}^{\ell} \Lambda \right)_{ij}} \int_{\sigma' \in S_{N}} \mathcal{K}_{\sigma'} \left(\theta_{1} \cdots \theta_{m} z^{R_{ij}^{\ell}} \Lambda \right)$$

$$= \frac{1}{f_{\Lambda}} \sum_{1 \leq i < j \leq N} \sum_{\ell = 0}^{S_{ij}^{\Lambda}} \sup \left(\sigma_{R_{ij}^{\ell}}^{\alpha} \Lambda \right) \frac{\mathcal{V}_{\Lambda_{ij}} \left(R_{ij}^{\ell} \Lambda \right)_{ij}}{f \left(R_{ij}^{\ell} \Lambda \right)_{ij}} \int_{\sigma' \in S_{N}} \mathcal{K}_{\sigma'} \left(\theta_{1} \cdots \theta_{m} z^{R_{ij}^{\ell}} \Lambda \right)$$

$$= \frac{1}{f_{\Lambda}} \sum_{1 \leq i < j \leq N} \sum_{\ell = 0}^{S_{ij}} \sup$$

Let us be more specific about some of these steps. In the fourth equality, we have introduced a factor $(1+\mathcal{K}_{ij})/2$ in order to generate two terms out of $\theta_1\cdots\theta_mz^{\Lambda}$ because the action of $B_{ij}+F_{ij}$ is defined on two terms (cf. eqs (38), (41) and (47)). We have thus generated an intermediate N=2 problem in order to use the N=2 results derived previously. The factor β is now reabsorbed in the coefficient $v_{\Lambda_{ij}\left(R_{ij}^{(\ell)}\Lambda\right)}$:

$$v_{\Lambda_{ij}\left(R_{ij}^{(\ell)}\Lambda\right)_{ij}} = 2\beta \left(|\Lambda_i - \Lambda_j| - \delta\ell\right), \qquad \ell \neq 0.$$
 (56)

Observe that the non-diagonal coefficients correspond to the cases where $\ell > 0$, but that the action of B+F also includes a diagonal contribution, to be analysed later. In the sixth equality, we reexpress the outcome in terms of a single term. However, the ordering of the variables may not be adequate for this to be a genuine supermonomial leading term. Some reordering may be required, forcing the appearance of extra factors $\mathcal{K}_{\sigma^a}\mathcal{K}_{\sigma^s}$ in the next equality. Moreover, reordering the anticommuting variables may generate a sign given by the factor $\mathrm{sgn}(\sigma^a)$. In the following step, we redefine the summation variable as follows: $\sigma' = \sigma \sigma^a \sigma^s$. In the final equality, we simply use the definition of supermonomials.

We are now in a position to compute the coefficient $\underline{v_{\Lambda\Omega}}$. Let $\#(\Omega,\Lambda)$ denote the number of distinct ways we can choose (i,j), i < j, such that $\Omega = R_{ij}^{(\ell)}\Lambda$. Note that such "symmetries" can only occur on the symmetric side, that is, if there is more than one possible choice of i (or j), then i > m (or j > m). Therefore, $\mathrm{sgn}(\sigma_{R_{ij}^{(\ell)}\Lambda}^a)$ is independent of the particular choice of the pair (i,j) leading to Ω , since the underlying degeneracy is independent of the antisymmetric side. The coefficient of m_{Ω} is thus

$$v_{\Lambda\Omega} = \frac{1}{f_{\Lambda}} \operatorname{sgn}(\sigma_{R_{ij}^{(\ell)}\Lambda}^{a}) \frac{v_{\Lambda_{ij}\left(R_{ij}^{(\ell)}\Lambda\right)_{ij}}}{f_{\left(R_{ij}^{(\ell)}\Lambda\right)_{ij}}} f_{\Omega} \#(\Omega,\Lambda).$$
(57)

The factor $\#(\Omega, \Lambda)$ comes from the different values of the pair (i, j) leading to the same superpartition Ω . From Remark 3, these distinct values all lead to the same coefficient $v_{\Lambda_{ij}\left(R_{ij}^{(\ell)}\Lambda\right)_{ij}}/f_{\left(R_{ij}^{(\ell)}\Lambda\right)_{ij}}$. Since it has also just been mentioned that these different choices do not affect the value of $\operatorname{sgn}(\sigma_{R_{ij}^{(\ell)}\Lambda}^a)$, $v_{\Lambda\Omega}$ is simply given by the coefficient of $m_{\overline{R_{ij}^{(\ell)}\Lambda}}$ in (55) (disregarding the sum) multiplied by $\#(\Omega, \Lambda)$.

Let us now evaluate explicitly the various factors entering in the expression of $v_{\Lambda\Omega}$, for the three cases treated separately. We first consider case III, that is, the case where $i,j\in\{m+1,\ldots,N\}$. If $\Lambda_i-\ell\neq\Lambda_j+\ell$, when going from Λ to Ω , the number of Λ_i and Λ_j decreases by 1, whereas the number of $\Lambda_i-\ell$ and $\Lambda_j+\ell$ increases by 1 (while the other Λ_k 's remain unaffected). Therefore, the ratio f_{Ω}/f_{Λ} depends only upon the multiplicity factors involving these parts, that is,

$$\frac{f_{\Omega}}{f_{\Lambda}} = \frac{n_{\omega^s}(\Lambda_i - \ell)! \, n_{\omega^s}(\Lambda_j + \ell)! \, n_{\omega^s}(\Lambda_i)! \, n_{\omega^s}(\Lambda_j)!}{n_{\lambda^s}(\Lambda_i - \ell)! \, n_{\lambda^s}(\Lambda_j + \ell)! \, n_{\lambda^s}(\Lambda_i)! \, n_{\lambda^s}(\Lambda_j)!}$$

$$= \frac{n_{\omega^s}(\Lambda_i - \ell)! \, n_{\omega^s}(\Lambda_j + \ell)! \, n_{\omega^s}(\Lambda_j)! \, n_{\omega^s}(\Lambda_j)!}{(n_{\omega^s}(\Lambda_i - \ell) - 1)! \, (n_{\omega^s}(\Lambda_j + \ell) - 1)! \, (n_{\omega^s}(\Lambda_i) + 1)! (n_{\omega^s}(\Lambda_j) + 1)!}$$

$$= \frac{n_{\omega^s}(\Lambda_i - \ell) n_{\omega^s}(\Lambda_j + \ell)}{(n_{\omega^s}(\Lambda_i) + 1) (n_{\omega^s}(\Lambda_j) + 1)}.$$
(58)

Finally, since in that case $\#(\Omega, \Lambda)$ is equal to $(n_{\omega^s}(\Lambda_i) + 1)(n_{\omega^s}(\Lambda_j) + 1)$ (the product of the number of Λ_i 's and Λ_j 's in the symmetric part of Λ), and since $f_{(R_{ij}^{(\ell)}\Lambda)_{i,j}} = 1$, we get

$$v_{\Lambda\Omega} = \operatorname{sgn}(\sigma_{R_{ij}^{(\ell)}\Lambda}^a) v_{\Lambda_{ij}(R_{ij}^{(\ell)}\Lambda)_{ij}} n_{\omega^s}(\Lambda_i - \ell) n_{\omega^s}(\Lambda_j + \ell), \qquad (59)$$

which agrees with the expression given in Theorem 2. In the case where $\Lambda_i - \ell = \Lambda_j + \ell$, when going from Λ to Ω , the number of Λ_i and Λ_j decreases by 1, whereas the number of $\Lambda_i - \ell$ increases by 2, that is,

$$\frac{f_{\Omega}}{f_{\Lambda}} = \frac{n_{\omega^s}(\Lambda_i - \ell) \left(n_{\omega^s}(\Lambda_i - \ell) - 1 \right)}{\left(n_{\omega^s}(\Lambda_i) + 1 \right) \left(n_{\omega^s}(\Lambda_j) + 1 \right)}.$$
 (60)

The number $\#(\Omega, \Lambda)$ is still $(n_{\omega^s}(\Lambda_i) + 1)(n_{\omega^s}(\Lambda_j) + 1)$, but now $f_{(R_{ij}^{(\ell)}\Lambda)_{ij}} = 2$, giving

$$v_{\Lambda\Omega} = \frac{1}{2} \operatorname{sgn}(\sigma_{R_{ij}^{(\ell)}\Lambda}^{a}) v_{\Lambda_{ij}(R_{ij}^{(\ell)}\Lambda)_{ii}} n_{\omega^{s}}(\Lambda_{i} - \ell) (n_{\omega^{s}}(\Lambda_{i} - \ell) - 1),$$

$$(61)$$

as announced.

Let us now consider case II with $\Lambda_i > \Lambda_j$. This case is similar to case III with $\Lambda_i - \ell \neq \Lambda_j + \ell$. The only difference is that since Λ_i now belongs to the antisymmetric sector, it does not influence the value of f_{Ω}/f_{Λ} . Therefore, in this case,

$$\frac{f_{\Omega}}{f_{\Lambda}} = \frac{n_{\omega^s}(\Lambda_j + \ell)}{n_{\omega^s}(\Lambda_j) + 1},\tag{62}$$

and

$$v_{\Lambda\Omega} = \operatorname{sgn}(\sigma_{R_{ij}^{(\ell)}\Lambda}^a) v_{\Lambda_{ij}(R_{ij}^{(\ell)}\Lambda)_{ij}} n_{\omega^s}(\Lambda_j + \ell).$$
(63)

The situation when $\Lambda_i < \Lambda_j$ is similar.

Finally, in case I, Λ_i and Λ_j both belong to the antisymmetric sector, and thus $f_{\Omega} = f_{\Lambda}$. This gives

$$v_{\Lambda\Omega} = \operatorname{sgn}(\sigma_{i,j,\ell}^a) \, v_{\Lambda_{ij} \left(R_{ij}^{(\ell)} \Lambda \right)_{ii}}. \tag{64}$$

4.3 The diagonal coefficients

To complete the study of the action of the Hamiltonian on the supermonomial basis, we now evaluate the diagonal coefficient explicitly. The full stCMS Hamiltonian reads

$$\bar{\mathcal{H}} = A + \beta(B+F), \quad \text{with} \quad A = \sum_{i=1}^{N} (z_i \partial_i)^2.$$
 (65)

The action of A on the supermonomial basis is diagonal:

$$A m_{\Lambda} = \left(\sum_{i=1}^{N} \Lambda_i^2\right) m_{\Lambda} \,. \tag{66}$$

There is also a contribution coming from the part $\beta(B+F)$ of the Hamiltonian. It can obtained from (55) for the case $\ell=0$ with

$$v_{\Lambda_{ij}\left(R_{ij}^{(0)}\Lambda\right)_{ij}} = v_{\Lambda_{ij}\Lambda_{ij}} = \beta|\Lambda_i - \Lambda_j|. \tag{67}$$

This value is independent of the sector – cf. eqs (40), (46) and (52)). The diagonal contribution of $\beta(B+F)m_{\Lambda}$ is thus

$$v_{\Lambda\Lambda} = \frac{1}{f_{\Lambda}} \sum_{1 \le i < j \le N} v_{\Lambda_{ij}\Lambda_{ij}} f_{\Lambda} = \beta \sum_{1 \le i < j \le N} |\Lambda_i - \Lambda_j|$$
$$= \beta \sum_{1 \le i < j \le N} (\Lambda_i^* - \Lambda_j^*) = \beta \sum_{k=1}^N (N + 1 - 2k) \Lambda_k^*.$$
(68)

The step in which we eliminate the alsolute value provides the technical reason why the eigenvalue is ultimately expressed in terms of the parts of Λ^* . The last equality is obtained as follows [3]. Let L be such that $\Lambda_L^* \neq 0$ but $\Lambda_{j>L}^* = 0$. We can write

$$\sum_{1 \le i < j \le N} (\Lambda_i^* - \Lambda_j^*) = \sum_{1 \le i \le L < j \le N} \Lambda_i^* + \sum_{1 \le i < j \le L} (\Lambda_i^* - \Lambda_j^*)$$

$$= \sum_{1 \le i \le L} (N - L) \Lambda_i^* + \sum_{1 \le k \le L} [(L - k) - (k - 1)] \Lambda_k^*$$

$$= \sum_{1 \le k \le L} (N + 1 - 2k) \Lambda_k^*. \tag{69}$$

To evaluate the double sum, we simply count the number of occurences of Λ_k^* with positive and negative signs: it arises N-k times with a plus sign and k-1 times with a minus sign. Note finally that the last sum can be extended from L to N without changing its value.

Combining the diagonal contribution of $\beta(B+F)$ to that of A yields

$$\varepsilon_{\Lambda} = \sum_{k=1}^{N} [\Lambda_k^2 + \beta(N+1-2k)\Lambda_k^*] = \sum_{k=1}^{N} [\Lambda_k^{*2} + \beta(N+1-2k)\Lambda_k^*]$$
 (70)

This is also the eigenvalue of the Jack superpolynomial \mathcal{J}_{Λ} , which is thus computed here without the help of Dunkl operators.

5 Determinantal expression for the Jack superpolynomials

5.1 A new definition of the Jack superpolynomials

Given that we have obtained a partial ordering on superpartitions, it is natural to ask whether we can replace Definition 1 of the Jack superpolynomials by a *stronger* definition involving the *weaker* ordering on superpartitions. That is, we would like to know whether the Jack superpolynomials can be characterized by the two conditions:

(i):
$$\mathcal{J}_{\Lambda} = m_{\Lambda} + \sum_{\Omega; \Omega < {}^{s}\Lambda} c_{\Lambda\Omega} m_{\Omega}$$

(ii): $\bar{\mathcal{H}} \mathcal{J}_{\Lambda} = \varepsilon_{\Lambda} J_{\Lambda}$. (71)

The interest of such a definition is that, given the triangular action of $\bar{\mathcal{H}}$ (see Corollary 6) and the fact that $\varepsilon_{\Lambda} \neq \varepsilon_{\Omega}$ if $\Omega <^s \Lambda$, we can obtain, as we will see later, a determinantal expression for \mathcal{J}_{Λ} . Therefore, there exist functions satisfying conditions (71). That we can indeed characterize the Jack superpolynomials using these conditions relies on the uniqueness of the Jack superpolynomials (cf. Definition 1), which we now establish.

Lemma 9. There exists a unique function \mathcal{J}_{Λ} satisfying the two conditions:

(i):
$$\mathcal{J}_{\Lambda} = m_{\Lambda} + \sum_{\Omega; \Omega^* < \Lambda^*} c_{\Lambda\Omega} m_{\Omega}$$
(ii):
$$\bar{\mathcal{H}} \mathcal{J}_{\Lambda} = \varepsilon_{\Lambda} J_{\Lambda} .$$
(72)

Proof: The existence of such a function will follow from the determinantal expression that we will present later on. The uniqueness is proved as follows. Let \mathcal{J}_{Λ} and $\bar{\mathcal{J}}_{\Lambda}$ be two functions satisfying

the conditions of the theorem. Then, if we suppose that $\mathcal{J}_{\Lambda} \neq \bar{\mathcal{J}}_{\Lambda}$, we have

$$\mathcal{J}_{\Lambda} - \bar{\mathcal{J}}_{\Lambda} = \sum_{\Omega: \Omega^* < \Lambda^*} d_{\Lambda\Omega} m_{\Omega} , \qquad (73)$$

for some coefficients $d_{\Lambda\Omega}$. Now, let $\Omega^{(1)} <^T \Omega^{(2)} <^T \cdots <^T \Omega^{(n)}$ be a total ordering (compatible with the ordering on superpartitions) of all the superpartitions $\Omega^{(i)}$, $i=1,\ldots,n$, such that $d_{\Lambda\Omega^{(i)}} \neq 0$. Since \mathcal{J}_{Λ} and $\bar{\mathcal{J}}_{\Lambda}$ are both eigenfunctions of $\bar{\mathcal{H}}$ with eigenvalue ε_{Λ} , we must have $\bar{\mathcal{H}}\left(\mathcal{J}_{\Lambda} - \bar{\mathcal{J}}_{\Lambda}\right) = \epsilon_{\Lambda}\left(\mathcal{J}_{\Lambda} - \bar{\mathcal{J}}_{\Lambda}\right)$, that is, from Corollary 6,

$$\bar{\mathcal{H}}\sum_{i=1}^{n}d_{\Lambda\Omega^{(i)}}m_{\Omega^{(i)}} = \sum_{i=1}^{n}d_{\Lambda\Omega^{(i)}}\left(\varepsilon_{\Omega^{(i)}}m_{\Omega^{(i)}} + \sum_{\Gamma < {}^{s}\Omega^{(i)}}v_{\Omega^{(i)}\Gamma}m_{\Gamma}\right) = \varepsilon_{\Lambda}\sum_{i=1}^{n}d_{\Lambda\Omega^{(i)}}m_{\Omega^{(i)}}.$$
 (74)

In the middle expression, the coefficient of $m_{\Omega^{(n)}}$ is simply equal to $\varepsilon_{\Omega^{(n)}}d_{\Lambda\Omega^{(n)}}$, since $\Omega^{(n)}$ dominates $\Omega^{(i)}$, $i=1,\ldots,n-1$, in the order on superpartitions. But in the last expression of (74), the coefficient of $m_{\Omega^{(n)}}$ is equal to $\varepsilon_{\Lambda}d_{\Lambda\Omega^{(n)}}$. We must thus have $\varepsilon_{\Lambda}d_{\Lambda\Omega^{(n)}}=\varepsilon_{\Omega^{(n)}}d_{\Lambda\Omega^{(n)}}$, with $d_{\Lambda\Omega^{(n)}}\neq 0$, that is we must have $\varepsilon_{\Lambda}=\varepsilon_{\Omega^{(n)}}$. But this is impossible because, from (73), $\Lambda^*>(\Omega^{(n)})^*$, which implies that $\varepsilon_{\Lambda}\neq\varepsilon_{\Omega^{(n)}}$. Therefore, a contradiction arises and \mathcal{J}_{Λ} cannot be different from $\overline{\mathcal{J}}_{\Lambda}$.

It is easy to see, from (34)(i), that if a function \mathcal{J}_{Λ} is such that $\mathcal{J}_{\Lambda} = m_{\Lambda} + \sum_{\Omega < s_{\Lambda}} c_{\Lambda\Omega} m_{\Omega}$, then it is also such that $\mathcal{J}_{\Lambda} = m_{\Lambda} + \sum_{\Omega; \Omega^* < \Lambda^*} e_{\Lambda\Omega} m_{\Omega}$, for some coefficients $e_{\Lambda\Omega}$. Therefore, the previous lemma (combined with the existence of functions satisfying conditions (71)) leads directly to the following theorem.

Theorem 10. The Jack superpolynomials can be defined by the two conditions:

(i):
$$\mathcal{J}_{\Lambda} = m_{\Lambda} + \sum_{\Omega; \Omega < {}^{s}\Lambda} c_{\Lambda\Omega} m_{\Omega}$$

(ii): $\bar{\mathcal{H}} \mathcal{J}_{\Lambda} = \varepsilon_{\Lambda} J_{\Lambda}$. (75)

5.2 Determinantal formulas

As mentioned earlier, an important offshoot of this new definition is that we can obtain determinantal expressions for the Jack superpolynomials. Indeed, because we are looking for Jack superpolynomials with a triangular expansion

$$\mathcal{J}_{\Lambda} = m_{\Lambda} + \sum_{\Omega < s_{\Lambda}} c_{\Lambda\Omega} m_{\Omega} , \qquad (76)$$

the fact that $\varepsilon_{\Omega} \neq \varepsilon_{\Lambda}$ if $\Omega <^{s} \Lambda$ combined with the triangular action (35)

$$\bar{\mathcal{H}} \, m_{\Lambda} = \varepsilon_{\Lambda} m_{\Lambda} + \sum_{\Omega < {}^{s}\Lambda} v_{\Lambda\Omega} \, m_{\Omega} \,, \tag{77}$$

leads to a determinantal expression.

Theorem 11. If $\Lambda^{(1)}, \Lambda^{(2)}, \ldots, \Lambda^{(n)} = \Lambda$ is a total ordering (compatible with the ordering on superpartitions) of all superpartitions $\leq^s \Lambda$, then the Jack superpolynomial \mathcal{J}_{Λ} is given by the following

6 CONCLUSION 17

determinant:

$$\mathcal{J}_{\Lambda} = c_{\Lambda} \begin{pmatrix}
m_{\Lambda^{(1)}} & m_{\Lambda^{(2)}} & \cdots & \cdots & m_{\Lambda^{(n-1)}} & m_{\Lambda^{(n)}} \\
\varepsilon_{\Lambda^{(1)}} - \varepsilon_{\Lambda^{(n)}} & v_{\Lambda^{(2)}\Lambda^{(1)}} & \cdots & \cdots & v_{\Lambda^{(n-1)}\Lambda^{(1)}} & v_{\Lambda^{(n)}\Lambda^{(1)}} \\
0 & \varepsilon_{\Lambda^{(2)}} - \varepsilon_{\Lambda^{(n)}} & \cdots & \cdots & v_{\Lambda^{(n-1)}\Lambda^{(2)}} & v_{\Lambda^{(n)}\Lambda^{(2)}} \\
\vdots & 0 & \ddots & \vdots & \vdots \\
\vdots & \vdots & \ddots & \ddots & \vdots \\
0 & 0 & \cdots & 0 & \varepsilon_{\Lambda^{(n-1)}} - \varepsilon_{\Lambda^{(n)}} & v_{\Lambda^{(n)}\Lambda^{(n-1)}}
\end{pmatrix}, (78)$$

where the constant of proportionality is

$$c_{\Lambda} = (-1)^{n-1} \prod_{i=1}^{n-1} \frac{1}{\varepsilon_{\Lambda^{(i)}} - \varepsilon_{\Lambda^{(n)}}}.$$
 (79)

For a proof that this determinant is in fact a non-zero eigenvector of $\overline{\mathcal{H}}$ with eigenvalue ε_{Λ} , the reader is referred to [11]. The proof found in this article can be applied to our case as well. Notice that in the determinantal expression, the relative ordering of the superpartitions not related by the partial ordering \geq^s is irrelevant.

It might be important to remark that, since the coefficients $v_{\Lambda\Omega}$ do not depend on the number of variables N, and since the differences

$$\varepsilon_{\Omega} - \varepsilon_{\Lambda} = \varepsilon_{\Omega^*} - \varepsilon_{\Lambda^*} = \sum_{k} \left((\Omega_k^{*2} - \Lambda_k^{*2}) - 2k\beta (\Omega_k^* - \Lambda_k^*) \right), \qquad |\Omega| = |\Lambda|, \tag{80}$$

also do not depend on the number of variables, the determinantal expression for \mathcal{J}_{Λ} does not depend on N.

For example, with

$$\bar{\mathcal{H}} m_{(3;1)} = (10 + 4 \beta N - 6 \beta) m_{(3;1)} + 2 \beta m_{(2;2)} + 8 \beta m_{(2;1,1)} + 2 \beta m_{(1;2,1)}
\bar{\mathcal{H}} m_{(2;2)} = (8 + 4 \beta N - 8 \beta) m_{(2;2)} + 4 \beta m_{(2;1,1)} + 2 \beta m_{(1;2,1)}
\bar{\mathcal{H}} m_{(2;1,1)} = (6 + 4 \beta N - 10 \beta) m_{(2;1,1)} + 6 \beta m_{(1;1,1,1)}
\bar{\mathcal{H}} m_{(1;2,1)} = (6 + 4 \beta N - 10 \beta) m_{(1;2,1)} + 12 \beta m_{(1;1,1,1)}
\bar{\mathcal{H}} m_{(1;1,1,1)} = (4 + 4 \beta N - 16 \beta) m_{(1;1,1,1)},$$
(81)

we have

$$\mathcal{J}_{(3;1)} = \frac{1}{64(3+5\beta)(1+\beta)^3} \begin{vmatrix}
m_{(1;1,1,1)} & m_{(1;2,1)} & m_{(2;1,1)} & m_{(2;2)} & m_{(3;1)} \\
-6-10\beta & 12\beta & 6\beta & 0 & 0 \\
0 & -4-4\beta & 0 & 2\beta & 2\beta \\
0 & 0 & -4-4\beta & 4\beta & 8\beta \\
0 & 0 & 0 & -2-2\beta & 2\beta
\end{vmatrix}$$

$$= m_{(3;1)} + \frac{\beta}{1+\beta} m_{(2;2)} + \frac{\beta(2+3\beta)}{(1+\beta)^2} m_{(2;1,1)} + \frac{\beta(1+2\beta)}{2(1+\beta)^2} m_{(1;2,1)} + \frac{3\beta^2}{(1+\beta)^2} m_{(1;1,1,1)} . \quad (82)$$

6 Conclusion

Although unravelling the natural generalization of the dominance ordering at the level of superpartitions was our original motivation for this work, we have obtained much more than that: we ended

up with a simple determinantal expression for the Jack superpolynomials. In a sense, the mere discovery of a determinantal expression for Jack superpolynomials ensures that the pivotal concepts underlying their construction, namely monomial symmetric superpolynomials and superpartitions (as well as their associated ordering) are the right ones.

There is a natural supersymmetric extension of the physical scalar product with respect to which the Jack polynomials are orthogonal. However, the Jack superpolynomials are not orthogonal with respect to this supersymmetric scalar product. The next step is thus to seek a general pattern for constructing linear combinations of Jack superpolynomials sharing the same Hamiltonian eigenvalue that would be orthogonal.

Also, we believe the present formalism to be extendable to the case of the Hi-Jack polynomials (or generalized Hermite polynomials) and their supersymmetric counterparts. We hope to report elsewhere on this subject.

NOTE ADDED

After this article got accepted for publication, we completed the program of constructing orthogonal Jack superpolynomials. These results will be presented in the article Jack polynomials in superspace.

ACKNOWLEDGMENTS

This work was supported by FCAR and NSERC. L.L. wishes to thank Luc Vinet for his financial support. P.D. would also like to thank the Fondation J.A Vincent for a student fellowship.

A Proofs of Identity 7 and 8

Proof of Identity 7: Expanding the factors z_{12} and z^{12} , Identity 7 is equivalent to

$$(r-s)\left(z_1^{r-s+1} - z_1^{r-s}z_2\right) + \sum_{\ell=1}^{r-s-1} 2(r-s-\ell)z_1^{r-s-\ell+1}z_2^{\ell} - \sum_{\ell=1}^{r-s-1} 2(r-s-\ell)z_1^{r-s-\ell}z_2^{\ell+1}$$

$$= (r-s)\left(z_1^{r-s+1} + z_1^{r-s}z_2\right) - 2\sum_{\ell=1}^{r-s} z_1^{r-s-\ell+1}z_2^{\ell}. \tag{83}$$

Now, doing simple transformations on the indices of summation, we have

$$(r-s)\left(z_{1}^{r-s+1}-z_{1}^{r-s}z_{2}\right)+\sum_{\ell=1}^{r-s-1}2(r-s-\ell)z_{1}^{r-s-\ell+1}z_{2}^{\ell}-\sum_{\ell=1}^{r-s-1}2(r-s-\ell)z_{1}^{r-s-\ell}z_{2}^{\ell+1}$$

$$=(r-s)\left(z_{1}^{r-s+1}-z_{1}^{r-s}z_{2}\right)+\sum_{\ell=1}^{r-s-1}2(r-s-\ell)z_{1}^{r-s-\ell+1}z_{2}^{\ell}-\sum_{\ell=2}^{r-s}2(r-s-\ell+1)z_{1}^{r-s-\ell+1}z_{2}^{\ell}$$

$$=(r-s)\left(z_{1}^{r-s+1}-z_{1}^{r-s}z_{2}\right)+2(r-s-1)z_{1}^{r-s}z_{2}-2z_{1}z_{2}^{r-s}-2\sum_{\ell=2}^{r-s-1}z_{1}^{r-s-\ell+1}z_{2}^{\ell}$$

$$=(r-s)\left(z_{1}^{r-s+1}+z_{1}^{r-s}z_{2}\right)-2\sum_{\ell=1}^{r-s}z_{1}^{r-s-\ell+1}z_{2}^{\ell},$$

$$(84)$$

REFERENCES 19

which proves (83) and, consequently, Identity 7.

Proof of Identity 8: Using Identity 7 twice (once with $z_1 \leftrightarrow z_2$), we have

$$(r-s)z^{12} \left(z_1^{r-s} + z_2^{r-s} \right) - 4 \sum_{\ell=1}^{r-s} z_1^{r-s-\ell+1} z_2^{\ell}$$

$$= z_{12} \left\{ (r-s) \left(z_1^{r-s} - z_2^{r-s} \right) + \sum_{\ell=1}^{r-s-1} 2(r-s-\ell) z_1^{r-s-\ell} z_2^{\ell} - \sum_{\ell=1}^{r-s-1} 2(r-s-\ell) z_1^{\ell} z_2^{r-s-\ell} \right\}.$$
(85)

Finally, after a sequence of simple transformations,

$$\sum_{\ell=1}^{r-s-1} 2(r-s-\ell)z_1^{r-s-\ell}z_2^{\ell} - \sum_{\ell=1}^{r-s-1} 2(r-s-\ell)z_1^{\ell}z_2^{r-s-\ell} \\
= \sum_{\ell=1}^{r-s-1} 2(r-s-\ell)z_1^{r-s-\ell}z_2^{\ell} - \sum_{\ell=1}^{r-s-1} 2\ell z_1^{r-s-\ell}z_2^{\ell} \\
= \sum_{\ell=1}^{r-s-1} 2(r-s-2\ell)z_1^{r-s-\ell}z_2^{\ell} \\
= \sum_{\ell=1}^{\lfloor \frac{r-s-1}{2} \rfloor} 2(r-s-2\ell)z_1^{r-s-\ell}z_2^{\ell} + \sum_{\ell=\lfloor \frac{r-s-1}{2} \rfloor+1}^{r-s-1} 2(r-s-2\ell)z_1^{r-s-\ell}z_2^{\ell} \\
= \sum_{\ell=1}^{\lfloor \frac{r-s-1}{2} \rfloor} 2(r-s-2\ell)z_1^{r-s-\ell}z_2^{\ell} - \sum_{\ell=1}^{r-s-1-\lfloor \frac{r-s-1}{2} \rfloor} 2(r-s-2\ell)z_1^{\ell}z_2^{r-s-\ell} \\
= \sum_{\ell=1}^{\lfloor \frac{r-s-1}{2} \rfloor} 2(r-s-2\ell)\left(z_1^{r-s-\ell}z_2^{\ell}-z_1^{\ell}z_2^{r-s-\ell}\right), \tag{86}$$

we obtain the Identity 8. Note that in the final step, when r-s is even, we have $r-s-1-\lfloor\frac{r-s-1}{2}\rfloor=\lfloor\frac{r-s-1}{2}\rfloor+1$. The extra case corresponding to $\ell=\lfloor\frac{r-s-1}{2}\rfloor+1=(r-s)/2$ can be ignored since it has a factor $(r-s-2\ell)=0$.

References

- F. Calogero, Ground state of one-dimensional N body system, J. Math. Phys. 10, 2197 (1969);
 Solution of a three body problem in one-dimension, J. Math. Phys. 10, 2191 (1969);
 Solution of the one-dimensional N body problems with quadratic and/or inversely quadratic pair potentials,
 J. Math. Phys. 12, 419 (1971).
- [2] J. Moser, Three integrable Hamiltonian systems connected with isospectral deformations, Surveys in applied mathematics, 235–258 (1976).
- [3] B. Sutherland, Quantum many body problem in one-dimension: ground state, J. Math. Phys. 12, 246 (1971); Exact results for a quantum many body problem in one-dimension, Phys. Rev. A4, 2019 (1971); Exact results for a quantum many body problem in one-dimension. II, Phys. Rev. A5, 1372 (1972).

REFERENCES 20

[4] M. A. Olshanetsky and A. M. Perelomov, Classical integrable finite dimensional systems related to Lie algebras, Phys. Rept. **71**, 313 (1981); Quantum integrable systems related to Lie algebras, Phys. Rept. **94** (1983) 313.

- [5] H. Jack, A class of symmetric polynomials with a parameter, Proc. Roy. Soc. Edinburgh Sect. A, 69, 1–18 (1970/1971); A surface integral and symmetric functions, Proc. Roy. Soc. Edinburgh Sect. A, 69, part 4, 347–364 (1972).
- [6] R. P. Stanley, Some combinatorial properties of Jack symmetric functions, Adv. Math., 77, 76-115 (1988).
- [7] P. Desrosiers, L. Lapointe and P. Mathieu, Supersymmetric Calogero-Moser-Sutherland models and Jack superpolynomials, Nucl. Phys. B 606, 547 (2001).
- [8] B. Sriram Shastry and B. Sutherland, Superlax pairs and infinite symmetries in the $1/r^2$ system, Phys. Rev. Lett. **70**, 4029 (1993), cond-mat/9212029.
- [9] I. G. Macdonald, Symmetric functions and Hall polynomials, 2ieme edition, The Clarendon Press Oxford University Press, (1995); Symmetric functions and orthogonal polynomials, Dean Jacqueline B. Lewis Memorial Lectures presented at Rutgers University, American Mathematical Society, (1998).
- [10] K. Sogo, Eigenstates of Calogero-Sutherland-Moser model and generalized Schur functions, J. Math. Phys. 35, 2282 (1994).
- [11] L. Lapointe, A. Lascoux and J. Morse, Determinantal formula and recursion for Jack polynomials, Electro. J. Comb. 7 467 (2000).
- [12] A. P. Polychronakos, Exchange operator formalism for integrable systems of particles, Phys. Rev. Lett. 69, 703 (1992), hep-th/9202057; J. A. Minahan and A. P. Polychronakos, Integrable systems for particles with internal degrees of freedom, Phys. Lett. B302 299 (1993), hep-th/9206046.