

## Renormalizable Abelian-Projected Effective Gauge Theory Derived from Quantum Chromodynamics II

Toru Shinohara<sup>1,†</sup>

<sup>1</sup> *Graduate School of Science and Technology, Chiba University, Chiba 263-8522, Japan*

### Abstract

In the previous paper[1], we derived the Abelian projected effective gauge theory as a low energy effective theory of the  $SU(N)$  Yang-Mills theory by adopting the maximal Abelian gauge. At that time, we have demonstrated the multiplicative renormalizability of the propagators for the diagonal gluon and the dual Abelian anti-symmetric tensor field. In this paper, we show the multiplicative renormalizability of the Green's functions also for the off-diagonal gluon. Moreover we complement the previous results by calculating the anomalous dimension and the renormalization group functions which are undetermined in the previous paper.

Key words: maximal Abelian gauge, Abelian dominance, renormalizability, non-Abelian gauge theory

PACS: 12.38.Aw, 12.38.Lg

---

<sup>†</sup> E-mail: [sinohara@cuphd.nd.chiba-u.ac.jp](mailto:sinohara@cuphd.nd.chiba-u.ac.jp), [sinohara@graduate.chiba-u.jp](mailto:sinohara@graduate.chiba-u.jp)

# 1 Introduction

Quark confinement and spontaneous breakdown of chiral symmetry are very important problems in the low energy Quantum Chromodynamics (QCD). It is difficult to analyze QCD in the low energy region due to strong coupling constant, while it is easy to analyze QCD in the high energy region due to the asymptotic freedom of coupling constant. Thus the effective theory is often used in order to investigate the phenomena in the low energy region of QCD. In fact, it is well-known that the problems mentioned above are qualitatively explained by dual Ginzburg-Landau (DGL) theory which describes the dual superconductivity due to the dual Meisner effects.[2, 3, 4] Therefore we expect that the DGL theory is a candidate of a low energy effective theory of QCD. However, the analytical derivation of DGL from QCD is not achieved so that we have tried to construct the interpolated effective theory connecting QCD and DGL in the series of our works[1, 5].

In the previous paper[1], which is referred to as (I) hereafter, we have derived Abelian projected effective gauge theory (APEGT) from the  $SU(N)$  Yang-Mills theory by making use of the maximal Abelian (MA) gauge[6] based on an idea of Abelian projection[3]. The APEGT is originally derived for the  $SU(2)$  gauge group in Ref. [5] and extended for the  $SU(N)$  gauge group in (I).

The original Yang-Mills theory includes the diagonal gluon, the off-diagonal gluon, the diagonal ghost and the off-diagonal ghost. According to the Abelian dominance[7], the diagonal components play the dominant roles in the low energy region of QCD, while the off-diagonal components hardly affect the phenomena in the low energy region. The mechanism of Abelian dominance can be understood by the fact that the off-diagonal fields become massive, while the diagonal fields remain massless or have smaller masses. We derived the dynamical mass generations of off-diagonal fields in Ref. [8] in agreement with numerical simulations on a lattice[9]. By introducing a dual Abelian anti-symmetric tensor fields together with two arbitrary parameters  $\rho$  and  $\sigma$  as a dual of the composite operator of off-diagonal gluons and by integrating out off-diagonal gluons and ghosts, we have obtained the APEGT which includes only the diagonal gluon and the dual Abelian tensor field. Moreover, by integrating out the diagonal gluon, we have arrived at the DGL-like effective theory as demonstrated in Refs. [1, 5].

In (I), we calculated the beta-function, the anomalous dimensions of the diagonal gluon and dual Abelian tensor fields, and the renormalization group (RG) functions of the gauge fixing parameter for the diagonal gluon and an arbitrary parameter  $\rho$  and demonstrated the multiplicative renormalizability for the diagonal gluon and the dual Abelian tensor field propagators. In this paper, moreover, we show the multiplicative renormalizability for the Green's functions including not only the diagonal fields but also off-diagonal gluons, and calculate the anomalous dimension of off-diagonal gluon and the RG functions of the gauge fixing parameter for the off-diagonal gluon and the remaining parameter  $\sigma$ . The results obtained here complement the previous results.

## 2 Modified maximal Abelian gauge

We construct the gauge fixing (GF) and the associated Faddeev-Popov (FP) ghost term for the MA gauge. To analyze the non-Abelian gauge theory in the MA gauge, we distinguish the color indices as follows:

$$\begin{cases} A, B, C, \dots & \rightarrow SU(N), \\ i, j, k, \dots & \rightarrow U(1)^{N-1} \text{ (diagonal)}, \\ a, b, c, \dots & \rightarrow SU(N)/U(1)^{N-1} \text{ (off-diagonal)}. \end{cases} \quad (1)$$

First, we define the MA gauge condition. The MA gauge is obtained by minimizing the functional  $R[A^U]$  with respect to the local gauge transformation  $U(x)$  of  $A_\mu^a$ . Here,  $R[A]$  is defined as the functional of off-diagonal gluons,

$$R[A] := \int d^4x \frac{1}{2} A_\mu^a(x) A^{\mu a}(x). \quad (2)$$

Then we obtain the differential form of the MA gauge condition,

$$F^a[A] := D_\mu A^{\mu a} = 0, \quad (3)$$

where we defined the covariant derivative with respect to the diagonal gluon  $A_\mu^i$  as

$$D_\mu \Phi^A := (\partial_\mu \delta^{AB} + g f^{AiB} A_\mu^i) \Phi^B, \quad (4)$$

for an arbitrary operator  $\Phi^A$ . The MA gauge condition partially fixes the color rotational symmetry from  $SU(N)$  to  $U(1)^{N-1}$ . In fact, the residual  $U(1)^{N-1}$  symmetry,

$$\begin{cases} \delta A_\mu^i = \partial_\mu \theta^i, \\ \delta \Phi^i = 0 \quad (\Phi^i = \phi^i, C^i, \bar{C}^i), \\ \delta \Phi^a = -g f^{abi} \Phi^b \theta^i \quad (\Phi^a = A_\mu^a, \phi^a, C^a, \bar{C}^a), \end{cases} \quad (5)$$

is not fixed by the MA gauge fixing condition. Here  $A_\mu$ ,  $\phi$ ,  $C$  and  $\bar{C}$  is gluon, Nakanishi-Lautrup field, ghost and antighost respectively. In order to complete the gauge fixing, we must fix the residual  $U(1)^{N-1}$  symmetry. In this paper, we employ the additional gauge fixing condition  $F^i[A_\mu] := \partial_\mu A^{i\mu} = 0$  for fixing the residual  $U(1)^{N-1}$  symmetry.

Next, we construct the GF+FP term for the MA gauge. In (I), we adopted the *naive* MA gauge fixing term

$$S_{\text{MA}} := - \int d^4x i \delta_B \left[ \bar{C}^a \left( D^\mu A_\mu^a + \frac{\alpha}{2} \phi^a \right) + \bar{C}^i \left( \partial^\mu A_\mu^i + \frac{\beta}{2} \phi^i \right) \right], \quad (6)$$

where  $\delta_B$  is the BRST transformation. However, the GF+FP term (6) is not appropriate from the viewpoint of multiplicative renormalizability of the theory. The reason is as follows. The divergent contribution proportional to a ghost-antighost self interaction term  $(f^{iab} \bar{C}^a C^b)^2$  is generated as a quantum correction due to the existence of the interaction  $f^{iad} f^{icb} \bar{C}^a C^b A^{uc} A_\mu^d$  in the MA gauge (6). However, the

GF+FP term (6) does not include such a term at the tree level so that the multiplicative renormalizability cannot be maintained in the off-diagonal gluon sector. Therefore, in order to preserve the multiplicative renormalizability, we should modify the GF+FP term to include such a term at the tree level. In this paper, we adopt the *modified* MA gauge fixing term[11, 8]

$$S_{\text{mMA}} := \int d^4x i \left[ \delta_{\text{B}} \bar{\delta}_{\text{B}} \left( \frac{1}{2} A_{\mu}^a A_{\mu}^a - \frac{\alpha}{2} i C^a \bar{C}^a \right) - \delta_{\text{B}} \left\{ C^i \left( \partial^{\mu} A_{\mu}^i + \frac{\beta}{2} \phi^i \right) \right\} \right], \quad (7)$$

where  $\bar{\delta}_{\text{B}}$  is the anti-BRST transformation. Integrating out  $\phi^i$  and  $\phi^a$ , we obtained the total gauge fixing term

$$\begin{aligned} S_{\text{mMA}} = \int d^4x \left\{ -\frac{1}{2\alpha} (D^{\mu} A_{\mu}^a)^2 - \frac{1}{2\beta} (\partial^{\mu} A_{\mu}^i)^2 - i g^2 f^{adi} f^{cbi} \bar{C}^a C^b A^{\mu c} A_{\mu}^d \right. \\ + i \bar{C}^a D^2 C^a + i g \bar{C}^a D^{\mu} (f^{abc} A_{\mu}^b C^c) + \frac{i}{2} g (D^{\mu} A_{\mu}^a) (f^{abc} \bar{C}^b C^c) \\ + \frac{\alpha}{8} g^2 f^{abe} f^{cde} \bar{C}^a \bar{C}^b C^c C^d + \frac{\alpha}{4} g^2 f^{abi} f^{cdi} \bar{C}^a \bar{C}^b C^c C^d \\ \left. + \frac{\alpha}{8} g^2 (f^{abc} \bar{C}^b C^c)^2 + i \bar{C}^i \partial^2 C^i + i \bar{C}^i \partial^{\mu} (g f^{ibc} A_{\mu}^b C^c) \right\}. \quad (8) \end{aligned}$$

This term includes the desired form ghost interaction term.<sup>1</sup> Moreover, we find that no interaction term with diagonal ghost  $C^i$  exists so that a diagonal ghost does not appear as an internal line in the perturbative loop calculation.

### 3 Composite operator and Abelian auxiliary field

Here, we consider the Green's function including Abelian composite operator which is made of off-diagonal fields and invariant under the  $U(1)$  gauge transformation (5). Then we introduce a dual Abelian tensor field  $B_{\mu\nu}^i$  as an auxiliary field corresponding to such a composite operator.

According to the Abelian dominance in the low energy region of QCD, an elementally off-diagonal field hardly affect the physics alone. However, the Abelian composite operator which is composed of off-diagonal fields can affect the low energy physics as well as elementary Abelian fields. There are four candidates of the Abelian composite operator made of off-diagonal fields with mass dimension 2, that is,  $f^{ibc} A_{\mu}^b A_{\nu}^c$ ,  $A_{\mu}^a A^{\mu a}$ ,  $f^{ibc} \bar{C}^b C^c$  and  $\bar{C}^a C^a$ . In this paper we focus on the two-rank anti-symmetric composite operator  $O_{\mu\nu}^i := f^{ibc} A_{\mu}^b A_{\nu}^c$  since this operator is very important to construct the dual Abelian gauge theory of Yang-Mills theory.<sup>2</sup> In order

<sup>1</sup>The GF+FP term (7) satisfies the charge conjugation invariance, global shift invariance for the diagonal ghost and global shift invariance for the diagonal antighost. See Ref [10] for more detail.

<sup>2</sup>Other three scalar composite operators are meaningful in the different sense. In the non-perturbative consideration by making use of effective potential, the dynamical mass of the off-diagonal fields are caused through the pair condensation of the off-diagonal ghost and antighost. However, in this paper, all calculations are based on the perturbative method and the non-zero masses of the off-diagonal fields are assumed.




Figure 1: The wavy line denotes the gluon and the circled cross denotes the composite source  $K_{\mu\nu}^i$ .

to consider the Green's functions with Abelian composite operator  $O^i$ , we introduce a source term  $S_K$  for  $O^i$  as well as  $S_J$  for elemental fields:

$$Z[J, K] = \int D\Phi \exp\{iS_{\text{YM}} + iS_{\text{mMA}} + iS_J + iS_K\}. \quad (9)$$

Where  $S_{\text{YM}} := -\frac{1}{4}F_{\mu\nu}^A F^{\mu\nu A}$  is Yang-Mills action and gauge fixing terms  $S_{\text{mMA}}$  have already been given in (7).  $S_J$  is a source term for the elementary fields,  $S_J := \int d^4x (A_\mu^i J^{\mu i} + A_\mu^a J^{\mu a})$ , and  $S_K$  is a source term for the composite operator:

$$S_K := \int d^4x \frac{i}{2} {}^*O_{\mu\nu}^i K^{\mu\nu i}, \quad (10)$$

where we define the Hodge dual of arbitrary two-rank anti-symmetric tensor as  ${}^*O_{\mu\nu} := \frac{1}{2}\epsilon_{\mu\nu\rho\sigma} O^{\rho\sigma}$ . However, we notice that the composite source term (10) is insufficient from the viewpoint of multiplicative renormalizability. Indeed, such a source term (10) generate the divergent contributions not only proportional to  $O_{\mu\nu}^i K^{\mu\nu i}$  but also proportional to  $\partial_\mu A_\nu^i K^{\mu\nu i}$  and  $K_{\mu\nu}^i K^{\mu\nu i}$ . (See Fig. 1) Therefore, in order to maintain the multiplicative renormalizability, we should re-define the composite source term as

$$S_K := \int d^4x \left( \frac{i}{2} {}^*Q_{\mu\nu}^i K^{\mu\nu i} + \frac{1}{4} K_{\mu\nu}^i K^{\mu\nu i} \right), \quad (11)$$

where we define new composite operator

$$Q_{\mu\nu}^i := \rho f_{\mu\nu}^i + \sigma f^{ibc} A_\mu^b A_\nu^c, \quad (12)$$

with new two parameters  $\rho$  and  $\sigma$ .<sup>3</sup> Then we can obtain the connected Green's functions including the Abelian composite operator  $Q_{\mu\nu}^i$  by differentiating  $W := -i \ln Z$  with respect to  $K_{\mu\nu}^i$ .

Now we try to define the effective action. The existence of the quadratic source term such as the last term of  $S_K$  prevents us from performing the Legendre transformation. Therefore we introduce a dual Abelian tensor field as an auxiliary field in order to put out quadratic term of  $K_{\mu\nu}^i$ . By making use of the identity:

$$\mathcal{N} \int DB_{\mu\nu}^i \exp i \int d^4x \left\{ -\frac{1}{4} (B_{\mu\nu}^i - {}^*Q_{\mu\nu}^i - K_{\mu\nu}^i)^2 \right\} \equiv 1, \quad (13)$$

<sup>3</sup>Three divergent contributions shown in Fig. 1 can be absorbed by renormalizing the composite source  $K_{\mu\nu}^i$  and two new parameters  $\rho$  and  $\sigma$ .

with an appropriate constant  $\mathcal{N}$ , we obtain

$$Z[J, K] = \mathcal{N} \int D\Phi DB_{\mu\nu}^i \exp\{iS_{\text{inv}} + iS_{\text{GF}} + iS_J + i \int d^4x \frac{1}{2} B_{\mu\nu}^i K^{\mu\nu i}\}, \quad (14)$$

where we defined

$$\begin{aligned} S_{\text{inv}} = \int d^4x \left\{ -\frac{1}{4} [D_\mu A_\nu^a - D_\nu A_\mu^a + g f^{abc} A_\mu^b A_\nu^c]^2 \right. \\ - \frac{1-\rho^2}{4} (f_{\mu\nu}^i)^2 - \frac{1-\rho\sigma}{2} g f_{\mu\nu}^i f^{ibc} A^{\mu b} A^{\nu c} - \frac{1-\sigma^2}{4} g^2 (f^{ibc} A_\mu^b A_\nu^c)^2 \\ \left. - \frac{1}{4} (B_{\mu\nu}^i)^2 - \frac{1}{4} \epsilon^{\mu\nu\rho\sigma} B_{\mu\nu}^i [\rho f_{\mu\nu}^i + \sigma g f^{ibc} A_\mu^b A_\nu^c] \right\}, \quad (15) \end{aligned}$$

with the Abelian field strength  $f_{\mu\nu}^i := \partial_\mu A_\nu^i - \partial_\nu A_\mu^i$ . It is easily confirmed that we can reproduce the original partition function (9) by integrating out  $B_{\mu\nu}^i$  in (14).

Now we can perform the Legendre transformation as

$$\Gamma[\bar{A}_\mu^i, \bar{A}_\mu^a, \bar{B}_{\mu\nu}^i] := W[J, K] - \int d^4x \left( \bar{A}_\mu^i J^{\mu i} + \bar{A}_\mu^a J^{\mu a} + \frac{1}{2} \bar{B}_{\mu\nu}^i K^{\mu\nu i} \right), \quad (16)$$

where we defined the background fields:

$$\bar{A}_\mu^A := \frac{\delta}{\delta J^{\mu A}} W[J, K], \quad \bar{B}_{\mu\nu}^i := \frac{\delta}{\delta K^{\mu\nu i}} W[J, K]. \quad (17)$$

We obtain one particle irreducible (1PI) Green's function by making use of the effective action  $\Gamma[\bar{A}_\mu^i, \bar{A}_\mu^a, \bar{B}_{\mu\nu}^i]$ .

## 4 Renormalization and RG functions

We introduce the multiplicative renormalization factors of the fields, parameters and coupling constant in order to absorb undesirable divergent contributions in the 1PI Green's functions. Especially, we pay attention to the 1PI Green's functions including the dual Abelian tensor field  $B_{\mu\nu}^i$  and determine the RG function of  $\sigma$  which is unfixed in (I).

In order to show the multiplicative renormalizability and determine the beta function, the anomalous dimensions and the RG functions we define the renormalized fields and parameters as

$$\begin{aligned} \bar{A}_\mu^i &= Z_a^{1/2} \bar{A}_{\text{R}\mu}^i, & \bar{A}_\mu^a &= Z_A^{1/2} \bar{A}_{\text{R}\mu}^a, & \bar{B}_{\mu\nu}^i &= Z_B^{1/2} \bar{B}_{\text{R}\mu\nu}^i, & g &= Z_g g_{\text{R}}, \\ \alpha &= Z_\alpha \alpha_{\text{R}}, & \beta &= Z_\beta \beta_{\text{R}}, & \rho &= Z_\rho \rho_{\text{R}}, & \sigma &= Z_\sigma \sigma_{\text{R}}. \end{aligned} \quad (18)$$

Here, at the one-loop level calculations, there is no need to take into account renormalization of each fluctuation field explicitly. Moreover the renormalization of the background ghost and antighost are also irrelevant because we consider the Green's functions including neither ghost nor antighost in this paper. We can determine each




Figure 2: The graphical representation of the Green's function including  $B_{\mu\nu}^i$  at the one-loop level. The zig-zag line denotes the dual Abelian tensor field. The lines labeled “ $i$ ” correspond to the diagonal fields.




Figure 3: Vacuum polarization graphs with respect to the off-diagonal gluon propagator at one-loop level.

renormalization constant defined here to absorb all divergent contributions calculated in the 1PI Green's function.

Now, we consider the Green's functions including  $B_{\mu\nu}^i$ . There are three divergent Green's functions  $\langle \bar{B}_{\mu\nu}^i \bar{B}_{\xi\eta}^j \rangle$ ,  $\langle \bar{B}_{\mu\nu}^i \bar{A}_\lambda^j \rangle$  and  $\langle \bar{B}_{\mu\nu}^i \bar{A}_\xi^a \bar{A}_\eta^b \rangle$ . The graphical representation is given in Fig. 2. The graphs (A), (B) and (C)s correspond to the first, second and third graphs in Fig. 1 respectively.

We have already calculated the vacuum polarization graphs (A) and (B) in (I) together with the vacuum polarization for the diagonal gluon propagator  $\langle A_\mu^i A_\nu^j \rangle$ . And we determined the renormalization constants  $Z_a$ ,  $Z_B$ ,  $Z_g$  and  $Z_\beta$  there. It is remarkable that the results in (I) paper are also valid here despite that we adopt the *modified* MA gauge in this paper which differs from the *naive* MA gauge in (I). Therefore the remaining renormalization constants to be determined in this paper are  $Z_A$ ,  $Z_\alpha$  and  $Z_\sigma$ . First,  $Z_A$  and  $Z_\alpha$  can be determined by requiring that the transverse and longitudinal mode of the off-diagonal gluon propagator are convergent. Then we can determine  $Z_\sigma$  by requiring the Green's function  $\langle \bar{B}_{\mu\nu}^i \bar{A}_\xi^a \bar{A}_\eta^b \rangle$  is convergent.

The divergent vacuum polarization graphs for the off-diagonal gluons are drawn in Fig. 3 and the divergent contributions to Green's function  $\langle \bar{B}_{\mu\nu}^i \bar{A}_\xi^a \bar{A}_\eta^b \rangle$  have already been shown in Fig. 2.

After straightforward calculations by making use of the dimensional regularization,

we obtain

$$Z_A = 1 + \frac{(g_R \mu^{-\epsilon})^2}{(4\pi)^2 \epsilon} \left[ \frac{17}{6} - \frac{\alpha_R}{2} - \beta_R + \left( \frac{5}{3} + \frac{1 - \alpha_R}{2} \right) (C_2(G) - 2) \right] + \dots, \quad (19)$$

$$Z_\alpha = 1 + \frac{(g_R \mu^{-\epsilon})^2}{(4\pi)^2 \epsilon} \left[ \frac{4}{3} - \alpha_R - \frac{3}{\alpha_R} + \left( \frac{5}{3} + \frac{1 - \alpha_R}{2} - \frac{3}{4} \alpha_R \right) (C_2(G) - 2) \right] + \dots, \quad (20)$$

$$Z_\sigma = 1 + \frac{(g_R \mu^{-\epsilon})^2}{(4\pi)^2 \epsilon} \left[ \frac{4}{3} + \frac{1 + \alpha_R}{2} \sigma_R^2 + \frac{5}{4} \alpha_R - \frac{1}{\alpha_R} + \frac{\beta_R}{4} - \frac{\beta_R}{2\alpha_R} \right. \\ \left. + \left( \frac{5}{12} + \frac{\alpha_R}{2} + \frac{1 + \alpha_R}{4} \sigma_R^2 \right) (C_2(G) - 2) \right] + \dots, \quad (21)$$

where  $\epsilon := (4 - d)/2$ .

Defining the anomalous dimension  $\gamma_\Phi$  for each field  $\Phi$  and the RG function  $\gamma_\chi$  for each parameter  $\chi$  as

$$\gamma_\Phi := \frac{1}{2} \mu \frac{\partial}{\partial \mu} \ln Z_\Phi, \quad \gamma_\chi := \mu \frac{\partial \chi_R}{\partial \mu} = -\chi_R \mu \frac{\partial}{\partial \mu} \ln Z_\chi, \quad (22)$$

we obtain the following results.

$$\beta(g_R) := \mu \frac{\partial g_R}{\partial \mu} = -g_R \mu \frac{\partial}{\partial \mu} \ln Z_g = -\frac{11}{3} C_2(G) \frac{g_R^3}{(4\pi)^2}, \quad (23)$$

$$\gamma_a(g_R) = \frac{1}{2} \mu \frac{\partial}{\partial \mu} \ln Z_a = -\frac{11}{3} C_2(G) \frac{g_R^2}{(4\pi)^2}, \quad (24)$$

$$\gamma_B(g_R) = \frac{1}{2} \mu \frac{\partial}{\partial \mu} \ln Z_B = \frac{1 + \alpha_R}{2} \sigma_R^2 C_2(G) \frac{g_R^2}{(4\pi)^2}, \quad (25)$$

$$\gamma_\rho(g_R) = -\rho_R \mu \frac{\partial}{\partial \mu} \ln Z_\rho \\ = -2\rho_R \left[ \frac{11}{6} + \frac{\sigma_R^2}{2} - 2\frac{\sigma_R}{\rho_R} + \frac{1 - \alpha_R}{2} \left( \frac{\sigma_R}{\rho_R} - \frac{\sigma_R^2}{2} \right) \right] C_2(G) \frac{g_R^2}{(4\pi)^2}, \quad (26)$$

$$\gamma_\beta(g_R) = -\beta_R \mu \frac{\partial}{\partial \mu} \ln Z_A = -2\gamma_a(g) \beta_R. \quad (27)$$

$$\gamma_A = -\frac{g_R^2}{(4\pi)^2} \left[ \frac{17}{6} - \frac{\alpha_R}{2} - \beta_R + \left( \frac{5}{3} + \frac{1 - \alpha_R}{2} \right) (C_2(G) - 2) \right], \quad (28)$$

$$\gamma_\alpha = \frac{2g_R^2}{(4\pi)^2} \left[ \frac{4}{3} \alpha_R - \alpha_R^2 - 3 + \alpha_R \left( \frac{5}{3} + \frac{1 - \alpha_R}{2} - \frac{3}{4} \alpha_R \right) (C_2(G) - 2) \right], \quad (29)$$

$$\gamma_\sigma = \frac{2g_R^2 \sigma_R}{(4\pi)^2} \left[ \frac{4}{3} + \frac{1 + \alpha_R}{2} \sigma_R^2 + \frac{5}{4} \alpha_R - \frac{1}{\alpha_R} + \frac{\beta_R}{4} - \frac{\beta_R}{2\alpha_R} \right. \\ \left. + \left( \frac{5}{12} + \frac{\alpha_R}{2} + \frac{1 + \alpha_R}{4} \sigma_R^2 \right) (C_2(G) - 2) \right]. \quad (30)$$

All anomalous dimensions and RG functions except for  $\gamma_B$ ,  $\gamma_\rho$  and  $\gamma_\sigma$  are completely independent of two parameters  $\rho$  and  $\sigma$  which were introduced in this paper together with the Abelian auxiliary field  $B_{\mu\nu}^i$ . Therefore the behavior of  $A_\mu^i$ ,  $A_\nu^a$ ,  $g$ ,  $\alpha$  and  $\beta$  are not affected by introduction of such an auxiliary field. Especially, beta function obtained here is exactly identical to that in the ordinary Lorentz gauge in which the  $SU(N)$  color rotational symmetry is unbroken. To the contrary,  $\gamma_B$ ,  $\gamma_\rho$  and  $\gamma_\sigma$  depend on two parameter  $\rho$  and  $\sigma$ .

In  $SU(2)$  case, we can put Casimir operator  $C_2(G)$  identical to 2. Then RG functions of each parameter are rewritten as

$$\gamma_\beta = \frac{44}{3}\beta_R \frac{g_R^2}{(4\pi)^2}, \quad (31)$$

$$\gamma_\alpha = -2 \left[ \left( \alpha_R - \frac{2}{3} \right)^2 + \frac{23}{9} \right] \frac{g_R^2}{(4\pi)^2}, \quad (32)$$

$$\gamma_\rho = -4\rho_R \left[ \frac{11}{6} + \frac{\sigma_R^2}{2} - 2\frac{\sigma_R}{\rho_R} + \frac{1 - \alpha_R}{2} \left( \frac{\sigma_R}{\rho_R} - \frac{\sigma_R^2}{2} \right) \right] \frac{g_R^2}{(4\pi)^2}, \quad (33)$$

$$\gamma_\sigma = 2\sigma_R \left[ \frac{4}{3} + \frac{1 + \alpha_R}{2}\sigma_R^2 + \frac{5}{4}\alpha_R - \frac{1}{\alpha_R} + \frac{\beta_R}{4} - \frac{\beta_R}{2\alpha_R} \right] \frac{g_R^2}{(4\pi)^2}. \quad (34)$$

Firstly, we consider the gauge fixing parameters  $\beta$  and  $\alpha$ . The RG flow of the Abelian gauge fixing parameter  $\beta$  has a fixed point at  $\beta = 0$ . It is similar to Landau gauge in the ordinary Lorentz gauge. To the contrary, the RG flow of the parameter  $\alpha$  does not have any fixed point because  $\gamma_\alpha$  is always negative for arbitrary real value of  $\alpha$ . The value of  $\alpha$  increases toward the infrared region. Therefore we cannot analyze a special point of  $\alpha$  differently from the case of  $\beta$ , so that we set  $\beta = 0$  and leave  $\alpha$  unfixed in the following discussion.

Next, we consider a parameter  $\rho$ . The RG flow of  $\rho$  does not have a fixed point at  $\rho = 0$  unless  $\sigma = 0$ . Of course, before the perturbative calculations, we can expect that such a trivial fixed point  $\rho = \sigma = 0$  exists since the case of  $\rho = \sigma = 0$  corresponds to the case in which Abelian auxiliary field has never been introduced. Because of the fact that a fixed point of  $\rho = 0$  does not exist when  $\sigma \neq 0$ , we understand that an introduction of  $\sigma$ -term must generate  $\rho$ -term. It is compatible with the argument in Sec. 3.

Finally, we consider a remaining parameter  $\sigma$ . The RG flow of  $\sigma$  has a trivial fixed point at  $\sigma = 0$ . However, we are not interested in such a trivial fixed point. At  $\beta = 0$ , we obtain

$$\gamma_\sigma = \sigma_R(1 + \alpha_R) \left[ \sigma_R^2 + \frac{5(\alpha_R - \alpha_1)(\alpha_R - \alpha_2)}{2(1 + \alpha_R)\alpha_R} \right] \frac{g_R^2}{(4\pi)^2}, \quad (35)$$

where  $\alpha_1 = (-8 - 2\sqrt{61})/15$  and  $\alpha_2 = (-8 + 2\sqrt{61})/15$ .  $\gamma_\sigma$  can be equal to 0 at

$$\sigma \equiv \sqrt{-\frac{5(\alpha_R - \alpha_1)(\alpha_R - \alpha_2)}{2(1 + \alpha_R)\alpha_R}}, \quad (36)$$

when  $\alpha_1 < \alpha_R < -1$  or  $0 < \alpha_R < \alpha_2$ . However this is not a true fixed point since  $\alpha$  is always unfixed in the MA gauge as mentioned above.

## 5 Conclusion

In (I) and in this paper, we have considered the possibility of the existence of the Abelian composite operator composed of off-diagonal gluons in the MA gauge. We have introduced the dual Abelian anti-symmetric tensor field  $B_{\mu\nu}^i$  into the  $SU(N)$  Yang-Mills theory as an auxiliary field together with two new parameters  $\rho$  and  $\sigma$ .

In (I), we demonstrated the multiplicative renormalizability of propagators for the diagonal gluon and the dual Abelian anti-symmetric tensor field. At that time, we calculated the beta-function, the anomalous dimensions of the diagonal gluon and dual Abelian anti-symmetric tensor field, and RG functions of the gauge parameter for diagonal gluon  $\beta$  and  $\rho$ . The results obtained in (I) are still valid here.

In addition to the demonstration in (I), in this paper, we have shown the multiplicative renormalizability for the Green's functions including not only diagonal fields but also off-diagonal gluons. We have calculated the anomalous dimension of the off-diagonal gluon and the RG functions of the gauge parameter for the off-diagonal gluon  $\alpha$  and parameter  $\sigma$ .

From the results obtained in this paper, we have found the following facts. First, the RG flow of the gauge fixing parameter  $\alpha$  for off-diagonal gluon does not have a fixed point anywhere. Next, an introduction of  $\sigma$  term must generate the  $\rho$ -term so that we should introduce  $\rho$ -term at the same time from the viewpoint of multiplicative renormalizability. Finally, the RG flow of a parameter  $\sigma$  has non-trivial fixed points depending on  $\alpha$ . The non-zero values of  $\rho$  and  $\sigma$  mean that the composite operator of off-diagonal gluons plays dominant role in the low energy region.

The results in this paper are based on the perturbative calculations so that it is valid in the ultraviolet region. However, these are important for the analysis in the infrared region in order to construct the low energy effective theory which is compatible to the original QCD.

## Acknowledgements

The author would like to thank Dr. Kei-Ichi Kondo for useful discussions and helpful advices.

## References

- [1] K.-I. Kondo and T. Shinohara, Prog. Theor. Phys. 105 649 (2001).
- [2] Y. Nambu, Phys. Rev. **D10**, 4262 (1974).
- [3] G. 't Hooft, Nucl. Phys. B 190 [FS3], 455-478 (1981).
- [4] S. Mandelstam, Phys. Report, 23, 245-249 (1976).
- [5] K.-I. Kondo, Phys. Rev. **D57**, 7467 (1998).

- [6] A. Kronfeld, M. Laursen, G. Schierholz and U.-J. Wiese, Phys. Lett. **B198**, 516 (1987).
- [7] Z. F. Ezawa and A. Iwazaki, Phys. Rev. **D25**, 2681 (1982).
- [8] K.-I. Kondo and T. Shinohara, Phys. Lett. **B491**, 263 (2000).
- [9] K. Amemiya and H. Suganuma, Phys. Rev. D 60, 114509 (1999).
- [10] T. Shinohara, T. Imai and K.-I. Kondo, hep-th/0105268, to be published in Intern. J. Mod. Phys. A.
- [11] K.-I. Kondo, Phys. Rev. **D58**, 105019 (1998).