

Gödel Type of Metrics in Various Dimensions

Metin Gürses*

*Department of Mathematics, Faculty of Sciences,
Bilkent University, 06800, Ankara, Turkey*

Atalay Karasu[†] and Özgür Sarioğlu[‡]

*Department of Physics, Faculty of Arts and Sciences,
Middle East Technical University, 06531, Ankara, Turkey*

(Dated: May 22, 2019)

Gödel type of metrics are introduced and used in producing charged dust solutions in various dimensions. The key ingredient is a $D - 1$ dimensional Riemannian geometry which is then employed in constructing solutions to Einstein Maxwell field equations with a dust distribution in D dimensions. The only essential field equation in the procedure turns out to be the source free Maxwell's equation in the relevant background. Similarly the geodesics of this type of metrics are described by the Lorentz force equation for a charged particle in the lower dimensional geometry. It is explicitly shown with several examples that Gödel type of metrics can be used in obtaining exact solutions to various supergravity theories and in constructing spacetimes that contain both closed timelike and closed null curves and that contain neither of these. Among the solutions that can be established using non-flat backgrounds, there exists a class which can be interpreted as black holes in a Gödel like universe.

PACS numbers: 04.20.Jb, 04.40.Nr, 04.50.+h, 04.65.+e

I. INTRODUCTION

Gödel metric [1] in general relativity is the solution of Einstein's field equations with homogeneous perfect fluid distribution having G_5 maximal symmetry [2]. This spacetime admits closed timelike and closed null curves but contains no closed timelike and null geodesics [3]. The Gödel universe is geodesically complete, it doesn't contain any singularities or horizons. There have been several attempts to generalize the Gödel metric in classical general relativity [4, 5, 6, 7]. The main goal of these works has been the elimination of closed timelike and closed null curves.

We call a metric in D dimensions as a *Gödel type of metric* if it can be written in the form $g_{\mu\nu} = h_{\mu\nu} - u_\mu u_\nu$ where $h_{\mu\nu}$ is a degenerate matrix of rank $D - 1$ and u^μ is a timelike unit vector. [We chose this name since Gödel's renowned metric can be cast like this.] Metrics of this form resemble the well known Kerr-Schild metrics of classical general relativity [8] which go like $g_{\mu\nu} = \eta_{\mu\nu} - \ell_\mu \ell_\nu$ for a null vector ℓ^μ and which we have recently used in constructing accelerated Kerr-Schild geometries for the Einstein Maxwell null dust

*Electronic address: gurses@fen.bilkent.edu.tr

[†]Electronic address: karasu@metu.edu.tr

[‡]Electronic address: sarioglu@metu.edu.tr

[9], Einstein Born Infeld null dust field equations [10], their extensions with a cosmological constant and respective zero acceleration limits [11].

Remarkably the very form of the Gödel type of metrics is reminiscent of the metrics used in Kaluza-Klein reductions in string theories [12]. However, as will be apparent in the subsequent sections, Gödel type of metrics have a number of characteristics that distinguish them from the Kaluza-Klein metrics. Here the background metric $h_{\mu\nu}$ is taken as positive definite whereas in the Kaluza-Klein case it must be locally Lorentzian. Moreover, opposite of what is done in the Kaluza-Klein mechanism, the Gödel type of metrics are used in obtaining a D dimensional theory starting from a $D - 1$ dimensional one. The D dimensional timelike vector u^μ is used in the construction of a Maxwell theory in D dimensions unlike the Kaluza-Klein vector potential which lives and defines a Maxwell theory in $D - 1$ dimensions. Even though Gödel type of metrics are akin to metrics employed in the Kaluza-Klein mechanism at face value, the applications we present here should make their real worth clear and should help in contrasting them with Kaluza-Klein metrics.

Gödel type of metrics also show up in supergravity theories in some dimensions. A special class of Gödel type of metrics is known to be the T-dual of the pp-wave metrics in string theory [13, 14, 15]. These metrics are all supersymmetric but contain closed timelike and closed null curves and thus violate causality [16, 17, 18, 19]. Recently there has been an attempt to remedy this problem by introducing observer dependent holographic screens [14, 20]. In [21], a new class of supergravity solutions have been constructed which locally look like the Gödel universe inside a domain wall made out of supertubes and which do not contain any closed timelike curves.

In this work, we consider Gödel type of metrics in a D dimensional spacetime manifold M . We show that in all dimensions the Einstein equations are classically equivalent to the field equations of general relativity with a charged dust source provided that a simple $D - 1$ dimensional Euclidean source free Maxwell's equation is satisfied. The energy density of the dust fluid is proportional to the Maxwell invariant F^2 . We next show that the geodesics of the Gödel type of metrics are described by solutions of the $D - 1$ dimensional Euclidean Lorentz force equation for a charged particle. We then present examples of spacetimes containing closed timelike and closed null curves which violate causality and examples of spacetimes without any closed timelike or closed null curves where causality is preserved. We show that the Gödel type of metrics we introduce provide exact solutions to various kinds of supergravity theories in five, six, eight, ten and eleven dimensions.

We next consider an interesting class of the Gödel type of metrics by taking a $D - 1$ dimensional non-flat background $h_{\mu\nu}$. We specifically consider the cases when the background $h_{\mu\nu}$ is conformally flat, an Einstein space and, as a subclass, a Riemannian Tangherlini solution. We explicitly construct such examples for $D = 4$ even though these can be generalized to dimensions $D > 4$ as well. When the background is an Einstein space, the corresponding source for the Einstein equations in D dimensions turns out to be a charged perfect fluid with pressure density $p = \frac{1}{2}(3 - D)\Lambda$ (so that $p > 0$ when $\Lambda < 0$) and energy density $\rho = \frac{1}{4}f^2 + \frac{1}{2}(D - 1)\Lambda$, where Λ is the cosmological constant and f^2 denotes the Maxwell invariant. We also discuss the existence of closed timelike and closed null curves in this class of spacetimes and explicitly construct geometries with and without such curves in $D = 4$. We show that when the background is a Riemannian Tangherlini space, the D dimensional solution turns out to describe a black hole depending on the parameters. We then finish off with our conclusions and a discussion of possible future work.

II. GÖDEL TYPE OF METRICS

Let M be a D dimensional manifold with a metric of the form

$$g_{\mu\nu} = h_{\mu\nu} - u_\mu u_\nu . \quad (1)$$

Here $h_{\mu\nu}$ is a degenerate $D \times D$ matrix with rank equal to $D - 1$. Throughout much of this work, we will further assume that $h_{0\mu} = 0$, $h_{ij} = \bar{\delta}_{ij}$, the $D - 1$ dimensional Kronecker delta symbol and $\partial_\alpha h_{\mu\nu} = 0$. We take Greek indices to run from $0, 1, \dots$ to $D - 1$ whereas the Latin indices range from 1 to $D - 1$. We also assume that u^μ is a timelike unit vector, i.e. $u_\mu u^\mu = -1$. (Our conventions are similar to the conventions of Hawking-Ellis [3].) The determinant of $g_{\mu\nu}$ is then $g = -u_0^2$ and moreover $u^\mu = -\frac{1}{u_0} \delta_0^\mu$. In what follows, we will also assume that $u_0 = 1$ and that u_μ is time independent, i.e. $\partial_0 u_\alpha = 0$.

With these assumptions, it is not hard to show that $u^\mu h_{\mu\nu} = 0$ and the inverse of the metric can be calculated as

$$g^{\mu\nu} = \bar{h}^{\mu\nu} + (-1 + \bar{h}^{\alpha\beta} u_\alpha u_\beta) u^\mu u^\nu + u^\mu (\bar{h}^{\nu\alpha} u_\alpha) + u^\nu (\bar{h}^{\mu\alpha} u_\alpha) . \quad (2)$$

Here $\bar{h}^{\mu\nu}$ is the $D - 1$ dimensional inverse of $h_{\mu\nu}$; i.e. $\bar{h}^{\mu\nu} h_{\nu\alpha} = \bar{\delta}_\alpha^\mu$ with $\bar{\delta}_\alpha^\mu$ denoting the $D - 1$ dimensional Kronecker delta: $\delta_\alpha^\mu = \bar{\delta}_\alpha^\mu + \delta_0^\mu \delta_\alpha^0$.

The Christoffel symbols can now be calculated as

$$\Gamma^\mu_{\alpha\beta} = \frac{1}{2} (u_\alpha f^\mu_{\beta} + u_\beta f^\mu_{\alpha}) - \frac{1}{2} u^\mu (u_{\alpha,\beta} + u_{\beta,\alpha}) \quad (3)$$

where we have used $f_{\alpha\beta} \equiv u_{\beta,\alpha} - u_{\alpha,\beta}$; $u_{\alpha,\beta} \equiv \partial_\beta u_\alpha$ and $f^\mu_{\nu} = g^{\mu\alpha} f_{\alpha\nu}$. We will also use a semicolon to denote a covariant derivative with respect to the Christoffel symbols given above; $u_{\alpha;\beta} \equiv \nabla_\beta u_\alpha$. Now one can easily show the following identities

$$u^\alpha u_{\alpha;\beta} = 0 , \quad (4)$$

$$u^\alpha f_{\mu\alpha} = 0 , \quad (5)$$

$$u^\alpha u_{\beta;\alpha} = 0 , \quad (6)$$

$$u_{\beta;\alpha} = \frac{1}{2} f_{\alpha\beta} , \quad (7)$$

$$u^\alpha_{;\alpha} = 0 , \quad (8)$$

$$\Gamma^\mu_{\mu\nu} = 0 . \quad (9)$$

The first identity is straightforward whereas the second one follows from the assumption that $u_0 = 1$ and $u^\mu = -\delta_0^\mu$. The next three identities all follow from the second one. Finally, since $\sqrt{-g} = 1$, one obtains the last identity. Notice that by (6), u^μ is tangent to a timelike geodesic curve and is a timelike Killing vector.

With the help of these identities, the Ricci tensor can be calculated as

$$R_{\mu\nu} = \frac{1}{2} f_\mu{}^\alpha f_{\nu\alpha} - \frac{1}{2} (u_\mu j_\nu + u_\nu j_\mu) + \frac{1}{4} f^2 u_\mu u_\nu \quad (10)$$

where we have used $f^2 \equiv f^{\alpha\beta} f_{\alpha\beta}$ and $j_\mu \equiv \partial_\alpha f_\mu{}^\alpha$. [Note that it is not possible to have $j_\mu = k u_\mu$ for a nontrivial constant k . Suppose the contrary is true, i.e. $j_\mu = k u_\mu$. Now by (5),

$$\partial_\alpha (u^\mu f_\mu{}^\alpha) = 0 .$$

However $u^\mu = -\delta_0^\mu$ and this gives $u^\mu j_\mu = 0$ which implies that $j_0 = k = 0$.] The Ricci scalar is then easily found as

$$R = \frac{1}{4} f^2 - u^\mu j_\mu . \quad (11)$$

Choosing $j_\mu = 0$, the Einstein tensor is simply given by

$$G_{\mu\nu} = \frac{1}{2} T_{\mu\nu}^f + \frac{1}{4} f^2 u_\mu u_\nu , \quad (12)$$

where

$$T_{\mu\nu}^f \equiv f_{\mu\alpha} f_\nu{}^\alpha - \frac{1}{4} g_{\mu\nu} f^2$$

is the Maxwell energy-momentum tensor for $f_{\mu\nu}$. Obviously, (12) implies that the Gödel type of metric $g_{\mu\nu}$ (1) is a solution of the charged dust field equations in D dimensions. The energy density of the dust fluid is just $\frac{1}{4} f^2$ then. Now one only needs to make sure that $j_\mu = 0$ is satisfied. However, this implies that

$$\partial_i f_{ij} = 0 , \quad (13)$$

with our choice of $h_{\mu\nu}$ and u_μ . Hence all that is left to solve is the flat $D - 1$ dimensional Euclidean source free Maxwell's equation. In covariant form (13) can also be written as

$$J^\nu \equiv \nabla_\mu f^{\mu\nu} = \frac{1}{2} f^2 u^\nu . \quad (14)$$

Indeed (14) will be useful for the remaining part of this work.

A few remarks regarding the positivity of energy and the character of the geodesics are in order at this point. For a timelike vector ξ^μ , one has $T_{\mu\nu}^f \xi^\mu \xi^\nu \geq 0$ by the very nature of $T_{\mu\nu}^f$ and since $f_{0\mu} = 0$, one has $f^2 = (f_{ij})^2 \geq 0$ as well. Hence it is readily seen that

$$G_{\mu\nu} \xi^\mu \xi^\nu = T_{\mu\nu}^f \xi^\mu \xi^\nu + \frac{1}{4} f^2 (u_\mu \xi^\mu)^2 \geq 0 ,$$

for all timelike ξ^μ and the weak energy condition is satisfied for spacetimes described by Gödel type of metrics.

As for the behavior of the geodesics, let us start by taking a geodesic curve Γ on M which is parametrized as $x^\mu(\tau)$. Using (3) and denoting derivative with respect to the affine parameter τ by a dot, the geodesic equation yields

$$\ddot{x}^\mu + f^\mu{}_\beta \dot{x}^\beta (u_\alpha \dot{x}^\alpha) - u^\mu \dot{x}^\alpha (u_{\alpha,\beta} \dot{x}^\beta) = 0 .$$

Noting that $u_{\alpha,\beta} \dot{x}^\beta = \dot{u}_\alpha$, writing $f^\mu{}_\beta$ explicitly via the inverse of the metric (2) and using (5), this becomes

$$\ddot{x}^\mu + u_\alpha \dot{x}^\alpha (\bar{h}^{\mu\sigma} + u^\mu \bar{h}^{\sigma\nu} u_\nu) f_{\sigma\beta} \dot{x}^\beta - u^\mu (\dot{u}_\alpha \dot{x}^\alpha) = 0 , \quad (15)$$

and contracting this with u_μ , one simply obtains

$$u_\mu \ddot{x}^\mu + \dot{u}_\mu \dot{x}^\mu = \frac{d}{d\tau} (u_\mu \dot{x}^\mu) = 0 .$$

Hence a constant of motion for the geodesic equation is

$$u_\mu \dot{x}^\mu = \dot{x}^0 + u_i \dot{x}^i = -e = \text{constant} . \quad (16)$$

Meanwhile setting the free index $\mu = i$ in (15), one also finds

$$\ddot{x}^i - e (\bar{h}^{i\sigma} f_{\sigma\beta} \dot{x}^\beta) = 0 ,$$

or simply

$$\ddot{x}^i = e f_{ij} \dot{x}^j , \quad (i = 1, 2, \dots, D-1) , \quad (17)$$

i.e. the $D-1$ dimensional Euclidean Lorentz force equation for a charged point particle of charge/mass ratio equal to e . [One could also obtain this by setting $\mu = 0$ in (15).] Moreover, contracting (17) further by $\dot{x}^i$, one obtains a second constant of motion

$$\dot{x}^i \dot{x}^i = \ell^2 = \text{constant} .$$

Since

$$g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu = h_{\mu\nu} \dot{x}^\mu \dot{x}^\nu - (u_\mu \dot{x}^\mu)^2 = \ell^2 - e^2 ,$$

one concludes that the nature of the geodesics necessarily depend on the sign of $\ell^2 - e^2$.

In retrospect, we have shown that the Gödel type of metric (1) solves the Einstein Maxwell dust field equations in D dimensions provided the flat $D-1$ dimensional Euclidean source free Maxwell's equation (13) holds. Moreover the geodesics of the Gödel type of metrics are described by the $D-1$ dimensional Euclidean Lorentz force equation (17).

A. A special solution to (13)

A solution to (13) is given by the simple choice

$$u_i = \frac{b}{2} J_{ij} x^j ,$$

where b is a real constant (we keep the $1/2$ factor for later convenience) and J_{ij} is fully antisymmetric with constant components that satisfy

$$J^k{}_j J^i{}_k = -\delta_j^i .$$

[Of course this is only possible when D is odd.] In this case, $f_{ij} = b J_{ij}$ and $f_{0\mu} = 0$ as before. Then

$$f_\mu{}^\alpha f_{\nu\alpha} = b^2 \bar{\delta}_{\mu\nu} = b^2 h_{\mu\nu} \quad (18)$$

and $f^2 = b^2(D-1)$. This in turn implies by (10) that (since $j_\mu = 0$)

$$R_{\mu\nu} = \frac{1}{2} b^2 g_{\mu\nu} + \frac{1}{4} b^2 (D+1) u_\mu u_\nu$$

and the Ricci scalar is $R = b^2(D-1)/4$. Hence the Einstein tensor can be written as

$$G_{\mu\nu} = \frac{1}{8} b^2 (5-D) g_{\mu\nu} + \frac{1}{4} b^2 (D+1) u_\mu u_\nu .$$

This in turn can be interpreted as coming from a perfect fluid source

$$G_{\mu\nu} = T_{\mu\nu} = p g_{\mu\nu} + (p + \rho) u_\mu u_\nu$$

by identifying the pressure p of the fluid as

$$p = \frac{1}{8} b^2 (5 - D)$$

and the mass-energy density ρ with

$$\rho = \frac{3}{8} b^2 (D - 1).$$

Notice that in this picture $p = 0$ when $D = 5$ and $p < 0$ when $D > 5$.

Alternatively, one can repeat this analysis by writing

$$h_{\mu\nu} = \frac{1}{b^2} f_\mu{}^\alpha f_{\nu\alpha} \quad \text{and} \quad u_\mu u_\nu = \frac{1}{b^2} f_\mu{}^\alpha f_{\nu\alpha} - g_{\mu\nu}$$

in (10). In this case

$$R_{\mu\nu} = \frac{D+1}{4} f_\mu{}^\alpha f_{\nu\alpha} - \frac{1}{4} f^2 g_{\mu\nu}$$

and the Ricci scalar is $R = b^2(D-1)/4$ as before. Now the Einstein tensor can be written in the form

$$G_{\mu\nu} = \frac{D+1}{4} \left[f_\mu{}^\alpha f_{\nu\alpha} - \frac{3}{2(D+1)} f^2 g_{\mu\nu} \right].$$

If one is to consider this as an Einstein Maxwell theory so that $G_{\mu\nu} \sim T_{\mu\nu}^f$, then

$$\frac{3}{2(D+1)} = \frac{1}{4}$$

which yields $D = 5$.

As a result, when $D = 5$ the special solution given above can either be thought of as describing a spacetime filled with dust or as a solution to the Einstein Maxwell theory. However, in general odd dimensions it can be considered as a solution of Einstein theory coupled with a perfect fluid source where the pressure $p < 0$ when $D > 5$.

Let us also take a look at the geodesics for the special solution described here. The $D-1$ dimensional Lorentz force equation (17) now takes the form

$$\ddot{\mathbf{x}} = \omega \mathbf{J} \dot{\mathbf{x}}, \tag{19}$$

where we have defined $\omega = eb$ and used matrix notation so that $\mathbf{x}$ denotes the $(D-1) \times 1$ column vector with entries x^i and $\mathbf{J}$ denotes the $(D-1) \times (D-1)$ matrix with components J_{ij} . Defining a second column vector $\mathbf{y}$ as $\mathbf{y} = \dot{\mathbf{x}}$, (19) can be integrated as

$$\mathbf{y} = e^{\omega\tau\mathbf{J}} \mathbf{x}_0,$$

for some constant column vector $\mathbf{x}_0$. Since $\mathbf{J}^2 = -\mathbf{I}_{D-1}$ (the $(D-1) \times (D-1)$ identity matrix and remember that D is odd) by construction, this can also be written as

$$\mathbf{y} = \dot{\mathbf{x}} = (\cos \omega\tau + \mathbf{J} \sin \omega\tau) \mathbf{x}_0,$$

or integrating once again

$$\mathbf{x}(\tau) = \frac{1}{\omega} (\sin \omega \tau - \mathbf{J} \cos \omega \tau) \mathbf{x}_0 + \mathbf{x}_1 \quad (20)$$

for a new constant column vector $\mathbf{x}_1$. Notice that

$$(\mathbf{x} - \mathbf{x}_1)^T (\mathbf{x} - \mathbf{x}_1) = \frac{1}{\omega^2} \mathbf{x}_0^T \mathbf{x}_0.$$

Defining a scalar as $\mathcal{R}^2 \equiv \mathbf{x}_0^T \mathbf{x}_0 / \omega^2$, one finds that $|\mathbf{x} - \mathbf{x}_1|^2 = \mathcal{R}^2$, which in general is the equation of a $(D - 2)$ dimensional sphere with radius $\mathcal{R}$ and center at $\mathbf{x}_1$.

The equation describing x^0 is given by (16) which reads

$$\dot{x}^0 + \frac{b}{2} \dot{\mathbf{x}}^T \mathbf{J} \mathbf{x} + e = 0$$

in matrix form. Using (20), this can be written as

$$\dot{x}^0 + \left(e + \frac{1}{2} b \omega \mathcal{R}^2 \right) + \frac{b}{2} (b_1 \sin \omega \tau + b_2 \cos \omega \tau) = 0$$

for some constants $b_1 \equiv \mathbf{x}_0^T \mathbf{x}_1$ and $b_2 \equiv \mathbf{x}_0^T \mathbf{J} \mathbf{x}_1$. Integrating this equation, one finally finds

$$x^0 = -e \left(1 + \frac{1}{2} b^2 \mathcal{R}^2 \right) \tau + \frac{1}{2e} (b_1 \cos \omega \tau - b_2 \sin \omega \tau) + b_3$$

for a new integration constant b_3 . Since the coefficient of the term linear in τ is not zero in general, x^0 can not be periodic. Hence we conclude that the geodesic curves move on $\mathbb{R} \times S^{D-2}$ and are complete for this special solution in odd dimensions.

[In fact, this result can also be generalized to the case when the rank of $\mathbf{J}$ is $2k$ where $2 \leq 2k \leq D - 1$ and $\mathbf{J}^2 = -\mathbf{I}_{2k}$, the $2k \times 2k$ identity matrix. Then $2k$ of the spatial coordinates will obey the Lorentz force equation

$$\ddot{x}^i = \omega J_{ij} \dot{x}^j, \quad (i = 1, 2, \dots, 2k),$$

whereas the remaining spatial ones will simply satisfy

$$\ddot{x}^i = 0, \quad (i = 2k + 1, \dots, D - 1).$$

Therefore the $2k$ dimensional part will be periodic in τ and will describe a $2k - 1$ dimensional sphere whereas the remaining parts will be linear in τ . Hence the geodesic curves will move on $\mathbb{R}^{D-2k} \times S^{2k-1}$ in this case.]

B. Spacetimes containing closed timelike curves

In this Subsection we give a simple solution which corresponds to a spacetime (1) that contains closed timelike or null curves. Here we take $D = 4$ for simplicity but what follows can easily be generalized to higher dimensions.

Obviously the simple choice $u_i = Q_{ij}x^j$, where Q_{ij} is fully antisymmetric with constant components ($i, j = 1, 2, 3$), solves (13). Now let $Q_{13} = Q_{23} = 0$ but $Q_{12} \neq 0$ for simplicity. Then

$$u_\mu dx^\mu = dt + Q_{12}(x^2 dx^1 - x^1 dx^2),$$

and employing the ordinary cylindrical coordinates (ρ, ϕ, z) this can be written as

$$u_\mu dx^\mu = dt - Q_{12} \rho^2 d\phi.$$

Using (1), this in turn implies that the line element $ds^2 = g_{\mu\nu} dx^\mu dx^\nu$ is

$$ds^2 = d\rho^2 + \rho^2 d\phi^2 + dz^2 - (dt - Q_{12} \rho^2 d\phi)^2.$$

Consider the curve $C = \{(t, \rho, \phi, z) \mid t = t_0, \rho = \rho_0, z = z_0\}$, where t_0, ρ_0 and z_0 are constants, in the manifold M . The norm of the tangent vector $v^\mu = \delta_\phi^\mu (\partial/\partial\phi)$ to this curve is then

$$v^2 \equiv v_\mu v^\mu = g_{\phi\phi} = \rho_0^2 (1 - (Q_{12})^2 \rho_0^2).$$

For a spacelike tangent vector, one has $v^2 > 0$, of course. The spacetime we are studying is obviously homogeneous and there passes a curve like C from each point of such a spacetime. Since ϕ is a periodic variable with $\phi = 0$ and $\phi = 2\pi$ identified, one then clearly finds that there exist closed timelike and null curves for $\rho_0 \geq 1/|Q_{12}|$ in this spacetime since then $v^2 \leq 0$. One can also show that there exists no closed timelike or null geodesics in this geometry.

C. Spacetimes without any closed timelike curves

In this Subsection we present a solution which describes a spacetime (1) that doesn't contain any closed timelike or null curves.

Now let $u_i = s(x^j) \omega_i$ where ω^i is a constant vector and s is a smooth function of the spatial coordinates x^j , ($i, j = 1, 2, \dots, D-1$). Hence $f_{ij} = (\partial_i s) \omega_j - (\partial_j s) \omega_i$ and (13) gives

$$\partial_i f_{ij} = (\nabla^2 s) \omega_j - (\partial_i \partial_j s) \omega_i = 0.$$

Then (13) is satisfied if one chooses $\nabla^2 s = 0$ and $\omega_i \partial_i s = \text{constant}$, which can further be set equal to zero.

Now let us take $D = 4$ specifically, but the following discussion can of course be generalized to $D > 4$ as well. As a simple example, choose $\omega_i = \delta_i^3$ above. Then any function s which is harmonic in the (x^1, x^2) variables will do. Now

$$u_\mu dx^\mu = dt + s(x^1, x^2) dx^3,$$

and using the cylindrical coordinates again, the line element becomes

$$ds^2 = d\rho^2 + \rho^2 d\phi^2 + dz^2 - (dt + s(\rho, \phi) dz)^2.$$

Consider the curve C and its tangent vector v^μ we used in Subsection IIB again. Now the norm of v^μ is

$$v^2 = g_{\phi\phi} = \rho_0^2,$$

and this is obviously always positive definite, $v^2 > 0$, i.e. v_μ is always spacelike. Hence there exists no closed timelike or null curves in this spacetime.

It is worth pointing out that the consideration of a curve of the form $\bar{C} = \{(t, \rho, \phi, z) \mid t = t_0, \rho = \rho_0, \phi = \phi_0, z \in [0, 2\pi)\}$, where t_0, ρ_0 and ϕ_0 are constants, and its tangent vector $\bar{v}^\mu = \delta_z^\mu (\partial/\partial z)$ gives

$$\bar{v}^2 = g_{zz} = 1 - (s(\rho_0, \phi_0))^2,$$

which at first sight indicates the existence of closed timelike or null curves in this geometry. On the other hand we don't confine ourselves to the small patch of spacetime where the z coordinate is on S^1 , we are interested in the universal covering of this patch and thus take z to be on the real line $\mathbb{R}$.

We thus conclude that the solutions we present correspond to spacetimes that contain both closed timelike and null curves and that contain neither of these depending on how one solves (13).

D. Einstein Maxwell 3-form field in D dimensions

Let us consider a theory that involves the coupling of a Maxwell 2-form field and a 3-form field with ordinary Einstein gravity in arbitrary $D > 4$ dimensions. The precise form of the field equations of such a theory can be determined once the underlying action is given, of course. However one can assume in general that the field equations will be of the form:

$$\nabla_\mu G^{\mu\nu\rho} = 0, \quad (21)$$

$$\nabla_\mu F^{\mu\nu} + c_1 G^{\nu\rho\sigma} F_{\rho\sigma} = 0, \quad (22)$$

$$R_{\mu\nu} = c_2 F_{\mu\rho} F_\nu{}^\rho + c_3 G_{\mu\rho\sigma} G_\nu{}^{\rho\sigma}, \quad (23)$$

at most.

To construct a solution to these equations, assume first that the metric $g_{\mu\nu}$ is of the form (1) and that the ordinary vector field $A^\mu = \lambda u^\mu$, where λ is a real constant. With this form of A_μ , $F_{\mu\nu} = \lambda f_{\mu\nu}$. Assume furthermore that one can take the 3-form field $G_{\mu\nu\rho}$ as

$$G_{\mu\nu\rho} = F_{\mu\nu} A_\rho + F_{\nu\rho} A_\mu + F_{\rho\mu} A_\nu \quad (24)$$

which then yields

$$G_{\mu\nu\rho} = \lambda^2 [f_{\mu\nu} u_\rho + f_{\nu\rho} u_\mu + f_{\rho\mu} u_\nu].$$

Then (21) implies that

$$\lambda^2 [f^{\mu\nu} u^\rho{}_{,\mu} + f^{\nu\rho}{}_{,\mu} u^\mu + f^{\rho\mu} u^\nu{}_{,\mu}] = 0$$

which is identically satisfied by our choice of $u^\mu = -\delta_0^\mu$.

Furthermore one can easily show, using (5), that

$$G^{\nu\rho\sigma} F_{\rho\sigma} = \lambda^3 f^2 u^\nu$$

independent of the dimension D and since $\nabla_\mu F^{\mu\nu} = \frac{\lambda}{2} f^2 u^\nu$, (22) holds provided

$$\lambda \left(\frac{1}{2} + c_1 \lambda^2 \right) = 0 \quad \text{or} \quad c_1 = -\frac{1}{2\lambda^2}$$

for a nontrivial λ .

Similarly one can calculate that

$$F_{\mu\rho} F_{\nu}{}^{\rho} = \lambda^2 f_{\mu\rho} f_{\nu}{}^{\rho} \quad \text{and} \quad G_{\mu\rho\sigma} G_{\nu}{}^{\rho\sigma} = \lambda^4 (f^2 u_{\mu} u_{\nu} - 2 f_{\mu\rho} f_{\nu}{}^{\rho})$$

which is again independent of the dimension D . Substituting these in (23) and using (10) with $j_{\mu} = 0$, one finds that

$$c_3 \lambda^4 = \frac{1}{4} \quad \text{and} \quad \frac{1}{2} = \lambda^2 (c_2 - 2c_3 \lambda^2)$$

which imply

$$c_2 = \frac{1}{\lambda^2} \quad \text{and} \quad c_3 = \frac{1}{4\lambda^4}.$$

Hence the Gödel type of metric (1), our choice of A_{μ} and our construction of the 3-form field $G_{\mu\nu\rho}$ (24) provide exact solutions to the field equations of the form (21), (22) and (23) which in general turn up in theories that involve the coupling of Einstein's gravity with both a 2-form and a 3-form field, as we show in some particular examples in the next Section.

III. SOLUTIONS OF SUPERGRAVITY THEORIES IN VARIOUS DIMENSIONS

In this Section, we use the results we have obtained so far in constructing solutions to some supergravity theories in dimensions $D \geq 5$.

A. Five dimensions

The Bosonic part of the minimal supergravity in $D = 5$ has the following field equations [17], [18]

$$R_{\mu\nu} = 2 (F_{\mu\alpha} F_{\nu}{}^{\alpha} - \frac{1}{6} g_{\mu\nu} F^2) \Leftrightarrow G_{\mu\nu} = 2 T_{\mu\nu}^F, \quad (25)$$

$$\nabla_{\mu} F^{\mu\nu} = \frac{1}{2\sqrt{3}} \eta^{\alpha\beta\gamma\mu\nu} F_{\alpha\beta} F_{\gamma\mu}, \quad (26)$$

where the Levi-Civita tensor η is given in terms of the Levi-Civita tensor density ϵ by $\eta^{\alpha\beta\gamma\mu\nu} = \epsilon^{\alpha\beta\gamma\mu\nu} / \sqrt{-g}$.

Let $g_{\mu\nu} = h_{\mu\nu} - u_{\mu} u_{\nu}$ (as in Section II) and $A_{\mu} = b u_{\mu}$, where b is a real constant. One then has $F^{\mu\nu} = b f^{\mu\nu}$.

Now let us concentrate on (26) first. By (14), $\nabla_{\mu} F^{\mu\nu} = \frac{b}{2} f^2 u^{\nu}$ and since $u^{\nu} = -\delta_0^{\nu}$, one finds that

$$\nabla_{\mu} F^{\mu\nu} = -\frac{b}{2} f^2 \delta_0^{\nu}.$$

Now when $\nu = i$, one of the first four indices of η on the right hand side of (26) has to be 0 and since $F_{0k} = 0$, (26) is satisfied identically in this case. When $\nu = 0$, $\nabla_{\mu} F^{\mu 0} = -\frac{b}{2} f^2$. On the other hand, the right hand side of (26) gives

$$\frac{1}{2\sqrt{3}} \epsilon^{\alpha\beta\gamma\mu 0} F_{\alpha\beta} F_{\gamma\mu} = \frac{b^2}{2\sqrt{3}} \epsilon^{ijkl} f_{ij} f_{kl}$$

since $\sqrt{-g} = 1$. So if one chooses f_{ij} to be further (anti) self dual in Euclidean $\mathbb{R}^4$, i.e. $f_{ij} = \pm \frac{1}{2} \epsilon_{ijkl} f_{kl}$ (in addition to the condition (13)), then the right hand side of (26) gives $\pm \frac{b^2}{\sqrt{3}} f^2$ and comparing with the left hand side, one finds $b = \mp \frac{\sqrt{3}}{2}$.

Paraphrasing, (26) is satisfied provided $b = \mp \frac{\sqrt{3}}{2}$, $f_{\mu\nu}$ satisfies (14) and f_{ij} is (anti) self dual in the Euclidean $\mathbb{R}^4$ space. Such an $f_{\mu\nu}$ can easily be constructed by choosing $u_i = J_{ij} x^j$ ($i, j = 1, 2, 3, 4$), where all components of the fully antisymmetric J_{ij} are constants with

$$J^k{}_j J^i{}_k = -\delta^i_j,$$

i.e. J_{ij} define an almost complex structure in $\mathbb{R}^4$. Moreover, for f_{ij} to be (anti) self dual, J_{ij} must itself be (anti) self dual in $\mathbb{R}^4$, i.e. $J_{ij} = \pm \frac{1}{2} \epsilon_{ijkl} J_{kl}$.

With the special choice of u_i as above,

$$T_{\mu\nu}^f = \frac{1}{4} f^2 u_\mu u_\nu \quad \text{and} \quad T_{\mu\nu}^F = b^2 T_{\mu\nu}^f.$$

Looking back at (25) and using (12) of Section II, one finds that $T_{\mu\nu}^f = \frac{1}{2(4b^2-1)} f^2 u_\mu u_\nu$. Comparing with the $T_{\mu\nu}^f$ above, this again yields $b = \mp \frac{\sqrt{3}}{2}$ as before. In fact this solution is supersymmetric [14], [15], [17], [18].

B. Six dimensions

In our conventions, the Bosonic part of the $D = 6, N = 2$ supergravity [22] reduces to the following field equations when all the scalars of the hypermatter ϕ^a and the 2-form field $B_{\mu\nu}$ in the theory are set to zero and when one assumes the dilaton φ to be constant with $\mu \equiv e^{\sqrt{2}\varphi}$:

$$R_{\mu\nu} = 2\mu F_{\mu\rho} F_\nu{}^\rho + \mu^2 G_{\mu\rho\sigma} G_\nu{}^{\rho\sigma}, \quad (27)$$

$$\nabla_\mu F^{\mu\nu} - \mu G^{\nu\rho\sigma} F_{\rho\sigma} = 0, \quad (28)$$

$$\nabla_\mu G^{\mu\nu\rho} = 0, \quad (29)$$

$$\frac{1}{3} \mu G_{\mu\nu\rho} G^{\mu\nu\rho} + \frac{1}{2} F_{\mu\nu} F^{\mu\nu} = 0. \quad (30)$$

Here all Greek indices run from 0 to 5 and $G_{\mu\nu\rho}$ is given by

$$G_{\mu\nu\rho} = F_{\mu\nu} A_\rho + F_{\nu\rho} A_\mu + F_{\rho\mu} A_\nu \quad (31)$$

and instead of a Yang-Mills field, we have taken an ordinary vector field A^μ to be present. Notice the striking resemblance of this theory with the model we described and solved in Subsection II D.

Now let $g_{\mu\nu} = h_{\mu\nu} - u_\mu u_\nu$ (as in Section II) and $A_\mu = \lambda u_\mu$, where λ is a real constant. Then (29) is satisfied identically since $u^\mu = -\delta_0^\mu$ for our choice. One also finds that with this A_μ ,

$$G_{\mu\nu\rho} G^{\mu\nu\rho} = -3\lambda^4 f^2 \quad \text{and} \quad F_{\mu\nu} F^{\mu\nu} = \lambda^2 f^2.$$

Hence (30) holds provided $\mu\lambda^2 = 1/2$. Since $\nabla_\mu F^{\mu\nu} = \frac{\lambda}{2} f^2 u^\nu$ for us and since

$$G^{\nu\rho\sigma} F_{\rho\sigma} = \lambda^3 f^2 u^\nu,$$

one again finds that (28) is satisfied when $\mu\lambda^2 = 1/2$. Finally, noting that

$$F_{\mu\rho} F_{\nu}{}^{\rho} = \lambda^2 f_{\mu\rho} f_{\nu}{}^{\rho}, \quad G_{\mu\rho\sigma} G_{\nu}{}^{\rho\sigma} = \lambda^4 (f^2 u_{\mu} u_{\nu} - 2f_{\mu\rho} f_{\nu}{}^{\rho})$$

and using (10), one finds that (27) is again satisfied when $\mu\lambda^2 = 1/2$.

Hence our Gödel type of metric (1) and choice of A_{μ} yield a class of exact solutions to $D = 6, N = 2$ supergravity theory. It should be further investigated to see whether this class of solutions preserve any supersymmetry.

C. Eight dimensions

The Bosonic part of the gauged $D = 8, N = 1$ supergravity theory coupled to n vector multiplets [23] has field equations which are very similar to the field equations of $D = 6, N = 2$ supergravity that we have examined in Subsection III B. Taking an ordinary vector field instead of a Yang-Mills field and setting the 2-form field B_{MN} equal to zero (as was done in Subsection III B), one again has

$$G_{MNP} = F_{MN} A_P + F_{NP} A_M + F_{PM} A_N, \quad (32)$$

similar to (31), where now capital Latin indices run from 0 to 7. We also set all the scalars in the theory to zero but assume that the dilaton σ is constant with $\mu \equiv e^{\sigma}$. These assumptions lead to the following field equations (see (26) of [23])

$$R_{MN} = 2\mu F_{MP} F_N{}^P + \mu^2 G_{MPS} G_N{}^{PS}, \quad (33)$$

$$\nabla_M F^{MN} = \mu G^{NPS} F_{PS}, \quad (34)$$

$$\nabla_M G^{MNP} = 0, \quad (35)$$

$$\frac{2}{9} \mu^2 G_{MNP} G^{MNP} + \frac{1}{3} \mu F_{MN} F^{MN} = 0. \quad (36)$$

which have the same form as (27), (28), (29) and (30), respectively.

Letting

$$g_{MN} = h_{MN} - u_M u_N$$

(as in Section II) and $A_M = \lambda u_M$ (with λ real), and following similar steps as in Subsection III B, it immediately follows that one obtains exact solutions to gauged $D = 8, N = 1$ supergravity with matter couplings provided $\mu\lambda^2 = 1/2$. Once again the conditions on u_M under which these solutions are supersymmetric should be studied further.

D. Ten dimensions

The following field equations can be obtained from a five dimensional action which is itself obtained by a Kaluza-Klein reduction of the type IIB supergravity theory with only a dilaton, a Ramond-Ramond 2-form gauge potential and a graviton. (The details of the reduction process, the corresponding splitting of the ten dimensional coordinates and the metric ansatz employed are explained in detail in [18] and we directly make use of the results

of that article here.)

$$\nabla_\mu F^{\mu\nu} = -\frac{1}{2} H^{\nu\rho\sigma} F_{\rho\sigma}, \quad (37)$$

$$\nabla_\mu H^{\mu\nu\rho} = 0, \quad (38)$$

$$H_{\mu\nu\rho} H^{\mu\nu\rho} = -3 F_{\mu\nu} F^{\mu\nu}, \quad (39)$$

$$G_{\mu\nu} = F_{\mu\alpha} F_\nu{}^\alpha - \frac{1}{4} g_{\mu\nu} F^{\alpha\beta} F_{\alpha\beta} + \frac{1}{4} \left(H_{\mu\alpha\beta} H_\nu{}^{\alpha\beta} - \frac{1}{6} g_{\mu\nu} H_{\rho\alpha\beta} H^{\rho\alpha\beta} \right). \quad (40)$$

Here all Greek indices run from 0 to 4.

Notice the striking resemblance of these equations with the equations of the $D = 6, N = 2$ supergravity theory of Subsection III B. We want to see whether our Gödel type of metric ansatz (1) and choice $A_\mu = \lambda u_\mu$, with λ a constant, solves equations (37), (38), (39), (40) above. We take the 2-form field B to be zero to that effect and following [18] find that $H_{\mu\nu\rho}$ is given by ($H = -A \wedge dA$)

$$H_{\mu\nu\rho} = -(F_{\mu\nu} A_\rho + F_{\nu\rho} A_\mu + F_{\rho\mu} A_\nu)$$

which already resembles the $G_{\mu\nu\rho}$ of Subsection III B. Following similar steps to what was done in Subsection III B, one can easily show that our Gödel type of metric ansatz (1) and choice of A_μ solve equations (37), (38), (39), (40) above provided that $\lambda^2 = 1$.

Following the discussion of [18], if one further assumes that u_μ is chosen in such a way that the 3-form field $H = -\star dA$, where $\star$ denotes Hodge duality with respect to the Gödel type of metric (1), and that the gauge field A is rescaled as $A \rightarrow 2A/\sqrt{3}$, this five dimensional solution can be further uplifted as the solution

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu + (dy + \frac{2}{\sqrt{3}} A_\mu dx^\mu)^2 + ds^2(\mathbb{T}^4), \quad (41)$$

$$\hat{H} = \frac{2}{\sqrt{3}} dA \wedge (dy + \frac{2}{\sqrt{3}} A) - \frac{2}{\sqrt{3}} \star dA, \quad (42)$$

of the type IIB supergravity theory. Here $ds^2(\mathbb{T}^4)$ is the metric on a flat four torus and y denotes one of the singled out extra dimensions. (See [18] for details.)

E. Eleven dimensions

The solution we gave in Subsection III A can also be uplifted to eleven dimensions as well. The field equations for the Bosonic part of $D = 11$ supergravity are as follows [24]:

$$R_{AB} = \frac{1}{12} \left(H_{ACDE} H_B{}^{CDE} - \frac{1}{12} H^2 g_{AB} \right), \quad (43)$$

$$\partial_A (\sqrt{-g} H^{ABCD}) = \frac{1}{2(4!)^2} \epsilon^{BCDMNKLPRST} H_{MNKL} H_{PRST}. \quad (44)$$

Here capital Latin indices run from 0 to 10. Now split the spacetime into $x^A = (x^\mu, x^m)$ where $\mu = 0, 1, 2, 3, 4$ of Subsection III A, $m = 5, 6, \dots, 10$ and let $u_A = (u_\mu, 0)$. With this choice of u_A , take the metric to be of Gödel type (1) as

$$g_{AB} = h_{AB} - u_A u_B. \quad (45)$$

Next define a 1-form field A as $A_A = bu_A$ where b is a real constant. Then $F = dA = bf$ where f has components $f_{\mu\nu}$ as in Subsection III A. Moreover one can also define a second 2-form $\mathcal{F}$ as

$$\mathcal{F} = \frac{2}{\sqrt{3}} (dx^5 \wedge dx^6 + dx^7 \wedge dx^8 + dx^9 \wedge dx^{10})$$

and the 3-form potential $\mathcal{G}$ as $\mathcal{G} = \mathcal{F} \wedge A$. Then the 4-form $\mathcal{H} = d\mathcal{G} = b\mathcal{F} \wedge f$. In component form, $\mathcal{H}$ can be written as

$$H_{ABCD} = b [\mathcal{F}_{AB} f_{CD} + \mathcal{F}_{AC} f_{DB} + \mathcal{F}_{AD} f_{BC} + \mathcal{F}_{CB} f_{DA} + \mathcal{F}_{DB} f_{AC} + \mathcal{F}_{CD} f_{AB}] .$$

Using the property that $\mathcal{F}_{AB} f^{BC} = 0$, one then obtains

$$H_{ACDE} H_B{}^{CDE} = 3b^2 [f^2 \mathcal{F}_{AC} \mathcal{F}_B{}^C + \mathcal{F}^2 f_{AC} f_B{}^C] .$$

Notice that the way $\mathcal{F}$ is constructed implies that $\mathcal{F}_{AC} \mathcal{F}_B{}^C = 0$ and $\mathcal{F}^2 = 8$. Substituting these in (43), one gets

$$R^\mu{}_\nu = 2b^2 [f^{\mu\sigma} f_{\nu\sigma} - \frac{1}{6} f^2 \delta_5{}^\mu{}_\nu]$$

where $\delta_5{}^\mu{}_\nu$ denotes the 5-dimensional Kronecker delta. This is exactly of the same form as (25) in $D = 5$. For the remaining field equation (44), first note that $\sqrt{-g} = 1$ and

$$\partial_A H^{ABCD} = \mathcal{F}^{BC} \partial_A f^{AD} + \mathcal{F}^{DB} \partial_A f^{AC} + \mathcal{F}^{CD} \partial_A f^{AB} \quad (46)$$

and the way $\mathcal{F}$ and f are constructed implies that only one of the terms in the right hand side of (46) survives, say for $B = \nu, C = 2a + 1, D = 2a + 2$ ($0 \leq \nu \leq 4; 2 \leq a \leq 4$). Then by (14) and $u^\mu = -\delta_0^\mu$, (44) is equivalent to

$$-\frac{b}{\sqrt{3}} f^2 \delta_0^\nu = \frac{1}{2(4!)^2} \epsilon^{\nu 2a+1 2a+2 MNKLPRST} H_{MNKL} H_{PRST} . \quad (47)$$

When $\nu = i$ ($1 \leq i \leq 4$), one of the last eight indices of ϵ on the right hand side of (47) must be a 0 and since $H_{0MNS} = 0$, (44) is satisfied identically in this case. When $\nu = 0$, the right hand side of (47) has nonzero contributions from terms of the form

$$\frac{1}{2(4!)^2} \epsilon^{0..ij..kl..} H_{ij..} H_{kl..} \quad (48)$$

where the dots indicate indices in the range 5 to 10. However the way $\mathcal{H}$ is constructed also implies that (48) is equal to [27]

$$\frac{1}{2(4!)^2} \left(\frac{4!}{2!2!} \right)^2 2 \left(\frac{2b}{\sqrt{3}} \right)^2 3 \epsilon^{ijkl} f_{ij} f_{kl}$$

which now must equal to $-\frac{b}{\sqrt{3}} f^2$ from the left hand side of (47). Remember that in $D = 5$, we chose f_{ij} to be (anti) self dual in Euclidean $\mathbb{R}^4$, which implies that

$$-\frac{b}{\sqrt{3}} = \pm \frac{b^2}{2}$$

or $b = \mp 2/\sqrt{3}$ for a solution.

Hence our Gödel type of metric (45) and choice of A^A and $\mathcal{F}$ yield a class of exact solutions to $D = 11$ supergravity theory. It should be further checked to find whether this class of solutions preserve supersymmetry.

IV. GÖDEL TYPE OF METRICS WITH $D - 1$ DIMENSIONAL NON-FLAT BACKGROUNDS

So far we have assumed that $h_{0\mu} = 0$ and $h_{ij} = \bar{\delta}_{ij}$, the $D - 1$ dimensional Kronecker delta symbol, in the Gödel type of metric (1). We have also taken $u_0 = 1$ and u_μ to be time independent with $\partial_0 u_\alpha = 0$. These assumptions simplified the calculation of the Ricci tensor (10) and we showed that for the Gödel type of metric (1) to be an exact solution to the Einstein Maxwell dust field equations in D dimensions, one had the $D - 1$ dimensional Euclidean source free Maxwell's equations (13) to solve.

Let us now examine how our analysis in Section II changes when the degeneracy of $h_{\mu\nu}$ is caused by taking $h_{k\mu} = 0$, where x^k is a fixed coordinate with $0 \leq k \leq D - 1$ (note that x^k does not necessarily have to be spatial), and by keeping the rest of $h_{\mu\nu}$, i.e. $\mu \neq k$ or $\nu \neq k$, dependent on all the coordinates x^α except x^k so that $\partial_k h_{\mu\nu} = 0$. Hence “the background” $h_{\mu\nu}$ can effectively be thought of as the metric of a $D - 1$ dimensional non-flat spacetime. Let u^μ be a timelike unit vector, $u_\mu u^\mu = -1$, as before but now assume that $u^\mu = -\delta_k^\mu$, ($u_k = 1$) and $\partial_k u_\mu = 0$.

One now finds that $u^\mu h_{\mu\nu} = 0$ and the inverse of the metric is given by (2) again. However the determinant of $g_{\mu\nu}$ is now different, $g = -h$, where h is the determinant of the $(D - 1) \times (D - 1)$ submatrix obtained by deleting the k^{th} row and the k^{th} column of $h_{\mu\nu}$. By defining the Christoffel symbols of $h_{\mu\nu}$ as

$$\gamma^\mu{}_{\alpha\beta} = \frac{1}{2} \bar{h}^{\mu\nu} [h_{\nu\alpha,\beta} + h_{\nu\beta,\alpha} - h_{\alpha\beta,\nu}], \quad (49)$$

one finds that the new Christoffel symbols of $g_{\mu\nu}$ are given by

$$\tilde{\Gamma}^\mu{}_{\alpha\beta} = \gamma^\mu{}_{\alpha\beta} + u^\mu u_\sigma \gamma^\sigma{}_{\alpha\beta} + \frac{1}{2} (u_\alpha f^\mu{}_\beta + u_\beta f^\mu{}_\alpha) - \frac{1}{2} u^\mu (u_{\alpha,\beta} + u_{\beta,\alpha}), \quad (50)$$

where $f_{\alpha\beta} \equiv u_{\beta,\alpha} - u_{\alpha,\beta}$ as before and we assume that the indices of u_μ and $f_{\alpha\beta}$ are raised and lowered by the metric $g_{\mu\nu}$; thus e.g. $f^\mu{}_\nu = g^{\mu\alpha} f_{\alpha\nu}$. By using a vertical stroke to denote a covariant derivative with respect to $h_{\mu\nu}$ so that

$$u_{\alpha|\beta} = u_{\alpha,\beta} - \gamma^\nu{}_{\alpha\beta} u_\nu, \quad (51)$$

(50) can be simply written as

$$\tilde{\Gamma}^\mu{}_{\alpha\beta} = \gamma^\mu{}_{\alpha\beta} + \frac{1}{2} (u_\alpha f^\mu{}_\beta + u_\beta f^\mu{}_\alpha) - \frac{1}{2} u^\mu (u_{\alpha|\beta} + u_{\beta|\alpha}). \quad (52)$$

Thus the ordinary commas in (3) have been replaced with vertical strokes and the Christoffel symbols (49) of $h_{\mu\nu}$ have been added to obtain the Christoffel symbols (52) of $g_{\mu\nu}$.

Notice that the definition of $f_{\alpha\beta}$ is left invariant if the ordinary derivatives are replaced by the covariant derivatives with respect to $h_{\mu\nu}$ (or $g_{\mu\nu}$ for that matter); i.e.

$$f_{\alpha\beta} = u_{\beta,\alpha} - u_{\alpha,\beta} = u_{\beta|\alpha} - u_{\alpha|\beta}. \quad (53)$$

To further remove any ambiguity, let us also denote a covariant derivative with respect to $g_{\mu\nu}$ by $\tilde{\nabla}_\mu$, thus

$$\tilde{\nabla}_\beta u_\alpha = u_{\alpha,\beta} - \tilde{\Gamma}^\nu{}_{\beta\alpha} u_\nu.$$

Using these preliminaries one can show the following:

$$u^\alpha \tilde{\nabla}_\beta u_\alpha = 0, \quad (54)$$

$$u^\alpha f_{\mu\alpha} = 0, \quad (55)$$

$$\tilde{\nabla}_\alpha u_\beta = \frac{1}{2} f_{\alpha\beta}, \quad (56)$$

$$u^\alpha \tilde{\nabla}_\alpha u_\beta = 0, \quad (57)$$

$$\tilde{\Gamma}^\nu{}_{\nu\mu} = \gamma^\nu{}_{\nu\mu}, \quad (58)$$

$$u^\alpha{}_{,\alpha} = 0, \quad (59)$$

$$u^\mu \gamma^\nu{}_{\mu\alpha} = 0, \quad (60)$$

$$\tilde{\nabla}_\alpha u^\alpha = 0. \quad (61)$$

The first one follows from $u_\mu u^\mu = -1$ whereas the second one holds since $u^\mu = -\delta_k^\mu$, $u_k = 1$ and $\partial_k u_\alpha = 0$. (56) is obtained by explicitly writing the left hand side in terms of (52), using (55), (51) and (53). (57) holds by (55) and (56). Since $\sqrt{-g} = \sqrt{|h|}$, (58) follows. Using $u^\mu = -\delta_k^\mu$, (59) holds trivially, and (60) is found by the assumptions $h_{k\mu} = 0$ and $\partial_k h_{\mu\nu} = 0$. Finally, (61) is obtained by using (59), (58) and (60) in the explicit expansion of the left hand side. Note again that by (57), u^μ is still a timelike Killing vector.

These identities help in the calculation of the Ricci tensor which turns out to be

$$\tilde{R}_{\mu\nu} = \hat{r}_{\mu\nu} + \frac{1}{2} f_\mu{}^\alpha f_{\nu\alpha} + \frac{1}{2} (u_\mu \tilde{j}_\nu + u_\nu \tilde{j}_\mu) + \frac{1}{4} f^2 u_\mu u_\nu, \quad (62)$$

where $f^2 = f^{\alpha\beta} f_{\alpha\beta}$ as before, $\tilde{j}_\mu \equiv f^\alpha{}_{\mu|\alpha}$ and $\hat{r}_{\mu\nu}$ is the Ricci tensor of $\gamma^\mu{}_{\alpha\beta}$. The Ricci scalar is now readily obtained as

$$\tilde{R} = \hat{r} + \frac{1}{4} f^2 + u^\mu \tilde{j}_\mu,$$

where $\hat{r}$ denotes the Ricci scalar of $\gamma^\mu{}_{\alpha\beta}$. [Note that

$$\hat{r} = g^{\alpha\beta} \hat{r}_{\alpha\beta} = \bar{h}^{\alpha\beta} \hat{r}_{\alpha\beta}$$

by using $u^\mu = -\delta_k^\mu$, (2) and (60) in the explicit calculation of $\hat{r}$.] Setting $\tilde{j}_\mu = 0$, the Einstein tensor is found to be

$$\begin{aligned} \tilde{G}_{\mu\nu} &= \hat{r}_{\mu\nu} - \frac{1}{2} g_{\mu\nu} \hat{r} + \frac{1}{2} T_{\mu\nu}^f + \frac{1}{4} f^2 u_\mu u_\nu, \\ &= \hat{r}_{\mu\nu} - \frac{1}{2} h_{\mu\nu} \hat{r} + \frac{1}{2} T_{\mu\nu}^f + \left(\frac{1}{4} f^2 + \frac{1}{2} \hat{r} \right) u_\mu u_\nu, \end{aligned} \quad (63)$$

where $T_{\mu\nu}^f$ denotes the Maxwell energy-momentum tensor for $f_{\mu\nu}$ as before.

Note that in fact

$$\tilde{j}_\mu = (g^{\alpha\beta} f_{\beta\mu})_{|\alpha} = (\bar{h}^{\alpha\beta} f_{\beta\mu})_{|\alpha}.$$

This follows by using (2), (55), $u^\mu = -\delta_k^\mu$ and the initial assumptions on $h_{\mu\nu}$. Hence $\tilde{j}_\mu = 0$ equivalently implies that $\bar{h}^{\mu\nu} \tilde{j}_\nu = 0$ or

$$\partial_\alpha (\bar{h}^{\alpha\mu} \bar{h}^{\beta\nu} \sqrt{|h|} f_{\mu\nu}) = 0. \quad (64)$$

Hence we find that the Einstein tensor of the $D - 1$ dimensional background $h_{\mu\nu}$ acts as a source term for the Einstein equations obtained for the D dimensional Gödel type of metric and that the curvature scalar of the background contributes to the energy density of the dust fluid provided that the $D - 1$ dimensional source free Maxwell equation (64) in the background holds.

A. Solutions with $D - 1$ dimensional conformally flat backgrounds

Let us now take the special fixed coordinate x^k as x^0 (i.e. $k = 0$) and let the background h_{ij} be conformally flat so that $h_{ij} = e^{2\psi}\bar{\delta}_{ij}$. Here Latin indices run from 1 to $D - 1$. If we denote the radial distance of $\mathbb{R}^{D-1}$ by $r = \sqrt{x^i x^i}$, take $\psi = \psi(r)$ and use a prime to denote derivative with respect to r , then one finds that (see e.g. [25])

$$\begin{aligned}\hat{r}_{ij} - \frac{1}{2} h_{ij} \hat{r} &= -\partial_i \partial_j \psi + (\partial_i \psi)(\partial_j \psi) + \bar{\delta}_{ij} \nabla^2 \psi \\ &= \bar{\delta}_{ij} \left(\psi'' + \frac{\psi'}{r} \right) + x_i x_j \left(\left(\frac{\psi'}{r} \right)^2 - \frac{\psi''}{r^2} + \frac{\psi'}{r^3} \right),\end{aligned}$$

and

$$\hat{r} = -2 e^{-2\psi} (2\nabla^2 \psi + (\nabla \psi)^2) = -2 e^{-2\psi} \left(\frac{2}{r^2} (r^2 \psi')' + (\psi')^2 \right),$$

for the special choice $D = 4$. [The discussion we present here can easily be generalized to $D > 4$ as well.]

If $\psi'' + \frac{\psi'}{r} = 0$, then one finds that $\psi = a \ln r + b$ for some constants a and b . Taking $b = 0$, this gives

$$\hat{r}_{ij} - \frac{1}{2} h_{ij} \hat{r} = \frac{a(a+2)}{r^4} x_i x_j.$$

If one chooses $a = -2$, then $h_{ij} = \frac{1}{r^4} \bar{\delta}_{ij}$ and now both $\hat{r}_{ij}$ and $\hat{r}$ vanish. One now has to solve (64) in this background to find the Gödel type of metric $g_{\mu\nu}$ which solves (63) with $\hat{r}_{\mu\nu} = \hat{r} = 0$. To construct such a solution, take $u_i = s(r) Q_{ij} x^j$ where Q_{ij} is fully antisymmetric with constant components. [Remember that $u_0 = 1$ now.] (64) implies that

$$\partial_i (r^2 f_{ij}) = \partial_i (r s' x^\ell (Q_{j\ell} x_i - Q_{i\ell} x_j) + 2 r^2 s Q_{ji}) = 0.$$

This can be further simplified as

$$(r^2 s'' + 6 r s' + 4 s) Q_{j\ell} x^\ell = 0,$$

and in general one obtains

$$u_i = \left(\frac{A}{r^4} + \frac{B}{r} \right) Q_{i\ell} x^\ell$$

for some real constants A and B .

Thus the line element corresponding the Gödel type of metric in $D = 4$ in this 3 dimensional conformally flat background

$$ds^2 = \frac{1}{r^4} (dx^2 + dy^2 + dz^2) - \left(dt + \left(\frac{A}{r^4} + \frac{B}{r} \right) Q_{i\ell} x^\ell dx^i \right)^2$$

solves the $D = 4$ Einstein charged dust field equations.

Let us further set $Q_{13} = Q_{23} = 0$ but $Q_{12} \neq 0$ for simplicity and write the resultant line element using cylindrical coordinates. [Note that $r^2 = \rho^2 + z^2$ then.] One finds

$$ds^2 = \frac{1}{r^4} (d\rho^2 + \rho^2 d\phi^2 + dz^2) - \left(dt - Q_{12} \frac{\rho^2}{r^4} (A + Br^3) d\phi \right)^2.$$

Employing the curve C of Subsection II B and its tangent vector v^μ , one finds that

$$v^2 = g_{\phi\phi} = \frac{\rho_0^2}{r_0^4} \left(1 - (Q_{12})^2 \frac{\rho_0^2}{r_0^4} (A + Br_0^3)^2 \right)$$

with $r_0^2 = \rho_0^2 + z_0^2$. Since v^2 is not positive definite in its full generality, we conclude that there exist closed timelike and closed null curves in this spacetime.

B. Solutions with $D - 1$ dimensional Einstein spaces as backgrounds (which are themselves conformally flat)

Let us again take the special fixed coordinate x^k as x^0 and the background h_{ij} as conformally flat so that $h_{ij} = e^{2\psi} \bar{\delta}_{ij}$. However let us now assume that $\hat{r}_{ij} = \Lambda h_{ij}$, where Λ denotes the cosmological constant; i.e. the background is a $D - 1$ dimensional Einstein space as well. This yields

$$\hat{r} = (D - 1)\Lambda \quad \text{and} \quad \hat{r}_{ij} - \frac{1}{2} h_{ij} \hat{r} = \left(\frac{3 - D}{2} \right) \Lambda e^{2\psi} \bar{\delta}_{ij}.$$

Substituting these in (63), one finds that when the background $h_{\mu\nu}$ is a $D - 1$ dimensional Einstein space, the Gödel type of metric $g_{\mu\nu}$ provides a solution to

$$\tilde{G}_{\mu\nu} = \left(\frac{3 - D}{2} \right) \Lambda g_{\mu\nu} + \frac{1}{2} T_{\mu\nu}^f + \left(\frac{1}{4} f^2 + \Lambda \right) u_\mu u_\nu,$$

which describes a charged perfect fluid source with

$$p = \frac{1}{2} (3 - D) \Lambda \quad \text{and} \quad \rho = \frac{1}{4} f^2 + \frac{1}{2} (D - 1) \Lambda.$$

[For $D \geq 4$, Λ must be negative in order to have a positive pressure density p .]

We now further assume that $\psi = \psi(z)$ and take $D = 4$ for simplicity. [One can again generalize the arguments we present here to $D > 4$.] Such a choice of ψ yields

$$\hat{r}_{ij} - \frac{1}{2} h_{ij} \hat{r} = -\partial_i \partial_j \psi + (\partial_i \psi)(\partial_j \psi) + \bar{\delta}_{ij} \nabla^2 \psi = (-\psi'' + (\psi')^2) \bar{\delta}_i^3 \bar{\delta}_j^3 + \psi'' \bar{\delta}_{ij},$$

where a prime denotes derivative with respect to z . One then has to solve $-\psi'' + (\psi')^2 = 0$ which yields $\psi = b - \ln|z + a|$ for some real constants a and b . Further demanding $\hat{r} = 3\Lambda$ fixes the constant b so that

$$h_{ij} = \frac{-2}{\Lambda(z + a)^2} \bar{\delta}_{ij}.$$

Thus for a physically acceptable background, it must be that $\Lambda < 0$.

Hence one now has to solve (64) in this background so that the Gödel type of metric $g_{\mu\nu}$ solves (63) in the form

$$\tilde{G}_{\mu\nu} = -\frac{1}{2}\Lambda g_{\mu\nu} + \frac{1}{2}T_{\mu\nu}^f + \left(\frac{1}{4}f^2 + \Lambda\right)u_\mu u_\nu,$$

i.e. the charged perfect fluid source has pressure density

$$p = -\frac{1}{2}\Lambda,$$

(and $p > 0$ when $\Lambda < 0$) and energy density

$$\rho = \frac{1}{4}f^2 + \frac{3}{2}\Lambda,$$

and $\Lambda < 0$ must be chosen properly so that $\rho > 0$.

To find a solution to (64), which simply takes the form

$$\partial_i((z+a)f_{ij}) = 0$$

in this background, let us use the ansatz $u_i = \bar{\delta}_i^3 s(x, y, z)$. [$u_0 = 1$ again.] Then

$$f_{ij} = \bar{\delta}_j^3 \partial_i s - \bar{\delta}_i^3 \partial_j s$$

and when the free index j is equal to 3, one finds

$$\partial_\ell((z+a)\partial_\ell s) = 0,$$

where the index ℓ runs over 1 and 2. When $j = \ell \neq 3$, one similarly obtains

$$\partial_3((z+a)\partial_\ell s) = 0,$$

which is easily integrated as

$$\partial_\ell s = \frac{c_\ell(x, y)}{z+a}$$

for some “integration constants” $c_\ell(x, y)$. Consistency with the $j = 3$ equation above further constrains c_ℓ to satisfy $\partial_\ell c_\ell = 0$. Hence letting $c_\ell = \partial_\ell c$ for a potential $c(x, y)$, one finds that

$$s(x, y, z) = \frac{c(x, y)}{z+a}$$

for a function $c(x, y)$ which is harmonic in the (x, y) variables.

Substituting these into the metric $g_{\mu\nu}$, one finds that the line element corresponding to this $D = 4$ example is given in the cylindrical coordinates as

$$ds^2 = \frac{-2}{\Lambda(z+a)^2} (d\rho^2 + \rho^2 d\phi^2 + dz^2) - \left(dt + \frac{c(x, y)}{z+a} dz\right)^2.$$

One then finds that the norm of the tangent vector v^μ to the curve C of Subsection II B is

$$v^2 = g_{\phi\phi} = \frac{-2\rho_0^2}{\Lambda(z_0+a)^2}.$$

One again sees that for v^2 to be positive definite Λ must be $\Lambda < 0$. [In that case the pressure density of the perfect fluid is also positive, $p > 0$.] Hence there exist no closed timelike or closed null curves in this spacetime provided $\Lambda < 0$.

C. $D = 3$ solutions with 2 dimensional backgrounds

In 2 dimensions regardless of what h_{ij} is, its Einstein tensor vanishes. [We take x^k as x^0 without any loss of generality.] In fact any 2 dimensional metric is conformally flat and can be put in the form $h_{ij} = e^{2\psi(x,y)} \bar{\delta}_{ij}$. In this case the Ricci scalar of h_{ij} is $\hat{r} = -2 e^{-2\psi} \nabla^2 \psi$ and the 3 dimensional Einstein equations for the Gödel type of metric simply becomes

$$\tilde{G}_{\mu\nu} = \frac{1}{2} T_{\mu\nu}^f + \left(\frac{1}{4} f^2 + \frac{1}{2} \hat{r} \right) u_\mu u_\nu. \quad (65)$$

So the only contribution of the 2 dimensional background is via its curvature scalar; the energy density of the charged dust fluid in 3 dimensions is

$$\rho = \frac{1}{4} f^2 + \frac{1}{2} \hat{r}$$

now. Of course one still has to solve the Maxwell's equation (64) which now simply reads

$$\partial_i (e^{-2\psi} f_{ij}) = 0,$$

and integrating this equation for the only nontrivial component f_{12} , one finds $f_{12} = c e^{2\psi}$ for some real constant c . Hence one simply has

$$f^2 = 2c^2, \quad T_{ij}^f = \frac{1}{2} c^2 e^{2\psi} \bar{\delta}_{ij} \quad \text{and} \quad \rho = \frac{1}{2} c^2 - e^{-2\psi} \nabla^2 \psi.$$

Therefore given ρ or equivalently $\hat{r}$ in $D = 2$, one can find the corresponding ψ , construct u_μ and thus find the Gödel type of metric $g_{\mu\nu}$ which solves the corresponding Einstein's equations (65) in $D = 3$.

D. Spacetimes with $D - 1$ dimensional Riemannian Tangherlini solutions as backgrounds

Let the $D - 1$ dimensional background metric $h_{\mu\nu}$ be the metric of an Einstein space,

$$\hat{r}_{\mu\nu} = \frac{2\Lambda}{3-D} h_{\mu\nu}, \quad (D \neq 3).$$

Then the full D dimensional Einstein tensor becomes

$$\tilde{G}_{\mu\nu} = \frac{1}{2} T_{\mu\nu}^f + \Lambda g_{\mu\nu} + \left(\frac{1}{4} f^2 + \frac{2\Lambda}{3-D} \right) u_\mu u_\nu, \quad (D \neq 3).$$

Hence our metric (1) solves the Einstein field equations with a charged dust source and a cosmological constant provided the source free Maxwell equation (64) holds. The energy density of the dust is

$$\rho = \frac{1}{4} f^2 + \frac{D-1}{3-D} \Lambda,$$

where Λ must be chosen so that $\rho > 0$.

Consider the line element corresponding to the $D-1$ dimensional Riemannian Tangherlini solution with a cosmological constant Λ

$$ds_{D-1}^2 = \zeta dt^2 + \frac{dr^2}{\zeta} + r^2 d\Omega_{D-3}^2,$$

where $\zeta = 1 - 2V$ with

$$V = \begin{cases} mr^{4-D} + \frac{\Lambda}{(D-3)(D-2)} r^2, & (D \geq 5) \\ m + \frac{\Lambda}{2} r^2, & (D = 4) \end{cases},$$

m is the constant mass parameter, $d\Omega_{D-3}^2$ is the metric on the $D-3$ dimensional unit sphere and we take the static limit so that all acceleration parameters vanish [11].

Let the special fixed coordinate x^k be x^{D-1} this time. Let us also assume that

$$u_\mu = u(r) \delta_\mu^0 + \delta_\mu^{D-1}.$$

Then

$$f_{\mu\nu} = (\delta_\mu^r \delta_\nu^0 - \delta_\mu^0 \delta_\nu^r) u',$$

where prime denotes derivative with respect to r . The only nontrivial component of (64) is obtained when $\beta = 0$ and in that case

$$(r^{D-3} u')' = 0,$$

which yields

$$u(r) = \begin{cases} ar^{4-D} + b, & (D \geq 5) \\ a \ln r + b, & (D = 4) \end{cases},$$

for some real constants a and b .

Substituting these in the Gödel type of metric (1), the D dimensional line element becomes

$$ds^2 = \zeta dt^2 + \frac{dr^2}{\zeta} + r^2 d\Omega_{D-3}^2 - (u(r) dt + dx^{D-1})^2.$$

For this solution one has

$$f^2 = \begin{cases} 2a^2(D-4)^2 r^{6-2D}, & (D \geq 5) \\ 2a^2/r^2, & (D = 4) \end{cases},$$

and one finds that the energy density of the dust diverges at $r = 0$. In the simple case $a = 0$, the Maxwell part of the full energy momentum tensor vanishes and one just has a dust source with

$$\rho = \frac{D-1}{3-D} \Lambda,$$

and for $\rho > 0$, Λ must be negative.

For $D = 4$, $\zeta = 0$ when $r^2 = (1 - 2m)/\Lambda$ and one has an event horizon at

$$r_H = \sqrt{\frac{1-2m}{\Lambda}} > 0 \quad \text{assuming} \quad \frac{1-2m}{\Lambda} > 0.$$

Similarly one can find conditions on the mass m and the cosmological constant Λ so that there exists an event horizon $r_H > 0$ which hides the singularity at $r = 0$ (which is present for $D \geq 5$). For example when $D = 5$, $\zeta = 0$ when

$$r^3 - \frac{3}{\Lambda}r + \frac{6m}{\Lambda} = 0, \quad (\Lambda \neq 0).$$

This cubic polynomial has at least one real root which can be identified as the event horizon $r_H > 0$ provided m and Λ satisfy certain inequalities. [When $\Lambda = 0$, the event horizon is at $r_H = 2m$ as expected.]

However the gist of the story is that our Gödel type of metric (1) provides a solution, which can have a black hole like singularity structure, to D dimensional Einstein Maxwell charged dust field equations with a cosmological constant.

E. Gödel metric

Finally we want to examine the original Gödel metric [1] to show how it has motivated us to christen the metrics studied in this work as Gödel type of metrics and how it takes its place among them.

The Gödel metric reads

$$ds^2 = -(dx^0)^2 + (dx^1)^2 - \frac{1}{2} e^{2x^1} (dx^2)^2 + (dx^3)^2 - 2 e^{x^1} dx^0 dx^2 \quad (66)$$

in our conventions. [In fact Gödel's original metric has an overall constant factor a^2 multiplying the line element. We have taken $a = 1$ here.]

Rearranging the terms, (66) can be further written as

$$ds^2 = (dx^1)^2 + \frac{1}{2} e^{2x^1} (dx^2)^2 + (dx^3)^2 - (dx^0 + e^{x^1} dx^2)^2, \quad (67)$$

which implies that the special fixed coordinate x^k is x^0 , the background $h_{\mu\nu}$ is non-flat and

$$u_\mu = \delta_\mu^0 + e^{x^1} \delta_\mu^2.$$

This simply gives

$$f_{\mu\nu} = e^{x^1} (\delta_\mu^1 \delta_\nu^2 - \delta_\mu^2 \delta_\nu^1)$$

and (64) yields

$$\partial_\alpha \left(\bar{h}^{\alpha\mu} \bar{h}^{\beta\nu} e^{2x^1} (\delta_\mu^1 \delta_\nu^2 - \delta_\mu^2 \delta_\nu^1) \right) = 0,$$

the only nontrivial component of which gives

$$\partial_1 (e^{2x^1} \bar{h}^{\beta 2}) = 0,$$

and this is identically satisfied. Hence $\tilde{j}_\mu = 0$ for the Gödel metric.

A straightforward calculation yields

$$\hat{r}_{\mu\nu} = -h_{\mu\nu} + \delta_\mu^3 \delta_\nu^3 \quad \text{and} \quad \hat{r} = -2,$$

which gives

$$\hat{r}_{\mu\nu} - \frac{1}{2} h_{\mu\nu} \hat{r} = \delta_\mu^3 \delta_\nu^3.$$

[In fact this shows that the background $h_{\mu\nu}$ describes a $\mathbb{R} \times AdS_2$ geometry.] Similarly it turns out that

$$f^2 = 4 \quad \text{and} \quad T_{\mu\nu}^f = g_{\mu\nu} + 2u_\mu u_\nu - 2\delta_\mu^3 \delta_\nu^3.$$

Substituting all of these in (63) finally yields

$$\tilde{G}_{\mu\nu} = \frac{1}{2} g_{\mu\nu} + u_\mu u_\nu,$$

which verifies the perfect fluid source of Gödel with $p = \rho = 1/2$, as expected [1].

Notice, however, that (67) is not the only way of rearranging the terms in (66). One could as well rewrite it as

$$ds^2 = (dx^0)^2 + (dx^1)^2 + (dx^3)^2 - \left(\sqrt{2} dx^0 + \frac{1}{\sqrt{2}} e^{x^1} dx^2 \right)^2. \quad (68)$$

The special fixed coordinate x^k is now x^2 and the background $h_{\mu\nu}$ is obviously flat in this new form. However $u_2 = e^{x^1}/\sqrt{2} \neq \text{constant}$, thus one can not study the Gödel metric (68) in the framework developed so far. A further analysis is needed which generalizes the assumption $u_k = 1$ to non-constant u_k . Work along this direction is in progress [26].

V. CONCLUSION

We have introduced and used Gödel type of metrics to find charged dust solutions to Einstein field equations in D dimensions. We started with a $D - 1$ dimensional Riemannian background (which could be taken as either flat or non-flat) and showed that solutions to D dimensional Einstein Maxwell theory with a dust source could be obtained provided source free Maxwell's equation is satisfied in the relevant background. The corresponding geodesics were also found to be described by the Lorentz force equation for a charged particle in the background geometry. We gave examples of spacetimes which contained closed timelike and closed null curves and others that contained neither of these. We used the Gödel type of metrics to find exact solutions to various kinds of supergravity theories.

In the case of non-flat backgrounds, we constructed explicit solutions for $D = 4$ when the background was taken as conformally flat, an Einstein space and a Riemannian Tangherlini solution. We showed that the Gödel type of metrics described a black hole depending on the parameters in the latter case. We also discussed the existence of closed timelike or closed null curves for conformally flat and Einstein space backgrounds.

It would be worth studying to see how much of supersymmetry is preserved in the solutions we have given to various supergravity theories here and to further seek whether Gödel type of metrics can be employed in finding new (possibly supersymmetric) solutions to others that we haven't considered. Throughout this work, we assumed the component of u_μ along the fixed special coordinate x^k to be constant. Another interesting point to investigate would be to generalize this assumption to non-constant u_k . One would expect to construct solutions to Einstein Maxwell dilaton 3-form field equations then. Work along these lines is in progress and the results will be published elsewhere [26].

VI. ACKNOWLEDGMENT

This work is partially supported by the Scientific and Technical Research Council of Turkey and by the Turkish Academy of Sciences.

-
- [1] K. Gödel, *Rev. Mod. Phys.* **21**, 447-450 (1949).
 - [2] D. Kramer, H. Stephani, M.A.H. MacCallum and E. Herlt, *Exact Solutions of Einstein's Field Equations*, (Cambridge, Cambridge University Press, 1980).
 - [3] S. Hawking and G.F.R. Ellis, *The Large Scale Structure of Space-Time* (Cambridge, Cambridge University Press, 1977).
 - [4] M.J. Rebouças and J. Tiomno, *Phys. Rev. D* **28**, 1251-1264 (1983).
 - [5] P. Kanti and C.E. Vayonakis, *Phys. Rev. D* **60**, 103519 (1999). [hep-th/9905032](#)
 - [6] A.E. Romano and C. Goebel, *Gen. Relat. Grav.* **35**, 1857-1863 (2003). [gr-qc/0208093](#)
 - [7] J.D. Barrow and M.P. Dabrowski, *Phys. Rev. D* **58**, 103502 (1998). [hep-th/9803048](#)
 - [8] R. Kerr and A. Schild, *Applications of Nonlinear Partial Differential Equations in Mathematical Physics*, Proceedings of Symposia on Applied Mathematics, (Amer. Math. Soc., Providence, R. I., 1965), Vol. XVII, p.199.
 - [9] M. Gürses and Ö. Sarioğlu, *Class. Quantum Grav.*, **19**, 4249-4261 (2002); *Erratum*, *ibid*, **20**, 1413 (2003). [gr-qc/0203097](#)
 - [10] M. Gürses and Ö. Sarioğlu, *Class. Quantum Grav.*, **20**, 351-358 (2003). [gr-qc/0207083](#)
 - [11] M. Gürses and Ö. Sarioğlu, *Some Further Properties of the Accelerated Kerr-Schild Metrics*, [gr-qc/0308020](#), (accepted for publication in *Gen. Relat. Grav.*)
 - [12] M.J. Duff, B.E.W. Nilsson and C.N. Pope, *Phys. Rept.*, **130**, 1 (1986).
 - [13] M.M. Caldarelli and D. Klemm, *Supersymmetric Gödel type universe in four dimensions*, [hep-th/0310081](#).
 - [14] E.K. Boyda, S. Ganguli, P. Hořava and U. Varadarajan, *Phys. Rev. D* **67**, 106003 (2003). [hep-th/0212087](#)
 - [15] T. Harmark and T. Takayanagi, *Nucl. Phys.* **B662**, 3-39 (2003). [hep-th/0301206](#)
 - [16] E.G. Gimon and A. Hashimoto, *Phys. Rev. Lett.*, **91**, 021601 (2003). [hep-th/0304181](#)
 - [17] J.P. Gauntlett, J.B. Gutowski, C.M. Hull, S. Pakis and H.S. Reall, *Class. Quantum Grav.* **20**, 4587-4634 (2003). [hep-th/0209114](#).
 - [18] C.A.R. Herdeiro, *Nucl. Phys.* **B665**, 189-210 (2003). [hep-th/0212002](#).
 - [19] C.A.R. Herdeiro, *Class. Quantum Grav.* **20**, 4891-4900 (2003). [hep-th/0307194](#)
 - [20] D. Brecher, P.A. DeBoer, M. Rozali and D.C. Page, *JHEP* **10**, 031 (2003). [hep-th/0306190](#)
 - [21] N. Drukker, B. Fiol and J. Simón, *Phys. Rev. Lett.*, **91**, 231601 (2003). [hep-th/0306057](#)
 - [22] H. Nishino and E. Sezgin, *Nucl. Phys.* **B278**, 353-379 (1986).
 - [23] A. Salam and E. Sezgin, *Phys. Lett.* **154B**, 37-42 (1985).
 - [24] E. Cremmer, B. Julia and J. Scherk, *Phys. Lett.* **76B**, 409-412 (1978).
 - [25] R.M. Wald, *General Relativity* (Chicago, The University of Chicago Press, 1984), see appendix D.
 - [26] M. Gürses, A. Karasu and Ö. Sarioğlu, in preparation.
 - [27] The factor 2 is inserted below to account for the $(ij) \rightleftharpoons (kl)$ interchange whereas the factor 3 is there to count the number of different choices for a , $2 \leq a \leq 4$.