On the Harmonic Oscillator Realisation of q-Oscillators

D. Gangopadhyay, A.P. Isaev
Laboratory of Theoretical Physics, JINR, Dubna, Russia

Abstract

The general version of the bosonic harmonic oscillator realisation of bosonic q-oscillators is given. It is shown that the currently known realisation is a special case of our general solution.

The investigation has been performed at the Laboratory of Theoretical Physics, JINR.

Recently, there has been much interest in quantum Lie algebras which first appeared in the investigations of the quantum inverse scattering problem while studying the Yang-Baxter equations¹. These quantum algebras can be considered as a "deformation" of the Lie algebra with the numerical deformation parameter s or $q = e^s$, such that the usual Lie algebra is reproduced in the limit $s \to 0$ or $q \to 1$. It has been shown that this structure essentially connects with quasi-triangular Hopf algebras and its generalisation to all simple Lie algebras has been given². There also exists the quantum generalisation of the Jordan-Scwinger mapping for $SU(2)_q$ algebra³. Moreover a q-oscillator realisation of many other quantum algebras has also been obtained^{4,5}. In Ref.4 a harmonic oscillator representation of the q-oscillators was also given. The motive of this paper is to show that the harmonic oscillator realisation of the q-oscillators admits a more general solution than the one currently in vogue⁴.

The basic equations characterising the q-deformed bosonic oscillator system are

$$aa^{\dagger} - qa^{\dagger}a = q^{-N} \quad ; \quad N^{\dagger} = N$$
 (1)

$$[N, a] = -a \; ; \; Na = a(N-1)$$
 (2)

$$[N, a^{\dagger}] = a^{\dagger} ; Na^{\dagger} = a^{\dagger}(N+1)$$
(3)

where a, a^{\dagger} are the annihilation and creation operators and N is the number operator.

Consider the case when q is complex. Then (1) implies

$$aa^{\dagger} - q^*a^{\dagger}a = (q^*)^{-N} \tag{4}$$

So from (1) and (2) we get

$$a^{\dagger}a = \frac{q^{-N} - (q^*)^{-N}}{q^* - q} \tag{5}$$

Multiplying (5) by a and then commuting a to the right in the right-hand side term we obtain

$$aa^{\dagger} = \frac{q^{-N-1} - (q^*)^{-N-1}}{q^* - q} \tag{6}$$

Substituting (5) and (6) in (1) then gives

$$q^{-N}(q^* - q^{-1}) = (q^*)^{-N}(q - (q^*)^{-1})$$
(7)

Now taking $q=|q|e^{i\alpha},q^*=|q|e^{-i\alpha}$ and putting these in (7) we have

$$e^{-i\alpha(N+1)}(|q| - \frac{1}{|q|}) = e^{i\alpha(N+1)}(|q| - \frac{1}{|q|})$$
(8)

Equation (8) has two solutions

$$|q| = \frac{1}{|q|}$$
 i.e. $|q| = 1$ (9a)

and

$$e^{-2i\alpha(N+1)} = 1 \quad i.e. \quad \alpha = \frac{\pi m}{N+1} \tag{9b}$$

with m being some integer. The second solution is not appropriate for us as we consider q as a number and not as an operator. Let us take the first solution (9a) viz. $q = e^{i\alpha}$. Then eqs.(5) and (6) can be written as

$$a^{\dagger}a = [N]$$
 , $aa^{\dagger} = [N+1]$ (10)

where $[x] = (q^x - q^{-x})/(q - q^{-1})$. It is straightforward to verify that (10) is indeed a solution of (1) even if q is real.

Let us address ourselves to determining the representation of the operators a and a^{\dagger} in terms of ordinary oscillators \hat{a} and \hat{a}^{\dagger} described by

$$[\hat{a}, \hat{a}^{\dagger}] = 1$$
 , $\hat{N} = \hat{a}^{\dagger} \hat{a} = \hat{a} \hat{a}^{\dagger} - 1$
$$[\hat{N}, \hat{a}] = -\hat{a} ; [\hat{N}, \hat{a}^{\dagger}] = a^{\dagger}$$
 (11a)

where \hat{N} is the usual number operator. We now find the solutions for a, a^{\dagger} and N satisfying equations (1), (2) and (3) together with

$$[\hat{N}, N] = 0 , [\hat{N}, a] = -a , [\hat{N}, a^{\dagger}] = a^{\dagger}$$
 (11b)

From (11b) one immediately has

$$N = \Phi(q, \hat{N}) , a = \hat{a}f(q, \hat{N})$$
 (12a)

with Φ and f some arbitrary functions at this moment. Reality of f and (12a) then give

$$a^{\dagger} = f(q, \hat{N})\hat{a}^{\dagger} \tag{12b}$$

Substituting (12) in (1) yields

$$f^{2}(q, \hat{N}+1)(\hat{N}+1) - qf^{2}(q, \hat{N})\hat{N} = q^{-\Phi(q, \hat{N})} = q^{-N}$$
(13)

With $q = e^s$ this means

$$\Phi(q, \hat{N}) = -\frac{1}{s} ln[f^2(q, \hat{N} + 1)(\hat{N} + 1) - qf^2(q, \hat{N})\hat{N}]$$
 (14)

Now from (2) and (3) we have

$$q^{-N}a = aq^{-N+1} (15a)$$

$$q^{-N}a^{\dagger} = a^{\dagger}q^{-N-1} \tag{15b}$$

Putting equation (13) in (15a) results in the functional equation

$$F(q, \hat{N})(\frac{1}{q} + q) - F(q, \hat{N} - 1) - F(q, \hat{N} + 1) = 0$$
 (16)

where $F(q, \hat{N}) = f^2(q, \hat{N})\hat{N}$. The same equation is also obtainable from (15b).

In order to solve eq.(16) for $F(q, \hat{N})$ note that

$$F(q, \hat{N}) \to \hat{N}$$
 (17)

for $s \to 0$ or $q \to 1$. This is simply because $f(q, \hat{N}) \to 1$ $(a \to \hat{a})$ when $q \to 1$.

Hence we have the following systems of equations:

$$(q + \frac{1}{q})F(q, N) - F(q, N - 1) - F(q, N + 1) = 0$$
 (18a)

$$F(1,N) = N , \Phi(1,\hat{N}) = \hat{N}$$
 (18b)

$$F(q, \hat{N} + 1) - qF(q, \hat{N}) = q^{-\Phi(q, \hat{N})} = q^{-N}$$
(18c)

The last of these equations is essentially equation (13). From (18c) we have

$$F(q,1) = qF(q,0) + q^{-\Phi(q,0)}$$
(19)

From (18a) and (19) we get

$$F(q,2) = q^{2}F(q,0) + (q+q^{-1})q^{-\Phi(q,0)}$$

A little algebra then leads to the general term

$$F(q, N) = q^{N} F(q, 0) + [N] q^{-\Phi(q, 0)}$$
(20)

It is readily verified that (20) satisfies (18a). Hence (20) is the solution of (18a) for arbitrary F(q,0) and $\Phi(q,0)$. Moreover, note that if $F \equiv \tilde{F}(q,N)$ is a solution of (18a), then $F \equiv \tilde{F}(q,-N)$ is also a solution.

It is by now obvious that we may write the general solution as

$$F(q,N) = \frac{q^N \Phi_1(q) - q^{-N} \Phi_2(q)}{q - q^{-1}}$$
(21)

where $\Phi_{1,2}$ are arbitrary functions with the restriction that $\Phi_{1,2}(1) = 1$. Then, using $F(q, \hat{N}) = f^2(q, \hat{N})\hat{N}$ we arrive at

$$f(q, \hat{N}) = \sqrt{\frac{q^{\hat{N}}\Phi_1 - q^{-\hat{N}}\Phi_2}{\hat{N}(q - q^{-1})}}$$

so that

$$a = \hat{a}\sqrt{\frac{q^{\hat{N}}\Phi_{1} - q^{-\hat{N}}\Phi_{2}}{\hat{N}(q - q^{-1})}}$$

$$a^{\dagger} = \sqrt{\frac{q^{\hat{N}}\Phi_{1} - q^{-\hat{N}}\Phi_{2}}{\hat{N}(q - q^{-1})}}\hat{a}^{\dagger}$$

$$N = \hat{N} - \frac{1}{s}ln\Phi_{2}$$
(22)

That the solutions (22) satisfy all the fundamental relations may be easily established. Choosing $\Phi_1 = \Phi_2 = 1$ gives the presently known realisation⁴.

Thus we prove that taking into account the additional conditions (11b), the representation (22) is the most general.

A similar analysis for fermionic q-oscillators leads to the known result $b = \hat{b}, b^{\dagger} = \hat{b}^{\dagger}$, and $M = \hat{M}$ after imposing the requirement $M = M^2$ for the number operator.

It is our pleasure to thank A.T.Filippov and J.Lukierski for enlightening discussions.

References

- Faddeev L.D., In: Les Houches Lectures 1982, North Holland, Amsterdam, 1984; Kulish P., Sklyanin E.K., Lecture Notes in Physics V151, Springer, Berlin, 1982, p.61; Kulish P., Reshetikin N.Yu., Zap. Nauk. Semin, 1981, LOMI 101, p.101; Jour. Sov. Math., 1983, 23 p.2435; Sklyanin E.K., Funk. Anal. Appl., 1982, 16 p.27; 1983, 17, p.34.
- [2] Drinfeld V.G., In: Proceedings of the International Congress of Mathematicians, Berkley, vol.1, 1986, p.198 (AMS,1986); Jimbo M., Lett.Math.Phys., 1985,10, p.63; ibid.,1986, 11, p.247; Comm. Math.Phys., 1987, 102, p.537; Woronowicz S.L., Comm.Math.Phys., 1987, 111, p.613; Manin Y.I., In: Quantum Groups and Non-Commutative Geometry, Centre de Recheres Mathematiques, Montreal, 1988.
- [3] Mac Farlane A. J. J. Phys., 1989, A22, p.4581; Biedenharn L.C. J. Phys.,
 1989, A22, p.L873; Sun C.P., Fu H.C., J. Phys., 1989, A22 p.L983.
- [4] Polychronakos A.P. A Classical Realisation of Quantum Algebras Univ. of Florida preprint HEP-89-23, 1989; Kulish P.P., Damaskinsky E.V., J.Phys.,1990, **A23**,p.L415; Floreanini E., Spiridonov V.P., Vinet L., Phys.Lett, 1990, **242**, p.383; Floreanini E., Spiridonov V.P., Vinet L., q-Oscillator Realisations of the Quantum Superalgebras $sl(m,n)_q$ and $osp(m,2n)_q$, preprint UCLA-90-TEP21.
- [5] Chaichian M., Kulish P., Phys. Lett., 1990, B234, p.72; Chaichian M.,
 Kulish P., Lukierski J. Phys. Lett., 1990, B237, p.401.