The Geometry of N=2 Strings with Torsion

C.M. Hull

Physics Department, Queen Mary and Westfield College, Mile End Road, London E1 4NS, U.K.

ABSTRACT

N=2 string theories are formulated in space-times with 2 space and 2 time dimensions. If the world-sheet matter system consists of 2 chiral superfields, the space-time is Kahler and the dynamics are those of anti-self-dual gravity. If instead one chiral superfield and one twisted chiral superfield are used, the space-time is a hermitian manifold with torsion and a dilaton. The string spectrum consists of a scalar, which is a potential K determining the metric, torsion and dilaton. The dynamics imply that the curvature with torsion is anti-self-dual, and an action is found for the potential K. It is argued that any N=2 sigma-model with twisted chiral multiplets in any dimension can be deformed to a conformally invariant theory if the lowest order contribution to the conformal anomaly vanishes. If there are isometries, more general geometries are possible in which the dilaton is the Killing potential for a holomorphic Killing vector.

In [1] a superstring with N=2 supersymmetry was shown to describe self-dual gravity in a Kahler space-time with signature (2,2) (or (4,0)). The world-sheet matter system consisted of two chiral scalar superfields Z^{α} ($\alpha = 1, 2$) satisfying

$$\bar{D}_{\pm}Z^{\alpha} = 0, \qquad D_{\pm}\bar{Z}^{\bar{\beta}} = 0 \tag{1}$$

where +,- are chiral spinor indices. (The superspace conventions are as in [2].) The lowest components of the superfield, $Z^{\alpha}|_{\theta=0}=z^{\alpha}$, are the bosonic complex coordinates of the space-time. The matter system is given by the N=2 sigma-model

$$S = \int d^2 \sigma d^4 \theta \, K(Z, \bar{Z}) \tag{2}$$

where K is the Kahler potential, so that the metric is

$$g_{\alpha\bar{\beta}} = \frac{\partial^2 K}{\partial z^\alpha \partial \bar{z}^{\bar{\beta}}} \tag{3}$$

For this to be a consistent string background, the sigma-model must be conformally invariant, which will be the case if the metric is Ricci-flat, or equivalently if the curvature is self-dual (or anti-self-dual)

$$*R_{\mu\nu\rho\sigma} \equiv \frac{1}{2} \epsilon_{\mu\nu}{}^{\lambda\tau} R_{\lambda\tau\rho\sigma} = \pm R_{\mu\nu\rho\sigma} \tag{4}$$

where $\mu, \nu \dots = 1, \dots 4$ are coordinate indices in a real coordinate system. As

$$R_{\alpha\bar{\beta}} = \partial_{\alpha}\partial_{\bar{\beta}}\log\det g_{\alpha\bar{\beta}} \tag{5}$$

the Ricci-flatness condition can be integrated to give

$$detg_{\alpha\bar{\beta}} = f(z)\bar{f}(\bar{z}) \tag{6}$$

for some holomorphic function f(z). With a suitable choice of coordinates, the right hand side can be set to -1 for (2,2) space-time signature or to +1 for (4,0)

signature. Writing $K = \tilde{K} + \phi$ where \tilde{K} is a background potential, this gives an equation for ϕ that can be derived from the Plebanski action (in the notation of [1])

$$\int \partial \phi \bar{\partial} \phi + \frac{1}{3} \phi \partial \bar{\partial} \phi \wedge \partial \bar{\partial} \phi \tag{7}$$

In [1], it was shown that the string spectrum consists of a scalar field ϕ whose effective dynamics was found from string scattering amplitudes to be governed by precisely the effective action (7), confirming that the N=2 string is a theory of self-dual gravity.

This is not the most general way in which conformal invariance can be achieved. More generally, one can introduce a dilaton Φ coupling through a supersymmetric Fradkin-Tseytlin term so that the condition for one-loop conformal invariance becomes

$$R_{\mu\nu} = -2\nabla_{\mu}\nabla_{\nu}\Phi\tag{8}$$

which implies

$$\nabla_{\alpha}\nabla_{\beta}\Phi = 0$$

$$\partial_{\alpha}\partial_{\bar{\beta}}\left(\log \det g_{\alpha\bar{\beta}} - 2\Phi\right) = 0$$
(9)

These imply that $\xi_{\alpha} = i\partial\Phi$ is a holomorphic Killing vector with Killing potential Φ , which satisfies

$$\Phi = \frac{1}{2} \log \det g_{\alpha \bar{\beta}} \tag{10}$$

in a suitable coordinate system [3]. In addition, there is a one-loop contribution to the Virasoro central charge proportional to $(\nabla \Phi)^2$, so that there are solutions in more than four dimensions. This gives field equations for ϕ , Φ which generalise the self-duality equations described above; some solutions with non-trivial dilaton are given in [3].

However, the Kahler sigma-model is not the most general matter system with (2,2) supersymmetry. The aim of this paper is to consider N=2 strings based

on the sigma-models of [2], using the results of [2-10]. The most general sigma-model with (2,2) supersymmetry off-shell has both chiral superfields $U^a, \bar{U}^{\bar{b}}$ $(a, b = 1, 2, ..., d_1)$ satisfying the constraints

$$\bar{D}_{\pm}U^{a} = 0, D_{\pm}\bar{U}^{\bar{b}} = 0 \tag{11}$$

and twisted chiral superfields $V^i, \bar{V}^{\bar{j}}$ $(i,j=1,2,\ldots,d_2)$ satisfying the constraints

$$D_{+}V^{i} = 0, \qquad \bar{D}_{-}V^{i} = 0, \qquad D_{-}\bar{V}^{\bar{j}} = 0, \qquad \bar{D}_{+}\bar{V}^{\bar{j}} = 0$$
 (12)

The action

$$S = \int d^2 \sigma d^4 \theta \, K(U, \bar{U}, V, \bar{V}) \tag{13}$$

defines a supersymmetric non-linear sigma-model with torsion on a target space of complex dimension $d_1 + d_2$ with coordinates $x^{\mu} = (u, \bar{u}, v, \bar{v})$ ($\mu = 1, ...2(d_1 + d_2)$) where u is the lowest component of the superfield U etc. The bosonic part of the component sigma-model action is

$$S = \frac{1}{2} \int d^2 \sigma \left(g_{\mu\nu} \partial_a x^{\mu} \partial^a x^{\nu} + b_{\mu\nu} \epsilon^{ab} \partial_a x^{\mu} \partial_b x^{\nu} \right) \tag{14}$$

where the metric $g_{\mu\nu}$ and anti-symmetric tensor $b_{\mu\nu}$ whose curl defines the torsion $H_{\mu\nu\rho} = \frac{1}{2} \partial_{[\mu} b_{\nu\rho]}$ are given by

$$g_{a\bar{b}} = K_{a\bar{b}}, \qquad g_{i\bar{j}} = -K_{i\bar{j}}$$

$$b_{a\bar{j}} = K_{a\bar{j}}, \qquad b_{i\bar{b}} = K_{i\bar{b}}$$

$$(15)$$

All other components of $g_{\mu\nu}$ and $b_{\mu\nu}$ not related to these by complex conjugation or symmetry vanish, and $K_{\mu\nu...\rho}$ denotes the partial derivative $\partial_{\mu}\partial_{\nu}...\partial_{\rho}K$. The geometry is that of a hermitian locally product space with two commuting complex structures $J^{\pm\mu}_{\nu}$; see [2] for details. In the special cases in which either $d_1 = 0$ or $d_2 = 0$, the torsion vanishes and the space is Kahler. It is useful to define the connections with torsion

$$\Gamma^{\pm\rho}_{\mu\nu} = \Gamma^{\rho}_{\mu\nu} \mp H^{\rho}_{\mu\nu} \tag{16}$$

where Γ is the usual Christoffel connection. The complex structures are each covariantly constant with respect to the corresponding connection: $\nabla^+ J^+ = 0$, $\nabla^- J^- = 0$ [2], so that both connections have holonomy $U(d_1 + d_2)$. It will also be useful to define the vectors v^{\pm} by

$$v_{\mu}^{\pm} = \pm J_{\mu\nu}^{\pm} J_{\rho\sigma}^{\pm} H^{\nu\rho\sigma} \tag{17}$$

and the U(1) parts of the two curvature tensors $R^{\pm}_{\mu\nu\rho\sigma}$ (defined as in [5])

$$C^{\pm}_{\mu\nu} = J^{\pm\rho\sigma} R^{\pm}_{\mu\nu\rho\sigma} \tag{18}$$

It follows from (15) that

$$v_a^{\pm} = \partial_a \log \det(g_{i\bar{j}})$$

$$v_i^{\pm} = \partial_i \log \det(g_{a\bar{b}})$$
(19)

The sigma-model will be conformally invariant at one-loop if there is a scalar $\Phi(x)$ such that

$$R_{\mu\nu} - H_{\mu}{}^{\rho\sigma} H_{\nu\rho\sigma} = -2\nabla_{\mu}\nabla_{\nu}\Phi$$

$$\nabla^{\nu} H_{\nu\rho\sigma} = 2H_{\nu\rho\sigma}\nabla^{\nu}\Phi$$
(20)

These are satisfied if [4,5,9]

$$\log \det(g_{a\bar{b}}) - \log \det(g_{i\bar{j}}) = f(u) + \bar{f}(\bar{u}) + g(v) + \bar{g}(\bar{v})$$
(21)

with

$$\Phi = -\frac{1}{2}\log \det(g_{a\bar{b}}) \tag{22}$$

for some holomorphic functions f, g. With a suitable choice of coordinates, the

right hand side of (21) can be set to zero. Then

$$v_{\mu}^{\pm} = -2\partial_{\mu}\Phi \tag{23}$$

and it follows from [5] that

$$C^{\pm}_{\mu\nu} = 0 \tag{24}$$

so that both connections Γ^{\pm} have $SU(d_1+d_2)$ holonomy and the first Chern class vanishes. Conversely, if (24) holds, then using $C_{ia}=i\partial_i v_a-i\partial_a v_i$ and (19), it can be shown that (21),(22) hold.

Thus if (24) is satisfied, the sigma-model will be conformally invariant at one loop, but there will in general be conformal anomalies at four loops and higher [12]. It is straightforward to show, using an argument similar to that of [15], that a deformation of K can be defined order by order in the sigma-model loop-counting parameter (\hbar or α') so that the beta-functions of the deformed sigma-model vanish. Although for general sigma-models this is not sufficient for conformal invariance [11], in this case it is possible to cancel the conformal anomaly to all orders by choosing the dilaton to be given again by (22), but now with the deformed metric appearing on the right hand side.

The equations (20) imply that

$$C \equiv -R + \frac{1}{3}H^2 - 4\nabla^2\Phi + 4(\nabla\Phi)^2$$
 (25)

is a constant. The resulting conformal field theory has central charge given by

$$c = 3(d_1 + d_2) + \frac{1}{4}\alpha'C + O(\alpha'^2)$$
(26)

However, for (2,2) geometries, the identity

$$\nabla^{\mu}v_{\mu}^{\pm} = (v^{\pm})^2 - \frac{2}{3}H^2 \tag{27}$$

implies that the quantity C given by (25) is given by

$$C = 2\nabla^{\mu}v_{\mu}^{\pm} = -\nabla^{2}\Phi = 0 \tag{28}$$

when (20) holds. In four dimensions, the definition (17) implies

$$v_{\mu}^{\pm} = \pm \frac{1}{3} \epsilon_{\mu\nu\rho\sigma} H^{\nu\rho\sigma} \tag{29}$$

so that $\nabla^{\mu}v_{\mu}^{\pm}=0$ and C vanishes identically. In this case, the sigma-model has in fact (4,4) supersymmetry [2], the central charge is given exactly by $c=3(d_1+d_2)$ and the theory is conformally invariant to all orders [14]. In other dimensions, integrating (28) and using the fact that C is both constant and a total divergence gives that C=0 for compact spaces, or for non-compact ones in which $\partial\Phi$ tends to zero sufficiently fast asymptotically. In other cases it can be non-zero, so there are quantum corrections to the central charge.

These geometries are the generalisations of the Ricci-flat Kahler spaces. As in the untwisted case, the general solution of (20) involves Killing vectors. Let

$$\xi_a = i\partial_a \left[\log \det(g_{i\bar{j}}) - 2\Phi \right]$$

$$\xi_i = i\partial_i \left[\log \det(g_{a\bar{b}}) - 2\Phi \right]$$
(30)

Then (20) implies that ξ_a and ξ_i are two commuting Killing vectors that are holomorphic with respect to both complex structures and which satisfy

$$\partial_a \xi^i = 0, \qquad \partial_i \xi^a = 0 \tag{31}$$

A Killing vector ξ^{μ} will leave the torsion $H_{\mu\nu\rho}$ invariant provided

$$H_{\mu\nu\rho}\xi^{\rho} = \partial_{[\mu}u_{\nu]} \tag{32}$$

for some u_{μ} . The equations (20) in addition imply that (30) satisfy (32) with

$$u_{a} = i\partial_{a} \left[\log \det(g_{a\bar{b}}) - 2\Phi \right]$$

$$u_{i} = i\partial_{i} \left[\log \det(g_{i\bar{j}}) - 2\Phi \right]$$
(33)

In general, C will be non-vanishing and the central charge gets quantum corrections, so that solutions in dimensions other than four are possible. The previous

case (21),(22) arises when the Killing vectors (30) both vanish.

Many geometries satisfying (21),(22) with vanishing ξ_i, ξ_a and leading to conformally invariant sigma-models are given implicitly by the twistor transform construction of [2] (these in fact have (4,4) supersymmetry and so do not need quantum deformations of K). An explicit compact example with $\Phi = 0$ and non-vanishing ξ_i, ξ_a was given in [13], while some non-compact examples both with and without Killing vectors were discussed in [3].

To construct a (2,2) string theory, it is necessary to choose c=6 to cancel the ghost contributions to the anomalies. The simplest case is that of backgrounds with $d_1 + d_2 = 2$ and vanishing ξ_i, ξ_a , so that C=0. This gives the Ricci-flat Kahler case if either d_1 or d_2 vanishes, while there will be non-vanishing torsion only if $d_1 = d_2 = 1$. Then the connections Γ^{\pm} both have SU(2) holonomy so that the model has (4,4) supersymmetry [8] and is perturbatively finite [14] and is in fact conformally invariant (as above). Both curvatures with torsion are anti-self-dual on both the first two and last two indices:

$$\frac{1}{2}\epsilon_{\mu\nu}{}^{\lambda\tau}R^{\pm}_{\lambda\tau\rho\sigma} = \frac{1}{2}R^{\pm}_{\mu\nu\lambda\tau}\epsilon^{\lambda\tau}{}_{\rho\sigma} = -R^{\pm}_{\mu\nu\rho\sigma}$$
 (34)

as can be checked using (24), $J_{[\mu\nu}J_{\rho\sigma]}=-\frac{1}{3}\epsilon_{\mu\nu\rho\sigma}$ and [7]

$$R^{+}_{\mu\nu\rho\sigma} = R^{-}_{\rho\sigma\mu\nu} \tag{35}$$

In this case the equation (21) reduces to the linear Laplace equation

$$K_{u\bar{u}} + K_{v\bar{v}} = 0 \tag{36}$$

(choosing coordinates so that the right hand side vanishes) which can be derived from the free effective action

$$\frac{1}{2} \int d^4x \, \partial_\mu K \partial^\mu K \tag{37}$$

This should also be the effective action that arises from considering the scattering amplitudes for such (2,2) strings; it would be interesting to check this explicitly.

To summarise, a (2,2) string theory can be constructed using d_1 chiral multiplets and d_2 twisted chiral multiplets, and the geometry is specified in terms of a potential K and a dilaton Φ . In the general situation, the gradient of the dilaton is given in terms of a Killing vector ξ_{α} . The case in which $\xi_{\alpha} = 0$ is easiest to analyse; in this case $d_1 + d_2 = 2$ and the space-time is four-dimensional. If $d_1 = 2$ or 0, then the geometry is Kahler, K is the Kahler potential and $\Phi = 0$. The dynamics require the Ricci tensor to vanish, which is equivalent to requiring the curvature to be anti-self-dual. The matter content is a scalar field governed by the Plebanski action (7). If $d_1 = d_2 = 1$, however, the geometry is no longer Kahler but is instead a complex geometry with torsion of the type introduced in [2] which is again given in terms of a potential K. The metric can have signature (2,2) or (4,0) and the dilaton is given by (22). The dynamics require the curvature with torsion to satisfy the anti-self-duality conditions (34). The string spectrum should again be the scalar field corresponding to the potential K, but now the dynamics is given by the free action (37). The target space theory should be a supersymmetric theory of self-dual supergravity in 2+2 dimensions; such theories have been studied in [16,17]. This exhausts the possibilities if the (2,2) world-sheet supersymmetry is to close off-shell. However, there remains the class of on-shell (2,2) models with two complex structures that do not commute and it would be interesting to understand the resulting string theory in this case also. Similar results can also be derived for for (2,1) strings and (2,0) strings. In particular, these involve a four-dimensional space-time for which the curvature with torsion has SU(2) holonomy, and so is selfdual [5-8]. This has particular relevance for the proposal of [18] relating the (2,1) string to the type IIB string and the D=11 membrane. This will be discussed in more detail elsewhere.

Acknowledgements: I would like to thank A. Sevrin for some helpful comments.

REFERENCES

- H. Ooguri and C. Vafa, Mod. Phys. Lett. A5 (1990) 1389; Nucl. Phys. 361 (191) 469.
- 2. S.J. Gates Jr., C.M. Hull and M. Roček, Nucl. Phys. **B248** (1984) 157.
- 3. E. Kiritsis, C. Kounnas and D. Lust, Int. J. Mod. Phys. A9 (1994) 1361.
- 4. T.H. Buscher, Phys. Lett. **159B** (1985) 127.
- 5. C.M. Hull, Nucl. Phys. **B267** (1986) 266.
- 6. C.M. Hull, Phys. Lett. 178B (1986) 357.
- C.M. Hull, in the Proceedings of the First Torino Meeting on Superunification and Extra Dimensions, edited by R. D'Auria and P. Fré, (World Scientific, Singapore, 1986).
- 8. C.M. Hull, in Super Field Theories (Plenum, New York, 1988), edited by H. Lee and G. Kunstatter.
- 9. M. Roček, in *Essays on Mirror Manifolds*, edited by S.-T. Yau, International Press, Hong Kong, 1992.
- 10. C.M. Hull and E. Witten, Phys.Lett. **160B** (1985) 398.
- 11. C.M. Hull and P.K. Townsend, Nucl. Phys. **B274** (1986) 349.
- J. Grundberg, A. Karlhede, U. Lindstrom, G. Theodoridis, Nucl. Phys. B282 (1987) 142; Class. Quant. Grav. 3L (1986) 129; I. Jack, Nucl. Phys. B371 (1992) 482.
- 13. M. Roček, K. Schoutens and A. Sevrin, Phys. Lett. **265B** (1991) 303.
- 14. P.S. Howe and G. Papadopoulos, Nucl. Phys. **B289** (1987) 264.
- 15. D. Nemeschansky and A. Sen, Phys. Lett. **178B** (1986) 385.

- S.J.Gates, Jr. and H.Nishino, Mod.Phys.Lett. A7, (1992) 2543; S.J. Gates, Jr., S.V.Ketov and H. Nishino, Phys.Lett. 307B (1993) 323; Phys.Lett. 297B (1993) 99; Nucl.Phys. B393 (1993) 149 and Phys.Lett. 307B (1993) 331.
- 17. E. Bergshoeff and E. Sezgin, Phys. Lett. **B292** (1992) 87; E. Sezgin, hep-th/9212092, in proceedings of Trieste Summer Workshop on Superstrings and Related Topics, 1992.
- 18. D. Kutasov and E. Martinec, hep-th/9602049.