## ULB-TH-98/17 ESI 644 (1998) hep-th/9812140

# The Wess-Zumino Consistency Condition For *p*-Form Gauge Theories

Marc Henneaux $^{a,b}$  and Bernard Knaepen<br/>  $^a$   $\ast$ 

<sup>a</sup> Physique Théorique et Mathématique, Université Libre de Bruxelles, Campus Plaine C.P. 231, B–1050 Bruxelles, Belgium

> <sup>b</sup> Centro de Estudios Científicos de Santiago, Casilla 16443, Santiago 9, Chile

#### Abstract

The general solution of the antifield-independent Wess-Zumino consistency condition is worked out for models involving exterior form gauge fields of arbitrary degree. We consider both the free theory and theories with Chapline-Manton couplings. Our approach relies on solving the full set of descent equations by starting from the last element down ("bottom").

<sup>\*</sup> henneaux@ulb.ac.be, bknaepen@ulb.ac.be

## 1 Introduction

p-form gauge theories are generalizations of electromagnetism in which the vector potential – a 1-form – is replaced by exterior forms of higher degree. They play an important rôle in supergravity and superstring theory.

The purpose of this paper is to derive the general solution of the antifieldindependent Wess-Zumino consistency condition [1] for free p-form theories – and theories with interactions that do not deform the gauge symmetry of the free case – as well as for theories involving Chapline-Manton couplings [2]. As is well known, the Wess-Zumino consistency condition is central to any gauge theory. For instance, at ghost number one, it constrains the candidate anomalies, while at ghost number zero, it determines the possible counterterms.

Counterterms and anomalies may actually depend also on the antifields [3, 4, 5, 6, 7, 8, 9]. Thus, one should in fact solve the Wess-Zumino consistency condition in the enlarged space containing these variables. However, as experience with the Yang-Mills theory indicates [10], it is useful to first work out the restricted, antifield-independent problem before tackling the full question. The complete solution of the Wess-Zumino consistency condition in the space of functions of the fields and the antifields will be given elsewhere [11].

The Wess-Zumino consistency condition takes the form

$$\gamma a + db = 0. \tag{1.1}$$

Here, a and b are local spacetime forms (for a precise definition, see below) while the differential d is the familiar exterior derivative operator acting on local forms. The differential  $\gamma$  is the BRST differential in the space of the fields, the ghosts and the ghosts of ghosts, whose explicit form depends on the theory at hand. It will be explicitly written down in the next section. Trivial solutions of (1.1) read  $a = \gamma m + dn$  for some local forms m and n. The physically distinct solutions of the Wess-Zumino consistency condition are obtained by quotienting out the trivial ones and are thus parametrized by the cohomology groups  $H(\gamma|d)$  of  $\gamma$  modulo d. The problem raised by the Wess-Zumino consistency condition is therefore a cohomological one.

Our method for investigating (1.1) follows closely the approach developed in [12, 13, 14, 15] for Yang-Mills gauge theories. In that approach, one solves (1.1) "from the bottom", i.e., one writes down the descent equations that follow from the Wess-Zumino consistency condition and determines the most general bottom of the descent. This leads to a generalization of the "transgression" formula for p-form theories, in which one associates with each non trivial last ghost of ghost a gauge-invariant curvature with opposite statistics.

Our main result is that the most general solution of (1.1) can be expressed in terms of exterior products and exterior derivatives of the basic form-variables, up to terms that descend trivially and up to trivial solutions. We say that a solution a descends trivially if it is equivalent to a strictly  $\gamma$ -closed solution, i.e.,  $a = a' + \gamma m + dn$ , with  $\gamma a' = 0$ . Thus, it is not necessary to assume that a and b in (1.1) depend only on the basic variables through the exterior forms they define and their exterior derivatives. One may allow a priori for a general dependence on the individual components of the basic variables and their independent derivatives of arbitrarily high (but finite) order. The fact that only exterior forms are relevant (up to solutions that descend trivially) emerges from the analysis and is not a restriction. This result generalizes the theorems established for 1-forms in [16, 17] and justifies in particular the usual methods followed in the literature for determining the anomalies.

Our paper is organized as follows. In Section 2, we introduce the various models considered in the article along with their respective BRST algebra. Then, in Section 3, we calculate for each model the cohomology  $H(\gamma)$  of the BRST differential. The knowledge of this cohomology is essential in order to solve to so-called descent equations, the study of which constitutes the core of our paper. In Section 4, we explicitly show how the analysis of the descent proceeds. In Section 5, we prove our main result, i.e., that any solution of (1.1) which descends non-trivially is an element of the exterior algebra generated by the form-fields, the ghosts and their exterior derivative (up to trivial solutions). Section 6 is then devoted to the explicit calculation of the auxiliary cohomologies  $E_k$  introduced in Section 4, which enable one to obtain the non-trivial solutions of the Wess-Zumino consistency condition. Finally, in Section 7, we exhibit the various counterterms and anomalies for the models defined in Section 2.

# 2 The models

## 2.1 Generalities

We consider an arbitrary system of exterior form gauge fields  $B^a_{\mu_1...\mu_{p_a}}$  of degree  $\geq 1$ . Each exterior form gauge field is accompanied by ghosts and ghosts of ghosts of decreasing form degree and increasing ghost number [18, 19, 20, 21, 5, 22],

$$C^a_{\mu_1\dots\mu_{p_a-1}},\dots,C^a_{\mu_1\dots\mu_{p_a-j}},\dots,C^a.$$
 (2.1)

The ghost number of the "first" ghosts  $C^a_{\mu_1...\mu_{p_a-1}}$  and their Grassmann parity are equal to 1. As one moves from one term to the next one to its right in (2.1), the Grassmann parity alternates and the ghost number increases by one unit up to  $p_a$ .

We denote by  $\mathcal{P}$  the algebra of spacetime forms with coefficients that are polynomials in the fields, ghosts, ghosts of ghosts and their derivatives. Thus, *a* belongs to  $\mathcal{P}$  if and only if

$$a = \frac{1}{q!} \alpha_{\nu_1 \dots \nu_q} ([B^a_{\mu_1 \dots \mu_{p_a}}], [C^a_{\mu_1 \dots \mu_{p_a-1}}], \dots, [C^a]) dx^{\nu_1} \wedge \dots \wedge dx^{\nu_q}$$
(2.2)

where the notation f([y]) means that f depends on y and its successive derivatives up to some finite order,

$$f = f([y]) \Longleftrightarrow f = f(y, \partial_{\mu}y, \dots, \partial_{\mu_1\dots\mu_k}y).$$
(2.3)

The form a is allowed to have components of various form degrees (there is a sum over q in (2.2)) but has no explicit dependence on the spacetime coordinates  $x^{\mu}$  since this is the case usually met in practice (although such a dependence can be handled by the present methods). From now on, we shall drop explicit reference to the wedge product in formulas like (2.2). It is in the algebra  $\mathcal{P}$  that the cohomological problem of computing the cohomology of  $\gamma$  modulo d will be analysed.

In both the free and interacting cases, the differential  $\gamma$  is first defined on the undifferentiated generators  $B^a_{\mu_1\dots\mu_{p_a}}, C^a_{\mu_2\dots\mu_{p_a}}, \dots, C^a$  of the algebra. It is then extended to the differentiated generators by requiring

$$\partial_{\mu}\gamma = \gamma\partial_{\mu} \tag{2.4}$$

which, together with  $\gamma(dx^{\mu}) = 0$ , implies the anticommutation relation

$$\gamma d + d\gamma = 0. \tag{2.5}$$

Finally, one extends the differential  $\gamma$  to the whole of  $\mathcal{P}$  by using the Leibnitz rule,

$$\gamma(ab) = (\gamma a)b + (-)^{\epsilon_a}a(\gamma b), \qquad (2.6)$$

where  $\epsilon_a$  is the Grassmann parity of a.

## 2.2 Free theory

In the free case, the differential  $\gamma$  is defined on the undifferentiated generators by the equations

$$\gamma B^{a}_{\mu_{1}...\mu_{p_{a}}} = \partial_{[\mu_{1}} C^{a}_{\mu_{2}...\mu_{p_{a}}]}, \qquad (2.7)$$

$$\gamma C^{a}_{\mu_{1}...\mu_{p_{a}-k_{a}}} = \partial_{[\mu_{1}} C^{a}_{\mu_{2}...\mu_{p_{a}-k_{a}}]}, \qquad (2.8)$$

$$\gamma C^a = 0 \tag{2.9}$$

 $(k_a = 1, \ldots, p_a - 1)$ . The field strengths or "curvatures" are given by

$$H^{a} = \frac{1}{(p_{a}+1)!} H^{a}_{\mu_{1}\dots\mu_{p_{a}+1}} dx^{\mu_{1}}\dots dx^{\mu_{p_{a}+1}} = dB^{a}, \qquad (2.10)$$

where

$$B^{a} = \frac{1}{p_{a}!} B^{a}_{\mu_{1}\dots\mu_{p_{a}}} dx^{\mu_{1}}\dots dx^{\mu_{p_{a}}}.$$
 (2.11)

and fulfill

$$\gamma(H^a_{\mu_1\dots\mu_{p_a+1}}) = 0. \tag{2.12}$$

One can rewrite the BRST variations in terms of forms and exterior derivatives. This yields

$$\gamma B^a + dC_1^a = 0, \qquad (2.13)$$

$$\gamma C_1^a + dC_2^a = 0, (2.14)$$

$$\gamma C^a_{p_a-1} + dC^a_{p_a} = 0, \qquad (2.15)$$

÷

$$\gamma C^a_{p_a} = 0. \tag{2.16}$$

In these equations,  $C_j^a$   $(j = 1, ..., p_a)$  are, up to numerical factors chosen to make the equations right, the  $(p_a - j)$ -forms whose components are respectively  $C_{\mu_1...\mu_{p_a-j}}^a$ .

It is sometimes convenient to adopt a more uniform notation that combines the ghosts and the fields, and to indicate explicitly the form degree and the ghost number. Thus one sets  $B^a \equiv B^{a(p_a,0)}$  and  $C_k^a \equiv B^{a(p_a-k,k)}$ . In these notations, the BRST equations are

$$\gamma B^{a\,(p_a-k,k)} + dB^{a\,(p_a-k-1,k+1)} = 0, \ \gamma B^{a\,(0,p_a)} = 0.$$
 (2.17)

 $(k=0,\ldots,p_a-1).$ 

Because the gauge transformations of *p*-form gauge theories whose couplings involve only the curvatures  $H^a_{\mu_1...\mu_{pa+1}}$  and their derivatives, or Chern-Simons couplings [23], are identical with those of the free theory, the above BRST transformations and field strength expressions encompass also these cases. The "free theories" cover therefore a larger class of models.

## 2.3 Chapline-Manton models

It has been proved in [24, 25] that the interactions between a set of exterior form gauge fields are severely constrained by the consistency requirement that the number of independent gauge symmetries should be unchanged when the interactions are switched on. This result complements the geometric analysis of [26] where it was shown that the non-abelian Yang-Mills construction cannot be generalized to *p*-forms viewed as connections for extended objects. [Topological field theory offers ways to bypass some of the difficulties [27], but will not be discussed here].

Among the few possible consistent interactions, the Chapline-Manton couplings are particularly interesting because the gauge algebra remains closed off-shell and the reducibility identities hold strongly even after the interactions are switched on. This is crucial here as it allows one to investigate the  $\gamma$ -cohomology without the antifields. In general, the new gauge algebra closes only on-shell and the reducibility identities become on-shell relations. This occurs for the celebrated Freedman-Townsend interaction [28] and its generalizations [25] (see also [29]). In that case,  $\gamma^2 \approx 0$  and it is meaningless to consider the strong cohomology of  $\gamma$  since  $\gamma$  is no longer a differential. One must either work on-shell, or, equivalently, introduce the antifields to recover nilpotency. In the Chapline-Manton models,  $\gamma^2 = 0$  strongly and without the antifields. One can thus consider the sub-problem of computing the cohomology of  $\gamma$  in the the algebra  $\mathcal{P}$ , which is an important step in the calculation of the full BRST cohomology. The Chapline-Manton interaction has been much studied in connection with the Green-Schwarz anomaly cancellation mechanism [30, 31]. Its construction follows a generic algebraic pattern discussed in [32]. Rather than discussing the general case, which would lead to non informative and uncluttered formulas, we shall consider four particular examples which illustrate the general construction.

The Chapline-Manton model is characterized by gauge-invariant curvatures  $H^a$  which differ from the free ones by terms that vanish with the coupling constant g,

$$H^{a} = dB^{a} + g\mu^{a} + O(g^{2})$$
(2.18)

The gauge transformations are

$$\delta_{\epsilon}B^{a} = d\epsilon^{a} + g\rho^{a} + O(g^{2}). \tag{2.19}$$

Here,  $\mu^a$  is given by a sum of exterior products of B's and dB's – which must match the form degree of  $dB^a$  – while  $\rho^a$  is given by a sum of exterior products of B's, dB's and  $\epsilon$ 's (linear in the  $\epsilon$ 's). The modified curvatures and gauge transformations must fulfill the consistency condition

$$\delta_{\epsilon} H^a = 0 \tag{2.20}$$

That is, the modified curvatures should be invariant under the modified gauge transformations. Furthermore, off-shell reducibility must be preserved, i.e.,  $\delta_{\epsilon}B^{a}$  should identically vanish for  $\epsilon^{a} = d\lambda^{a} + \theta^{a}$  for some appropriate  $\theta^{a}(\epsilon, B, dB, g)$ . The Lagrangian is a function of the curvatures and their derivatives,  $\mathcal{L} = \mathcal{L}([H^{a}_{\mu_{1}...\mu_{p_{a}+1}}])$  and is thus automatically gauge-invariant. To completely specify the model, it is thus necessary to give, besides the field spectrum, the modified curvatures and gauge transformations fulfilling (2.20). In many cases, the curvatures are modified by the addition of Chern-Simons forms of same degree, but this is not the only possibility as the example 3 below indicates. We shall set in the sequel the coupling constant g equal to one.

#### 2.3.1 Model 1

The first example contains one *p*-form, denoted by  $A \equiv A^{(p,0)}$ , and one (p+1)-form, denoted by  $B \equiv B^{(p+1,0)}$ . The new field strengths are

$$F = dA + B, \ H = dB \tag{2.21}$$

while the modified gauge transformations take the form

$$\delta_{\epsilon,\eta}A = d\epsilon - \eta, \qquad (2.22)$$

$$\delta_{\epsilon,\eta}B = d\eta \tag{2.23}$$

where  $\epsilon$  is a (p-1)-form and  $\eta$  a *p*-form. The gauge transformations are abelian and remain reducible off-shell since the choice of gauge parameters  $\epsilon = d\rho + \sigma$ ,  $\eta = d\sigma$  clearly leads to no variations of the fields. The BRST transformations of the undifferentiated generators are

$$\gamma A^{(p-k,k)} + dA^{(p-k-1,k+1)} + B^{(p-k,k+1)} = 0$$
(2.24)

for the A-variables, and

$$\gamma B^{(p+1-k,k)} + dB^{(p-k,k+1)} = 0, \qquad (2.25)$$

$$\gamma B^{(0,p+1)} = 0 \tag{2.26}$$

 $(k = 0, \dots, p)$  for the *B*-ones. One has

$$\gamma F = 0 = \gamma H. \tag{2.27}$$

This model describes in fact a massive (p+1)-form. Indeed, one can use the gauge freedom of B to set A = 0. Once this is done, one is left with the Lagrangian for a massive (p+1)-form.

#### 2.3.2 Model 2

The second example contains an abelian 1-form  $A \equiv A^{(1,0)}$  and a 2*r*-form  $B \equiv B^{(2r,0)}$  (r > 0). The field strengths are

$$F = dA, \ H = dB + F^r A \tag{2.28}$$

with  $F^r \equiv FF \cdots F$  (r times). The gauge transformations read

$$\delta_{\epsilon,\eta}A = d\epsilon \tag{2.29}$$

$$\delta_{\epsilon,\eta}B = d\eta - F^r \epsilon \tag{2.30}$$

and clearly leave the curvatures invariant.

The BRST transformations are

$$\gamma A^{(1,0)} + dA^{(0,1)} = 0, \qquad (2.31)$$

$$\gamma A^{(0,1)} = 0, \qquad (2.32)$$

$$\gamma B^{(2r,0)} + dB^{(2r-1,1)} + F^r A^{(0,1)} = 0, \qquad (2.33)$$

$$\gamma B^{(2r-k,k)} + dB^{(2r-k-1,k+1)} = 0, \qquad (2.34)$$

$$\gamma B^{(0,2r)} = 0 \tag{2.35}$$

(k = 1, ..., 2r - 1).

### 2.3.3 Model 3

Let A, B and C be respectively 1-, 2- and 3-forms. Define the curvatures through

$$F = dA + B, \ H = dB, \ G = dC + AdB + (1/2)B^2.$$
 (2.36)

The gauge transformations are  $\delta_{\epsilon,\Lambda,\mu}A = d\epsilon - \Lambda$ ,  $\delta_{\epsilon,\Lambda,\mu}B = d\Lambda$  and  $\delta_{\epsilon,\Lambda,\mu}C = d\mu - \epsilon dB - \Lambda B$ , where  $\epsilon$ ,  $\Lambda$  and  $\mu$  are respectively 0-, 1- and 2-forms. The gauge algebra is non-abelian and one easily verifies that the gauge transformations are off-shell reducible.

The BRST differential is defined by

$$\gamma A^{(1,0)} + dA^{(0,1)} + B^{(1,1)} = 0, \qquad (2.37)$$

$$\gamma A^{(0,1)} + B^{(0,2)} = 0, \qquad (2.38)$$

$$\gamma B^{(2,0)} + dB^{(1,1)} = 0, \qquad (2.39)$$

$$\gamma B^{(1,1)} + dB^{(0,2)} = 0, \qquad (2.40)$$

$$\gamma B^{(0,2)} = 0,$$
 (2.41)

$$\gamma C^{(3,0)} + dC^{(2,1)} + A^{(0,1)}H + B^{(1,1)}B^{(2,0)} = 0, \qquad (2.42)$$

$$\gamma C^{(2,1)} + dC^{(1,2)} + \frac{1}{2}B^{(1,1)}B^{(1,1)} + B^{(0,2)}B^{(2,0)} = 0, \qquad (2.43)$$

$$\gamma C^{(1,2)} + dC^{(0,3)} + B^{(1,1)}B^{(0,2)} = 0, \qquad (2.44)$$

$$\gamma C^{(0,3)} + \frac{1}{2} B^{(0,2)} B^{(0,2)} = 0 \qquad (2.45)$$

This example arises in some formulations of massive supergravity in 10 dimensions [33, 34].

#### 2.3.4 Model 4

Our final example is the original Chapline-Manton model, involving a Yang-Mills connection  $A^a$  and a 2-form B. We assume the gauge group to be SU(N) for definiteness although the analysis proceeds in the same way for any other compact group. The curvatures are

$$F = dA + A^2 \tag{2.46}$$

$$H = dB + \omega_3 \tag{2.47}$$

where  $\omega^{(3,0)}(A, dA)$  is the Chern-Simons 3-form

$$\omega_3 = \frac{1}{2} \left[ tr(AdA + \frac{2}{3}A^3) \right]. \tag{2.48}$$

The BRST differential reads

$$\gamma A + DC = 0, \qquad (2.49)$$

$$\gamma C + C^2 = 0, \qquad (2.50)$$

$$\gamma B + \omega_2 + d\eta = 0, \qquad (2.51)$$

$$\gamma \eta + \omega_1 + d\rho = 0, \qquad (2.52)$$

$$\gamma \rho + \frac{1}{3} tr C^3 = 0. (2.53)$$

Here, the one-form  $\omega_1$  and the two-form  $\omega_2$  are related to the Chern-Simons form  $\omega_3$  through the descent,

$$\gamma\omega_3 + d\omega_2 = 0, \quad \omega_2 = tr(CdA), \quad (2.54)$$

$$\gamma \omega_2 + d\omega_1 = 0, \quad \omega_1 = tr(C^2 A),$$
 (2.55)

$$\gamma \omega_1 + d(\frac{1}{3}trC^3) = 0. (2.56)$$

The full BRST cohomology of this model was worked out in [35], so we shall only illustrate here some aspects of the descent equation that were not explicitly discussed in that paper.

# **3 BRST** cohomology $H^*(\gamma)$

### 3.1 Free Theory

In order to compute the general solution of the Wess-Zumino consistency condition, one needs to know  $H(\gamma)$ , i.e., the general solution of  $\gamma a = 0$  modulo coboundaries  $(a = \gamma b)$ . We start with the free models. The cohomology  $H(\gamma)$  for this case was given in [36] but without giving all the details. This is done here.

**Theorem 3.1** The cohomology of  $\gamma$  is given by,

$$H(\gamma) = \mathcal{I} \otimes \mathcal{C},\tag{3.1}$$

where  $\mathcal{C}$  is the algebra generated by the "last", undifferentiated ghosts of ghosts  $C_{p_a}^a \ (\equiv B^{a(0,p_a)})$ , and  $\mathcal{I}$  is the algebra generated by the fields strength components  $H^a_{\mu_1...\mu_{p_a+1}}$  and their derivatives. **Proof:** One follows the standard method which consists in separating the

**Proof:** One follows the standard method which consists in separating the variables into three sets obeying respectively  $\gamma x^i = 0$ ,  $\gamma y^{\alpha} = z^{\alpha}$ ,  $\gamma z^{\alpha} = 0$ . The variables  $y^{\alpha}$  and  $z^{\alpha}$  form "contractible pairs" and the cohomology is generated by the (independent) variables  $x^i$  [37]. In our case, the  $x^i$  are given by the fields strength components, their derivatives and the last (undifferentiated) ghosts of ghosts.

To arrive at the appropriate decomposition, we split the generators of  $\mathcal{P}$ , which are the successive derivatives  $\partial_{\alpha_1...\alpha_k} B^{a(l,p_a-l)}_{\mu_1...\mu_l}$  of the fields and the ghosts, into irreducible tensors under the full linear group GL(n). Since the  $\partial_{\alpha_1...\alpha_k} B^{a(l,p_a-l)}_{\mu_1...\mu_l}$  are completely symmetric in  $\alpha_1...\alpha_k$  and completely antisymmetric in  $\mu_1...\mu_l$ , they transform under GL(n) as the variables of the tensor product representation symbolically denoted by



in [38].

Convenient generators for the irreducible spaces corresponding to diagrams 1) and 2) are respectively,

$$\partial_{(\alpha_1\dots\alpha_k} B^{a(l,p_a-l)}_{[\mu_1)_1\dots\mu_l]_2} \quad \text{and} \quad \partial_{\alpha_2\dots\alpha_k} H^{a(l+1,p_a-l)}_{\alpha_1\mu_1\dots\mu_l}, \tag{3.2}$$

with  $H^{a(l,p_a-l+1)}_{\mu_1...\mu_l} = \partial_{[\mu_1} B^{a(l-1,p_a-l+1)}_{\mu_2...\mu_l]}$ . In particular,  $H^{a(p_a+1,0)}_{\mu_1...\mu_{p_a+1}} \equiv H^a_{\mu_1...\mu_{p_a+1}}$ . Here, [] and () mean respectively antisymmetrization and symmetrization; the subscript indicates the order in which the operations are done. The generators  $\partial_{\alpha_2...\alpha_k} H^{a(l+1,p_a-l)}_{\alpha_1\mu_1...\mu_l}$  are obtained by first symmetrizing  $\partial_{\alpha_1...\alpha_k} B^{a(l,p_a-l)}_{\mu_1...\mu_l}$  with respect to  $\mu_1, \alpha_2, ..., \alpha_k$  and then antisymmetrizing with respect to the  $\mu$ 's and  $\alpha_1$ . In this last step, the terms with  $\partial_{\mu_1} \partial_{\alpha_1}$  yield zero.

A direct calculation shows that

$$\gamma B^{a(l,p_a-l)}_{\mu_1\dots\mu_l} = H^{a(l,p_a-l+1)}_{\mu_1\dots\mu_l}, \qquad (3.3)$$

$$\gamma \partial_{(\alpha_1 \dots \alpha_k} B^{a(l,p_a-l)}_{[\mu_1)_1 \dots \mu_l]_2} = c \partial_{\alpha_1 \dots \alpha_k} H^{a(l,p_a-l+1)}_{\mu_1 \dots \mu_l}$$
(3.4)

with  $c = \frac{k+l}{l(k+1)}$  and  $2 \le l \le p$ , and

$$\gamma H^{a(p_a+1,0)}_{\mu_1\dots\mu_{p_a+1}} = 0, \qquad (3.5)$$

$$\gamma \partial_{\alpha_1 \dots \alpha_k} H^{a(p_a+1,0)}_{\mu_1 \dots \mu_{p_a+1}} = 0.$$
(3.6)

Similarly, the relations involving the last ghosts of ghosts are

$$\gamma B_{\mu_1}^{a(1,p_a-1)} = \partial_{\mu_1} B^{a(0,p_a)}, \qquad (3.7)$$

$$\gamma \partial_{(\alpha_1 \dots \alpha_k} B^{a(1,p_a-1)}_{\mu_1)} = \partial_{\alpha_1 \dots \alpha_k \mu_1} B^{a(0,p_a)}, \qquad (3.8)$$

$$\gamma B^{a(0,p_a)} = 0. (3.9)$$

All the generators are are now split according to the rule recalled at the beginning of the subsection: the z's are the  $B^{a(l,p_a-l)}_{\mu_1\dots\mu_l}$  with  $1 \leq l \leq p$ , and their derivatives  $\partial_{(\alpha_1\dots\alpha_k}B^{a(l,p_a-l)}_{[\mu_1)_1\dots\mu_l]_2}$  corresponding to the diagram of Young type 1). The y's are their  $\gamma$ -variations, i.e., the  $H^{a(l,p_a-l+1)}_{\mu_1\dots\mu_l}$  and their derivatives  $(2 \leq l \leq p)$ , and the derivatives  $\partial_{\alpha_1\dots\alpha_k\mu_1}B^{a(0,p_a)}$  of the last ghosts of ghosts. Of course, any identity verified by the y's is also verified by the z's since they correspond to identical Young tableaux and hence transform in the same representation of GL(n). There are thus as many independent y's as there are independent z's. Finally, the x' are the "left-over" variables, i.e., the curvatures  $H^{a(p_a+1,0)}_{\mu_1\dots\mu_{p_a+1}} \equiv H^a_{\mu_1\dots\mu_{p_a+1}}$ , their successive derivatives and the last, undifferentiated ghosts of ghosts  $B^{a(0,p_a)}$ .

The cohomology is therefore generated by

$$B^{a(0,p_a)}, \ H^{a(p_a+1,0)}_{\mu_1\dots\mu_{p+1}} \text{ and } \partial_{\alpha_1\dots\alpha_k} H^{a(p_a+1,0)}_{\mu_1\dots\mu_{p+1}}.$$
 (3.10)

This ends the proof of the theorem. As shown recently [39], the same method applies to the calculation of the  $\gamma$ -cohomology of gauge fields with more general symmetry structure [40]. Note that the generators are not independent but restricted by the Bianchi identity  $dH^{a(p_a+1,0)} = 0$ .  $\Box$ .

## **3.2** Chapline-Manton model 1

The cohomology  $H^*(\gamma)$  of the Chapline-Manton model can be worked out as in the free case, by exhibiting explicitly the contractible part of the algebra. This contractible part typically gets larger with the coupling: some cocycles are removed from  $H^*(\gamma)$ . This happens for the examples 1, 3 and 4.

In the absence of couplings, the  $\gamma$ -cohomology for the first model would be given, as the previous subsection indicates, by polynomials  $P([F_{\mu_1...\mu_{p+1}}], [H_{\mu_1...\mu_{p+2}}], A^{(0,p)}, B^{(0,p+1)})$  in the components of the curvatures and their derivatives as well as in the last, undifferentiated ghosts. When the coupling is turned on, however, some of these "x"-variables become contractible pairs and get cancelled in cohomology. Specifically, it is the last ghosts of ghosts that disappear. To see this, one first makes the same redefinitions of generators for the *B*-sector as in the free case. The *x*, *y* and *z*-variables are taken as before, except that the last, undifferentiated ghost of ghost  $B^{(0,p+1)}$  counts now as a *y* (see below).

In the A-sector, one takes for the x's the improved field strength components  $F_{\mu_1...\mu_{p+1}}$  and their derivatives. The z-variables are taken as before, i.e., are the derivatives of  $A_{\mu_1...\mu_l}^{(l,p-l)}$  (l > 0) of Young symmetry-type 1) and the yvariables are just defined to be their  $\gamma$ -variations. This is an invertible change of generators provided one includes as well the last, undifferentiated ghost of ghost  $A^{(0,p)}$ , as in the free case. But this variable counts now as a z since it is no longer closed. And it actually "kills"  $B^{(0,p+1)}$  since  $\gamma A^{(0,p)} + B^{(0,p+1)} = 0$ . Thus, the variables  $A^{(0,p)}$  and  $B^{(0,p+1)}$ , which were previously x-variables, form now a contractible pair and disappear as announced.

The Bianchi identities for the new field strengths read

$$dF = H, \ dH = 0.$$
 (3.11)

They can be used to express the *H*-components and their derivatives in terms of the components  $F_{\mu_1...\mu_{p+1}}$  and their derivatives, which thus completely generate the cohomology.

To summarize, the  $\gamma$  cohomohology is given by

**Theorem 3.2** For the Chapline-Manton model (2.21), (2.22), (2.23), the cohomology  $H(\gamma)$  is given by the polynomials in the improved field strength components  $F_{\mu_1...\mu_{p+1}}$  and their derivatives,

$$\gamma\omega = 0 \Leftrightarrow \omega = \frac{1}{q!} \omega_{\nu_1 \dots \nu_q} ([F_{\mu_1 \dots \mu_{p+1}}]) dx^{\nu_1} \dots dx^{\nu_q}.$$
(3.12)

In particular, there is no cohomology at non-vanishing ghost number.

The situation is very similar to the discussion of the gauged principal U(1) sigma model [41] (see also [42] in this context).

### 3.3 Chapline-Manton model 2

In this case, the  $\gamma$ -cohomology is unchanged compared with the free case (in terms of the improved, gauge-invariant curvatures). The last ghosts remain in cohomology because  $A^{(0,1)}$  is still  $\gamma$ -closed, so the mechanism of the previous subsection is not operative. We leave it to the reader to make the change of generators necessary to prove the following theorem.

**Theorem 3.3** The cohomology of  $\gamma$  for the Chapline-Manton model (2.28), (2.29), (2.30), is given by

$$H(\gamma) = \tilde{\mathcal{I}} \otimes \mathcal{C} \tag{3.13}$$

where C is the algebra generated by the last, undifferentiated ghosts  $A^{(0,1)}$ and  $B^{(0,2r)}$ , and where  $\tilde{\mathcal{I}}$  is the agebra generated by the gauge invariant field strength components  $F_{\mu\nu}$ ,  $H_{\mu_1...\mu_{2r+1}}$  and their derivatives.

Note the new form of the Bianchi identities on the curvatures

$$dF = 0, \ dH = F^{r+1}.$$
 (3.14)

### 3.4 Chapline-Manton model 3

The discussion of the third example proceeds to a large extent like that of the first one. The last ghosts of ghosts  $A^{(0,1)}$  and  $B^{(0,2)}$  form a contractible pair and disappear in cohomology; the improved last ghost of ghost

$$\tilde{C}^{(0,3)} = C^{(0,3)} - \frac{1}{2}A^{(0,1)}B^{(0,2)}$$
(3.15)

remains. Thus one has

**Theorem 3.4** The cohomology of  $\gamma$  for the Chapline-Manton model (2.36) is given by

$$H(\gamma) = \tilde{\mathcal{I}} \otimes \tilde{\mathcal{C}} \tag{3.16}$$

where  $\tilde{\tilde{\mathcal{I}}}$  is the algebra generated by the gauge invariant field strength components  $F_{\mu\nu}$ ,  $G_{\mu\nu\rho\sigma}$  and their derivatives, and where  $\tilde{\mathcal{C}}$  is the algebra generated by the last, improved ghost of ghost  $\tilde{C}^{(0,3)} \equiv C^{(0,3)} - \frac{1}{2}A^{(0,1)}B^{(0,2)}$ .

Again, note the new form of the Bianchi identities,

$$dF = -H, \ dH = 0, \ dG = -FH$$
 (3.17)

which enable one in particular to express H in terms of the derivatives of F.

### 3.5 Chapline-Manton model 4

The  $\gamma$ -cohomology for this model is explicitly given in [35]. It is given by the theorem:

**Theorem 3.5** The cohomology of  $\gamma$  for the Chapline-Manton model (2.46), (2.47) is given by

$$H(\gamma) = \mathcal{J} \otimes \mathcal{D} \tag{3.18}$$

where (i)  $\mathcal{J}$  is the algebra of the invariant polynomials in the Yang-Mills curvature components and their covariant derivatives, as well as in the components of the gauge invariant curvature H and their derivatives; and (ii)  $\mathcal{D}$ is the algebra generated by the "primitive forms"  $trC^5$ ,  $trC^7$ , ...,  $trC^{2N-1}$ .

We recall that the Lie algebra cohomology for SU(N) is generated by the primitive forms  $trC^3$ ,  $trC^5$ , ... up to  $trC^{2N-1}$  [43, 44]. The primitive form  $trC^3$  is removed from the cohomology of  $\gamma$  because it is exact when the coupling to the 2-form is introduced: the last ghost of ghost  $\rho$  kills  $trC^3$  in cohomology. The Bianchi identity reads

$$DF = 0, \ dH = trF^2.$$
 (3.19)

## 4 Descent equation and lifts of $\gamma$ -cocycles

### 4.1 The descent equation

Our method for solving the Wess-Zumino consistency condition is that developped in [12, 13, 15] for the Yang-Mills case. In that approach, one analyses the  $\gamma$ -cocycles that can appear as "bottoms" of the descent equations [45]. We shall first briefly summarize the main ideas. These depend only on the generic properties of  $\gamma$  and d and not on their specific forms. We shall then apply them to the models at hand.

To a given solution  $a^{(p,q)}$  of the Wess-Zumino consistency condition

$$\gamma a^{(p,q)} + da^{(p-1,q+1)} = 0 \tag{4.1}$$

(where p and q denote respectively the form degree and the ghost number), one can associate another solution of the Wess-Zumino consistency condition, namely,  $a^{(p-1,q+1)}$ . Indeed, the triviality of d ("algebraic Poincaré lemma" [46]) implies<sup>1</sup>

$$\gamma a^{(p-1,q+1)} + da^{(p-2,q+2)} = 0 \tag{4.2}$$

for some  $a^{(p-2,q+2)}$ . There are ambiguities in the choice of  $a^{(p-1,q+1)}$  given the class  $[a^{(p,q)}]$  of  $a^{(p,q)}$  in  $H^{(p,q)}(\gamma|d)$ , but it is easy to verify that the map  $\partial: H^{(p,q)}(\gamma|d) \to H^{(p-1,q+1)}(\gamma|d)$  is well defined.

The map  $\partial$  is in general not injective. There are non trivial classes of  $H^{*,*}(\gamma|d)$  that are mapped on zero through the descent. For instance, if one iterates  $\partial$ , one gets from  $a^{(p,q)}$  a chain of cocycles in  $H^{*,*}(\gamma|d)$ ,  $[a^{(p,q)}] \mapsto \partial[a^{(p,q)}] \in H^{(p-1,q+1)}(\gamma|d) \mapsto \partial^2[a^{(p,q)}] \in H^{(p-2,q+2)} \mapsto \cdots \mapsto \partial^k[a^{(p,q)}] \in H^{(p-k,q+k)} \mapsto 0$  which must eventually end on zero since there are no forms of negative form degree. The equations defining the successive images of  $[a^{(p,q)}]$  are

$$\gamma a^{(p,q)} + da^{(p-1,q+1)} = 0 \tag{4.3}$$

$$\gamma a^{(p-1,q+1)} + da^{(p-2,q+2)} = 0 \tag{4.4}$$

<sup>&</sup>lt;sup>1</sup>In the space of x-independent forms, the cohomology of d is actually not trivial, even in form degree < n (the case relevant to the descent). Indeed, the constant forms are in the cohomology. But these never get in the way because  $\gamma a^{(p-1,q+1)}$  and the successive terms in the descent have strictly positive ghost number.

$$\gamma a^{(p-k,q+k)} + da^{(p-k-1,q+k+1)} = 0 \tag{4.5}$$

$$[a^{(p-k-1,q+k+1)}] = 0 (4.6)$$

and are known as the "descent equations" [45]. Since  $a^{(p-k-1,q+k+1)}$  is trivial in  $H^{(p-k-1,q+k+1)}(\gamma|d)$ , i.e.,  $a^{(p-k-1,q+k+1)} = \gamma b^{(p-k-1,q+k)} + db^{p-k,q+k+1)}$ , one may redefine  $a^{(p-k,q+k)} \rightarrow a^{(p-k,q+k)} - db^{(p-k-1,q+k)} = a'^{(p-k,q+k)}$  in such a way that we have  $\gamma a'^{(p-k,q+k)} = 0$ , i.e.,  $a'^{(p-k-1,q+k+1)} = 0$ . Conversely, if  $a^{(p-k,q+k)}$ is annihilated by  $\gamma$ , then  $\partial [a^{(p-k,q+k)}] = 0$ . Thus, the last non-trivial element  $a^{(p-k,q+k)}$ , or "bottom", of the descent is a  $\gamma$ -cocycle that is not exact in  $H^{*,*}(\gamma|d)$ . The non-injectivity of  $\partial$  follows precisely from the existence of such cocycles.

The length of the descent associated with  $[a^{(p,q)}]$  is the integer k for which  $\partial^k[a^{(p,q)}]$  is the last non-trivial cocycle occuring in the chain. One says that a descent is non trivial if it has length  $\geq 1$ . The idea of [12, 13] is to classify the elements of  $H^{*,*}(\gamma|d)$  according to the length of the associated descent.

In order to achieve this, one must determine the possible bottoms, i.e., the elements of  $H(\gamma)$  which are not trivial in  $H(\gamma|d)$  and which can be lifted k times.

## **4.2** Lifts of elements of $H(\gamma)$ - An example

The difficulty in the analysis of the lift is that contrary to the descent which carries no ambiguity in cohomology, the lift is ambiguous because  $H(\gamma)$  is not trivial. Furthermore, for the same reason, the lift can be obstructed, i.e., given  $a \in H(\gamma)$ , there may be no descent (i) which has a as bottom; and (ii) which starts with a solution b of the Wess-Zumino consistency condition such that db = 0 (while any descent ends always with an a such that  $\gamma a = 0$ ). The "first" b may be such that  $db \neq 0$  or even  $db \neq \gamma$ (something). In that case, there is no element c above b such that  $\gamma c + db = 0$  (while there is always an element e below a such that  $\gamma a + de = 0$ , namely e = 0: the descent effectively stops at a but is not obstructed at a).

In this subsection, we shall illustrate these features on a specific example: that of a free 1-form A and a free 2-form B, with BRST algebra

$$\gamma A^{(1,0)} + dA^{(0,1)} = 0, \ \gamma A^{(0,1)} = 0$$
(4.7)

$$\gamma B^{(2,0)} + dB^{(1,1)} = 0, \ \gamma B^{(1,1)} + dB^{(0,2)} = 0, \ \gamma B^{(0,2)} = 0.$$
 (4.8)

The curvatures are F = dA and H = dB, with  $\gamma F = \gamma H = 0$ .

Consider the  $\gamma$ -cocycle  $A^{(0,1)}B^{(0,2)}$ . It has form-degree zero and ghost number three. The descent that ends on this bottom has length one, and not the maximum length three. Indeed, the  $\gamma$ -cocycle  $A^{(0,1)}B^{(0,2)}$  can be lifted once, since there exists  $a \in \mathcal{P}$  such that  $\gamma a + d(A^{(0,1)}B^{(0,2)}) = 0$ . One may take  $a = A^{(1,0)}B^{(0,2)} + A^{(0,1)}B^{(1,1)}$ . Of course, a has form-degree one and ghost number two. If one tries to lift the given  $\gamma$ -cocycle  $A^{(0,1)}B^{(0,2)}$  once more, one meets an obstruction. Namely, there is no b such that  $\gamma b + da = 0$ . This is because da is in the same  $\gamma$ -class as  $FB^{(0,2)}$ , which is non-trivial, i.e., which cannot be written as a  $\gamma$ -variation. It is easy to verify that one cannot remove the obstruction by adding to a a  $\gamma$ -cocycle (which would not change  $\gamma a$ ). This provides an example of a  $\gamma$ -cocycle for which the lift is obstructed after one step.

Consider now the  $\gamma$ -cocycle  $\frac{1}{2}F(B^{(0,2)})^2$  with ghost number four and formdegree two. This cocycle can be lifted a first time, for instance  $FB^{(1,1)}B^{(0,2)}$  is above it,

$$\gamma[FB^{(1,1)}B^{(0,2)}] + d[\frac{1}{2}F(B^{(0,2)})^2] = 0.$$
(4.9)

It can be lifted a second time to  $\frac{1}{2}(B^{(1,1)})^2 + FB^{(2,0)}B^{(0,2)}$ . However, if one tries to lift it once more, one meets apparently the obstruction  $FHB^{(0,2)}$ , since the exterior derivative of  $\frac{1}{2}(B^{(1,1)})^2 + FB^{(2,0)}B^{(0,2)}$  differs from the  $\gamma$ -cocycle  $FHB^{(0,2)}$  by a  $\gamma$ -exact term. It is true that  $FHB^{(0,2)}$  is a non-trivial  $\gamma$ -cocycle. However, the obstruction to lifting three times  $\frac{1}{2}F(B^{(0,2)})^2$  is really absent. What happens is that we made a "wrong" choice for the term above  $\gamma$ -cocycle  $\frac{1}{2}F(B^{(0,2)})^2$  and should have taken not  $FB^{(1,1)}B^{(0,2)}$ , but rather, a term that differs from it by an appropriate  $\gamma$ -cocycle. This is because  $FHB^{(0,2)}$  is in fact the true obstruction to lifting twice the  $\gamma$ -cocycle  $A^{(0,1)}HB^{(0,2)}$ . Thus if one replaces (4.9) by

$$\gamma [FB^{(1,1)}B^{(0,2)} - A^{(0,1)}HB^{(0,2)}] + d[\frac{1}{2}F(B^{(0,2)})^2] = 0, \qquad (4.10)$$

which is permissible since  $\gamma(A^{(0,1)}HB^{(0,2)}) = 0$ , one removes the obstruction to lifting further  $\frac{1}{2}F(B^{(0,2)})^2$ . This shows that the obstructions to lifting k times a  $\gamma$ -cocycle are not given by elements of  $H(\gamma)$ , but rather, by elements of  $H(\gamma)$  that are not themselves obstructions of shorther lifts. The ambiguity in the choice of the lifts plays accordingly a crucial rôle in the analysis of the obstructions. In fact, the given  $\gamma$ -cocycle  $\frac{1}{2}F(B^{(0,2)})^2$  is actually trivial in  $H(\gamma|d)$ 

$$\frac{1}{2}F(B^{(0,2)})^{2} = d\left[\frac{1}{2}A^{(1,0)}(B^{(0,2)})^{2} + A^{(0,1)}B^{(1,1)}B^{(0,2)}\right] 
+ \gamma\left[A^{(1,0)}B^{(1,1)}B^{(0,2)} + \frac{1}{2}A^{(0,1)}(B^{(1,1)})^{2} + A^{(0,1)}B^{(2,0)}B^{(0,2)}\right]$$
(4.11)

and therefore, its lift can certainly never be obstructed.

## **4.3** Lifts of elements of $H(\gamma)$ - The first two steps

To control the features that we have just illustrated, it is necessary to introduce new differential algebras [12, 13]. Let  $E_0 \equiv H(\gamma)$ . We define a map  $d_0: E_0 \to E_0$  as follows:

$$d_0[a] = [da] \tag{4.12}$$

where [] is here the class in  $H(\gamma)$ . This map is well defined because  $\gamma da = -d\gamma a = 0$  (so da is a  $\gamma$ -cocycle) and  $d(\gamma m) = -\gamma(dm)$  (so d maps a  $\gamma$ -coboundary on a  $\gamma$ -coboundary). Now,  $d_0$  is a derivation and  $d_0^2 = 0$ , so it is a differential. Cocycles of  $d_0$  are elements of  $H(\gamma)$  that can be lifted at least once since  $d_0[a] = 0 \Leftrightarrow da + \gamma b = 0$  for some b, so b descends on a. By contrast, if  $d_0[a] \neq 0$ , then a cannot be lifted and, in particular, a is not exact in  $H(\gamma|d)$  (if it were,  $a = \gamma m + dn$ , one would have  $da = -\gamma dm$ , i.e., da = 0 in  $H(\gamma)$ ). Let  $F_0$  be a subspace of  $E_0$  supplementary to  $Ker d_0$ . One has the isomorphism (as vector spaces)

$$E_0 \simeq Ker \, d_0 \oplus F_0. \tag{4.13}$$

The next step is to investigate cocycles that can be lifted at least twice. In order to be liftable at least once, these must be in  $\operatorname{Ker} d_0$ . Among the elements of  $\operatorname{Ker} d_0$ , those that are in  $\operatorname{Im} d_0$  are not interesting, because they are elements of  $H(\gamma)$  that are trivial in  $H(\gamma|d)$  ( $[a] = d_0[b] \Leftrightarrow a = db + \gamma m$ ). Thus the relevant space is  $E_1 \equiv H(d_0, E_0)$ . One has

$$Ker \, d_0 \simeq Im \, d_0 \oplus E_1 \tag{4.14}$$

One then defines the differential  $d_1: E_1 \to E_1$ 

$$d_1[[a]] = [[db]] \tag{4.15}$$

where b is defined through  $da + \gamma b = 0$  – recall that  $d_0[a] = 0$  – and where [[a]] is the class of [a] in  $E_1$ . It is easy to see that (4.15) provides a well-defined differential in  $E_1^2$ .

If  $[[a]] \in E_1$  is a  $d_1$ -cocycle, then it can be lifted at least twice since [[db]] = 0 in  $E_1$  means  $db = du + \gamma$  (something) with  $\gamma u = 0$ . Thus one has  $da + \gamma b' = 0$  with b' = b - u and  $db' = \gamma$  (something). If on the contrary,  $d_1[[a]] \neq 0$ , then the corresponding elements in  $H(\gamma)$  cannot be lifted twice,  $d_1[[a]]$  being the obstruction to the lift. More precisely, the inequality  $d_1[[a]] \neq 0$  in  $E_1$  means  $[db] \neq d_0[c]$  in  $E_0$ . Thus, db cannot be written as a  $\gamma$ -variation, even up to the exterior derivative of a  $\gamma$ - closed term (ambiguity in the definition of b).

Analogous to the decomposition (4.13), one has

$$E_1 \simeq Ker \, d_1 \oplus F_1 \tag{4.16}$$

where  $F_1$  is a subspace of  $E_1$  supplementary to  $Ker d_1$ . The elements in  $Im d_1$  are trivial in  $H(\gamma|d)$  and thus of no interest from the point of view of the Wess-Zumino consistency condition.

To investigate the (non-trivial)  $\gamma$ -cocycle that can be lifted at least three times, one defines

$$E_2 = H(d_1, E_1) \tag{4.17}$$

and the differential  $d_2$  through

$$d_2: E_2 \to E_2, \ d_2[[[a]]] = [[[dc]]]$$

$$(4.18)$$

where the triple brackets denote the classes in  $E_2$  and where c is defined through the successive lifts  $da + \gamma b = 0$ ,  $db + \gamma c = 0$  (which exist since  $d_1[[a]] = 0$ ). It is a straightforward exercise to verify that  $d_2$  is well-defined in  $E_2$ , i.e., that the ambiguities in b and c play no rôle in  $E_2$ . Furthermore, a  $\gamma$ -cocycle a such that  $d_0[a] = 0$  (so that  $[[a]] \in E_1$  is well-defined) and  $d_1[[a]] = 0$  (so that  $[[[a]]] \in E_2$  is well-defined) can be lifted a third time if and only if  $d_2[[[a]]] = 0$ . Indeed, the relation  $d_2[[[a]]] = 0$  is equivalent to [[[dc]]] = 0, i.e.  $dc = \gamma u + dv + dw$ , with  $\gamma v = 0$  (this is the  $d_0$ -term) and

<sup>&</sup>lt;sup>2</sup>Proof:  $d_0[a] = 0 \Rightarrow da + \gamma b = 0 \Rightarrow \gamma db = 0$ . Hence, db is a  $\gamma$ -cocycle, which is clearly annihilated by  $d_0, d_0[db] = [d^2b] = 0$ . Furthermore, the class of db in  $E_1$  does not depend on the ambiguity in the definition of b, since if b is replaced by b + dm + u with  $\gamma u = 0$ , then db is replaced by db + du which is equivalent to db in  $E_1$  (the class of du in  $E_0$  is equal to  $d_0[u]$  since  $\gamma u = 0$ , and this is zero in  $E_1$ ). The derivation property is also easily verified,  $d_1(ab) = (d_1a)b + (-1)^{\epsilon_a}ad_1b$ .

 $\gamma w + dt = 0, \ \gamma t = 0$  (this is the  $d_1$ -term). Thus, by redefining b as b - t and c as c - v - w, one gets,  $dc_{Redefined} = \gamma u$ .

## 4.4 Lifts of elements of $H(\gamma)$ - General theory

One can proceed in the same way for the next lifts. One finds in that manner a sequence of spaces  $E_r$  and differentials  $d_r$  with the properties

- 1.  $E_r = H(E_{r-1}, d_{r-1}).$
- 2. There exists an antiderivation  $d_r: E_r \to E_r$  defined by  $d_r[[\ldots [X] \ldots ]] = [[\ldots [db] \ldots ]]$  for  $[[\ldots [X] \ldots ]] \in E_r$  where  $[[\ldots [db] \ldots ]]$  is the class of the  $\gamma$ -cocycle db in  $E_r$  and where b is defined through  $dX + \gamma c_1 = 0$ ,  $dc_1 + \gamma c_2 = 0$ , ...,  $dc_{r-1} + \gamma b = 0$ . Similarly,  $[[\ldots [X] \ldots ]]$  denotes the class of the  $\gamma$ -cocycle X in  $E_r$  (assumed to fulfill the successive conditions  $d_0[X] = 0$ ,  $d_1[[X]] = 0$  etc ... so as to define an element of  $E_r$ ).
- 3.  $d_r^2 = 0$ .
- 4. A  $\gamma$ -cocycle X can be lifted r times if and only if  $d_0[X] = 0$ ,  $d_1[[X]] = 0$ ,  $d_2[[[X]]] = 0$ , ...,  $d_{r-1}[[\ldots [X] \ldots ]] = 0$ . If  $d_r[[\ldots [X] \ldots ]] \neq 0$ , the  $\gamma$ -cocycle X cannot be lifted (r+1) times and is not exact in  $H(\gamma|d)$ .
- 5. A necessary and sufficient condition for an element Y in  $H(\gamma)$  to be exact in  $H(\gamma|d)$  is that there exists a k such that  $d_i[\ldots[Y]\ldots] = 0$ ,  $(i = 1, 2, \ldots, k - 1)$  and  $[\ldots[Y]\ldots] = d_k[\ldots[Z]\ldots]$ . This implies in particular  $d_j[\ldots[Y]\ldots] = 0$  for all j's.
- 6. Conversely, if a  $\gamma$ -cocycle Y fulfills  $d_i[\ldots[Y]\ldots] = 0$  for i = 0, 1, ..., k-1and  $d_k[\ldots[Y]\ldots] \neq 0$ , then, it is not exact in  $H(\gamma|d)$ . The condition is not necessary, however, because there are elements of  $H(\gamma)$  that are non trivial in  $H(\gamma|d)$  but which are annihilated by all  $d_i$ 's. This is due to the fact that there are no exterior form of degree higher than the spacetime dimension. We shall come back to this point below.

The meaning of the integer k for which  $Y = d_k Z$  in item 5 (with  $Y \neq d_i$ (something) for i < k) is as follows (we shall drop the multiple brackets when no confusion can arise). If the  $\gamma$ -cocycle a is in  $Im d_0$ , then  $a = db + \gamma c$ ,

where b is also a  $\gamma$ -cocycle. If a is a non-zero element of  $E_1$  in the image of  $d_1$ , then again  $a = db + \gamma c$ , but b is now not a cocycle of  $\gamma$  since a would then be in  $Im d_0$  and thus zero in  $E_1$ . Instead, one has  $\gamma b + d\beta = 0$  where  $\beta$  is a cocycle of  $\gamma$  ( $\gamma\beta = 0$ ) which is not trivial in  $H(\gamma|d)$ . More generally, k characterizes the length of the descent below b in  $a = db + \gamma c$ ,  $\gamma b + d\beta = 0$  etc.

The proof of items 1 to 4 proceeds recursively. Assume that the differential algebras  $(E_i, d_i)$  have been constructed up to order r - 1, with the properties 2 through 4. Then, one defines the next space  $E_r$  as in 1. Let x be an element of  $E_r$ , and let X be one of the  $\gamma$ -cocycles such that the class  $[[\dots [X] \dots ]]$  in  $E_r$  is precisely x. Since X can be lifted r times, one has a sequence  $dX + \gamma c_1 = 0$ , ...,  $dc_{r-1} + \gamma b = 0$ . The ambiguity in X is  $X \to X + \gamma a + du_0 + du_1 + \dots + du_{r-1}$ , where  $u_0$  is a  $\gamma$ -cocycle (this is the  $d_0$ -exact term),  $u_1$  is the first lift of a  $\gamma$ -cocycle (this is the  $d_1$ -exact term) etc. Setting  $u = u_0 + u_1 + \dots + u_{r-1}$ , one sees that the ambiguity in X is of the form  $X \to X + \gamma a + du$ . On the other hand, the ambiguity in the successive lifts takes the form  $c_1 \to c_1 + m_1$ , where  $m_1$  is a  $\gamma$ -cocycle that can be lifted r-1 times,  $c_2 \to c_2 + n_1 + m_2$ , where  $n_1$  descends on  $m_1$  and  $m_2$  is a  $\gamma$ -cocycle that can be lifted r-2 times, ..., and finally  $b \to b + a_1 + a_2 + \dots + a_{r-1} + a_r$ , where  $a_1$  descends (r-1) times, on  $m_1$ ,  $a_2$  descends (r-2) times, on  $m_2$ , etc, and  $a_r$  is a  $\gamma$ -cocycle.

The element  $X_r \equiv db$  is clearly a cocycle of  $\gamma$ , which is annihilated by  $d_0$  and the successive derivations  $d_k$  because  $dX_r = 0$  exactly and not just up to  $\gamma$ -exact terms. The ambiguity in the successive lifts of X plays no rôle in the class of  $X_r$  in  $E_r$ , since it can (suggestively) be written  $db \rightarrow db + d_{r-1}m_1 + d_{r-2}m_2 + \cdots + d_1m_{r-1} + d_0a_r$ . Thus, the map  $d_r$  is well-defined as a map from  $E_r$  to  $E_r$ . It is clearly nilpotent since  $dX_r = 0$ . It is also a derivation, because one may rewrite the lift equations for X as  $\tilde{\gamma}(X + c_1 + c_2 + \ldots + b) = d_r X$  where

$$\tilde{\gamma} = \gamma + d. \tag{4.19}$$

Let Y be another element of  $E_r$  and  $e_1$ ,  $e_2$ , ... $\beta$  its successive lifts. Then,  $\tilde{\gamma}(Y + e_1 + e_2 + \ldots + \beta) = d_r Y$ . Because  $\tilde{\gamma}$  is a derivation, one has  $\tilde{\gamma}[(X + c_1 + \ldots + \beta)(Y + e_1 + \ldots + \beta)] = (d_r X)Y + (-1)^{\epsilon_X} X d_r Y +$  forms of higher form-degree, which implies  $d_r(XY) = (d_r X)Y + (-1)^{\epsilon_X} X d_r Y$ :  $d_r$  is also an odd derivation and thus a differential. This establishes properties 2 and 3. To prove property 4, one observes that X can be lifted once more if and only if one may choose its successive lifts so that db is  $\gamma$ -exact. This is equivalent to stating that  $d_r X$  is zero in  $E_r$ . Properties 5 and 6 are rather obvious: if a is a  $\gamma$ -cocycle which is exact in  $H(\gamma|d)$ ,  $a = db + \gamma c$ , then  $a = d_k m$  where k is the length of the descent associated with  $\gamma b + dn = 0$ , which has bottom m.

As shown in [12, 13], the above construction may be elegantly captured in an exact couple [47]. The detailed analysis of this exact couple and the proof of the above results using directly the powerful tools offered by this couple may be found in [13, 14, 15].

One has, for each r, the vector space isomorphisms

$$E_r \simeq Ker \, d_r \oplus F_r \simeq Im \, d_r \oplus E_{r+1} \oplus F_r \tag{4.20}$$

where  $F_r$  is a subspace supplementary to  $Ker d_r$  in  $E_r$ . Thus

$$E_0 \simeq \bigoplus_{k=0}^{k=r-1} F_k \bigoplus_{k=0}^{k=r-1} Im \, d_k \oplus E_r \tag{4.21}$$

Because there is no form of degree higher than the spacetime dimension,  $d_n = 0$  ( $d_n a$  has form-degree equal to FormDeg(a) + n + 1). Therefore,  $E_n = E_{n+1} = E_{n+2} = \dots$  This implies

$$E_0 \simeq \bigoplus_{k=0}^{k=n-1} F_k \bigoplus_{k=0}^{k=n-1} Im \, d_k \oplus E_n.$$

$$(4.22)$$

The elements in any one of the  $F_k$ 's are non trivial bottoms of the descent which can be lifted exactly k times. All the elements above them in the descent are also non trivial solutions of the Wess-Zumino consistency condition. The elements in  $Im d_k$  are bottoms which are trivial in  $H(\gamma|d)$  and which define therefore trivial solutions of the Wess-Zumino consistency condition. Finally, the elements in  $E_n$  are bottoms that can be lifted all the way up to form degree n. These are non trivial in  $H(\gamma|d)$ , since they are not equal to  $d_im$  for some i and m. The difference between the elements in  $\oplus F_k$  and those in  $E_n$  is that the former ones cannot be lifted all the way up to form-degree n: one meets an obstruction before, which is  $d_k a$  (if  $a \in F_k$ ). By contrast, the elements in  $E_n$  can be lifted all the way up to form degree n. This somewhat unpleasant distinction between  $\gamma$ -cocycles that are non-trivial in  $H(\gamma|d)$  will be removed below, where we shall assign an obstruction to the elements of  $E_n$  in some appropriate higher dimensional space.

In order to solve the Wess-Zumino consistency condition, our task now is to determine explicitly the spaces  $E_r$  and  $F_r$ .

## 5 Covariant Poincaré lemma – Small algebra

To that end, we first work out the cohomology of  $d_0$  in  $E_0 \equiv H(\gamma)$ . Let u be a  $\gamma$ -cocycle. Without loss of generality, we may assume that u takes the form

$$u = \sum P_I \omega^I \tag{5.1}$$

where the  $\omega^{I}$  form a basis of the algebra generated by the last, non trivial (if necessary, improved) ghosts of ghosts in the cohomology (as well as  $trC^{5}$ ,  $trC^{7}$  etc for the fourth Chapline-Manton model), and where the  $P_{I}$ 's are polynomials in the (improved) field strength components and their (covariant) derivatives, with coefficients that involve  $dx^{\mu}$ . The  $P_{I}$ 's are called "gauge-invariant polynomials". A direct calculation using the fact that the *d*-variation of the last ghosts (and  $trC^{5}$ ,  $trC^{7}$  etc) is  $\gamma$ -exact yields  $du = \sum (dP_{I})\omega^{I} + \gamma v'$ . This is  $\gamma$ -exact if and only if  $dP_{I} = 0$ .

Now, if  $P_I = dR_I$  where  $R_I$  is also a gauge invariant polynomial, then u is d-exact modulo  $\gamma$ ,  $u = da + \gamma b$ , with  $\gamma a = 0$ . Conversely, if  $u = da + \gamma b$  with  $\gamma a = 0$ , then  $P_I$  is d-exact in the space of invariant polynomials. Thus, the class of u (in  $E_0$ ) is a non trivial cocycle of  $d_0$  if and only if  $P_I$  is a non trivial cocycle of the *invariant* cohomology of d. We give below the relevant "covariant Poincaré lemma" for each of the models of this paper.

Since we are interested in lifts of  $\gamma$ -cocycles from form-degree k to formdegree k + 1, we shall investigate the *d*-invariant cohomology only in formdegree strictly smaller than the spacetime dimension n. This will be assumed throughout the remainder of this section. [In form-degree n, there is clearly further invariant cohomology since any invariant *n*-form is annihilated by d, even when it cannot be written as the d of an invariant form].

#### 5.1 Free case

The calculation of the invariant cohomology of d is given in the appendix A of [36], so we just recall the result.

**Theorem 5.1** The invariant cohomology of d for the free theory is given by the polynomials in the exterior forms  $H^a$ .

This is a direct generalization of the result established for 1-forms in [16, 17]. Note that the cohomology contains in particular the constants and

the constant forms. These latter can be eliminated by imposing Lorentz invariance.

The theorem implies, according to the general analysis of the descent equation given above, that the only bottoms u ( $\gamma u = 0$ ) that can be lifted at least once can be expressed in terms of exterior products of the curvature forms  $H^a$  and the last ghosts of ghosts (up to trivial redefinitions). Out of the infinitely many generators of  $H(\gamma)$ , only  $H^a$  and  $B^{a(0,p_a)}$  survive in  $E_1$ .

Because the objects that survive the first step in the lift can be expressed in terms of forms, it is convenient to introduce the so-called "small algebra"  $\mathcal{A}$  generated in the exterior product by the exterior forms  $B^{a(k,p_a-k)}$  and  $dB^{a(k,p_a-k)}$   $(k = 0, ..., p_a)$ . This algebra is stable under  $\gamma$  and d. If one denotes by  $E_0^{small}$  the cohomology of  $\gamma$  in the small algebra, one finds

$$E_0^{small} \equiv H(\gamma, \mathcal{A}) \simeq \mathcal{B} \tag{5.2}$$

where  $\mathcal{B}$  is the subalgebra of  $\mathcal{A}$  generated by the curvatures  $H^a$  and the last ghosts of ghosts  $B^{a(0,p_a)}$ .

One defines  $E_1^{small}$  as  $H(d_0^{small}, E_0^{small})$ , where  $d_0^{small}$  is the restriction of  $d_0$  to  $E_0^{small}$ . Because  $dH^a = 0$  and  $dB^{a(0,p_a)} = \gamma$ (something), the restriction  $d_0^{small}$  identically vanishes. Thus

$$E_1^{small} \simeq E_0^{small} \simeq \mathcal{B}. \tag{5.3}$$

What is the relationship between  $E_1^{small}$  and  $E_1$ ? These two spaces are in fact isomorphic,

$$E_1 \simeq E_1^{small}.\tag{5.4}$$

Indeed, let q be the map from  $E_1^{small}$  to  $E_1$  that assigns to a cohomological class in  $E_1^{small}$  its cohomological class in  $E_1$  ( $a \in E_1^{small} \simeq \mathcal{B}$  fulfills  $\gamma a = 0$ and  $d_0 a = 0$  and thus defines of course an element of  $E_1$ ). It follows from the above theorem that the map q is surjective since any class in  $E_1$  possesses a representative in the small algebra. The map q is also injective because there is no non trivial class in  $E_1^{small}$  that becomes trivial in  $E_1$ . If the smallalgebra  $\gamma$ -cocycle  $r = \sum P_I \omega^I$  with  $P_I, \omega^I \in \mathcal{B}$  can be written as  $r = du + \gamma t$ where u and v are in the big algebra and u is a  $\gamma$ -cocycle, then r is actually zero. This can be easily seen by setting the derivatives of the ghosts and of the field strength components equal to zero.

The differentials  $d_1^{small}$  and  $d_1$  are mapped on each other in this isomorphism,  $d_1q = qd_1^{small}$ . It then follows that the next cohomological spaces

 $E_k$  and  $E_k^{small}$  are also equal,  $E_2 \equiv H(d_1, E_1) \simeq E_2^{small} \equiv H(d_1^{small}, E_1^{small})$ ,  $E_3 \simeq E_3^{small}$  etc (but of course,  $E_0 \neq E_0^{small}$ ). There is thus no loss of generality in investigating in the small algebra the solutions of the Wess-Zumino consistency condition that descend non trivially.

## 5.2 Chapline-Manton models

The small algebra  $\mathcal{A}$  is also relevant to the Chapline-Manton models because the invariant cohomology can be computed in it without loosing any cohomological class. Indeed, one has

**Theorem 5.2** Let P be a gauge invariant polynomial. If P is closed, then P is the sum of a closed, gauge invariant polynomial belonging to the small algebra and of the exterior derivative of an invariant polynomial,

$$dP = 0 \Leftrightarrow P = Q + dR, \ Q \in \mathcal{A}, \ dQ = 0 \tag{5.5}$$

(with P, Q and R all gauge-invariant). Furthermore, if Q is d-exact in the algebra of gauge-invariant polynomials, Q = dS with S gauge-invariant, one may assume that S is in the small algebra (and gauge-invariant). Therefore, the invariant cohomology of d is to be found in A.

Note that while the conditions  $Q \in \mathcal{A}$  and Q = dS (with S gauge-invariant) imply Q = 0 in the free case, this is no longer true here.

We shall prove the theorem for the specific case of the second model. The proof proceeds in the same way for the other models. Introduce a grading N that counts the number of derivatives of the B field. According to this grading P and d split as

$$P = P_k + P_{k-1} + \dots + P_1 + P_0, \quad d = D_1 + D_0, \tag{5.6}$$

with

$$N(P_i) = i, \quad N(D_i) = i.$$
 (5.7)

The differential  $D_1$  takes derivatives only of the *B*-field, the differential  $D_0$  takes derivatives only of the *A*-field. Because *P* is gauge-invariant, the *B*-field enters *P* only through the components of *dB* and their derivatives. Furthermore, even though the  $P_i$ 's with i < k may involve the components

 $A_{\mu}$ 's and their symmetrized derivatives,  $P_k$  depends on A only through the  $F_{\mu\nu}$  and their derivatives.

The equation dP = 0 yields  $D_1P_k = 0$  at the highest value of the N-degree. According to the results for the free case, this implies  $P_k =$  $D_1R_{k-1} + m_k$  where  $R_{k-1}$  is a polynomial in the components of dB and their derivatives, while  $m_k$  is a polynomial in the form dB, both with coefficients in the components of F and their derivatives (which fulfill  $D_1 F_{\mu\nu} = 0$ ). Covariantize  $R_{k-1}$  and  $m_k$  by completing dB into H. This only introduces terms of lower N-degree. We denote the covariant objects by r and m, respectively. One has  $P_k = (dr + m)_k$  and  $P = dR_{k-1} + m_k + more$ , where "more" is an invariant polynomial of maximum N-degree strictly smaller than k. The invariant polynomial m - which exists only if k = 1 or 0 since  $H^2 = 0$  - is of order k in the exterior form H. It must be closed by itself since there can be no compensation between  $D_0m$  and  $D_1(more)$ , which is necessarily of lower degree in the components of H and their derivatives. It follows from  $D_0m = 0$  that  $m = \mu(F, H) + ds$ , where  $\mu$  is a polynomial in the forms F and H and where s is an invariant polynomial (use again the results for the free case and Hdp = -d(Hp) + more). Thus one can get rid of  $P_k$  by adding to P terms of the form (5.5) of the theorem. By repeating the argument at the successive lower degrees, one reaches the desired conclusion.

To prove the second part of the theorem, one first observes that if dQ = 0, then Q(F, H) does not involve in fact H, Q = Q(F) (see subsection 5.2.2 below). Assume then that Q = dU, where U is a gauge-invariant polynomial, U = U([H], [F]). By expanding U according to the N-degree,  $U = U_0 + U_1 + \ldots + U_l$ , one finds at higher order  $D_1U_l = 0$ , which implies as above  $U_l = D_1R_{l-1} + m_l$  where  $m_l$  is a polynomial in the form dB. One can remove  $D_1R_{l-1}$  from  $U_l$  by substracting  $dR_{l-1}$  from U, which does not modify Q. Thus, only  $m_l$ , which is present for l = 1 or l = 0, is relevant. By repeating the argument, one finally arrives at

$$U = Ha([F]) + b([F])$$
(5.8)

The condition Q = dU implies da = 0 and thus  $a = d\nu([F]) + \rho(F)$  where  $\rho(F)$  is a polynomial in the form F. The term  $Hd\nu([F])$  is irrelevant since it can be absorbed into b([F]) up to a d-exact term. Thus,  $U = H\rho(F) + b'([F])$ . The condition Q(F) = dU reads now Q(F) = k(F) + db'([F]) where k(F) is a polynomial in F and implies db' = 0 because db'([F]) necessarily involves

one derivative of  $F_{\lambda\mu}$  if it is not zero. But then, again, one can drop b' from U, which proves the second assertion.

It follows from this theorem that there is no restriction in investigating the invariant *d*-cohomology in the small algebra. Elements of  $H(\gamma)$  that can be lifted at least once necessarily belong to  $\mathcal{A}$  up to trivial terms. There is no restriction in the investigation of the next lifts either because again  $E_1^{small} = E_1$ . If a  $\gamma$ -cocycle  $a \in \mathcal{A}$  can be written as  $a = du + \gamma v$  where uand v are in the big algebra and  $\gamma u = 0$ , then one may find u' and v' in  $\mathcal{A}$ such that  $a = du' + \gamma v'$  (with  $\gamma u' = 0$ ). This follows from the second part of the theorem. Obstructions to lifts within  $\mathcal{A}$  are not removed by going to the big algebra.

#### 5.2.1 Model 1

For the first Chapline-Manton model discussed above, the invariant cohomology of d is trivial. Indeed, in the algebra generated by F and H, the differential d takes the contractible form dF = H, dH = 0. Thus

$$E_1 \equiv H(d_0, E_0^{small}) = 0 \tag{5.9}$$

where  $E_0^{small}$  is the algebra generated by F and H.

#### 5.2.2 Model 2

In the algebra generated by the gauge-invariant curvatures, d takes the form

$$dF = 0, \ dH = F^{r+1}.$$
 (5.10)

Since  $H^2 = 0$ , any element in this algebra is of the form

$$a = \alpha(F) + \beta(F)H \tag{5.11}$$

where  $\alpha(F)$  and  $\beta(F)$  are polynomials in F. The condition that *a* is closed implies  $\beta(F)F^{k+1} = 0$ , which forces  $\beta(F)$  to vanish. Furthermore  $a \equiv \alpha(F)$ is exact if it is in the ideal generated by  $F^{r+1}$ . Thus, we have the theorem:

**Theorem 5.3** The invariant cohomology of d for the Chapline-Manton model 2 is the quotient of the algebra generated by the F's by the ideal generated by  $F^{r+1}$ .

#### 5.2.3 Model 3

For the third model, d is given by (3.17). By redefining the curvature G as

$$G_M = G - \frac{F^2}{2}$$
(5.12)

this can be brought to the form

$$dF = -H, \ dH = 0, \ dG_M = 0 \tag{5.13}$$

from which it follows that:

**Theorem 5.4** For the third model, the invariant cohomology of d is given by the polynomials in the variable  $G_M = G - F^2/2$ .

#### 5.2.4 Model 4

The invariant polynomials in the small algebra are the polynomials in the gauge-invariant curvature H of the 2-form and in the "fundamental" invariants  $trF^2$ ,  $trF^3$ , ...  $trF^N$  for SU(N) (this is a basis for the SU(N) symmetric polynomials). These polynomials are closed, except H, which fulfills  $dH = trF^2$ . Hence, H and  $trF^2$  do not appear in the cohomology.

**Theorem 5.5** For the fourth model, the invariant cohomology of d is given by the polynomials in  $trF^3$ ,  $trF^4$ , ...  $trF^N$ .

### 5.3 Universal algebra $\mathcal{U}$

The small algebra  $\mathcal{A}$  involves only exterior forms, exterior products and exterior derivatives. It does "remember" the spacetime dimension since its generators are not free: any product of generators which form-degree exceeding the spacetime dimension vanishes.

It is useful to drop this relation and to work in the algebra freely generated by the potentials, the last ghosts of ghosts and their exterior derivatives with the sole condition that these commute or anti-commute (graded commutative algebra) but without imposing any restriction on the maximally allowed form degree [13, 48]. This algebra is called the universal algebra and denoted by  $\mathcal{U}$ . In this algebra, the cohomology of d is trivial in all form-degrees and the previous theorems on the invariant cohomology of d are also valid in formdegree  $\geq n$ . Furthermore, one can sharpen the condition for a cocycle in  $H(\gamma)$  to be non trivial in  $H(\gamma|d)$ .

**Theorem 5.6** A necessary and sufficient condition for  $X \in H(\gamma)$  to be nontrivial in  $H(\gamma|d)$  is that there exists r such that  $d_rX \neq 0$ . That is, the lift of X must be obstructed at some stage. (For the equation  $d_rX \neq 0$  to make sense,  $d_iX$  must vanish for i < r. Also, we denote again  $X \in E_0$  and its representative in  $E_r$  by the same letter).

Proof: The decomposition of  $E_n$  is now non-trivial since da does not necessarily vanish even when a is a n-form. Thus,  $d_n$  is not necessarily zero and the procedure of lifting can be pursued above form-degree n. Suppose that one does not encounter an obstruction in the lifting of X. That is, one can go all the way up to ghost number zero, the last two equations being  $dc_k + \gamma b = 0$  (with b of ghost number zero) and db = 0 (so b lifts to zero). Then, one can write b = dm since the cohomology of d is trivial in any form-degree in the universal algebra  $\mathcal{U}$  (except for the constants, which cannot arise here since b involves the fields). The triviality of the top-form bimplies the triviality in  $H(\gamma|d)$  of all the elements below it. Thus, a necessary condition for the bottom to be non trivial in  $H(\gamma|d)$  is that one meets an obstruction in the lift at some stage. The condition is also clearly sufficient.

One can summarize our results as follows

**Theorem 5.7** (Generalized "transgression" lemma) Let  $X \in E_0$  be a nontrivial element of  $H(\gamma|d)$ . Then there exists an integer r such that  $d_iX = 0$ , i < r and  $d_rX = Y \neq 0$ . The element Y is defined through the chain  $dX + \gamma c_1 = 0, ..., dc_{r-1} + \gamma c_r = 0, dc_r + \gamma c_{r+1} = Y$ , where the elements  $c_i \in \mathcal{U}$  (i = 1, r + 1) are chosen so as to go all the way up to  $c_{r+1}$ . One has  $\gamma Y = 0$  and Y should properly be viewed as an element of  $E_r$  (reflecting the ambiguities in the lift). One calls the obstruction Y to a further lift of Xthe (generalized) "transgression" of X. The element X and its transgression have opposite statistics.

This is the direct generalization of the analysis of [13] to the case of *p*-forms. "Primitive elements" of  $E_0$  are those that have form-degree zero and for which the transgression has ghost number zero, i.e., they are the elements that can be lifted all the way up to ghost number zero ("that can be transgressed"). We refer to [13, 43, 49] for more background information applicable to the Yang-Mills case.

Because the space  $E_n$  and the next ones can be further decomposed in the universal algebra,

$$E_n \simeq Imd_n \oplus E_{n+1} \oplus F_n, \ E_{n+1} \simeq etc$$
 (5.14)

where the decomposition for a given  $\gamma$ -cocycle ultimately ends at form-degree equal to the ghost number, one has

$$E_0 \simeq \bigoplus_{k=0}^{\infty} F_k \bigoplus_{k=0}^{\infty} Imd_k.$$
(5.15)

# 6 Results

We can now compute the spaces  $E_k$ .

## 6.1 Free case

Let  $0 < p_1 < p_2 < \ldots < p_M$  be the form degrees of the gauge potentials  $B^a$ . We denote by  $B_1^{a_1}$  the forms of degrees  $p_1$ ,  $B_2^{a_2}$  the forms of degree  $p_2$  etc.

The first non-vanishing differential (in  $E_0^{small}$ ) is  $d_{p_1}$  so that  $E_0^{small} = E_1 = E_2 = \dots = E_{p_1}$ . Any bottom in  $E_0^{small}$  can be lifted at least  $p_1$  times. In  $E_1$ , the differential  $d_{p_1}$  acts as follows

$$d_{p_1}B_1^{a_1(0,p_1)} = H_1^{a_1}, \ d_{p_1}H_1^{a_1} = 0$$
(6.1)

in the sector of the forms of degree  $p_1$  and

$$d_{p_1}B_k^{a_k(0,p_k)} = 0, \ d_{p_1}H_k^{a_k} = 0, \ k > 1$$
(6.2)

in the other sectors. The form of the differential  $d_{p_1}$  makes explicit the contractible part of  $(E_{p_1}, d_{p_1})$ . The variables  $B_1^{a_1(0,p_1)}$  and  $H_1^{a_1}$  are removed from the cohomology, so that  $E_{p_1+1}$  is isomorphic to the algebra generated by the curvatures  $H_k^{a_k}$  of form-degree  $> p_1 + 1$  and the last ghosts of ghosts of ghost number  $> p_1$ .

A subspace  $F_{p_1}$  complementary to  $\operatorname{Ker} d_{p_1}$  is easily constructed. In fact, a monomial in  $B_1^{a_1(0,p_1)}$  and  $H_1^{a_1}$  is defined by a tensor  $f_{a_1\dots a_k b_1\dots b_m}$  which is symmetric (respectively antisymmetric) in  $a_1, \ldots, a_k$  and antisymmetric (respectively symmetric) in  $b_1, \ldots, b_m$  if the last ghosts of ghosts are commuting (respectively anticommuting). Its irreducible components can be of two Young-symmetry types, one of which must be zero if the monomial is to be annihilated by  $d_{p_1}$ . The space  $F_{p_1}$  can be taken to be the space generated by the monomials of this symmetry type (not annihilated by  $d_{p_1}$ ), tensored by the algebra generated by the curvatures and last ghosts of ghosts of higher degree. Together with their successive lifts, the elements in  $F_{p_1}$  provide all the non-trivial solutions of the Wess-Zumino consistency condition which are involved in descents whose bottoms can be lifted exactly  $p_1$  times.

Similarly, one finds that the next non-vanishing differential is  $d_{p_2}$ . The generators  $B_2^{a_2(0,p_2)}$  and  $H_2^{a_2}$  drop from the cohomology of  $d_{p_2}$  while those of higher degree remain. A space  $F_{p_2}$  can be constructed along exactly the same lines as the space  $F_{p_1}$  above and characterizes the solutions of the Wess-Zumino consistency condition involved in descents whose bottoms can be lifted exactly  $p_2$  times.

More generally, the non-vanishing differentials are  $d_{p_k}$ . They are defined (in  $E_{p_k}$ , which is isomorphic to the algebra generated by the curvatures of form-degree  $> p_{k-1}+1$  and the last ghosts of ghosts of ghost number  $> p_{k-1}$ ) through

$$d_{p_k} B_k^{a_k(0,p_k)} = H_k^{a_k}, \ d_{p_k} H_k^{a_k} = 0$$
(6.3)

and

$$d_{p_k} B_j^{a_j(0,p_j)} = 0, \ d_{p_k} H_j^{a_j} = 0, \ j > k.$$
(6.4)

The generators  $B_k^{a_k(0,p_k)}$  and  $H_k^{a_k}$  disappear in cohomology. The subspace  $F_{p_k}$  is again easily constructed along the previous lines. Together with their successive lifts, the elements in  $F_{p_k}$  provide all the non-trivial solutions of the Wess-Zumino consistency condition which are involved in descents whose bottoms can be lifted exactly  $p_k$  times.

The discussion can be illustrated in the case of the simple model with one free 1-form  $A^{(1,0)}$  and one free 2-form  $B^{(2,0)}$  considered in subsection 4.2. The space  $E_0^{small}$  is isomorphic to the space of polynomials in the curvatureforms F, H and the last ghosts of ghosts  $A^{(0,1)}$ ,  $B^{(0,2)}$ . The differential  $d_0^{small}$ vanishes so  $E_1 = E_0^{small}$ . One finds next  $d_1 A^{(0,1)} = F$ ,  $d_1 F = 0$ ,  $d_1 B^{(0,2)} = 0$ and  $d_1 H = 0$ . The space  $E_2$  is isomorphic to the space of polynomials in  $B^{(0,2)}$  and H. One may take for  $F_1$  the space of polynomials linear in  $A^{(0,1)}$ . These can be lifted exactly once, their lifts being linear in  $A^{(1,0)}$  and  $A^{(0,1)}$ ,

$$a \in F_1 \Leftrightarrow a = A^{(0,1)} \sum (B^{(0,2)})^l F^k H^m \quad (m = 0 \text{ or } 1)$$
 (6.5)

Then, one gets

$$da + \gamma b = 0 \tag{6.6}$$

with

$$b = \sum \left( A^{(1,0)} (B^{(0,2)})^l F^k H^m + l A^{(0,1)} B^{(1,1)} (B^{(0,2)})^{l-1} F^k H^m \right).$$
(6.7)

They cannot be lifted a second time since the obstruction  $d_1 a = \sum (B^{(0,2)})^l F^{k+1} H^m$  does not vanish. The above *a*'s and *b*'s are the most general solutions of the Wess-Zumino consistency condition involved in descents of length 1.

The differential  $d_2$  in  $E_2$  is given by  $d_2 B^{(0,2)} = H$ ,  $d_2 H = 0$ . Because  $H^2 = 0$ , one may take for  $F_2$  the space of polynomials in  $B^{(0,2)}$  only. For those, the descent reads,

$$\begin{aligned} \alpha &= (B^{(0,2)})^l \quad , \quad \gamma \alpha = 0 \\ \beta &= l B^{(1,1)} (B^{(0,2)})^{l-1} \quad , \quad d\alpha + \gamma \beta = 0 \\ \lambda &= l B^{(2,0)} (B^{(0,2)})^{l-1} + \frac{l(l-1)}{2} (B^{(0,2)})^{l-2} (B^{(1,1)})^2 \quad , \quad d\beta + \gamma \lambda = 0 \ (6.8)^{l-1} \end{aligned}$$

The elements of the form  $\alpha$ ,  $\beta$  or  $\lambda$  are the most general solutions of the Wess-Zumino consistency condition involved in descents of length 2. With the solutions involved in descents of length 1 and those that do not descend (i.e., which are strictly annihilated by  $\gamma$ ), they exhaust all the (antifield-independent) solutions of the Wess-Zumino consistency condition.

A straightforward consequence of our discussion is the following theorem, which will prove useful in [11].

**Theorem 6.1** Let  $\omega$  be a  $\gamma$ -cocycle that takes the form

$$\omega = \alpha (H_s^{a_s} \ B_s^{a_s,(0,p_s)}) \beta (B_k^{a_k(0,p_k)}, H_k^{a_k}), \ k > s$$
(6.9)

where  $\alpha$  vanishes if  $H_s^{a_s}$  and  $B_s^{a_s(0,p_s)}$  are set equal to zero (no constant term) and fulfills

$$d_{p_s}\alpha = 0 \tag{6.10}$$

(the first potential obstruction in the lift of  $\omega$  is absent). Then,  $\omega$  is trivial in  $H(\gamma|d)$ .

The proof is direct, one has  $\alpha = d_{p_s}\mu$  for some  $\mu(H_s^{a_s}, B_s^{a_s(0,p_s)})$  since  $d_{p_s}$  is acyclic in the space of the  $\alpha(H_s^{a_s}B_s^{a_s,(0,p_s)})$  with no constant term. Thus  $\omega$  is  $d_{p_s}$ -exact,  $\omega = d_{p_s}(\mu\beta)$ :  $\omega$  is the first obstruction to the further lift of  $\mu\beta$  and as such, is trivial.

The theorem applies in particular when  $\alpha$  is an arbitrary polynomial of degree > 0 in the curvatures  $H_s^{a_s}$ .

## 6.2 Chapline-Manton models

#### 6.2.1 Model 1

The analysis is trivial in this case since there is no non trivial descent. All solutions of the Wess-Zumino consistency condition can be taken to be strictly annihilated by  $\gamma$ , i.e., can be taken to be in  $E_0$  ( $E_1 = 0$ ). They are thus completely described by Theorem **3.2** (from which one must remove the *d*-exact terms  $d\alpha([F])$ ).

#### 6.2.2 Model 2

The second model is more interesting. The algebra  $E_0^{small}$  has generators H, F,  $A^{(0,1)}$  and  $B^{(0,2r)}$ . One may take for  $F_0^{small}$  the space of elements of the form  $HQ(F, A^{(0,1)}, B^{(0,2r)})$  for which  $d_0(HQ(F, A^{(0,1)}, B^{(0,2r)})) = F^{r+1}Q \neq 0$ . These  $\gamma$ -cocycle do not lift at all. The space  $E_1$  is isomorphic to the algebra generated by F,  $A^{(0,1)}$  and  $B^{(0,2r)}$ , with the relation  $F^{r+1} = 0$ . Thus, it is not a free algebra contrary to the situation encountered in the free case. The differential  $d_1$  is non trivial and given by

$$d_1 A^{(0,1)} = F, \ d_1 F = 0, \ d_1 B^{(0,2r)} = 0$$
 (6.11)

when r > 1, which we shall assume at first. Because F is subject to the relation  $F^{r+1} = 0$ , the cohomological space  $E_2 \equiv H(d_1, E_1)$  is isomorphic to the algebra generated by  $B^{(0,2r)}$  and  $\mu(A, F)$  with

$$\mu(A,F) = -A^{(0,1)}F^r \tag{6.12}$$

The next differentials  $d_2$ ,  $d_3$  ... vanish up to  $d_{2r-1}$ . So,  $E_2 = E_3 = \ldots = E_{2r-1}$ . One has

$$d_{2r-1}B^{(0,2r)} = \mu(A,F), \ d_{2r-1}\mu(A,F) = 0.$$
(6.13)

Thus  $E_{2r} = 0$ .

One can take for  $F_1$  the space of polynomials of the form  $(B^{(0,2r)})^l Q_l(F)$  $A^{(0,1)}$  where  $Q_l$  is a polynomial in F of degree strictly less than r. Similarly, one may take for  $F_{2r-1}$  the space of polynomials in  $B^{(0,2r)}$  (with no constant piece). We leave it to the reader to write down explicitly the lifts of these elements. Note in particular that  $\mu(A, F)$  does not appear in any of the spaces  $F_k$ . This is because it is now trivial. In the free case,  $\mu(A, F)$  is an element of  $F_1$  and is the bottom of a non-trivial descent of length two. The coupling to the 2-form makes it disappear from the cohomology. At the same time, the cocycle  $F^{r+1}$ , which is in the invariant cohomology of d in the free case, has now become d-exact in the space of invariant polynomials. Also, while  $B^{(0,2r)}$  can be transgressed all the way up to H in the free case, its lift stops now at ghost number one with  $\mu$ .

The situation for r = 1 is similar, the two steps corresponding to the differentials  $d_1$  and  $d_{2r-1}$  being combined in one, so that the space  $E_2$  vanishes. The easiest way to see this is to observe that  $H(d_1, E_1)$  (with  $d_1 A^{(0,1)} = F$ ,  $d_1F = 0$  and  $d_1B^{(0,2)} = \mu(A, F)$  for r = 1) is isomorphic to  $H(D, E_0)$  with  $DA^{(0,1)} = F$ , DF = 0,  $DH = F^{r+1}$ ,  $DB^{(0,2)} = \mu(A, F) + H$ . Indeed, one may view the generator H as Koszul generator for the equation  $F^{r+1} = 0$ . The change of variable  $H \to H' = H + \mu$  brings then D to the manifestly contractible form.

#### 6.2.3 Model 3

The third model is essentially a combination of the first model in the (A, B)sector and of the free model for the improved 3-form  $C_M = C - AB - \frac{1}{2}AdA$ , with curvature  $G_M = dC_M$  and improved last ghost of ghost  $\tilde{C}^{(0,3)}$  (3.15). Only  $d_0$  and  $d_3$  are non trivial. The details are left as an exercise.

#### 6.2.4 Model 4

In the absence of coupling between the 2-form and the Yang-Mills field, the non trivial differentials are

$$d_2 B^{(0,2)} = H, \ d_2 H = 0 \tag{6.14}$$

 $(B^{(0,2)} \equiv \rho)$  and

$$d_3 tr C^3 = tr F^2, \ d_3 tr F^2 = 0 \tag{6.15}$$

$$d_5 tr C^5 = tr F^3, \ d_5 tr F^3 = 0 \tag{6.16}$$

$$d_{2N-1}trC^{2N-1} = trF^N ag{6.18}$$

(see [13]). We have written explicitly only the action of the non trivial  $d_k$ 's on the contractible pairs. The last ghost of ghost  $B^{(0,2)}$  is non trivial and can be lifted twice;  $trC^3$  is non trivial and can be lifted three times;  $trC^5$  is non trivial and can be lifted five times; more generally,  $trC^{2k+1}$  is non trivial and can be lifted (2k + 1) times.

÷

When the coupling is turned on, the variables  $\rho$  and  $trC^3$  disappear from the cohomology. It follows that all the solutions of the Wess-Zumino consistency condition that were previously above  $trC^3$  (or above a polynomial involving  $trC^3$ ) become trivial. This is the Green-Schwarz anomaly cancellation mechanism [30]. At the same time, the differential  $d_0$  becomes non trivial, as for the previous Chapline-Manton models. One has

$$d_0 H = tr F^2, \ d_0 tr F^2 = 0 \tag{6.19}$$

which shows that  $trF^2$  disappears from the invariant cohomology, as already pointed out above. The other differentials (6.16) through (6.18) remain unchanged. The cohomology  $H(\gamma|d)$  is given in [35].

## 7 Counterterms and anomalies

We finally summarize our results by giving explicitly the antifield-independent counterterms and anomalies, i.e.,  $H^{(n,0)}(\gamma|d)$  and  $H^{(n,1)}(\gamma|d)$ . These can be of two types: (i) the ones that descend trivially ("type A"); these can be assumed to be strictly annihilated by  $\gamma$  and are described by  $H(\gamma)$  up to trivial terms; and (ii) the ones that lead to a non-trivial descent ("type B"); these can be assumed to be in the small algebra modulo solutions of the previous type. For small ghost number, it turns out to be more convenient to determine the solutions of "type B" directly from the obstructions sitting above them rather than from the bottom. That this procedure, which works in the universal algebra, yields all the solutions, is guaranteed by our general analysis.

## 7.1 Counterterms and anomalies of type A

a) The counterterms that lead to a trivial descent involve in general the individual components of the gauge-invariant field strengths and their derivatives and generically cannot be expressed as exterior products of the forms F or H. They are the gauge-invariant polynomials introduced above and read explicitly

$$a = a([H^a])d^nx (7.1)$$

for the free models $^3$ ,

$$a = a([F])d^nx (7.2)$$

for the first Chapline-Manton models,

$$a = a([F], [H])d^nx$$
 (7.3)

for the second Chapline-Manton models and

$$a = a([F], [G])d^nx (7.4)$$

for the third Chapline-Manton models (with the condition  $a \neq db$  in all cases which is equivalent to the condition that the variational derivatives of a with respect to the fields do not identically vanish).

For the fourth Chapline-Manton model, the counterterms are invariant functions of  $F^a_{\mu\nu}$  and their covariant derivatives, as well as of [H],

$$a = P_I([F^a], [H])d^n x,$$
 (7.5)

where  $P_I$  is invariant under the adjoint representation of SU(N).

We have assumed that the spacetime forms  $dx^{\mu}$  occur only through the product  $dx^0 dx^1 \cdots dx^{n-1} \equiv d^n x$  as this is required by Lorentz-invariance.

<sup>&</sup>lt;sup>3</sup>Recall that the "free models" encompass in fact all models having the gauge symmetries of the free theory since the present analysis depends only on the form of the gauge transformations and not on the specific Lagrangian. So the results for the "free models" cover general Lagrangians  $L([H^a])$  or models with Chern-Simons interactions.

b) The anomalies that lead to a trivial descent are sums of terms of the form  $a = P C d^n x$  where P is a gauge-invariant polynomial and C is a last ghost of ghost of ghost number one, which must be non trivial in  $H(\gamma)$ . These anomalies exist only for a free theory with 1-forms and in the second Chapline-Manton models since only in these cases are there non trivial, last ghosts of ghosts of ghost number one. One has explicitly

$$a = P_A([H^a])B^{A(0,1)}, (7.6)$$

where A runs over the 1-forms (free models) or

$$a = P([F], [H])A^{(0,1)},$$
(7.7)

(second CM models). In both cases a will be trivial if P = dR where R is an invariant polynomial or if  $P_A = P_A(H^a)$  with  $P_A H^A = 0$  in the first case and  $a = \mu$  in the second case.

The existence of such anomalies - which generically cannot be expressed as exterior products of curvatures and ghosts - was pointed out in [50] for Yang-Mills gauge models with U(1) factors.

## 7.2 Counterterms of type B

The solutions that lead to a non trivial descent can be assumed to be in the small algebra, i.e., can be expressed in terms of exterior product of the fields, the ghosts (which are all exterior forms) and their exterior derivatives (modulo solutions of type A). If a is a non-trivial solution of the Wess-Zumino consistency condition with ghost number zero, then  $da \neq 0$  (in the universal algebra). Since a has ghost number zero, it is the top of the descent and dais the obstruction to a further lift. Because da is a  $\gamma$ -cocycle, it is a gaugeinvariant polynomial. It must, in addition, be d-closed but not d-exact in the space of gauge-invariant polynomials since otherwise, a could be redefined to be of type A. Therefore, da is an element of the invariant cohomology of d and it will be easier to determine a directly from the obstruction darather than from the bottom of the descent because one knows the invariant cohomology of d.

#### 7.2.1 Free models

In the free case, any polynomial P(H) in the curvatures  $H^a$  is d-closed and thus d-exact,

$$P(H) = dQ(H,B) \tag{7.8}$$

where Q is linear in the forms  $B^a$ ,

$$Q = R_a(H^b)B^a, (7.9)$$

One may in fact assume that Q involves only the potentials  $B^a$  of the curvatures of smaller form-degree occuring in P. To searched-for solution of the Wess-Zumino consistency condition of which P is the obstruction to a further lift is of course just Q. These are the familiar Chern-Simons terms, which exist provided one can match the spacetime dimension n with a polynomial in the curvatures  $H^a$  and the forms  $B^a$ , linear in  $B^a$ .

The whole descent associated with Q is generated through the "Russian formula" [45, 32]

$$P = \tilde{\gamma}Q(H,\tilde{B}) \tag{7.10}$$

$$\tilde{\gamma} = d + \gamma \tag{7.11}$$

$$\tilde{B}^a = B^{a(p_a,0)} + B^{a(p_a-1,1)} + \dots + B^{a(0,p_a)}$$
(7.12)

which follows from the "horizontability condition" [32]

$$\tilde{\gamma}\tilde{B}^a = H^a. \tag{7.13}$$

By expanding (7.10) according to the ghost number, one gets the whole tower of descent equations. The bottom takes the form  $R_a(H^b)B^{a(0,p_a)}$  and is linear in the last ghosts of ghosts associated with the forms of smaller form degree involved in P. That the bottoms should take this form could have been anticipated since these are the only bottoms with the right degrees that can be lifted all the way to form-degree n. The non-triviality of the bottom implies also the non-triviality of the whole tower.

It is rather obvious that the Chern-Simons terms are solutions of the Wess-Zumino consistency condition. The main result here is that these are the only solutions that descend non trivially (up to solutions of type A).

#### 7.2.2 Chapline-Manton model 1

There is in this case no non trivial solution of type B since there is no non trivial descent.

#### 7.2.3 Chapline-Manton model 2

One may proceed as for the free theory. The polynomial P must be taken in the invariant cohomology of d and so is a polynomial in the curvatures F with  $F^{r+1}$  identified with zero. This leads, as in the free theory, to the Chern-Simons terms

$$a = F^m A, (7.14)$$

except that  $F^{r}A$  is now absent because it can be brought to class A up to exact terms. These Chern-Simons terms are available in all odd dimensions.

#### 7.2.4 Chapline-Manton model 3

 $\tilde{\gamma}$ 

In this case, the obstruction P is a polynomial in the improved field strength  $G_M$ . One has  $P = dQ(G_M, C_M)$  where

$$Q(G_M, C_M) = R(G_M)C_M$$
(7.15)

is linear in the improved potential  $C_M = C - AB - \frac{1}{2}AdA$ . The Chern-Simons solution Q exists only in spacetime dimension 4k - 1. As in the free case, the whole descent associated with Q is generated through the Russian formula:

$$P = \tilde{\gamma}Q(G_M, \tilde{C}_M) \tag{7.16}$$

$$= d + \gamma \tag{7.17}$$

$$\tilde{C}_M = C_M + E_M + L_M + \tilde{C}^{(0,3)},$$
(7.18)

with  $E_M = C^{(2,1)} - \frac{1}{2}A^{(1,0)}B^{(1,1)} - \frac{1}{2}dAA^{(0,1)} - BA^{(0,1)}$  and  $L_M = C^{(1,2)} - \frac{1}{2}AB^{(0,2)} - \frac{1}{2}A^{(0,1)}B^{(1,1)}$ . This follows from the "horizontability condition"

$$\tilde{\gamma}\tilde{C}_M = G_M. \tag{7.19}$$

#### 7.2.5 Chapline-Manton model 4

Again, one finds as solutions the familiar higher order Yang-Mills Chern-Simons not involving  $trF^2$  or  $\omega_3$ . These are available in odd dimensions > 3.

### 7.3 Anomalies of type B

The anomalies a of type B can themselves be of two types. They can arise from an obstruction that lives one dimension higher or from an obstruction that lives two dimensions higher. In the first case, the obstruction da has form degree n + 1 and ghost number 1. This case is only possible for the free models with 1-forms and the second Chapline-Manton model, since there is no  $\gamma$ -cohomology in ghost number one for the other models. In the other case, the anomaly can be lifted once,  $da + \gamma b = 0$ . The obstruction db to a further lift is then a (n + 2)-form of ghost number 0.

In the first case, the obstruction da reads

$$da + \gamma(\text{something}) = P_A(H)B^{A(0,1)} \tag{7.20}$$

(we consider explicitly the free case, the second CM model being handled similarly). The right-hand side of (7.20) is necessarily the  $d_1$  of something. Indeed, it cannot be the  $d_k$  (k > 1) of something, say m, since this would make m trivial: the first obstruction to the lift of m would have to vanish and m involves explicitly the variables of the 1-form sector (see theorem **6.1** above). This implies

$$P_A(H)B^{A(0,1)} = C_{AB}(H)H^A B^{B(0,1)}, \quad C_{AB}(H) = -C_{BA}(H)$$
(7.21)

so that  $P_A(H)B^{A(0,1)} = d_1(\frac{1}{2}C_{AB}(H)B^{A(0,1)}B^{B(0,1)})$ . One thus needs at least two 1-forms to construct such solutions. If  $C_{AB}(H)$  involves the curvatures  $H^A$  of the 1-forms, it must be such that (7.21) is not zero. The anomaly following from (7.20) is

$$a = C_{AB}(H)B^{A(1,0)}B^{B(0,1)}, (7.22)$$

and the associated descent is generated through

$$C_{AB}(H)H^{A}B^{B(0,1)} = \tilde{\gamma}(\frac{1}{2}C_{AB}(H)\tilde{B^{A}}\tilde{B^{B}})$$
(7.23)

In the second case and for the free theory, the obstruction  $P \in H^{inv}(d)$ is a polynomial in  $H^a$  of form-degree n + 2, which can be written P = dQwhere Q is linear in the potentials associated with the curvatures of lowest degree occuring in P. The solution a and the descent are obtained from the Russian formula (7.10), exactly as for the counterterms,

$$a = R_a(H^b)C^a. (7.24)$$

They are linear in the ghosts and exist only if there are forms of degree > 1 which are the only ones that can occur in P since otherwise a is either trivial or of type A. Indeed, if variables from the 1-form sector occur in P, then  $P = d_1 a$  (if P is non trivial) and the descent has only two steps. But this means that a is the bottom of the descent and is really of type A.

There is no solution of this type for the Chapline-Manton model 1 because of the lack of a non-trivial descent.

For the Chapline-Manton model 2, there is again no anomaly that can be lifted once because there is no element belonging to an  $F_k$  yielding upon lifting the appropriate obstruction.

For the Chapline-Manton model 3, solutions descending from polynomials  $P(G_M)$  in two dimensions higher exist only in spacetime dimensions equal to 4k - 2. They are given by

$$a = Q(G_M)L_M, (7.25)$$

with  $L_M$  defined in Section 7.2.4.

Finally, for the Chapline-Manton model 4, one has all the anomalies of the SU(N) Yang-Mills theory, except those involving the cocycle  $trC^3$  and its lifts since they are now trivial.

## 8 Conclusions

In this paper, we have derived the general solution of the antifield-independent Wess-Zumino consistency condition for models involving p-forms. We have justified in particular why one may assume that the solutions can be expressed in terms of exterior products of the fields, the ghosts (which are all exterior forms) and their exterior derivatives, when these solutions occur in non trivial descents. This is not obvious to begin with since there are solutions that are not expressible in terms of forms (those that descend trivially) and justify the usual calculations made for determining the anomalies. Once one knows that the solutions involved in non trivial descents can be expressed in terms of forms (up to solutions that descend trivially), one can straightforwardly determine their explicit form in ghost numbers zero and one. This was done in the last section where all counterterms and anomalies have been classified. The counterterms for the free models are either strictly gauge invariant and given by (7.1) or of the Chern-Simons type (when available) and given by (7.9). The counterterms for the Chapline-Manton models are also either strictly invariant (Eqs. (7.2), (7.3), (7.4), (7.5)) or of the Chern-Simons type. These Chern-Simons solutions exist for all the models except the first one in appropriate dimensions (see Eqs. (7.14), (7.15) respectively and Section **7.2.5**).

The anomalies may also be either strictly annihilated by  $\gamma$ , or lead to a non-trivial descent. The first type generalizes the anomalies of Dixon and Ramon Medrano [50] and exist in the free case and the second CM model. The more familiar anomalies with a non trivial descent are analyzed in Section **7.4** and listed in Eqs. (7.22), (7.24), (7.25) and below (7.25).

The method applies also to other values of the ghost number, which are relevant in the analysis of the antifield-dependent cohomology.

As we have shown in [36], the natural appearance of exterior forms holds also for the characteristic cohomology: all higher order conservation laws are naturally expressed in terms of exterior products of field strengths and duals to the field strengths. It is this property that makes the gauge symmetrydeforming consistent interactions for p-form gauge fields expressible also in terms of exterior forms and exterior products [25, 11].

# Acknowledgements

The authors are grateful to Glenn Barnich, Friedemann Brandt, Michel Dubois-Violette, Christiane Schomblond, Michel Talon, Claude Viallet and André Wilch for useful discussions. B. K. is "Aspirant du Fonds National de la Recherche Scientifique" (Belgium). MH thanks the Erwin Schrödinger Institute for hospitality while this paper was completed. This work has been partly supported by I.I.S.N.-Belgium (convention 4.4505.86) and the "Ac-

tions de Recherche Concertées" of the "Direction de la Recherche Scientifique - "Communauté francaise de Belgique".

# References

- [1] J. Wess and B. Zumino, *Phys. Lett.* **37** B (1971) 95.
- [2] G.F. Chapline and N.S. Manton, *Phys. Lett.* B120 (1983) 105; H. Nicolai and P.K. Townsend, *Phys. Lett.* 98B (1981) 257; A. H. Chamseddine, *Nucl. Phys.* B185 (1981) 403; *Phys. Rev.* D24 (1981) 3065; E. Bergshoeff, M. de Roo, B. de Wit and P. van Nieuwenhuizen, *Nucl. Phys.* B195 (1982) 97.
- [3] C. Becchi, A. Rouet and R. Stora, Commun. Math. Phys. 42 (1975) 127; Ann. Phys. (NY) 98 (1976) 287.
- [4] J. Zinn-Justin, Renormalisation of gauge theories in Trends in elementary particle theory, Lecture notes in Physics n<sup>0</sup>37 (Springer, Berlin 1975); Quantum Field Theory and Critical Phenomena, 3<sup>d</sup> Edition, Clarendon Press (Oxford: 1995).
- [5] I.A. Batalin and G.A. Vilkovisky, *Phys. Lett.* **102** B (1981) 27; *Phys. Rev.* D **28** (1983) 2567.
- [6] B. L. Voronov and I. V. Tyutin, *Theor. Math. Phys.* 50 (1982) 218; 52 (1982) 628.
- [7] C. Itzykson and J.-B. Zuber, "Quantum Field Theory", Mc Graw Hill (New York: 1980).
- [8] O. Piguet and S. P. Sorella, "Algebraic Renormalization", Lecture Notes in Physics vol. m28, Springer Verlag (Berlin: 1995).
- [9] S. Weinberg, "The Quantum Theory of Fields", volume 2, Cambridge University Press (Cambridge: 1996).
- [10] G. Barnich and M. Henneaux, *Phys. Rev. Lett.* **72** (1994) 1588; G. Barnich, F. Brandt and M. Henneaux, *Commun. Math. Phys.* **174** (1995) 93.

- [11] M. Henneaux and B. Knaepen, *Counterterms and anomalies for p-form gauge theories*, in preparation.
- [12] M. Dubois-Violette, M. Talon and C. Viallet, Phys. Lett. 158 B (1985) 231.
- [13] M. Dubois-Violette, M. Talon and C. Viallet, Commun. Math. Phys. 102 (1985) 105.
- [14] M. Dubois-Violette, M. Talon and C. Viallet, Ann. Inst. Henri Poincaré 44 (1986) 103.
- [15] M. Talon, "Algebra of anomalies", Cargese Summer Inst. Jul 15-31 (1985) 433.
- [16] F. Brandt, N. Dragon and M. Kreuzer, *Phys. Lett.* B231 (1989) 263; *Nucl. Phys.* B332 (1990) 250.
- [17] M. Dubois-Violette, M. Henneaux, M. Talon and C. M. Viallet, Phys. Lett. B289 (1992) 361.
- [18] W. Siegel, *Phys. Lett.* **93** B (1980) 170.
- [19] I.A. Batalin and E.S. Fradkin, *Phys. Lett.*, **122B** (1983) 157.
- [20] J. Thierry-Mieg, Nucl. Phys. B335 (1990) 334.
- [21] L. Baulieu and J. Thierry-Mieg, Nucl. Phys. **B228** (1983) 259.
- [22] M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems*, Princeton University Press (Princeton: 1992).
- [23] S. Deser, R. Jackiw and S. Templeton, *Phys. Rev. Lett.* 48 (1982) 975.
- [24] M. Henneaux, *Phys. Lett.* **368B** (1996) 83.
- [25] M. Henneaux and B. Knaepen, *Phys. Rev.* D 56 (1997) 6076.
- [26] C. Teitelboim, *Phys. Lett.* **167B** (1986) 63.
- [27] L. Baulieu, On Forms with Non-Abelian Charges and Their Dualities, hep-th/9808055.

- [28] D. Freedman and P. K. Townsend, Nucl. Phys. B177 (1981) 282.
- [29] F. Brandt and N. Dragon, Nonpolynomial gauge invariant interactions of 1-form and 2-form gauge potentials, in Theory of Elementary Particles, pp. 149-154, H. Dorn, D. Lüst, G. Weigt (eds.) (Wiley-VCH, Weinheim, 1998), hep-th/9709021.
- [30] M. B. Green and J. H. Schwarz, *Phys. Lett.* **149B** (1984) 117.
- [31] L. Baulieu, *Phys. Lett.* **167** B (1986) 56.
- [32] L. Baulieu, Algebraic construction of gauge invariant theories, in Perspectives in Particles and Fields, Cargèse 1983, M. Levy, J.-L. Basdevant, D. Speiser, J. Weyers, M. Jacob and R. Gastmans eds, NATO ASI Series B126, Plenum Press, New York (1983).
- [33] L. Romans, *Phys. Lett.* **B 169** (1986) 374.
- [34] E. Bergshoeff, M. de Roo, G. Papadopoulos, M. B. Green and P. K. Townsend, Nucl. Phys. B 470 (1996) 113.
- [35] M. Henneaux B. Knaepen and C. Schomblond, Lett. Math. Phys. 42 (1997) 337.
- [36] M. Henneaux, B. Knaepen and C. Schomblond, Commun. Math. Phys. 186 (1997) 137.
- [37] D. Sullivan, "Infinitesimal Computations in Topology", Publ. I.H.E.S. 47 (1977) 269.
- [38] M. Hamermesh, *Group Theory*, Addison Wesley (1962).
- [39] J. A. Garcia and B. Knaepen, Couplings between Generalized Gauge Fields, hep-th/9807016, to apear in Phys. Lett. B.
- [40] T. Curtright, *Phys. Lett.* **165B** (1985) 304.
- [41] M. Henneaux and A. Wilch, *Phys. Rev.* D 58 (1998) 025017.
- [42] C. Bizdadea and S. O. Saliu, Int. J. Mod. Phys. A 11 (1996) 3523.

- [43] W. Greub, S. Halperin and R. Vanstone, *Connections, curvature and cohomology*, vol. III, Academic Press (New York: 1976).
- [44] J. L. Koszul, Bull. Soc. Math. Fr. 78 (1950) 65.
- [45] R. Stora, in New Developments in QFT and Statistical Mechanics, M. Levy and P. Mitter eds, Plenum Press (New York: 1977); R. Stora, in Recent Progress in Gauge Theories, Lehmann G. et al eds, Plenum Press (New York: 1984); B. Zumino, in Relativity, Groups and Topology II, B. S. De Witt and R. Stora eds, North Holland (Amsterdam: 1984).
- [46] A.M. Vinogradov, Sov. Math. Dokl. 19 (1978) 1220; M. De Wilde, Lett. Math. Phys. 5 (1981) 351; W.M. Tulczyjew, Lecture Notes in Math. 836 (1980) 22; P. Dedecker and W.M. Tulczyjew, Lecture Notes in Math. 836 (1980) 498; T. Tsujishita, Osaka J.of Math. 19 (1982) 311; L. Bonora and P. Cotta-Ramusino, Commun. Math. Phys. 87 (1983) 589; P.J. Olver, Applications of Lie Groups to Differential Equations, Graduate Texts in Mathematics, volume 107, Springer Verlag (New York: 1986); R.M. Wald, J. Math. Phys. 31 (1990) 2378; L.A. Dickey, Contemp. Math. 132 (1992) 307. F. Brandt, N. Dragon and M. Kreuzer, Nucl. Phys. B332 (1990) 224; M. Dubois-Violette, M. Henneaux, M. Talon and C. M. Viallet, Phys. Lett. B267 (1991) 81.
- [47] S. Mac Lane, *Homology*, Springer (New York: 1963).
- [48] L. Bonora, P. Cotta-Ramusino, M. Rinaldi and J. Stasheff, Commun. Math. Phys. 112 (1987) 237.
- [49] H. Cartan, in Colloque de Topologie (Bruxelles 1950), Masson (Paris: 1951)
- [50] J. Dixon and M. Ramon Medrano, *Phys. Rev.* D 22 (1980) 429.